

UNITED STATES DEPARTMENT OF COMMERCE

R. P. LAMONT, Secretary

BUREAU OF FISHERIES

HENRY O'MALLEY, Commissioner

REPORT
OF THE
UNITED STATES
COMMISSIONER OF FISHERIES

FOR THE FISCAL YEAR 1928

WITH

APPENDIXES

Part II



UNITED STATES
GOVERNMENT PRINTING OFFICE
WASHINGTON : 1930

CONTENTS

	Page
CHECK LIST OF THE FISHES AND FISHLIKE VERTEBRATES OF NORTH AND MIDDLE AMERICA NORTH OF THE NORTHERN BOUNDARY OF VENEZUELA AND COLOMBIA. By David Starr Jordan, Barton Warren Evermann, and Howard Walton Clark. (Document No. 1055. Issued February, 1930)-----	1-670

DEPARTMENT OF COMMERCE

BUREAU OF FISHERIES

HEADQUARTERS STAFF, 1927-28

Commissioner

HENRY O'MALLEY

Deputy Commissioner.—LEWIS RADCLIFFE.

Assistants in Charge of Divisions:

Office.—IRVING H. DUNLAP.

Fish Culture.—GLEN C. LEACH.

Inquiry Respecting Food Fishes.—ELMER HIGGINS.

Fishery Industries.—OSCAR E. SETTE.

Alaska Service.—WARD T. BOWER.

Architect and Engineer.—GEORGE A. SCHNEIDER.

Accountant.—CHARLES W. SCUDDER.

Superintendent Central Station and Aquaria.—L. G. HARRON.

Editor.—ANNA B. BROWN.

CHECK LIST OF THE FISHES AND FISHLIKE VERTEBRATES OF NORTH AND MIDDLE AMERICA NORTH OF THE NORTHERN BOUNDARY OF VENEZUELA AND COLOMBIA ¹

By

DAVID STARR JORDAN, BARTON WARREN EVERMANN, and
HOWARD WALTON CLARK

INTRODUCTION

The present memoir comprises a list of all the fishes and fishlike vertebrates known to occur in the waters of north and middle America; more specifically all of continental America north of the Isthmus of Panama, and the outlying islands including the West Indies, the Greater and the Lesser Antilles, Greenland, Iceland, and the islands off the Pacific coast of Central America, Mexico, the United States, and Alaska. The salt-water species on the northern coasts of Colombia and Venezuela have been included. Those of the Galapagos, the Sea of Okhotsk, and the west coast of Kamchatka are not included. Oceanic species in general within the 1,000-fathom curve have been admitted, although it is not always easy to draw the line.

The majority of the species included will be found described in *Fishes of North and Middle America*, a 4-volume work by Jordan and Evermann published in 1896 to 1900 as Bulletin 47 of the United States National Museum. No other general work restricted to this continent and containing descriptions of all known species has been written. On that work was based the Check List of 1896 by Jordan and Evermann, of which the present Check List may be regarded as a revision.

In the present list are given, in as natural sequence as our present knowledge permits, (1) the names of all species and subspecies that the authors admit as valid, (2) as many of the vernacular or common names for each species as have been readily obtainable, (3) the known geographic distribution of each species, (4) the reference to the original description of the species, and (5) the reference to every real synonym. If a species has been described as new but once, there will, of course, be but one reference (as, for example, *Photocorynus spiniceps* on p. 509); if it has been described as new six times there will be six references (as, for example, *Eupomotis gibbosus* on p. 302). Thus, this work is really a check list of all the scientific names that have ever been applied to any American fish.

As 4,139 species and subspecies are admitted as valid, and as there are more than 3,000 real synonyms, it appears that each American fish has been described as new about two times on an average.

The first attempt at a Synopsis of the Fishes of North America was published by David Humphreys Storer in the *Memoirs of the Ameri-*

¹ Appendix X to the Report of the U. S. Commissioner of Fisheries for 1928. B. F. Doc. 1055.

can Academy of Arts and Sciences of Boston in 1846. That work enumerates 739 species distributed in 221 genera. As the fishes of the Rocky Mountain region, Mexico, the Pacific coast, and the West Indies were then scarcely known, this list was very incomplete and the nomenclature naturally was imperfect.

After an elaborate survey of the Pacific coast fish fauna in 1880, David Starr Jordan and the late Charles Henry Gilbert published a Synopsis of the Fishes of North America (1882), including all species known north of the Mexican boundary. The 1,340 species names in that synopsis were arranged in 487 genera; a few other species appeared in the appendix.

Jordan's Catalogue of the Fishes Known to Inhabit the Waters of North America North of the Tropic of Cancer (1887) was based on this synopsis and includes a number of additional species.

In the present memoir we have followed implicitly the rules of nomenclature laid down by the International Commission for Zoological Nomenclature, of which the senior author has been a member since 1908. One single case now under discussion (*Carcharias*) has not been disposed of fully as yet, and the view recommended by Jordan is adopted here tentatively. An important rule set forth by Jordan and Evermann in 1895 has not been acted upon. This rule reads: "We regard all generic names as different unless actually spelled alike." In accordance with decisions of the commission, another rule, which we have followed, is this: "No change is to be made in current nomenclature unless the case is fairly clear; as, for example, a matter of priority." In general we have followed current practice rather than accept questionable changes. One doubtful opinion should not be exchanged for another. The original orthography of all names (misprints aside) has been retained. Some supposed misprints (*Cypsilurus*, *Pachynathus*, etc.) have been referred to the International Commission, whose verdicts have been followed.

In accordance with the decision of the International Commission, we have disregarded (unless revived by later authors) the polynomial generic names of Gronow, Klein, Browne, and other able early writers, who failed to adopt the Linnæan code simply because they had not heard of it. This decision rests not on consideration of justice but upon that of permanence.

The most serious problem in taxonomy is that of the assignment of generic types to the genera of authors who had no conception of types. The "rule of the first reviser" was adopted by the commission at Boston in 1908, and the present writers have endeavored to follow it.

By far the most complete work on North American fishes is that by Jordan and Evermann, mentioned above—the Fishes of North and Middle America. In that work the authors recognized 1,113 genera, 3,263 species, and 133 subspecies—a total of 3,396 species and subspecies. The check list of Fishes and Fishlike Vertebrates of North and Middle America, by Jordan and Evermann, based upon the above and published in 1896, enumerates 3,100 species arranged in 1,053 genera, these being about double those in the Jordan catalogue of 1887, the additions being mainly from Mexico, the West Indies, and the deep seas.

The late modern period in systematic ichthyology is from 1880 to the present time and is especially signalized by the explorations of

the deep sea, of most of the continents, and of fossil beds. The unexampled explorations of Eigenmann in South America belong to this period, as also the records of Boulenger of the fishes of South Africa and the varied researches of Regan in the rich material of the British Museum. In this period the faunas of Australia, Japan, and Siberia have been studied critically and successfully. Our knowledge of fossil fishes is assuming definite form through the work of Woodward, Zittel, Hay, Errol Ivor White, and others who have given vertebrate paleontology, as related to ichthyology, a constructive turn.

We may again emphasize the distinction between the ancient and modern periods in animal taxonomy. Beginning with 1758, the date of Linnæus's *Systema Naturæ*, for 60 years a genus was usually regarded merely as a convenient pigeonhole into which species were thrown. With Cuvier's *Règne Animal* (1817) came the conception of arrangement "according to organization," or comparative anatomy. With Darwin's *Origin of Species*, 40 years later (1859), zoologists began, though slowly at first, to realize that a classification must be more than an inventory; that its basis must be genetic, and that the problems involved in natural grouping are vastly more complex than even great morphologists like Cuvier and Agassiz had realized. To paraphrase a sentence of Elliott Coues, "genera and species are but larger and smaller twigs of a tree, which we try to arrange as nearly as possible in accordance with nature's ramifications." This view, applied to taxonomy, involves the extensive subdivision of accepted genera, many of which were in the highest degree unnatural. Subdivision as practiced in the present paper may go too far in some cases, but our material must be analyzed first before we can group its parts in natural synthesis. No system of naming can progress beyond the knowledge on which it rests. We may emphasize here a statement attributed to Aristotle that "one should never expect a degree of accuracy which the subject matter does not permit."

Each tendency in evolution, large or small, diverging from the main current, should properly stand as a separate genus.

The problem of subspecies among fishes is especially incomplete and confused. Most of the marine species are clearly definable and not often subject to intergradations. Among the fresh-water forms, all sorts of modifications occur. A species, in general, is a definable group of related individuals that have run the gauntlet of life and have endured. Among our river fishes we find species of different types. There are numerous cases of species of compact though often wide distribution, in which all individuals are, to all appearances, similar. Examples of this type are the common perch (*Perca fluviatilis*), the common sucker (*Catostomus commersonii*), and the minnows (*Ericymba buccata* and *Exoglossum maxillingua*). Other forms, which we have elsewhere called geminate or twin species, may be distinct enough, having a few tangible characters of difference in spite of general resemblance. These, for the most part, are set apart from each other by barriers, their distinctness proportional to the efficiency of the barrier. Their origin may be due to the crossing of these barriers by individuals from the mass of their kind, thus acquiring a different environment with a different type of selection. Illustrations of geminate species set off by isolation may be found in abundance, as among the trout of western America, and other species,

especially in the Panama region, where many geminate species are found on each side of the Isthmus, which has split its species from their mates since Miocene times. The Miocene fossil fishes of California are largely geminate in origin, twins or parents of species now appearing on the coast. In most cases such forms are clearly distinct but to all appearance ancestral, the existing distinctions apparently due to varying incidents of selection in varying environment. Geminate species rarely inhabit the same region, though exceptions are known, attributable to reinvasion.

In many cases the distinctions between geminate forms are not complete, intergradations occurring along the line of the barriers or at different sections in climatic range. Types thus connected are known as subspecies. Subspecies are often nascent species; but in most cases a large amount of material is necessary before subspecies can be separated from geminate species. Among fresh-water fishes, each river basin shows some features of distinction, large or small, as does each oceanic island with regard to marine fishes. In most families of American river fishes studies of species have not been extensive enough to establish facts of intergradation. While in birds, reptiles, and mammals the subspecies are usually well defined and named, among the fishes this is still largely a matter of guesswork. Many authors base the recognition of subspecies not on actual intergradation but on guesses based on degrees of probability, a practice we have not followed. It has seemed to us better to regard related forms as distinct species until intergradation has been shown.

We find also among river fishes, and sometimes among marine forms (as in Lutianus), occasional individuals that to all appearance are hybrids. These are marked by a distinct blending of specific traits. They are extremely rare, however, often known only from a single example. A few hybrids are known or suspected among certain minnows, sunfishes, and darters. The best-known hybrids thus far recorded in America occur among the Centrarchidæ, or sunfishes, in which group they have been studied critically by Mr. Hubbs. Two common species (*Apomotis cyanellus* and *Eupomotis gibbosus*) figure particularly as probable parents. Hybrids may give rise to distinct species, but no example of this is known among our fishes.

Among fishes there are also occasional monstrosities, freaks, or "mutations," in which certain traits may be absent or exaggerated. Whether such qualities may be inherited is not known. If so, it is likely that they would soon be swamped by breeding with the mass unless especially protected by segregation. It does not appear that any species of fishes have arisen by mutation or from hybridization, though the latter possibility is greater than the former.

We may observe that compilations such as the present paper should constitute an exercise in modesty. Such work can never be made absolutely complete nor free from errors. It deals with the observations and thoughts of great minds, as also with a degree of ignorance, carelessness, and perversity, by which names have been multiplied without corresponding extension of ideas. Synonymy (the use of different names for the same group) in general represents the lack of exact knowledge or else a failure (often unavoidable) to deal justly with the work of others. It is, therefore, as has been said by Doctor Coues, often "a burden and disgrace to science."

Only those who have had experience in cataloguing, in a similar way, the species in some large group of animals or plants can appreciate the difficulties involved. In the first place, in order to make the list complete by the inclusion of all valid names and all real synonyms it was necessary to make a careful search of all ichthyological literature. It is too much to expect that we have been entirely successful in this search; some names probably have been overlooked. The verification of the thousands of references as to the publication, volume, page, date, spelling, and type locality, was even more exacting and arduous; but most difficult of all were the multitude of taxonomic problems involved. In many cases existing knowledge is not adequate for final settlement. It is not improbable that some of our decisions as to the taxonomic relationships of certain species and the validity of others may not meet the approval of all of our coworkers, but that is inevitable where knowledge is incomplete.

The present senior author, as a lifelong teacher of biology, may be pardoned a personal reference. Himself a student of Agassiz and Gill of the earlier period, he has had the honor of being also a friend, to some extent a protégé, of Baird, Günther, Vaillant, Lütken, Hilgendorf, and Cope. In addition, he has numbered among his own students many of those active in systematic ichthyology during the late modern period. Among these are James Francis Abbott, Willis Stanley Blatchley, Charles Harvey Bollman, Alembert Winthrop Brayton, Charles Victor Burke, Howard Walton Clark*, Herbert Edson Copeland, Frank Cramer, George Bliss Culver, Bradley Moore Davis, Mary Cynthia Dickerson, Charles Lincoln Edwards, Carl H. Eigenmann, Rosa Smith Eigenmann, Barton Warren Evermann, Bert Fesler, Morton William Fordice, Henry Weed Fowler, Charles Henry Gilbert, James Zaccheus Gilbert, David Kop Goss, Ulysses Simpson Grant IV, Arthur White Greeley, Joseph Grinnell, Harold Hannibal, Edmund Heller, Albert William Christian Theodore Herre, Jennie Horning, Carl Leavitt Hubbs, Elizabeth G. Hughes, Oliver Peebles Jenkins, Eric Knight Jordan, Philip Henry Kirsch, Richard Crittenden McGregor, Ernest Alexander McGregor, Charles Leslie McKay, Seth Eugene Meek, Charles William Metz, Heraclio R. Montalban, George Sprague Myers, Edward William Nelson, Robert Newland, Masamitsu Oshima, Keinosuke Otaki, Charles J. Pierson, Cora D. Reeves, Robert Earl Richardson, Cloudsley Rutter*, Norman Bishop Scofield, Alvin Seale, Michitaro Sindo, Tage Skogsberg, Robert Evans Snodgrass, John Otterbein Snyder, Edwin Chapin Starks, Joseph Swain, Shigeo Tanaka, Lionel William Wiedey, Wilbur Wilson Thoburn, Joseph C. Thompson, William Francis Thompson, Deogracias Villamin Villadolid, Yojiro Wakiya, Frank Walter Weymouth, Thomas Marion Williams, and Albert Jefferson Woolman.*

Those marked with an asterisk have also been students of Doctor Evermann, as have also the following: Frederick Morton Chamberlain, Ulysses Orange Cox, Samuel Frederick Hildebrand, Chancey Juday, William Converse Kendall, Clarence Hamilton Kennedy, William J. Moenkhaus, John Treadwell Nichols, Lewis Radcliffe, Josiah Thomas Scovell, and Alfred Cleveland Weed.

Many of these have made important contributions to our knowledge of fishes.

The preparation of this Check List has extended over a period of more than 20 years. It was begun in 1906 by the senior authors,

who worked upon it from time to time, as their other duties permitted, until 1925, when they were joined by the junior author, who thereafter gave most of his time to the work until its completion. The senior authors also have devoted much of their time to it since 1925.

During the progress of the work the authors have endeavored to consult and utilize every record bearing on the fish fauna of America. In addition to this they have received invaluable assistance and suggestions from virtually every present-day student of American fishes, among whom special mention should be made of the late Dr. Charles Henry Gilbert and Prof. John Otterbein Snyder of Stanford University, Dr. William Converse Kendall of the United States Bureau of Fisheries, Dr. Carl Leavitt Hubbs of the University of Michigan, George Sprague Myers of Stanford University, Barton Appler Bean of the United States National Museum, William Francis Thompson of the International Fisheries Commission, Dr. Arthur W. Henn of the Carnegie Museum, John Treadwell Nichols and Dr. E. W. Gudger of the American Museum of Natural History, and many others. To all these we express our indebtedness and grateful appreciation. For clerical assistance of one kind or another, particularly in looking up and verifying references, we are greatly indebted to Miss Jessie E. Drayton, Harry J. Christoffers, and Henry D. Aller, of the United States Bureau of Fisheries. Without the assistance so generously given by these and others, this memoir could hardly have been completed.

DAVID STARR JORDAN.
BARTON WARREN EVERMANN.
HOWARD WALTON CLARK.

STANFORD UNIVERSITY,
CALIFORNIA ACADEMY OF SCIENCES,
January 4, 1928.

Class I. LEPTOCARDII. Lancelets

Order A. CIRROSTOMI. Cirrostomes

Family 1. BRANCHIOSTOMIDÆ. Lancelets

Genus 1. BRANCHIOSTOMA Costa.

Branchiostoma Costa, Cenni Zoologici Napol., 49, 1834 (*B. lubricum* Costa=*lanceolatus* Pallas).

Amphioxus Yarrell, Hist. Brit. Fishes, 1836, 468 (*Limax lanceolatus* Pallas).

Amphioxides ² Gill, Amer. Nat., XXIX, 1895, 458 (*Branchiostoma pelagicum* Gunther).

1. *Branchiostoma virginia* ³ Hubbs. *Virginia lancelet*.
Shallow waters, burying itself in the sand, from Chesapeake Bay to Florida; Beaufort, Cape Lookout, etc.
Branchiostoma virginia Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 105, 8, July 6, 1922, Sewall's Point, Va.
2. *Branchiostoma florida* Hubbs. *Florida lancelet*.
Coasts of Florida; Tampa, Pensacola, Cape Romans, St. Martin's Reef, Snapper Banks, etc.
Branchiostoma florida Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 105, 7, July 6, 1922, Tampa Bay, Fla.
3. *Branchiostoma bermuda* Hubbs. *Bermuda lancelet*.
Bermuda Islands.
Branchiostoma bermuda Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 105, 9, July 6, 1922, Bermuda.
4. *Branchiostoma caribæum* ⁴ Sundevall. *West Indian lancelet*.
West Indies.
Branchiostoma caribæum Sundevall, Vet. Akad. Förhandl., XI, 1853, 12, St. Thomas
5. *Branchiostoma californiense* Gill. *California lancelet*; *California amphioxus*.
Pacific coast of America from San Diego Bay southward to San Luis Gonzales Bay, Gulf of California; reported also from Panama. Abundant in shallow water, buried in the sand.
Branchiostoma californiense Gill, in Andrews, Studies Biol. Lab. Johns Hopkins Univ., V, 1893, 235, San Diego.

Family 2. EPIGONICHTHYIDÆ

Genus 2. ASYMMETRON Andrews.

Asymmetron Andrews, Studies Biol. Lab. Johns Hopkins Univ., V, 1893, 237 (*A. lucayanum* Andrews).

6. *Asymmetron lucayanum* Andrews. *Bahama lancelet*; *Bahama amphioxus*.
West Indies; Bemini and Nassau, Bahamas; Culebra Island; Humacao, Porto Rico; south to Brazil. Common.
Asymmetron lucayanum Andrews, Studies Biol. Lab. Johns Hopkins Univ., V, 1893, 237, Bemini, Bahamas.

² Based on larvæ showing no buccal cirri, found in the open sea near Hawaii

³ Until the recent studies of Hubbs, this species was wrongly identified with the European form (*B. lanceolatum*), which, however, does not occur in America

⁴ Represented on the coast of Brazil (Rio Janeiro, etc.) by *B. platæ* Hubbs.

Class II. CYCLOSTOMI. Lampreys

Order B. HYPEROTRETA. Hagfishes

Family 3. MYXINIDÆ. Hagfishes

Genus 3. MYXINE Linnæus.

Myxine Linnæus, Syst. Nat., ed. X, 1758, 650 (*M. glutinosa* L.).

Gastrobranchus Bloch, Ichth., XII, 1797, 51, tab. 413 (*G. cæcus* Bloch).

Murænoblenna Lacépède, Hist. Nat. Poiss., V, 1803, 647 (*M. olivacea* Lacépède).

Anopsus Rafinesque, Anal. Nat., 1815, 93 (*M. olivacea* Lacépède).

7. *Myxine limosa* Girard. Hagfish; Borer; Sucker; Slimefish.

Newfoundland, south to Cape Cod; Grand Manan Island, New Brunswick; replacing in the western Atlantic the European *M. glutinosa* L.; not abundant.

Myxine limosa Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 224, Grand Manan.

Family 4. EPTATRETIDÆ. Borers

Genus 4. POLISTOTREMA Gill.

Polistotrema Gill, Proc. U. S. Nat. Mus., IV, 1881, 30 (*Gastrobranchus dombey* Cuvier).

8. *Polistotrema stoutii* (Lockington). California hagfish; Lamperina.

Coast of California; common in Monterey Bay; Eel River, Humboldt Co., Calif., and other streams of northern California; San Francisco Bay.

Bdellostoma stoutii Lockington, Amer. Nat., XII, December, 1878, 793, Eel River, Calif.

9. *Polistotrema deani* Evermann and Goldsborough.

Alaska to Santa Barbara, Calif., in deep water.

Polistotrema deani Evermann and Goldsborough, Bull. U. S. Bur. Fish., XXVI, 1906, 225, fig. 1, Albatross station 4238 in Behm Canal off Nose Point in 229 to 231 fathoms.

Polistotrema curtiss-jamesi Townsend and Nichols, Bull. Amer. Mus. Nat. Hist., LII, 1925, 4, off California, lat. 34° to 36° in 440 to 585 fathoms.

Order C. HYPEROARTIA. Lampreys

Family 5. PETROMYZONIDÆ. Lampreys

Genus 5. PETROMYZON Linnæus.

Petromyzon (Artedi) Linnæus, Syst. Nat., ed. X, 1758, 230 (*P. marinus* L.).

Ammocetes Duméril in Cuvier, Règne Anim., ed. I, 1817, 119, variant spelling of *Ammocetus*, a name given to a larval form (*branchialis*).

Bathymyzon Gill, Proc. U. S. Nat. Mus., VI, 1883, 25 (*Bathymyzon bairdii* Gill); probably an old example that has lost teeth.

Oceanomyzon Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 461 (*O. wilsoni* Fowler); like the preceding, supposed by Creaser and Hubbs to be an old individual from the Gulf Stream.

10. *Petromyzon marinus* Linnæus. Great sea lamprey; Lamprey eel; Sucker; Lamprey; River lamprey; Green lamprey; Shad lamprey; Nine-eyes.

Atlantic coasts of Europe and America, southward in America to Chesapeake Bay; abundant northward; ascending streams to spawn.

Petromyzon marinus Linnæus, Syst. Nat., ed. X, 1758, 230, European seas.

Petromyzon americanus LeSueur, Trans. Amer. Phil. Soc. Phila., I, 1818, 383, coast of Massachusetts.

Petromyzon nigricans LeSueur, loc. cit., 385, Massachusetts; small example from fresh water.

Ammocetes bicolor LeSueur, loc. cit., 386, Northampton, Mass. (larva).

? *Petromyzon* (*Bathymyzon*) *bairdii* Gill, Proc. U. S. Nat. Mus., VI, 1883, 254, Gulf Stream, north lat. 40° 02', west long. 68° 50' 30'', in 547 fathoms.

? *Oceanomyzon wilsoni* Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 461; open sea.

11. *Petromyzon marinus dorsatus* Wilder. *Lake lamprey; Mud lamprey*.
Landlocked in Cayuga Lake and other lakes of western and central New York.
Petromyzon marinus dorsatus Wilder, in Jordan and Gilbert, Synop. Fishes North Amer., 1883, 869, Cayuga Lake, N. Y.
- Genus 6. *ICHTHYOMYZON* Girard. *River lampreys*.
Ichthyomyzon Girard, Pac. R. R. Surv., X, 1858, 381 (*Petromyzon argenteus* Kirtland).
Scolecospoma Girard, loc. cit., 385 (*Ammocetes concolor* Kirtland); larva of *argenteus*.
12. *Ichthyomyzon concolor* (Kirtland). *Silver lamprey*.
Great Lakes; upper Mississippi Valley; tributaries of Hudson Bay; south to Arkansas and Louisiana.
Petromyzon argenteus Kirtland, Zool. Ohio, 170, 1838, 197, Big Miami River, Ohio; name preoccupied.
Ammocetes concolor Kirtland, Jour. Bost. Soc. Nat. Hist., III, 1840, 473, plate, Mahoning River; Scioto River; larva.
Petromyzon bdelium Jordan, Cat. Fishes North Amer., 1885, 4; after Kirtland.
Ichthyomyzon castaneus Girard, Pac. R. R. Surv., X, 1858, 381, Galena, Minn.
Ichthyomyzon hirudo Girard, loc. cit., 382, Fort Smith, Ark.
- Genus 7. *REIGHARDINA* Creaser and Hubbs.
Reighardina Creaser and Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 120, 1922, 4 (*Ichthyomyzon fossor* Reighard and Cummins = *Ammocetes unicolor* DeKay).
13. *Reighardina unicolor* (DeKay).
Ammocetes unicolor DeKay, N. Y. Fauna: Fishes, 1842, 383, pl. 79, fig. 250; Lake Champlain.
Ammocetes borealis Agassiz, Lake Superior, 1850, 252, Michigan River, Mich.
Ichthyomyzon fossor Reighard and Cummins, Occ. Papers Mus. Zool. Univ. Mich., 31, 1916, 1 to 12, pl. 1, 2, Michigan.
- Genus 8. *LETHENTERON* Creaser and Hubbs.
Lethenteron Creaser and Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 120, 1922, 6 (*Lampetra wilderi* Gage = *Petromyzon appendix* DeKay).
14. *Lethenteron appendix* (DeKay). *Small black lamprey; Brook lamprey*.
New York to Iowa, in tributaries of the Great Lakes and of the Ohio and Mississippi, south to Virginia.
Petromyzon nigrum Rafinesque, Ichth. Ohiensis, 1820, 84, falls of Ohio River; name preoccupied.
Petromyzon appendix DeKay, N. Y. Fauna: Fishes, 1842, 381, pl. 64, fig. 211, Providence; Hudson Bay.
Lampetra wilderi Gage, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 13, Cayuga Lake, N. Y.
- Genus 9. *TETRAPLEUODON* Creaser and Hubbs.
Tetrapleurodon Creaser and Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 120, 1922, 9 (*Lampetra spadicea* Bean).
15. *Tetrapleurodon spadiceus* (Bean). *Mexican lamprey*.
Known only from the Rio Lerma, near Guanajuato, and from Lago de Chapala, Mexico.
Lampetra spadicea Bean, Proc. U. S. Nat. Mus., X, 1887, 374, Pl. XX, fig. 6, Rio Lerma, Guanajuato, Mexico.
- Genus 10. *ENTOSPHEUS* Gill.
Entosphenus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 331 (*Petromyzon tridentatus* Gairdner).

16. *Entosphenus tridentatus* (Gairdner). *Three-toothed lamprey; Western lamprey*.
Pacific coast of America, from southern California to Unalaska; ascending streams even to the headwaters of the Columbia for spawning purposes.
Petromyzon tridentatus Gairdner, in Richardson, Fauna Bor.-Amer., 1836, 293, falls of the Willamette River, Oreg.
Petromyzon lvidus Girard, Pac. R. R. Surv., X, 1858, Pt. 4, Fishes, 379, Willamette River, Oreg.
Petromyzon astori Girard, loc. cit., 380, Astoria, Oreg.
Ammocetes cibarius Girard, loc. cit., 383 (larva).
17. *Entosphenus ciliatus* (Ayres).
Streams of northern California.
Petromyzon ciliatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 43, San Francisco.
Petromyzon epihexodon Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 331, Fort Reading, Calif.
- Genus 11. LAMPETRA (Bonnaterre) Gray. *Brook lampreys*.
Lampetra Bonnaterre, Enc. Meth., 1788; no paging; no type, used for all.
Lampreda Rafinesque, Anal. Nat., 1815, 94 (no species named; no description).
Præcus Rafinesque, loc. cit. (no species named); no description.
Lampetra Gray, Proc. Zool. Soc. Lond., 1851, 236 (*Petromyzon fluviatilis* L.).
18. *Lampetra ayresii* (Günther). *Western lamprey*.
Pacific coast of North America from the Fraser River southward to the Sacramento; very close to *L. fluviatilis* of Europe.
Petromyzon plumbeus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 27, San Francisco; name preoccupied.
Petromyzon ayresii Günther, Cat., VIII, 1870, 505, San Francisco; substitute for *plumbeus* Ayres, preoccupied.
19. *Lampetra borealis* (Girard). *Arctic lamprey*.
Streams of northern Alaska and Kamchatka.
Petromyzon borealis Girard, Pac. R. R. Surv., X, 1858, Pt. 4, Fishes, 377, Great Slave Lake; not *Ammocetes borealis* Agassiz.
Ammocetes aureus Bean, Proc. U. S. Nat. Mus., IV, 1881, 159, Yukon River.
- Genus 12. OKKELBERGIA Creaser and Hubbs.
Okkelbergia Creaser and Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 120, 1922, 8 (*Petromyzon lamotteni* LeSueur).
20. *Okkelbergia lamotteni* (LeSueur).
Petromyzon lamotteni LeSueur, New York Fauna: Fishes, 1842, 382, pl. 79, fig. 249; no locality.
Ammocetes æpyptera C. C. Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 327 (type redescribed by Fowler, *ibid.*, 1901, 328; 1907, 466).

Class III. ELASMOBRANCHII. (Selachii, Plagiostomata, Chondropterygii, Antacca). Sharks; Skates and rays; Chimæras

Subclass SELACHII. Sharks

Order D. CESTRACIONTES

Suborder PROARTHRI

Family 6. HETERODONTIDÆ. Bullhead sharks

- Genus 13. GYROPLEURODUS Gill.
Gyroleurodus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 331 and 489 (*Cestracion francisci* Girard).
Tropidodus Gill, loc. cit., 489 (*Cestracion pantherinus* Valenciennes = *Cestracion quoyi* Fréminville).
21. *Gyroleurodus francisci* (Girard). *Bullhead shark*.
Coast of California; abundant south of Point Conception.
Cestracion francisci Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 196, Monterey Bay, Calif.

Order E. NOTIDANI. (Diplospondyli)

Suborder OPISTHARTHRI

Family 7. HEXANCHIDÆ. Cow sharks

Genus 14. HEXANCHUS Rafinesque.

Hexanchus Rafinesque, Caratteri, etc., 1810, 14 (*Squalus griseus* Bonnaterre).

Notidanus Cuvier, Règne Anim., ed. 1, 1817, 128 (*Squalus griseus* L.).

Monopterhinus Blainville, Faune Française, 1828, 77 (*Squalus griseus* Gmelin).

22. *Hexanchus griseus* (Bonnaterre). Cow shark; *Caffa bota*.

Mediterranean to Scotland; Cuba.

Squalus griseus Bonnaterre, Ichth. R. 1788, 9; Mediterranean; after *le grisé* Bonnaterre.

Squalus griseus Gmelin, Syst. Nat., I, 1795, 1789; after Broussonet.

23. *Hexanchus corinus* Jordan and Gilbert. Shovel-nosed shark.

Pacific coast of America; Monterey Bay to Puget Sound.

Hexanchus corinus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880 (December 21), 352, Neah Bay, Wash., and Soquel, Calif.

Genus 15. NOTORYNCHUS Ayres.

Notorynchus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 73 (*Notorynchus maculatus* Ayres).

24. *Notorynchus maculatus*^a Ayres.

Pacific coast of the United States, from Monterey Bay northward to Washington; common northward, especially in Humboldt and San Francisco Bays.

Notorynchus maculatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 72, San Francisco Bay.

Notorhynchus borealis Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 150, Nisqually, Wash.

Order F. EUSELACHII

Suborder GALEI. True sharks

Family 8. SCYLLIORHINIDÆ. Cat sharks

Genus 16. SCYLLIORHINUS Blainville. Roussettes.

Catulus Valmont, Dict. Raisonné, IV, 1768, 451 (*C. saxatilis* Valmont=*Squalus stellaris* L.); non binominal.

Catulus Smith, Proc. Zool. Soc. Lond., 1837, 85 (*Scyllium capense* Smith); name preoccupied by *Catulus* Knipphof, 1759, a genus of insects.

Scylliorhinus Blainville, Jour. Phys., 1816, 263 (*Squalus canicula* L., as restricted); spelled *Scylliorhinus* by Garman.

Scyllium Cuvier, Règne Anim., ed. I, 1817, 124 (*Squalus canicula* L., as restricted).

25. *Scylliorhinus retifer* (Garman).

Gulf Stream in deep water; occasionally off South Atlantic coast.

Scyllium retiferum Garman, Bull. Mus. Comp. Zool., VIII, 1881, 233, off coast of Virginia in deep water.

26. *Scylliorhinus cephalus* (Gilbert).

Gulf of California and southward in deep water.

Catulus cephalus Gilbert, Proc. U. S. Nat. Mus., XV, 1891, 541, deep water near the Revillagigedo Islands and in the Gulf of California.

Genus 17. CEPHALOSCYLLIUM Gill.

Cephaloscyllium Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 407, 412 (*Scyllium laticeps* A. Duméril).

^a Regarded by Garman as identical with *N. platycephalus* (Tenore) of the Mediterranean.

27. *Cephaloscyllium uter* (Jordan and Gilbert). *Swell shark*.
 Monterey to San Diego; very abundant in Santa Barbara channel.
Catulus uter Jordan and Gilbert, in Jordan and Evermann, Fishes North
 and Mid. Amer., 1896, 25; Monterey, San Diego, and Santa Barbara
 Channel.

Genus 18. *APRISTURUS* Garman.

Apristurus Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI,
 1913, 96 (*Scylliorhinus indicus* Brauer).

28. *Apristurus brunneus* (Gilbert).
 Gulf of California and northward in deep water; Monterey Bay; Puget
 Sound; Hood Canal.
Catulus brunneus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 542, Gulf of
 California.

29. *Apristurus profundorum* (Goode and Bean).
 North Atlantic, in deep water.
Scylliorhinus profundorum Goode and Bean, Oceanic Ichth., 1896, 17, Gulf
 Stream.

Genus 19. *PRISTIURUS* (Bonaparte) Müller and Henle.

Pristiurus Bonaparte, Prospetto del Systema d'Ittiologia Generale, 1831,
 121 (*Galeus melanostomus* Rafinesque); name only; defined in Icono-
 graphia della Fauna Italica, 1841 (no paging).

Pristiurus Müller and Henle, Mag. Nat. Hist., III, Jan., 1838, 341 (*Galeus
 melanostomus* Rafinesque).

Pristidurus Bonaparte, Mem. Soc. Sci. Neufchâtel, II, 1839, 11, variant
 spelling.

30. *Pristiurus xaniurus* (Gilbert).
 Pacific coast of southern and Lower California; abundant in deep water.
Catulus xaniurus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 540, off
 southern and Lower California in 184 to 684 fathoms.

31. *Pristiurus aræ* Nichols.
 Off the coast of Florida.
Pristiurus aræ Nichols, Amer. Mus. Novit., No. 256, March 12, 1927, 1,
 off the coast of Florida.

Family 9. *GINGLYMOSTOMIDÆ*. Nurse sharks

Genus 20. *GINGLYMOSTOMA* Müller and Henle.

Ginglymostoma Müller and Henle, Wiegmann's Archiv, I, 1837, 396
 (*Squalus cirratus* Gmelin).

32. *Ginglymostoma cirratum* (Gmelin). *Nurse shark; Gata*.
 West Indies and the west coast of Mexico; Cape Verde Islands; occasional
 on our South Atlantic coast.
Squalus cirratus Gmelin, Syst. Nat., I, 1492, 1788, American seas; after
 Broussonet.
Squalus punctatus Bloch and Schneider, Syst. Ichth., 1801, 134, Cuba;
 after Gata Hispanus of Parra
Squalus punctulatus Bloch and Schneider, loc. cit., 549, Cayenne.
Squalus argus Bancroft, Zool. Jour., V, 1832, 82, West Indies.
Ginglymostoma fulvum Poey, Memorias, II, 1861, 342, Habana; young.
Ginglymostoma caboverdianus Capello, Jour. Sci. Math. Phys. Nat. Lisbon,
 1867, 167, Cape Verde Islands.

Family 10. *PSEUDOTRIAKIDÆ*

Genus 21. *PSEUDOTRIAKIS* Capello.

Pseudotriakis Capello, Jour. Sci. Math. Phys. Nat. Lisbon, IV, 1867, 321
 (*Pseudotriakis microdon* Capello).

33. *Pseudotriakis microdon* Capello. *Peixe carago*.
 Portugal; Amagansett, Long Island.
Pseudotriakis microdon Capello, Jour. Sci. Math. Phys. Nat. Lisbon, 1868,
 321, pl. 5, fig. 1, Portugal.

Family 11. CARCHARIIDÆ. Gray sharks

Genus 22. CYNIAS Gill.

Cynias Gill, Proc. U. S. Nat. Mus., XXVI, 1903, 960 (*Squalus canis* Mitchell); embryos without placental attachment.

Mustelus Valmont de Bomare, Dict. Raisonné, ed. II, 1768, 786 (*Galeus asterias* Valmont = *Squalus canis* Mitchell; not given as a scientific name but as Latin vernacular).

34. *Cynias canis* ⁶ (Mitchill). *Dog shark; Dogfish; Smooth dogfish; Blue dog.* Cape Cod to Cuba; abundant on the Atlantic coast of both continents. *Squalus canis* Mitchell, Trans. Lit. Phil. Soc. N. Y., 1815, 486, New York. *Mustelus asterias* Cloquet, Dict. Raisonné, 1820, 407, Europe. *Mustellus stellatus* Risso, Europ. Mérid., III, 1826, 1126, Nice. *Mustelus plebejus* Bonaparte, Fauna Italica, Pesci, Fasci, Tome III, 1834, 43, pl. 132, Italy.

35. *Cynias lunulatus* (Jordan and Gilbert). *Gato.*

West coast of Mexico; common in Gulf of California.

Mustelus lunulatus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 108, Mazatlan, Mexico.

Genus 23. MUSTELUS Linck.

Mustelus Linck, Magazin für Neueste aus Physik und Naturgeschichte, 1790, 31 (*Squalus mustelus* Linnæus); not of Valmont de Bomare.

Galeus ⁷ Rafinesque, Caratteri, 1801, 13 (*Squalus mustelus* L., as restricted by Jordan and Gilbert).

Mustelus Leach, Observ. Squalus, 1813, 62 (*Squalus mustelus* L.).

Mustelus Cuvier, Règne Anim., ed. I, 1817, 128 (*Squalus mustelus* Cuv.).

Pleuracromylon Gill, Proc., Acad. Nat. Sci. Phila., XVI, 1864, 148, (*Mustelus lævis* Risso); embryo attached by a placenta.

36. *Mustelus mustelus* (Linnæus). *Smooth hound; Boce dulce.*

Southern Europe; occasional in America, lately taken on coast of New Jersey.

Squalus mustelus Linnæus, Syst. Nat., ed. X, 1758, 235, Europe.

Mustelus lævis Risso, Europ. Mérid., 1826, 327, Nice; and of European writers generally.

Galeus mustelus Rafinesque, Caratteri, 1910.

37. *Mustelus californicus* Gill.

California, north to Cape Mendocino.

Mustelus californicus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 148, San Francisco.

38. *Mustelus dorsalis* Gill.

Panama and neighboring waters, north to Gulf of California.

Mustelus dorsalis Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 149, Panama.

Mustelus mento Cope, Proc. Amer. Phil. Soc., XVII, 1877, 47, Pacific Ocean at Pacasmayo, Peru.

Genus 24. RHINOTRIACIS Gill.

Rhinotriacis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 486 (*Rhinotriacis henlei* Gill).

⁶ The European synonymy of *Mustelus mustelus* and *Cynias canis* is in a state of great confusion, as few of the earlier writers recognized any difference between the two species externally very much alike.

⁷ The genus *Galeus* of Rafinesque was based on species with spiracles, differing from *Squalus* by the presence of an anal fin. One new species is described, *Galeus melastomus*, later type of Bonaparte's genus *Pristiurus*; but on the same page and the next page Rafinesque speaks of other species—*Galeus vulpecula*, *G. catulus*, and *G. mustelus*. It is plain that *Squalus galeus* L. was his intended type, after the general custom of writers who elevated species to the rank of genus, but he does not mention that species in the paper nor in his Indices, and as *Squalus galeus* was therefore not available as type, Jordan and Gilbert selected *Squalus mustelus* for that purpose, thus making *Galeus* Rafinesque a synonym of *Mustelus* Leach, rather than to displace *Pristiurus*. Apparently this procedure was allowable, unless some earlier author had chosen either *Pristiurus melastomus* or *Scylliorhinus catulus* (*canicula*) for that purpose.

39. *Rhinotriakis henlei* Gill.
Coast of California from Humboldt Bay to Monterey; rather rare.
Rhinotriakis henlei Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 486, San Francisco.
- Genus 25. **TRIAKIS** Müller and Henle.
Triakis Müller and Henle, Mag. Nat. Hist., II, 1838, 36 (*Triakis scyllium*); corrected to *Triakis* by authors.
40. *Triakis semifasciatum* Girard. *Cat shark*; *Leopard shark*.
Cape Mendocino to San Diego; common.
Triakis semifasciatum Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 196, Presidio de San Francisco.
Mustelus felis Ayres, Proc. Calif. Acad. Nat. Sci., 1854, 17, San Francisco.
- Genus 26. **GALEORHINUS** Blainville.
Galeorhinus Blainville, Bull. Soc. Philom., 1816, 121 (*Squalus galeus* L.; restricted by Gill).
Eugaleus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 148 (*Squalus galeus* L.); the first species named by Blainville being *Squalus mustelus*, for which reason Garman uses *Galeorhinus* as the name of the genus here called *Mustelus*.
41. *Galeorhinus zyopterus* Jordan and Gilbert. *Oil shark*; *Soup-fish shark*.
Coast of southern and Lower California, from San Francisco to Cedros Island; very abundant.
Galeorhinus zyopterus Jordan and Gilbert, Synop. Fishes North Amer., 1883, 871, San Pedro, Calif.
- Genus 27. **TRIÆNODON** Müller and Henle.
Triænodon Müller and Henle, Plagiostomen, 1841, 55 (*Carcharias obesus* Rüppell).
42. *Triænodon obesus* (Rüppell).
Tropical Pacific; reported from Cocos Island.
Carcharias obesus Rüppell, Neue Wirbel., Fisch., 1835, 64, pl. 18, fig. 2, Djidda, Arabian Coast of Red Sea.
- Genus 28. **GALEOCERDO** Müller and Henle
Galeocerdo Müller and Henle, Plagiostomen, 1838, 59 (*Galeocerdo tigrinus* Müller and Henle).
Boreogaleus Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 411 (*Squalus arcticus* Faber).
43. *Galeocerdo arcticus* (Faber). *Spotted shark*; *Leopard shark*; *Tiger shark*; *Tigrone*.
Tropical seas; occasionally to Iceland, Cape Cod, and San Diego.
Squalus arcticus Faber, Fische Islands, 1829, 17, Iceland and neighboring seas.
Galeocerdo tigrinus Müller and Henle, Plagiostomen, 1838, 59, Pondicherry.
Galeus maculatus Ranzani, De Novis Speciebus Piscium, Dissert. Prima, in Novi Com. Acad. Boron., IV, 1840, 68, Brazil.
- Genus 29. **HEMIGALEUS** Bleeker.
Hemigaleus Bleeker, Verh. Bot. Gen., XIV, 1852, 45 (*Hemigaleus microstoma* Bleeker); not *Hemigalea*, an earlier genus of mammals.
44. *Hemigaleus pectoralis* Garman.
Coast of New England, probably in the Gulf Stream.
Hemigaleus pectoralis Garman, Bull. Mus. Comp. Zool., XLIV, 1906, 203, "Aquarial Gardens," the fishes in which were taken off Massachusetts and Rhode Island.
- Genus 30. **SCOLIODON** Müller and Henle.
Scoliodon Müller and Henle, Wiegmann's Archiv, I, 1837, 398 (*Scoliodon laticaudus* Müller and Henle).
45. *Scoliodon terræ-novæ* (Richardson). *Sharp-nosed shark*; *Newfoundland shark*.
Cape Cod to Florida; very common southward along the Atlantic coast.
Squalus (*Carcharias*) *terræ-novæ* Richardson, Fauna Bor.-Amer., III, 1836, 289, Newfoundland; an error, probably for Florida.
Squalus punctatus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 483, New York; not of Bloch and Schneider.

46. *Scoliodon lalandi* (Muller and Henle).
West Indies to Brazil.
Carcharias lalandi Muller and Henle, Plagiostomen, 1838, 30, Rio Janeiro; Martinique; Guadeloupe.
47. *Scoliodon porosus* (Poey).
Cuba.
Squalus porosus Poey, Memorias, II, 1861, 339, tab. 19, L. 11, 12, Habana.
48. *Scoliodon longurio* (Jordan and Gilbert).
Pacific coast of Mexico.
Carcharias longurio Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 106, Mazatlan, Mexico.
- Genus 31. *APRIONODON* Gill.
Aprion Muller and Henle, Plagiostomen, 1838-1841, 31 (*Carcharias isodon* Muller and Henle); preoccupied.
Aprionodon Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 411 (*Aprionodon punctatus* Gill; not *Squalus punctatus* Mitchell = *Carcharias isodon* Muller and Henle).
49. *Aprionodon isodon* (Müller and Henle).
Atlantic Ocean; recorded from New York, Virginia, and Cuba; probably rare.
Carcharias isodon Muller and Henle, Plagiostomen, 1838, 32, New York.
Carcharias punctatus Gunther, Cat., VIII, 1870, New York; not *Squalus punctatus* Mitchell, which is *Scoliodon terre-novæ*.
- Genus 32. *CARCHARIAS* Rafinesque. Gray shark; Requén.
Carcharias Rafinesque, Caratteri, 1810, 10 (*Squalus carcharias* L. intended, but *Carcharias taurus* the only species mentioned).
Carcharhinus Blainville, Bull. Soc. Philom., 1816, 264 (*Squalus commersonii* Blainville, apparently the same as *Carcharias lamia* Rafinesque).
Carcharias Cuvier, Règne Anim., ed. I, 1817, 125 ("Squalus carcharias," *Carcharias lamia* intended); not of Rafinesque.
Eulamia Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 401 (*Carcharias lamia* Rafinesque; *Squalus commersonii* Blainville being intended = *Carcharias lamia* Rafinesque was *Carcharodon carcharias* Rafinesque, the specific name being a substitute for *Squalus carcharias* L.).
Platyodon Gill, loc. cit. (*Carcharias menisorrhah* Muller and Henle).
Isogomphodon Gill, loc. cit. (*Carcharias oxyrhynchus* Muller and Henle).
Lamiopsis Gill, loc. cit. (*Carcharias temminckii* Muller and Henle).
Isoplagiodon Gill, loc. cit. (*Carcharias sorrah* Muller and Henle).
50. *Carcharias commersonii* (Blainville). Cub shark; Requén; Requén;
Lamia; Common gray shark.
Tropical parts of the Atlantic; common northward to the Florida Keys and in the Mediterranean; rarely straying northward to New York in the Gulf Stream.
Squalus commersonii Blainville, Bull. Soc. Philom., 1816, 121, Mediterranean.
Carcharias lamia Risso, Europ. Mérid., III, 1826, 19, Nice; not of Rafinesque.
Squalus longimanus Poey, Memorias, II, 1861, 338, Cuba.
51. *Carcharias lamiella* Jordan and Gilbert. Bay shark.
San Diego Bay and southward along the Mexican coast.
Carcharias lamiella Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 110, San Diego; regarded by Garman as identical with *C. commersonii* of the Atlantic.
52. *Carcharias platyrhynchus* (Gilbert).
Magdalena Bay, Lower California, to Galapagos Islands and the Revillagigedo Islands
Eulamia platyrhynchus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 543, Clarion Island; Magdalena Bay; regarded by Garman as identical with *C. commersonii*.
53. *Carcharias platyodon* (Poey). Cub shark.
Cuba to Texas; abundant in the Gulf of Mexico
Squalus platyodon Poey, Memorias, II, 1861, 336, Habana.
Squalus obtusus Poey, loc. cit., 337, Habana.

54. *Carcharias milberti* Valenciennes. *Small blue shark; White shark.*
Cape Cod to Florida; not rare; recorded from the Mediterranean.
Carcharias (Prionodon) milberti Valenciennes, in Muller and Henle, Plagiostomen, 1838, 38, New York.
Carcharias cæruleus DeKay, N. Y. Fauna: Fishes, 1842, 49, New York.
Lamna caudata DeKay, loc. cit., 354, Brenton Reef, R. I.
55. *Carcharias obscurus* (LeSueur). *Dusky shark.*
North and Middle Atlantic; frequently on our coast.
Squalus obscurus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 223, New York.
Carcharias falcipinnis Lowe, Proc. Zool. Soc. Lond., 1839, 90, Madeira.
Prionodon obvelatus Valenciennes, Poiss. Îles Canaries, 1839, 103, Canaries.
56. *Carcharias azureus* Gilbert and Starks.
Carcharias azureus Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 11, Pl. II, fig. 5, Panama.
57. *Carcharias nicaraguensis* (Gill and Bransford). *Nigrone; Fresh-water shark.*
Lake Nicaragua (landlocked); also in the Bay of Panama.
Eulamia nicaraguensis Gill and Bransford, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 190, Lake Nicaragua.
58. *Carcharias henlei* Valenciennes.
Coast of Brazil, northward to Guiana.
Carcharias henlei Valenciennes, in Müller and Henle, Plagiostomen, 1838, 46, Cayenne.
Carcharias porosus Ranzani, De Novis Speciebus Piscium, Dissert. Prima., in Novi Com. Acad. Bonon., IV, 1840, 70, Brazil.
59. *Carcharias cerdale* (Gilbert).
Abundant at Panama.
Carcharhinus cerdale Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., III, 1898, 2746, Panama; united by Garman with *C. menisorrhah* from the East Indies.
60. *Carcharias falciformis* Bibron. *Cazon de playa (wharf) shark.*
Cuba and neighboring waters.
Carcharias falciformis Bibron, in Muller and Henle, Plagiostomen, 1838, 47, Cuba.
Squalus tiburo Poey, Memorias, II, 1861, 331, Habana.
61. *Carcharias remotus* Valenciennes.
West Indies and Brazil.
Carcharias remotus Valenciennes, in Duméril, Hist. Nat. Poiss., I, 1865, 374.
Platypodon perezii Poey, Anal. Soc. Española Hist. Nat., V, 194, pl. 9, figs. 2, 3, Cuba.
62. *Carcharias acronotus* (Poey).
Cuba.
Squalus acronotus Poey, Memorias, II, 1861, 335, Cuba.
63. *Carcharias velox* (Gilbert).
Panama.
Carcharhinus velox Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2747, Panama.
64. *Carcharias oxyrhynchus* Müller and Henle.
Surinam.
Carcharias oxyrhynchus Müller and Henle, Plagiostomen, 1838, 41, Surinam.
65. *Carcharias limbatus* Müller and Henle. *Carconetta.*
Tropical seas north to Florida; a stray example taken at Woods Hole, Mass.
Carcharias (Prionodon) limbatus Müller and Henle, Plagiostomen, 1838, 49, Martinique.
Isogomphodon maculipinnis Poey, Repertorio, I, 1866, 191, Cuba.
Carcharias (Prionodon) millerii Steindachner, Sitzungs. Kais. Akad. Wiss. Wien, LVI, 1867, 356, West Indies.
Carcharias microps Lowe, Proc. Zool. Soc. Lond., 1840, 38, Madeira.
Prionodon cucuri Castelnau, Anim. Amer. Sud. Poiss., 1855, 99, Bahia.

66. *Carcharias æthalarus* Jordan and Gilbert.
Mazatlan to Panama.
Carcharias æthalarus Jordan and Gilbert, Proc., U. S. Nat. Mus., V, 1882, 104, Mazatlan; regarded by Garman as inseparable from *C. limbatus*.
67. *Carcharias natator* (Meek and Hildebrand).
Gulf of Panama.
Carcharhinus natator Meek and Hildebrand, Marine Fishes of Panama, Field Mus. Nat. Hist., Zool., XV, Pt. I, Dec. 20, 1923, 40, pl. 1, fig. 1, Panama City fish market.
68. *Carcharias fronto* Jordan and Gilbert. *Tiburón*.
Pacific coast of Mexico.
Carcharias fronto Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 102, Mazatlan. Regarded by Garman as inseparable from *C. milberti*.
- Genus 33. **HYPOPRION** Müller and Henle.*
Hypoprion Müller and Henle, Plagiostomen, 1838, 34 (*Carcharias maculoti* Müller and Henle).
Hypoprionodon Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 401 (*Carcharias hemiodon* Valenciennes).
69. *Hypoprion brevirostris* Poey.
West Indies, north to Charleston and Pensacola.
Hypoprion brevirostris Poey, Repertorio, II, 1868, 451, Tab. 4, Cuba.
70. *Hypoprion signatus* Poey.
Cuba.
Hypoprion signatus Poey, Synop., 1868, 452, Cuba.
Hypoprion longirostris Poey, Enumeratio, 1876, 199, Cuba.
- Genus 34. **PRIONACE**† Cantor.
Galeus Valmont de Bomare, Dict. Raisonné, I, 1768, 371, "Cagnot bleu, *Galeus glaucus* Valmont"; Latin vernacular.
Prionodon Muller and Henle, Plagiostomen, 1838, 36 (*Squalus glaucus* L.); name preoccupied.
Prionace Cantor, Malayan Fishes, 1850, 399, substitute for *Prionodon* (*Squalus glaucus* Linnæus).
Cynocephalus Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 401 (*Squalus glaucus* L.); after Klein, whose names are not binomial.
71. *Prionace glauca* (Linnæus). *Great blue shark*.
A large shark of the warm seas, occasionally taken on our coasts, both Atlantic and Pacific.
Squalus glaucus Linnæus, Syst. Nat., ed. X, 1758, 235, Seas of Europe.
Squalus cæruleus Blainville, Faune Française, 1828, 91, Mediterranean.
Squalus hirundinaceus Valenciennes, in Müller and Henle, Plagiostomen, 1838, 37, Brazil.

Family 12. SPHYRNIDÆ. Hammer-headed sharks

- Genus 35. **RENICEPS** Gill.
Reniceps Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 403 (*Squalus tiburo* Linnæus).
72. *Reniceps tiburo* (Linnæus). *Shovelhead shark*; *Bonnethead shark*.
Atlantic Ocean, abundant from Long Island southward; Pacific Ocean ranging from Mazatlan and Bay of Panama to China.
Squalus tiburo Linnæus, Syst. Nat., ed. X, 1758, 234, America.
- Genus 36. **PLATYSQUALUS** Swainson.
Platysqualus Swainson, Class. Anim., II, 1839, 318 (*Zygæna tudes* Cuvier, in Valenciennes); misnamed *Zygæna tiburo*.

* Müller and Henle's Systematische Beschreibung der Plagiostomen was published from 1838 to 1841. Garman, probably correctly, gives this and other genera of sharks the date of 1841. Some of them appeared earlier in periodicals.

† The reasons for not accepting the names of Valmont de Bomare in scientific nomenclature are given by Jordan and Evermann, Genera of Fishes, I, p. 27. These names (*Galeus*, *Vulpecula*, *Catulus*, and *Mustelus*) are binomial only by accident, and not accepted by the author himself as generic in his list of genera (1791). These are, however, accepted by Garman (Plagiostomia, 1913).

73. *Platysqualus tudes* (Cuvier).
Gulf of California to Bay of Panama; West Indies; Mediterranean Sea and Indian Ocean.
Zygæna tudes Cuvier, in Valenciennes, Mém. Mus. Hist. Nat., IX, 1822, 225, Nice; after *Pantouffier* of Risso.
- Genus 37. *SPHYRNA* Rafinesque.
Sphyrna Rafinesque, Indice, 1810, 60 (*Squalus zygæna* Linnæus).
Cestrorhinus Blanville, Jour. Phys., 1816, 264 (*Squalus zygæna* Linnæus).
Zygæna Cuvier, Règne Anim., ed. I, 1817, 127 (*Squalus zygæna* Linnæus); name preoccupied.
Sphyrichthys Thienemann, Lehrbuch der Zoologie, III, 1828, 408 (*Squalus zygæna* Linnæus).
Eusphyræ Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 412 (*Squalus zygæna* Linnæus).
Cestracion Garman, Plagiostomen, 1813, 133 (*Squalus zygæna* L.); after Klein; not binomial; not of Cuvier.
74. *Sphyrna zygæna* (Linnæus). *Hammerhead shark; Hammerhead; Shovelnose shark.*
Cape Cod and Point Conception southward; Bay of Panama and Mazatlan in Pacific; and across to Hawaii and Japan; abundant.
Squalus zygæna Linnæus, Syst. Nat., ed. X, 1758, 234, Europe; America.
Zygæna malleus Valenciennes, Mém. Mus. Hist. Nat., IX, 1822, 223, pl. 11, fig. 1.
Zygæna lewini Lord, in Griffith, Animal Kingdom, X, 1834, 640, pl. 50, New Holland.
Zygæna subarcuata Storer, Proc. Bost. Soc. Nat. Hist., III, 1848, 71, Cape Cod.

Family 13. RHINEODONTIDÆ. Whale sharks

- Genus 38. *RHINEODON* Smith.
Rhincodon Smith, Zool. Jour. Lond., No. XVI, 1829, 443 (*R. typicus* Smith); misprint for *Rhineodon*
Micristodus Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 177 (*M. punctatus* Gill).
Rhinodon and *Rhincodon* of authors.
75. *Rhineodon typicus* Smith. *Whale shark; Mhor; Chagrín; Tiburon ballenas; Tint reu; East Indian basking shark.*
Coast of Florida; Gulf of California; Cape of Good Hope; Japan to China; India; rare but widely diffused.
Rhincodon typicus Smith, Zool. Jour. Lond., No. XVI, 1829, 443, 444, Table Bay, Cape of Good Hope.
Micristodus punctatus Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 177, Gulf of California.

Family 14. ALOPIIDÆ. Thresher sharks

- Genus 39. *ALOPIAS* Rafinesque.
Vulpecula Valmont de Bomare, Dict. Raisonné, III, 1768, 740 (*V. marina* Valmont); Latin vernacular not intended as a generic name.
Alopias Rafinesque, Caratteri, 1810, 12 (*Alopias macrourus* Rafinesque = *Alopecias vulpes* Gunther).
Alopecias Müller and Henle, Plagiostomen, 1838, 74, amended orthography.
76. *Alopias vulpinus* (Bonnaterre). *Thresher; Fox shark; Swingletail; Longtail shark; Fish shark; Sea fox; Sea ape; Pez zorro; Pesca pavone.*
Mediterranean, Atlantic, and Pacific.
Vulpecula marina Valmont, Dict. Raisonné, III, 740; not eligible.
Squalus vulpinus Bonnaterre, Tableau Encycl. Ichth., 1788, 9, Mediterranean; after Pennant.
Squalus vulpes Gmelin, Syst. Nat., I, 1789, 1496, Mediterranean; after Pennant.
Alopias macrourus Rafinesque, Caratteri, 1810, 12, Sicily.
Galeus vulpecula Rafinesque, Caratteri, 1810, 15; passing notice.

Family 15. ODONTASPIDÆ. Sand sharks

Genus 40. ODONTASPIS Agassiz.

Odontaspis Agassiz, Poiss. Fossiles, III, 1836, 87 (*Squalus ferox* Risso).

77. *Odontaspis littoralis* (Mitchill). Sand shark; Shovelnose shark; Horned dogfish; Picked dogfish; Dogfish; Bone-dog; Skittle-dog; Hoe.
Atlantic coast between Cape Cod and Cape Hatteras; regarded by Garman as not separable from *Carcharias taurus* Rafinesque of the Mediterranean.
Squalus americanus Mitchill, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 483, New York; not of Shaw.
Squalus littoralis and *Squalus macrodus* Mitchill, Amer. Month. Mag., II, 1818, 328, New York.
Carcharias griseus Ayres, Bost. Jour. Nat. Hist., IV, 1843, 288, Long Island.

Family 16. LAMNIDÆ. Mackerel sharks; Man-eaters

Genus 41. LAMNA Cuvier. Porbeagles.

Lamna Cuvier, Règne Anim., ed. I, 1817, 126 (*Squalus cornubicus* Gmelin).

Lamia Risso, Europ. Mérid., III, 1826, 123 (*Squalus cornubicus* Gmelin); name preoccupied.

Selanonius Fleming, Brit. Anim., 1828, 169 (*Selanonius walkeri* Fleming = *Squalus cornubicus* Gmelin).

78. *Lamna nasus* (Bonnaterre). Porbeagle; Mackerel shark; Salmon shark.
North Atlantic and North Pacific; abundant in central and southeast Alaska.
Squalus nasus Bonnaterre, Tableau Encycl., Ichth., 1788, 10, shores of Cornwall.
Squalus cornubicus Gmelin, Syst. Nat., I, 1789, 1497, shores of Cornwall.
Squalus pennanti Walbaum, Artedi Pisc., 1792, 517, Cornwall; after Pennant.
Squalus monensis Shaw, Gen. Zool., V, 1804, 350, Anglesea.
Selanonius walkeri Fleming, Brit. Anim., 1828, 169, Lochfyne, Argyleshire.

Genus 42. ISURUS Rafinesque.

Isurus Rafinesque, Caratteri, 1810, 11 (*Isurus oxyrinchus* Rafinesque).

Oxyrhina Agassiz, Poiss. Fossiles, III, 1836, 276 (*Isurus spallanzani* Rafinesque = *Isurus oxyrinchus* Rafinesque).

Isoropsis Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 398, 409 (*I. dekayi* Gill).

79. *Isurus punctatus* (Storer). Mackerel shark.
Atlantic coast of United States; common.
Lamna punctata Storer, Rept. Fishes Mass., 1839, 185, pl. 3, fig. 2, Massachusetts.
Oxyrhina dekayi Gill, Cat. Fish. E. Coast N. Amer., Suppl. Proc. Acad. Nat. Sci. Phila., XIII, 1861, 60, after Storer; name misprint for *dekayi*.
80. *Isurus tigris* (Atwood). Mackerel shark; Mackerel porbeagle; Dentuda.
Cape Cod to West Indies; not generally common except in the Gulf of Mexico.
Lamna punctata DeKay, N. Y. Fauna: Fishes, 1842, 352, New York; not *Squalus punctatus* Mitchill.
Isoropsis dekayi Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 409; after DeKay; not *Oxyrhina dekayi* Gill, 1861.
Carcharias tigris Atwood, Proc. Bost. Soc. Nat. Hist., XII, 1869, 268, Provincetown, Mass.

81. *Isurus oxyrinchus* Rafinesque. Mackerel shark; Pesce tondo; Cane de mare.
Mediterranean Sea and neighboring parts of the Atlantic; occasional on the Atlantic coast of America.
Isurus oxyrinchus Rafinesque, Caratteri, 1810, 12, Palermo.
Isurus spallanzani Rafinesque, Indice, 1810, 60, Sicily; after Spallanzani.
Oxyrhina gomphodon Müller and Henle, Plagiostomen, 1838, 68, Atlantic Ocean.
Squalus rostratus Macri, Mem. Ac. Sci. Napoli, I, 1819, 55, Tab. 1, fig. 2, Naples.

Genus 43. *CARCHARODON*¹⁰ Smith. *Man-eater sharks; Great white sharks.*
Carcharodon Andrew Smith, in Müller and Henle, Mag. Nat. Hist., II,
 second article, January, 1838, 37 (*Carcharodon capensis* Smith=
Squalus carcharias Linnæus).

82. *Carcharodon carcharias* (Linnæus). *Man-eater shark; Great white shark.*
 Temperate and tropical parts of Atlantic and Pacific; occasionally on our
 coasts, north to New York and Monterey Bay. A dangerous shark.
Squalus carcharias Linnæus, Syst. Nat., ed. X, 1758, 235, Europe; after
 Artedi.
Carcharias lamia Rafinesque, Indice, 1810, 44, after Linnæus; not of
 Risso and later authors.
Carcharodon verus Agassiz, Poiss. Fossiles, III, 1838, 91.
Carcharodon rondeleti Müller and Henle, Plagiostomen. 1838, 70, Mediter-
 ranean Sea and Atlantic Ocean.
Carcharodon Smith, in Bonaparte, Mem. Soc. Sci. Neufchâtel. II, 1839, 9
 (errata), New York and Monterey Bay
Carcharias atwoodi Storer, Proc. Bost. Soc. Nat. Hist., III, 1848, 72,
 Provincetown.
Carcharodon capensis Smith, Ills. Zool. S. Africa, Pisces, 1849, pl. 4,
 Cape of Good Hope.

Family 17. CETORHINIDÆ. Basking sharks

Genus 44. *CETORHINUS* Blainville.

Cetorhinus Blainville, Bull. Soc. Philom., 1810, 169 (*Squalus gunneri*
 Blainville=*Squalus maximus* Gunner).
Selache Cuvier, Règne Anim., ed. I, 1817, 129 (*Squalus maximus* Gunner).
Polyprosopus Couch, Hist. Brit. Fishes, I, 1862, 67 (*Squalus rashleighanus*
 Couch=*Squalus maximus* Gunner).

83. *Cetorhinus maximus* (Gunner). *Basking shark; Pêlerin; Elephant shark;*
Bone shark; Sailfish; Sunfish; Hoe-mother; Capidoli or oilfish.
 Arctic seas, straying southward to Portugal, Virginia, and California.
Squalus maximus Gunner, Trondhjem Skrift., III, 1765, 33, coast of
 Norway.
Squalus gunneri Blainville, Bull. Sci. Philom., 1810, 256; after Gunner.
Squalus pelegrinus Blainville, loc. cit., 257, Europe.
Squalus homianus Blainville, loc. cit., 257, Europe.
Cetorhinus shavianus Blainville, loc. cit., 264, Europe.
Squalus isodus Saverino Macri, Mem. della R. Ac. Sci. Napoli, I, 1819, 55,
 pls. 1 and 2, Naples.
Squalus elephas LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 343, New
 Jersey.
Squalus rashleighanus Couch, Trans. Linn. Soc., XIV, 1825, 91, Cornwall.
Squalus cetaceus Gronow, Cat. Fishes, 1854, 6, Norway.
Polyprosopus macer Couch, Hist. Brit. Fishes, I, 1877, 68, England.

¹⁰ As to the authorship of the name *Carcharodon*, we have this information from the late Allan R. McCulloch, of the Australian Museum:

"The name *Carcharodon* is not used by Smith. The earliest reference to it is that of Müller and Henle, as quoted above. These authors saw Smith's specimens (p. 33) and perhaps heard his paper read at the Zoological Society in September, 1837. They, therefore, quoted the name as of Smith. But Smith's paper was not published until February 13, 1838, he having read Müller and Henle's paper about a month previously, in which *Carcharodon* appeared, and he evidently withdrew it with names he refers to in his paper." (In lit. June 28, 1920.)

Order G. TECTOSPONDYLII.¹¹ Tectospondylous sharks

Suborder SQUALOIDEI. (Sharks without anal fin)

Family 18. SQUALIDÆ. Dogfishes

Genus 45. SQUALUS Linnæus.

Squalus (Artedi) Linnæus, Syst. Nat., ed. X, 1758, 233 (includes all sharks; *Squalus acanthias* Linnæus made type by subsequent writers, following Rafinesque, 1810).

Acanthorhinus Blainville, Jour. Phys., 1816, 263 (*Squalus acanthias* Linnæus).

Acanthias Risso, Europ. Mérid., III, 1826, 131 (*Squalus acanthias* Linnæus).

84. *Squalus acanthias* Linnæus. *Spiny dogfish; Cod shark; Codfish shark; Thornback shark; Picked dogfish; Bone-dog; Dogfish; Skittle-dog.*

North Atlantic, on both coasts and about Cuba; abundant along shores of Northern and Middle States.

Squalus acanthias Linnæus, Syst. Nat., ed. X, 1758, 233, coast of Europe.

Acanthias vulgaris Risso, Europ. Mérid., III, 1826, 131, Nice.

Acanthias americanus Storer, Synop. N. Amer. Fishes, in Mem. Am. Acad., II, 1846, 506, Massachusetts.

85. *Squalus suckleyi* (Girard). *California dogfish; Dogfish.*

Pacific coast of America, from Santa Barbara to the Aleutian Islands; abundant in southeast and middle Alaska, through to Kamchatka.

Spinax (*Acanthias*) *suckleyi* Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 196, Fort Steilacoom, Wash.

Genus 46. CENTROSCYMNUS Bocage and Capello.

Centroscymnus Bocage and Capello, Proc. Zool. Soc. Lond., 1864, 263 (*C. celolepis* Bocage and Capello).

86. *Centroscymnus celolepis* Bocage and Capello.

Coast of Portugal and neighboring parts of the Atlantic; Gloucester, Mass., and Nova Scotia Banks.

Centroscymnus celolepis Bocage and Capello, Proc. Zool. Soc. Lond., 1864, 263, Portugal.

Genus 47. CENTROSCYLLIUM Müller and Henle.

Centroscyllium Müller and Henle, Plagiostomen, 1838, 191 (*Spinax fabricii* Reinhardt).

87. *Centroscyllium fabricii* (Reinhardt).

Greenland seas; Gloucester and the Nova Scotia Banks.

Spinax fabricii Reinhardt, Dansk. Vidensk. Selsk. Forhandl., III, 1828, XVI, Greenland.

Genus 48. ETMOPTERUS Rafinesque.

Etmopterus Rafinesque, Caratteri, 1810, 14 (*Etmopterus aculeatus* Rafinesque=*Squalus spinax* L.).

Acanthidium Lowe, Proc. Zool. Soc. Lond., 1839, 91 (*Acanthidium pusillum* Lowe, as restricted by Jordan and Evermann, 1896; restricted by Garman, 1912, to *A. calcea*, a species of *Deania* Jordan and Snyder).

88. *Etmopterus hillianus* (Poey).

West Indies; Cuba; St. Kitts.

Spinax hillianus Poey, Memorias, II, 1861, 340, Cuba.

Family 19. SOMNIOSIDÆ. Sleeper sharks

Genus 49. SOMNIOSUS LeSueur.

Somniosus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 222 (*Somniosus brevipinna* LeSueur=*Squalus microcephalus* Bloch and Schneider.)

Lamargus Müller and Henle, Plagiostomen, 1838, 93 (*Squalus borealis* Scoresby=*Squalus microcephalus* Bloch and Schneider).

Leiodon Wood, Proc. Bost. Soc. Nat. Hist., II, 1847, 174 (*Leiodon echinatum* Wood=*Squalus microcephalus* Bloch and Schneider).

Rhinocymnus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 264 (*Scymnus rostratus* Risso).

¹¹ We group together under this name those sharks that, like the rays, lack the anal fin and that have also simplified or degenerate vertebrae, in which the hard material is arranged in rings without radiation.

89. *Somniosus microcephalus* (Bloch and Schneider). *Sleeper shark; Nurse shark.*
 Arctic seas, south to Cape Cod, Oregon, San Francisco Bay, and France.
Squalus microcephalus Bloch and Schneider, Syst. Ichth., 1801, 135, northern seas.
Somniosus brevipinna LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 222, Massachusetts.
Squalus borealis Scoresby, Arct. Reg., I, 1820, 538, pl. 15, Arctic.
Squalus norvegicus Blainville, Faune Française, 1828, 61, Norway.
Squalus glacialis Faber, Natur der Fische Islands, 1829, 23, Iceland.

Family 20. ECHINORHINIDÆ. Bramble sharks

Genus 50. ECHINORHINUS Blainville.

Echinorhinus Blainville, Bull. Soc. Philom., 1816, 121 (*Squalus spinosus* Gmelin).

Goniodus Agassiz, Poiss. Fossiles, III, 1836, 183 (*Squalus spinosus* Gmelin).

90. *Echinorhinus brucus* (Bonnaterre). *Bramble shark.*
 Atlantic coasts of Europe, America, and Africa; scarce in the western Atlantic.
Squalus brucus Bonnaterre, Tableau Encycl., Ichth., II, 1788, 1500, "the ocean."
Squalus spinosus Gmelin, Syst. Nat., 1789, 1500, "the ocean."
Echinorhinus obesus Smith, Ills. Zool. So. Africa, Fishes, 1845, Pl. I, Cape of Good Hope.

Suborder SQUATINOIDEI

Family 21. SQUATINIDÆ. Angel sharks

Genus 51. SQUATINA Duméril.

Squatina Valmont de Bomare, Dict. Raisonnée, I, 1768, 117; not eligible.

Rhina Klein, Neuer Schauplatz, II, 1776, 587; non-binominal.

Squatina Duméril, Zool. Analyt., 1806, 102 (*Squatina angelus* Duméril = *Squalus squatina* Linnæus).

Rhina Rafinesque, Caratteri, 1810, 14 (*Squalus squatina* L.)

91. *Squatina duméril*¹² LeSueur. *Angelfish; Monkfish.*
 Atlantic coast of United States; Newport, R. I.
Squatina duméril LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 225, pl. 10, probably Florida.
92. *Squatina squatina* (Linnæus). *Monkfish; Angelfish; Squato; Shark ray; Monkeyfish; Kingston; Fiddlefish; Little bullhead shark.*
 Atlantic on both coasts, north to southern England and Cape Cod.
Squalus squatina Linnæus, Syst. Nat., ed. X, 1758, 233, coasts of Europe.
Squatina lævis Cuvier, Règne Anim., ed. I, 1817, 131; after Linnæus.
Squatina vulgaris Risso, Ichth. Nice, 1810, 45, Nice.
Squatina fimbriata Müller and Henle, Plagiostomen, 1838, 101, Mediterranean Sea.
Squatina oculata Bonaparte, Faun. Ital., Pesc., XXVIII, 1840, Fasci. 147, pl. 143, Italy.
Squatina lewis Couch, Trans. Linn. Soc. Lond., XIV., 1825, 90, Cornwall.
93. *Squatina californica* Ayres. *California angel shark.*
 Pacific coast of America, from southeastern Alaska southward; not rare on coast of California.
Squatina californica Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 29, San Francisco.

¹² As this species has not been distinguished generally from its near relative, *Squatina squatina* L., it is possible that all western Atlantic records belong to it.

Order H. BATOIDEI. Rays

Suborder SARCURA. Thick-tailed rays

Family 22. PRISTIDÆ. Sawfishes

Genus 52. PRISTIS Linck.

Pristis Klein, Neuer Schauplatz, VII, 1779, 403; non-binominal.

Pristis Linck, Mag. Neueste Physik und Naturgeschichte, 1790, 31 (*Squalus pristis* L.).

Pristis Latham, Trans. Linn. Soc. Lond., II, 1794, 276 (*Squalus pristis* L.).

94. *Pristis pectinatus* Latham. Common sawfish; *Pez sierra*.

Tropical Atlantic, north to West Indies and Florida; occasionally at Cape May and Grassy Sound.

Pristis pectinatus Latham, Trans. Linn. Soc. Lond., II, 1794, 278, "in the ocean."

Pristis granulosa Bloch and Schneider, Syst. Ichth., 1801, 352, Habana; after Parra.

Pristis mississippiensis Rafinesque, Ichth. Ohiensis, 1820, 86, lower Mississippi.

Pristis megalodon Duméril, Elasmobranchs, 1870, 476, pl. 9, Cayenne.

Pristis acutirostris Duméril, loc. cit., 479, Martinique.

Pristis occa Duméril, loc. cit., 479, locality unknown.

95. *Pristis zephyreus*¹⁸ Jordan and Starks. *Pez de espada*.

Tropical seas entering rivers, north to Mazatlan on the Pacific coast.

Pristis zephyreus Jordan and Starks, Fishes of Sinaloa, in Proc. Calif. Acad. Sci., V, 1895, 383, Mazatlan, Mexico; regarded by Garman as inseparable from *Pristis microdon* Latham.

Family 23. RHINOBATIDÆ. Guitarfishes

Genus 53. RHINOBATOS Linck. Guitarfishes.

Rhinobatus Klein, Neuer Schauplatz, II, 1776, 593 (*Rhinobatos seu = Squatina raja*; non-binominal).

Rhinobatos Linck, Magazin Neuestes, 1790, 32; type not specified.

Rhinobatus Bloch and Schneider, Syst. Ichth., 1801, 353 (*Rhinobatus rhinobatus*).

Syrrhina Muller and Henle, Plagiostomen, 1838, 113 (*Rhinobatus columna = Rhinobatos rhinobatus*).

Glaucostegus Bonaparte, Catalogo Metodico, 1846, 14 (*Rhinobatus rhinobatus*).

96. *Rhinobatos horkeli* (Müller and Henle).

Brazil north to Windward Islands; St. Eustatius.

Rhinobatus horkeli Müller and Henle, Plagiostomen, 1838, 122, pl. 41.

97. *Rhinobatos lentiginosus* (Garman). *Guitarro*.

Atlantic from Charleston, S. C., to Yucatan.

Rhinobatus lentiginosus Garman, Bull. Mus. Comp. Zool., VI, 1880, 168, coast of Florida.

98. *Rhinobatos percellens* (Walbaum). *Fiddlerfish*; *Puraque*.

West Indies to Jamaica and southern Brazil.

Raja percellens Walbaum, Artedi Pisc., 1792, 525, Brazil; after *Puraque* Marcgrave.

Rhinobatus electricus Bloch and Schneider, Syst. Ichth., 1801, 356, Brazil.

Rhinobatus undulatus Ölfers, Torpedo, 1831, 22, Brazil.

Rhinobatus glaucostictus Ölfers, loc. cit., 1831, 22, Brazil.

Rhinobatus marcgravi Henle, Ueber Narcine, 1834, 34, Brazil.

99. *Rhinobatos stellio* (Jordan and Rutter).

West Indies.

Rhinobatus stellio Jordan and Rutter, Proc., Acad. Nat. Sci. Phila., XLIX, 1897, 91, Kingston, Jamaica.

¹⁸ The African species, *P. perroteti* Valenciennes, probably does not occur in America.

100. *Rhinobatos spinosus* (Günther).
Mexico.
Rhinobatos spinosus Günther, Cat., VIII, 1870, 518, Mexico.
101. *Rhinobatos productus* (Ayres). *Guilartfish*.
San Francisco to San Diego.
Rhinobatos producta Ayres, in Girard, Proc., Acad. Nat. Sci. Phila., VII, 1854, 196, Presidio of Monterey, Calif.
102. *Rhinobatos glaucostigma* (Jordan and Gilbert). *Guilartro*.
Gulf of California and southward.
Rhinobatos glaucostigma Jordan and Gilbert, Proc. U. S. Nat. Mus., VI, 1883, 210, Mazatlan.
103. *Rhinobatos leucorhynchus* (Günther).
Panama and vicinity.
Rhinobatos leucorhynchus Günther, Proc. Zool. Soc. Lond., 1866, 604, Panama.
Genus 54. ZAPTERYX Jordan and Gilbert.
Zapteryx Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 53 (*Platyrrhina exasperata* Jordan and Gilbert).
104. *Zapteryx exasperatus* (Jordan and Gilbert).
San Diego Bay; abundant.
Platyrrhina exasperata Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 32, San Diego.
Trigonorrhina alveata Garman, Bull. Mus. Comp. Zool., VI, 1880, 169, San Diego.
105. *Zapteryx xyster* Jordan and Evermann.
Panama.
Zapteryx xyster Jordan and Evermann, Fishes North and Mid. Amer., 1896, 65, Panama.

Family 24. PLATYRRHINIDÆ

- Genus 55. PLATYRRHINOIDIS Garman.
Platyrrhinoidis Garman, Proc. U. S. Nat. Mus., III, 1880, 522 (*Platyrrhina triseriata* Jordan and Gilbert).
106. *Platyrrhinoidis triseriatus* (Jordan and Gilbert).
Coast of California, from Point Conception southward.
Platyrrhina triseriata Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 36, Santa Barbara, Calif.

Family 25. RAJIDÆ. Skates and rays

- Genus 56. RAJA Linnæus.
Raja (Artedi) Linnæus, Syst. Nat., ed. X, 1758, 231 (*Raja batis* Linnæus).
Diptyrus Rafinesque, Caratteri, 1810, 16 (*Raja batis* L.).
Platopterus Rafinesque, Anal. Nat. 1815, 93 (*Raja batis* L.).
Dasybatus Blainville, Jour. Phys. 1816, 260 (*communis*).
Propterygia Otto, Nova Acta Acad. Cæs. Leop. Carol. Nat. Curios., 1824, 777 (*hyposticta*; monstrous example with fins not adnate to head).
Læviraja Bonaparte, Faun. Ital., XXV, 1839, 130 (*oxyrhynchus*).
Hieroptera Fleming, Edinburgh New Philos. Jour., XXVI, 1841, 236 (*H. abredonensis* Flem. = *R. clavata* L.).
Batis Bonaparte, Cat. Metod., 1846, 12 (*radula*; no description).
Raja of various authors; merely change of spelling.
Amblyraja Malm, Göteborg og Bohuslän's Fauna, 1877, 607 (*Raja circularis* Malm = *Raja radiata* Donovan).
Lencoraja Malm, loc. cit., 609 (*Raja fullonica* L.).
Alpharaca Leigh-Sharpe,¹⁴ Jour. Morph., XXXIX, 1924, 568 (*Raja circularis* Couch).

¹⁴ Mr. Leigh-Sharpe (Jour. Morphology, XXXIX, December, 1924) divides the genus *Raja* into nine "pseudogenera" on the morphology of the genital organs. To settle the question of the standing of pseudogenera, Jordan (in Copeia, 1925) accepts all as subgenera. Most of those not named above (*Epsilon-raja*, *Zelara*, *Etaria*, *Thetaria*, and *Iotaria*), outside of our limits, may prove valid genera.

- Betaraia* Leigh-Sharpe, loc. cit., 568 (*Raja clavata* Linnaeus, a synonym of *Heiroptera* Fleming).
- Gammaraia* Leigh-Sharpe, loc. cit., 568 (*Raja batis* Linnaeus, a synonym of *Raja* Linnaeus).
- Deltaraia* Leigh-Sharpe, loc. cit., 568 (*Raja radiata* Linnaeus, a synonym of *Amblyraja* Malm).
107. *Raja laevis* Mitchill. *Barndoor skate; Winier skate; Peck-nosed skate.*
Massachusetts to Florida, common northward.
Raja laevis Mitchill, Amer. Month. Mag., II, 1817, 327, New York; name perhaps preoccupied by eligible authors (Gronow, 1763; Valmont, 1765; Duhamel, 1782).
Raja stabuliformis Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI, 1913, 341, Pl. XXII, fig. 2, Pl. XLIV, figs. 4-6, New England coast (substitute for *laevis*, regarded as preoccupied, a matter to be questioned).
108. *Raja radiata* Donovan.
Northern Iceland, southward on both coasts.
Raja radiata Donovan, Hist. Brit. Fish., V, 1820, pl. 114, Great Britain.
109. *Raja scabrata* Garman.
Raja scabrata Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI, 1913, 340, pl. 21, pl. 44, figs. 1-3, Massachusetts Bay. Long confused with *Raja radiata*.
110. *Raja granulata* Gill.
Banks of Newfoundland.
Raja granulata Gill, in Goode and Bean, Bull. Essex Inst., XI, 1879, 28, La Have Bank.
111. *Raja fyllæ* Lütken.
Davis Straits, Greenland.
Raja fyllæ Lütken, Vid. Medd. Naturh. Foren. Kjøbenhavn, 1887, 1, pl. 1, Davis Straits.
112. *Raja eglanteria* Bosc. *Skate; Brier ray; Possum ray; Sea possum; Bob-tailed skate; Clear-nosed skate; Summer skate.*
Cape Cod, southward to Florida; rather common.
Raja eglanteria Bosc, in Lacépède, Hist. Nat. Poiss., II, 1800, 104, Charleston, S. C.
Raja desmarestia LeSueur, Jour. Acad. Nat. Sci. Phila., IV, 1824, 100, Florida.
Raja chantenay LeSueur, loc. cit., 106, America.
Raja americana DeKay, N. Y. Fauna: Fishes, 1842, 368, pl. 66, fig. 215, Staten Island.
113. *Raja erinacea* Mitchill. *Hedgehog ray; Common skate; Little skate; Tobacco; Prickly skate; Summer skate.*
Nova Scotia to the Carolinas; very abundant from Virginia to Maine.
Raja erinacea Mitchill, Amer. Jour. Sci. Arts, IX, 1825, 290, pl. 6, New York.
114. *Raja diaphanes* Mitchill. *Clear ray.*
East Coast of United States.
Raja ocellata Mitchill, Trans. Lit. Phil. Soc. N. Y., 1, 1803, 877, New York; not of Russell.
Raja diaphanes Mitchill, loc. cit., 478, New York.
115. *Raja senta* Garman.
Banks of Newfoundland to Cape Cod; in deep water.
Raja senta Garman, Proc. U. S. Nat. Mus., VIII, 1885, 43, Cape Cod Bay; La Have Bank.
116. *Raja microtrachys* Osburn and Nichols.
Off west coast of Lower California.
Raja microtrachys Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1917, 142, fig. 1, off Guadalupe Island in deep water.
117. *Raja ackleyi* Garman.
Yucatan Banks
Raja ackleyi Garman, Bull. Mus. Comp. Zool., VIII, 1881, 234, Yucatan Banks.

118. *Raja ornata* Garman.
Coast of South Carolina and Florida.
Raja ornata Garman, Bull. Mus. Comp. Zool., VIII, 1881, 235, off South Carolina.
119. *Raja plutonia* Garman.
Deep water off South Carolina.
Raja plutonia Garman, Bull. Mus. Comp. Zool., VIII, 1881, 236, off South Carolina.
120. *Raja rhina* Jordan and Gilbert.
Monterey to Alaska; not rare.
Raja rhina Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 251, Monterey; San Francisco.
121. *Raja parmifera* Bean.
Common on coast of Alaska; Bristol Bay.
Raja parmifera Bean, Proc. U. S. Nat. Mus., IV, 1881, 157, Unalaska.
122. *Raja aleutica* Gilbert.
Unalaska, Aleutian Islands.
Raja aleutica Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1895), 397, pl. 21, Sannak Pass, Aleutian Islands, in 81 fathoms.
123. *Raja rosispinis* Gill and Townsend.
Bering Sea.
Raja rosispinis Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 231, Bering Sea.
Raja obtusa Gill and Townsend, loc. cit., 231, Bering Sea.
124. *Raja interrupta* Gill and Townsend.
Raja interrupta Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 232, Bering Sea.
125. *Raja binoculata* Girard. *Big skate of California.*
Pacific coast; abundant from Monterey to Sitka.
Raja binoculata Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 196, San Francisco.
Raja coopers Girard, Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 372, Shoalwater Bay, Wash.
126. *Raja stellulata* Jordan and Gilbert.
Coast of California and northward to Unalaska
Raja stellulata Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 133, Monterey Bay.
127. *Raja inornata* Jordan and Gilbert. *Common skate (California.)*
Coast of California; very abundant from San Diego to Cape Mendocino.
Raja inornata Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 457, San Francisco.
Raja jordani Garman, loc. cit., VIII, 1885, 43, San Francisco.
Raja inornata inermis Jordan and Gilbert, loc. cit., IV, 1881, 74, Santa Barbara.
128. *Raja montereyensis* Gilbert.
Monterey Bay.
Raja montereyensis Gilbert, Proc., U. S. Nat. Mus., XLVIII, 1915, 307, Monterey Bay off Santa Cruz, Calif., in 26 to 28 fathoms
129. *Raja equatorialis* Jordan and Bollman.
West coast of Colombia, between Panama and Galapagos Islands.
Raja equatorialis Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 150, off Colombia.
130. *Raja trachura* Gilbert.
Santa Barbara Islands; south of Shumagin Islands, Central Alaska.
Raja trachura Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 539, off Santa Barbara, at Albatross station 2923 in 822 fathoms.
131. *Raja abyssicola* Gilbert.
Queen Charlotte Island, British Columbia, in deep water.
Raja abyssicola Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1895), 396, pl. 20, off Queen Charlotte Island, at Albatross station 3342 in 1,588 fathoms.

Genus 57. DACTYLOBATUS Bean and Weed.

Dactylobatus Bean and Weed, Proc. U. S. Nat. Mus., XXVI, 1909, 459
(*D. armatus* Bean and Weed).

132. Dactylobatus armatus Bean and Weed.

Deep water off South Carolina.

Dactylobatus armatus Bean and Weed, Proc. U. S. Nat. Mus., XXXVI, 1911, 459, off Charleston in 270 fathoms.

Suborder NARCACIONTES**Family 26. TORPEDINIDÆ. Electric rays****Genus 58. TETRANARCE**¹⁵ Gill. *Great torpedo*.

Tetranarce Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 387 (*Torpedo occidentalis* Storer); misprinted *Tetronarce*.

Gymnotorpedo Fritsch, Arch. Anal. Phys., 1886, 365 (*Torpedo occidentalis* Storer).

133. Tetranarce occidentalis (Storer). *Electricfish; Crampfish; Torpedo; Numbfish.*

Atlantic coast of United States, Cape Cod to Cuba; not very common.

Torpedo occidentalis Storer, Amer. Jour. Sci. Arts, XLV, 1843, 165, pl. 3, Massachusetts. Garman regards this species as inseparable from *Torpedo nobiliana* Bonaparte of the Mediterranean.

134. Tetranarce californica (Ayres). *California torpedo; Crampfish.*

Coast of California; not recorded from south of Monterey; recently reported from Cape Flattery, Wash.

Torpedo californica Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 74, San Francisco.

Genus 59. NARCINE Henle.

Narcine Henle, Ueber *Narcine*, 1834, 31 (*Torpedo brasiliensis* von Ölfers).

Syraxia Jourdain, in Bonaparte, Fauna Italica, fasc. 160, 1841 (*Narcine indica* Henle).

Cyclonarce Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 387 (*Rana timlei* Schneider).

Gonionarce Gill, loc. cit., 387 (*Narcine indica* Henle).

135. Narcine brasiliensis (Ölfers). *Torpedofish; Small electric ray; Trembler.*

West Indies and Brazil, occasionally northward to Key West and the coast of North Carolina.

Torpedo brasiliensis von Ölfers, *Torpedo*, 1831, 19, tab. 2, fig. 4, Brazil.
Torpedo bancrofti Griffith, Anim. King., X, 1834, 649, Pl. XXXIV, Jamaica.

Narcine nigra Duméril, Revue Zool., 272, 1852, Brazil.

Torpedo pictus Gronow, Cat. Fishes, 13, 1854, Antilles.

Narcine brasiliensis corallina Garman, Bull. Mus. Comp. Zool., VIII, 1881, 234, Florida.

Narcine umbrosa Jordan, Proc. U. S. Nat. Mus., VII, 1884, 105, Key West.

136. Narcine entemedor Jordan and Starks. *Entemedor.*

Pacific coast of tropical America; Mazatlan, La Paz, Panama.

Narcine entemedor Jordan and Starks, Fishes of Sinaloa, Proc. Calif. Acad. Sci., V, 1895, 387, Mazatlan, Mexico.

Genus 60. DISCOPYGE Heckel.

Discopyge Heckel, in Tschudi, Fauna Peruana, 1846, 32 (*D. tschudii* Tschudi).

137. Discopyge ommata Jordan and Gilbert.

Panama; rare.

Discopyge ommata Jordan and Gilbert, Proc. U. S. Nat. Mus., XII, 1889, 151, Panama.

¹⁵ In the Old World genus *Torpedo* Forskål, as then understood, the spiracles are provided with a fringe of filaments, at least in joints. In *Tetranarce* these are absent. The following is the synonymy of *Torpedo*: *Torpedo* Forskål, Fauna Arabica, 1775, 16, in part (*Raja torpedo* L., confused with *Malapterurus electricus*, the electric catfish of the Nile).

Narcacion Klein, Neuer Schauplatz, IV, 1777, 726, nonbinomial (*Raja torpedo* L.).
Fimbriatorpedo Fritsch, loc. cit., 365 (*Torpedo marmorata* Risso = *Raja torpedo* L.).

Suborder MASTICURA. Whip-tailed rays

Family 27. DASYATIDÆ. Stingrays

Genus 61. PASTINACHUS Rüppell.

Pastinachus Rüppell, Atlas, Reise Nord Africa, 1828, 82 (*Raja sephen* Forskål); tail rounded above, with a keel below.

Hypolophus Müller and Henle, Plagiostomen, 1838, 170 (*Raja sephen* Forskål).

138. *Pastinachus centrourus* (Mitchill). *Stingaree*; *Stingray*; *Devilfish*; *Pigeon stingray*; *Clam-cracker*.

Coast of Maine to Cape Hatteras, abundant; the same or a related species in the Mediterranean.

Raja centroura Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 479, New York.

139. *Pastinachus schmardæ* (Werner).

West Indies.

Dasyatis schmardæ Werner, Zool. Jahrb., XXI, 1904, 298, Jamaica.

140. *Pastinachus torrei* (Garman). *Sabina*; *Lebisa*.

West Indies.

Dasyatis torrei Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI, 1913, 386, Tunas de Zaza, Cuba.

Genus 62. DASYATIS Rafinesque. *Stingrays*

Dasybatus Klein, Neuer Schauplatz, I, 1775, 191; not binomial.

Dasyatis Rafinesque, Caratteri, 1810, 16 (*Dasyatis ujo* Rafinesque = *Raja pastinaca* L.).

Uroxis Rafinesque, Indice, 1810, 61 (*U. ujo* Rafinesque).

Trigonobatus Blainville, Jour. Phys., 1816, 261 (*T. vulgaris* Blainville = *Pastinaca* L.).

Trygon Adanson, in Cuvier, Règne Anim., ed. I, 1817, 135 (*Raja pastinaca* L.).

Pastinaca Swainson, Class. Anim., 1839, 319 (*Pastinaca olivacea* Swainson).

Pastinaca DeKay, N. Y. Fauna: Fishes, 1842, 372 (*Raja pastinaca* L.).

Dasibatis Garman, corrected spelling.

141. *Dasyatis longus* (Garman).

Gulf of California to Panama: Galapagos.

Trygon longa Garman, Bull. Mus. Comp. Zool., VI, 1880, 170, Acapulco, Mexico.

142. *Dasyatis hastatus* (DeKay). *Whip stingray*; *Kit*; *Stingaree*.

West Indies to Brazil, north to Florida, North Carolina and Rhode Island.

Pastinaca hastata DeKay, N. Y. Fauna: Fishes, 1842, 373, pl. 65, fig. 214, Rhode Island.

143. *Dasyatis guttatus* (Schneider).

Surinam to Brazil; recorded from Granada.

Raja guttata Schneider, Syst. Ichth., 1801, 361, no locality given.

Trygon gymnura J. Müller, in Erdmann's Reise um die Erde, 1830, 25, taf. 13, Brazil.

Raja tuberculata Lacépède, Hist. Nat. Poiss., II, 1800, 106, pl. 4, fig. 1, Cayenne; not of Bonnatere.

Genus 63. AMPHOTISTIUS Garman.

Amphotistius Garman, Plagiostomia, Mem. Mus. Comp. Zool., 1913, 375 (*Trygon sinensis* Steindachner); tail with a distinct fold above and below.

144. *Amphotistius say* (LeSueur). *Southern stingray*.

New Jersey to Brazil, common in Florida; occasionally northward to New York.

Raja say LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 42, New Jersey.

145. *Amphotistius sabinus* (LeSueur).
Streams and estuaries of Florida; abundant on both coasts.
Trygon sabina LeSueur, Jour. Acad. Nat. Sci. Phila., IV, 1824, 109, Florida.
Trygon osteosticta Müller, in Erdmann's Reise um die Erde, 1830, 25, pl. 14, Brazil.
146. *Amphotistius dipterurus* (Jordan and Gilbert).
Bay of San Diego and southward; locally abundant.
Dasybatis dipterurus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 31, San Diego; identified by Garman with *Dasyatis brevis* Garman, from Peru.
- Genus 64. **PTEROPLATEA** Müller and Henle.
Pteroplatea Müller and Henle, Plagiostomen, 1838, 168 (*Raja altavela* L.)
147. *Pteroplatea altavela* (Linnæus)
Tropical Atlantic, occasionally northward to Mediterranean and in Gulf Stream; Beaufort, N. C.; Woods Hole, Mass.; taken at Penzance Point in Buzzards Bay (Nichols).
Raja altavela Linnæus, Syst. Nat., ed. X, 1758, 232, no locality.
Dasyatis attavilla Rafinesque, Indice, 1810, 49, Sicily.
Pteraplatea canariensis Valenciennes, Ichth. Canaries, 1843, 100, pl. 23, fig. 9, Canary Islands.
Pteroplatea valenciennesi A. Duméril, Elasmobranches, 1865, 612, no locality.
Pteroplatea vaillanti Rochebrune, Bull. Sci. Philom., IV, 1880, 159, no locality.
148. *Pteroplatea micrura* (Schneider). *Butterfly ray*.
Rhode Island to Brazil.
Raja micrura Schneider, Syst. Ichth., 1801, 360, Surinam.
Raja macrura LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 41, Rhode Island.
149. *Pteroplatea marmorata* Cooper.
Coast of California, from Point Conception southward to Cedros Island; common.
Pteroplatea marmorata Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 112, fig. 25, San Diego.
150. *Pteroplatea crebripunctata* Peters. *Manta rara colorado*; *Manta rara*.
Gulf of California, southward along the west coast of Mexico; common.
Pteroplatea crebripunctata Peters, Monatsber. Akad. Wiss. Berlin, 1869, 703, Mazatlan.
Pteroplatea rara Jordan and Starks, in Jordan, Fishes of Sinaloa, Proc. Calif. Acad. Sci., Ser. 2, V, 1895, 390, Mazatlan, Mexico.
- Genus 65. **UROBATUS** Garman.
Urobatis Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI, 1913, 401 (*Leobatus sloani* Blainville).
151. *Urobatis sloani* (Blainville). *Maid*.
West Indies, generally common; occasionally northward; once (perhaps doubtfully) recorded from New Jersey.
Leobatus sloani Blainville, Bull. Sci. Philom., 1816, 121, Jamaica; after *Pastinaca marina* Sloan.
Raja jamaicensis Cuvier, Règne Anim., ed. I, 1817, 137, Jamaica.
Trygonobatus torpedinus Desmarest, Prem. Dec. Ichth., I, 1823, 6, Cuba.
152. *Urobatis vermiculatus* (Garman).
West Indies.
Urobatis sloani vermiculatus Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI, 1913, 802, Progreso, Yucatan.
153. *Urobatis halleri* (Cooper). *Round stingray*.
Coast of California, from Point Conception southward.
Urolophus halleri Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 95, San Diego.

154. *Urobatis umbrifer* (Jordan and Starks).
West coast of Mexico.
Urolophus umbrifer Jordan and Starks, in Jordan, Fishes Sinaloa, Proc. Calif. Acad. Sci., Ser. 2, V, 1895, 389, Mazatlan, Mexico; regarded by Garman as inseparable from *Urobatis halleri*.
155. *Urobatis nebulosus* (Garman).
Gulf of California to Panama.
Urolophus nebulosus Garman, Proc. U. S. Nat. Mus., VIII, 1885, 41, Colima, Mexico.
156. *Urobatis concentricus* Osburn and Nichols.
Gulf of California.
Urobatis concentricus Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 144, fig. 2, Esteban Island, Gulf of California.
- Genus 66. *UROTRYGON* Gill.
Urotrygon Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 173 (*Urotrygon mundus* Gill).
157. *Urotrygon mundus* Gill.
West coast of Mexico.
Urotrygon mundus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 173, Panama.
Urolophus asterias Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 579, Mazatlan; Panama.
Urolophus rogersi Jordan and Starks, in Jordan, Fishes of Sinaloa, Proc. Calif. Acad. Sci., Ser. 2, V, 1895, 388, Mazatlan.
158. *Urotrygon goodei* (Jordan and Bollman).
Panama and vicinity.
Urolophus goodei Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 151, near Panama.
159. *Urotrygon aspidurus* (Jordan and Gilbert).
Panama.
Urolophus aspidurus Jordan and Gilbert, Bull. U. S. Fish Com., IV, 1881, 307, Panama.

Family 28. AETOBATIDÆ. Eagle rays

- Genus 67. *STOASODON* Cantor.
Aetobatis Müller and Henle, Plagiostomen, 1841, 180 (*Raia narinari* Euphrasen); not *Aetobatus* Blainville.
Stoasodon Cantor, Cat. Malayan Fishes, 1850, 434 (*Raia narinari* Euphrasen).
Goniobatis Agassiz, Proc. Bost. Soc. Nat. Hist., VI, 1859, 385 (*flagellum*).
160. *Stoasodon narinari* (Euphrasen). *Spotted stingray*; *Bishop ray*; *Obispo*; *Lady ray*.
Tropical seas; north on the Atlantic coast to Virginia; Pacific coast, Gulf of California to Panama.
Raia narinari Euphrasen, Vedensk. Akad. Nya Handl., XI, 1790, 217, Brazil.
Raia flagellum Bloch and Schneider, Syst. Ichth., 1801, 361, Coromandel.
Myliobatis eeltenkei Rüppell, Neu. Wirb., 1835, 70, Red Sea.
Goniobatis macroptera McClelland, Calcutta Jour. Nat. Hist., I, 1841, 60, Bengal.
Aetobatus laticeps Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1865, 137, California.
- Genus 68. *PTEROMYLÆUS* Garman.
Pteromyllæus Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI, 1913, 437 (*Myliobatis asperimus* Gilbert).
161. *Pteromyllæus asperimus* (Gilbert).
Panama.
Myliobatis asperimus Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2754, Panama.

Genus 69. AETOBATUS ¹⁶ Blainville. *Eagle rays*.

Aetobatus Blainville, Jour. Phys., LXXXIII, 1816, 261 (*Raja aquila* as *Aetobatus vulgaris* Blainville).

Myliobatis Duméril in Cuvier, Règne Anim., ed. I, 1817, 137 (*Raja aquila*).

Holorhinus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 331 (*Rhinoptera vespertilio* Girard = *Myliobatus californicus* Gill).

162. Aetobatus frémouvillii (LeSueur). *Stingaree; Sharp-nosed ray; Long-tailed stingray*.

Cape Cod to Brazil.

Myliobatis fremouvillii LeSueur, Jour. Acad. Nat. Sci. Phila., IV, 1824, 111, Rhode Island.

Myliobatis bispinosus Storer, Proc. Bost. Soc. Nat. Hist., I, 1841, 53, Massachusetts.

Myliobatis acuta Ayres, loc. cit., I, 1842, 65, Brookhaven, Long Island, N. Y.

163. Aetobatus goodiei (Garman).

Central America to Brazil; Rio Grande do Sul.

Myliobatis goodiei Garman, Proc. U. S. Nat. Mus., VIII, 1885, 39, Central America.

164. Aetobatus californicus (Gill). *Califorma stingray; Batfish*.

California from Cape Medocino southward.

Rhinoptera vespertilio Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 137, Tomales Bay; not *Myliobatus vespertilio* Bleeker.

Myliobatis californicus Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1865, 137; after Girard.

Family 29. RHINOPTERIDÆ**Genus 70. RHINOPTERA** Kuhl.

Rhinoptera Kuhl in Cuvier, Règne Anim. ed. II, II, 1829, 401 (*Myliobatis marginatus* Geoffroy).

Zygobatis Agassiz, Poiss. Fossiles, III, 1836, 79 (*Z. jussieuvi* Agassiz).

Myliorhina Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1865, 136 (*Rhinoptera lalandi* Muller and Henle).

Micromesurus Gill, loc. cit., 136 (*Rhinoptera adspersa* Muller and Henle).

165. Rhinoptera bonasus Mitchill. *Cowfish; Cow-nosed ray*.

Cape Cod to Florida.

Rhinoptera bonasus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 479, New York; description inadequate.

Raja quadriloba LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 44, New Jersey.

166. Rhinoptera steindachneri Evermann and Jenkins. *Gabilan*.

Gulf of California.

Rhinoptera steindachneri Evermann and Jenkins, Proc. U. S. Nat. Mus., XIV, 1891, 130, pl. 1, fig. 1, Guaymas, Sonora, Mexico.

167. Rhinoptera encenadæ Rosa Smith.

Todos Santos Bay, Lower California.

Rhinoptera encenadæ Rosa Smith, Proc. U. S. Nat. Mus., IX, 1886, 220, Enseñada de Todos Santos, Lower California. Jaw only.

Family 30. MOBULIDÆ. Sea devils**Genus 71. MOBULA** Rafinesque.

Mobula Rafinesque, Indice, 1810, 61 (*M. auriculata* = *Squalus edentulus* Brunnich).

Apterurus Rafinesque, Indice, 1810, 62 (*A. fabroni* Rafinesque).

Cephalopterus Risso, Ichth. Nice, 1810, 14 (*Raja giorna* Lacépède); not of Geoffroy St. Hilaire, 1809, a genus of birds.

Dicerobatus Blainville, Bull. Soc. Philom., 1816, 116 (*Raja mobular* Lacépède).

Pterocephala Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 321, (*Raja giorna* Lacépède).

¹⁶ Müller and Henle correctly referred *Aetobatus* Blainville to the type of *Raja aquila*, spelling the name, however, as *Aetobatis*. They then created a new genus, *Aetobatis* for *Raja narinari*, an arrangement without warrant.

168. *Mobula hypostoma* (Bancroft).

Jamaica

Cephalopterus hypostomus Bancroft, Proc. Comm. Zool. Soc., 1830, 134, Jamaica.169. *Mobula oliersi* (Muller and Henle). *Small devilfish.*

Cape Lookout, N. C., to Brazil.

Cephalopterus oliersi Muller and Henle, Abhandl. Akad. Wiss. Berlin, 1834, 311, Brazil; regarded by Garman as identical with *M. thypostoma*.

Genus 72. CERATOBATIS Boulenger.

Ceratobatis Boulenger, Ann. Mag. Nat. Hist., XX, 1897, 227 (*C. robertsi* Boulenger).170. *Ceratobatis robertsi* Boulenger.

Jamaica.

Ceratobatis robertsi Boulenger, Ann. Mag. Nat. Hist., XX, 1897, 227, Jamaica.

Genus 73. MANTA Bancroft.

Manta Bancroft, Zool. Jour., IV, 1828-29, 444 (*Cephalopterus manta* Bancroft=*Raja birostris* Walbaum, 1792).*Ceratoptera* Muller and Henle, Syst. Besch., Plagiostomen, 1838, 186 (*Cephaloptera vampyrus* Mitchill).*Brachioptilon* Newman, Zoologist, 1849, 74 (*B. hamiltoni*).*Diabolichthys* Holmes, Proc. Elliott Soc. Nat. Hist., I, 1856, 39 (*D. ellioti* Holmes).171. *Manta birostris* (Walbaum). *Sea devil; Devilfish; Manta; Blanketfish.*

Tropical waters of America; north to New Jersey and San Diego; not rare on the Florida coast.

Raja birostris Walbaum, Artedi Pisc., 1792, 535, after *Diabolus marinus* Willughby, etc.*Raja manatia* Schneider, Svst. Ichth., 1801, 364, tropical America.*Raja fimbriata* Lacépède, Hist. Nat. Poiss., IV, 1803, 677, Gulf Stream.*Cephalopterus vampyrus* Mitchill, Ann. Lyc. Nat. Hist. N. Y., I, 1823, 23, Delaware Bay.*Cephalopterus giorna* LeSueur, Jour. Acad. Nat. Sci. Phila., IV, 1824, 115, Georgia.*Cephaloptera manta* or *Manta americana* Bancroft, Zool. Jour., IV, 1828, 444, Jamaica.*Cephaloptera johni* Muller and Henle, Syst. Besch. Plagiostomen, 1838, 186, West Indies.*Diabolichthys ellioti* Holmes, Proc. Elliott Soc. Nat. Hist., I, 1856, 39, Charleston, S. C.172. *Manta hamiltoni* (Newman).San Diego; Gulf of California to Panama, not rare; not known to be distinct from *Manta birostris*.*Brachioptilon hamiltoni* Newman, Zoologist, 1849, 74, Gulf of California.

Subclass HOLOCEPHALI. Chimæras

Order I. CHIMÆEROIDEI. Chimæroids

Family 31. CHIMÆRIDÆ. Chimæras

Genus 74. CHIMÆRA Linnæus. *Elephantfishes.**Chimæra* Linnæus, Syst. Nat., ed. X, 1758, 236 (*C. monstrosa* Linnæus).173. *Chimæra monstrosa* Linnæus.

Deep waters of the Atlantic; off Portugal and off the American coast from Cape Cod northward.

Chimæra monstrosa L., Syst. Nat., ed. X, 1758, 236, Atlantic.*Chimæra mediterranea* Risso, Europ. Mérid., III, 1826, 168, Nice.*Chimæra cristata* Faber, Natur. der Fische Islands, 1829, 45, Iceland.*Callorhynchus centrina* and *Callorhynchus atlanticus* Gronow, Cat. Fishes, 15, 16, 1854.

Genus 75. BATHYALOPEX Collett.

Bathyalopez Collett, Arch. Math. Natur., XXXIII, 1901, 125 (*Chimæra mirabilis* Collett).

174. Bathyalopez plumbeus (Gill).

Gulf Stream in deep water.

Chimæra plumbea Gill, Bull. Phil. Soc. Wash., Dec., 1877, 182, La Have Bank, lat. 42° 40' N., long. 63° 23' W., in 350 fathoms.

175. Bathyalopez abbreviatus (Gill).

Chimæra abbreviata Gill, Proc. U. S. Nat. Mus., VI, 1883, 254, at 40° N., long. 67° W., in 1,290 fathoms.

Genus 76. PSYCHICHTHYS Fowler.

Psychichthys Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 419 (*Hydrolagus waiteri* Fowler).

176. Psychichthys affinis (Capello).

Open sea, from Cape Cod to Portugal.

Chimæra affinis Capello, Jour. Soc. Math., Lisbon, IV, 1868; 314, Portugal in deep water.

Genus 77. HYDROLAGUS Gill.

Hydrolagus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 331 (*Chimæra collei* Lay and Bennett).

177. Hydrolagus collei (Lay and Bennett). *Ratfish; Elephantfish.*

Pacific coast, Monterey to Alaska; abundant in cold water at no great depth.

Chimæra collei Lay and Bennett, Beechey's Voy., Zool., 1839, 71, North Pacific.

Family 32. RHINOCHIMÆRIDÆ**Genus 78. HARRIOTTA** Goode and Bean.

Harriotta Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1895, 471 (*H. raleighana* Goode and Bean).

Antelochimæra Tanaka, Jour. Coll. Sci., Imp. Univ. Tokyo, XXVIII, art. 8, 1909, Japan (*A. chætirhamphus* Tanaka).

178. Harriotta raleighana Goode and Bean.

Deep water off the north Atlantic coast of the United States.

Harriotta raleighana Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894 (Jan. 26, 1895), 472, pl. 19, Gulf Stream at 39° N., 70° W.

179. Harriotta curtis-jamesi Townsend and Nichols.

Deep water off Lower California.

Harriotta curtis-jamesi Townsend and Nichols, Bull. Amer. Mus. Nat. Hist., LII, 1925, 6, fig. 2, off Lower California, lat. 26°, in 645 fathoms.

Class IV. PISCES. True fishes**Subclass ACTINOPTERI****Order J. GLANIOSTOMI****Family 33. ACIPENSERIDÆ. Sturgeons****Genus 79. ACIPENSER**¹⁷ Linnaeus. *Sturgeons.*

Acipenser Linnaeus, Syst. Nat., ed. X, 1758, 237 (*A. sturio* L.).

Ichthocolla Geoffroy, Descr. de 719 Plantes, etc., 1767, 399 (presumably *A. huso* L.).

Sterleta Guldenstadt, Nov. Com. Sci. Petropol., XVI, 1772, 533 (*A. ruthenus* L.).

Sturio Rafinesque, Indice, 1810, 41 (*Sturio vulgaris* Rafinesque = *A. sturio* L.).

Sterletus Rafinesque, Ichth. Ohiensis, 1820, 80 (*A. serotinus* Rafinesque = *A. fulvescens* Rafinesque).

¹⁷ This group ultimately may need subdivision into distinct genera, but none of the many proposed is adequately defined.

Helops Brandt and Ratzeburg, Medizinische Zoologie, II, 1833, 3, 349 (*A. stellatus* Pallas = *A. halops* Pallas).
Sterletus Brandt and Ratzeburg, loc. cit., 349 (*A. ruthenus* L.).
Sturio Muller, Anab. Myxinoidea, 1835, 77 (*A. sturio* L.).
Antaceus Heckel and Fitzinger, Ann. Wiener Mus., I, 1836, 293 (*A. schypa* Eichwald).
Lioniscus Heckel and Fitzinger, loc. cit., 370 (*A. glabra* Fitzinger).
Sterletus Bonaparte, Cat. Metod, 1845, 21 (*A. vultharius* L.).
Ellops Gistel, Naturg. Thierreichs, IX, 1848 (*A. helops* Pallas).
Schüpa Brandt, Bull. Acad. Sci. Petersb., VII, 1850, 113 (*A. schypa* Eichwald).

180. *Acipenser transmontanus* Richardson. *White sturgeon; Oregon sturgeon; Sacramento sturgeon.*

Pacific coast of North America, from Monterey to Alaska; once abundant, now nearly extinct; largest of the sturgeons, reaching a weight of 1,000 pounds. A record sturgeon, *Acipenser transmontanus*, taken at Astoria, weighed 1,900 pounds.

Acipenser transmontanus Richardson, Fauna Bor.-Amer., III, 1836, 278, Fort Vancouver.

Acipenser brachyrhynchus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 15, San Francisco.

Acipenser aleutensis Fitzinger and Heckel, Ann. Wien. Mus. Naturg., 1836, Aleutian Islands, after Pallas.

Acipenser caryi Duméril, Nouv. Arch. Mus., III, 1867, 169, San Francisco.

Acipenser ayresi Duméril, loc. cit., 171, San Francisco.

Acipenser putnami Duméril, loc. cit., 178, San Francisco.

181. *Acipenser oxyrhynchus* Mitchill. *Common sturgeon.*

New York to Carolina, once abundant along the shores; very close to the common sea sturgeon of Europe (*Acipenser sturio* L.), but with fewer lateral plates and rougher ossification; the two perhaps identical.

Acipenser oxyrhynchus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1814, 462, New York.

Acipenser mitchilli Duméril, Nouv. Arch. Mus., III, 1867, New York.

Acipenser kennicotti Duméril, loc. cit., James River.

Acipenser girardi Duméril, loc. cit., Maryland.

Acipenser macrorhinus Duméril, loc. cit., New York.

Acipenser milberti Duméril, loc. cit., New York.

Acipenser bairdi Duméril, loc. cit., Maryland.

Acipenser storeri Duméril, loc. cit., Boston.

Acipenser holbrooki Duméril, loc. cit., Charleston.

Acipenser lecontei Duméril, loc. cit., New York.

182. *Acipenser brevirostris* LeSueur. *Short-nosed sturgeon.*

Cape Cod to Florida.

Acipenser brevirostrum LeSueur, Trans. Amer. Phil. Soc., I, 1818, 290, Atlantic coast.

Acipenser microrhynchus Duméril, Hist. Nat. Poiss., II, 1870, 105 and 164, New York.

Acipenser lesueurii Valenciennes, in Duméril, loc. cit., 105 and 166, Atlas, pl. 16, figs. 1 and 1a, New York.

Acipenser dekayi Duméril, loc. cit., 105 and 168, New York.

183. *Acipenser acutirostris* Ayres. *Green sturgeon.*

Pacific coast of America, ascending rivers from Monterey northward; not rare.

Acipenser acutirostris Ayres, Proc., Calif. Acad. Nat. Sci., I, 1854, 14, San Francisco.

Acipenser medirostris Ayres, loc. cit., 15, San Francisco.

Acipenser (Antaceus) agassizii Duméril, Nouv. Arch. Mus. Paris, III, 1867, 181, Pl. XI, figs. 2, 2a, 2b, San Francisco.

Acipenser (Antaceus) alexandri Duméril, loc. cit., 183, Pl. XV, figs. 1, 1a, 1b, San Francisco.

Acipenser (Antaceus) oligopeltis Duméril, loc. cit., 184, Pl. XV, figs. 2, 2a, 2b, San Francisco.

184. *Acipenser fulvescens* Rafinesque. *Lake sturgeon; Ohio sturgeon; Red sturgeon.*

Mississippi Valley, Great Lakes and northward, ascending rivers, but not entering the sea; varying much with age.

Acipenser fulvescens Rafinesque, Amer. Month. Mag., I, August, 1817, 288, Great Lakes.

Acipenser rubicundus LeSueur, Trans. Amer. Phil. Soc., I, 1818, 388, Lakes Ontario, Erie, and all the upper lakes.

Acipenser maculosus LeSueur, loc. cit., 393, Ohio River; young.

Acipenser serotinus Rafinesque, Ichth. Ohiensis, 1820, 80, Ohio River.

Acipenser ohiensis Rafinesque, loc. cit., 81, Ohio River.

Acipenser macrostomus Rafinesque, loc. cit., 81, Ohio River.

Dinectus truncatus Rafinesque, loc. cit., 82; erroneous, based on a drawing by Audubon.

Acipenser rupertianus Richardson, Fauna Bor.-Amer., III, 1836, 311, Albany River, Rupert Land.

Acipenser laevis Agassiz, Lake Superior, 1850, 267; Lake Superior.

Acipenser carbonarius Agassiz, loc. cit., 271, Lake Superior.

Acipenser rhynchus Agassiz, loc. cit., 276, Lake Superior.

Acipenser koppelis Gunther, Cat. Fishes, VIII, 1870, 341, Mississippi.

Acipenser (Huso) copei Duméril, Hist. Nat. Poiss., 1870, 105 and 108, supposedly upper Missouri.

Acipenser (Huso) rauchi Duméril, loc. cit., 105 and 118, Osage River, Mo.

Acipenser (Huso) richardsoni Duméril, loc. cit., 105 and 120, upper Mississippi.

Acipenser (Huso) anasimos Duméril, loc. cit., 105 and 122, Missouri River at St. Louis.

Acipenser (Huso) paranasimos Duméril, loc. cit., 105 and 124, Huntsville, Ala.

Acipenser (Huso) anthracinus Duméril, loc. cit., 105 and 126, Lake Erie.

Acipenser (Huso) megalaspis Duméril, loc. cit., 105 and 133, Lake Champlain.

Acipenser (Huso) lamarii Duméril, loc. cit., 105 and 139, Mississippi River or Lake Erie.

Acipenser (Huso) atelaspis Duméril, loc. cit., 105 and 141, Saskatchewan River.

Acipenser (Huso) rafinesquii Duméril, loc. cit., 105 and 143, Ohio River.

Acipenser (Huso) rosarium Duméril, loc. cit., 105 and 152, Atlas, pl. 17, figs. 2 and 2a, upper Mississippi or Lake Erie.

Acipenser (Huso) platyrhinus Duméril, loc. cit., 105 and 154, upper Mississippi or Lake Erie.

Acipenser (Huso) kirilandi Duméril, loc. cit., 105 and 161, Lake Erie.

Acipenser (Huso) nertinianus Duméril, loc. cit., 105 and 162, Michigan.

Acipenser (Antaceus) cincinnatus Duméril, loc. cit., 220 and 228, Ohio River.

Acipenser (Antaceus) buffalo Duméril, loc. cit., 220 and 231, Lake Erie, Buffalo.

Genus 80. *SCAPHIRHYNCHUS* Heckel. *Shovelnose sturgeons.*

Scaphirhynchus Heckel, Ann. Wien. Mus. Naturg., I, 1835, 71 (*S. rafinesquii* Heckel = *Acipenser platyrhynchus* Rafinesque); not *Scaphorhynchus* Maximilian, a genus of birds.

Scaphirhynchops Gill, Trans. Amer. Phil. Soc., V, 1863, 12, 178 (*Acipenser platyrhynchus* Rafinesque).

185. *Scaphirhynchus platyrhynchus* (Rafinesque). *Shovelnose sturgeon; Swichtail; Hackleback; Sand sturgeon.*

Mississippi Valley; common in the larger streams.

Acipenser platyrhynchus Rafinesque, Ichth. Ohiensis, 1820, 80, Ohio River.

Genus 81. *PARASCAPHIRHYNCHUS* Forbes and Richardson.

Parascaphirhynchus Forbes and Richardson, Bull. Ill. Lab. Nat. Hist., VII, 1905, 38 (*P. albus* Forbes and Richardson).

186. *Parascaphirhynchus albus* Forbes and Richardson. *White shovelnose sturgeon; White sturgeon.*

Mississippi River; rare.

Parascaphirhynchus albus Forbes and Richardson, Bull. Ill. Lab. Nat. Hist., VII, 1905, 38, Mississippi River at Grafton, Ill.

Order K. SELACHOSTOMI

Family 34. POLYODONTIDÆ. Paddlefishes

Genus 82. POLYODON Lacépède.

- Polyodon* Lacépède, Bull., Soc. Philom., 1797, 49 (*P. folium* Lacépède).
Spathularia Shaw, Gen. Zool., V, 1804, 362 (*S. reticulata*=*Squalus spathula* Walbaum).
Platirostra LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817 (1818), 228 (*P. edentula*).
Plamirostra Rafinesque, Ichth. Ohiensis, 1820, 83 (*P. edentula*).
Proceros Rafinesque, loc. cit., 87 (*P. maculatus*).

187. *Polyodon spathula* (Walbaum). *Paddlefish*; *Spoonbill cat*; *Duckbill cat*; *Spadefish*.
 Mississippi Valley and south in the larger streams, reaching a weight of 173 pounds; rare in Lake Erie.
Squalus spathula Walbaum, Artedi Pisc., 1792, 522; after *Spathula* Rosier, Jour. Phys., 1774, 384, pl. 2; no locality given.
Polyodon feuille Lacépède, Hist. Nat. Poiss., I, 1798, 402, locality unknown.
Polyodon folium Bloch and Schneider, Syst. Ichth., 1801, 457; after Lacépède.
Spatularia reticulata Shaw, Gen. Zool., V, 1804, 362, after Lacépède.
Platirostra edentula LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 227, Ohio River.
Acipenser lagenarius Rafinesque, Ichth. Ohiensis, 1820, 86, Ohio River.
Proceros maculatus Rafinesque, loc. cit., 87, Mississippi River at St. Genevieve, Mo.
Proceros vittatus Rafinesque, loc. cit., 1820, 86, 87, Lake Ontario.

Order L. HOLOSTEI

Suborder GINGLYMODI

Family 35. LEPISOSTEIDÆ. Gar pikes

Genus 83. LEPISOSTEUS Lacépède. Gar pikes.

- Lepisosteus* Lacépède, Hist. Nat. Poiss., V, 1803, 331 (*L. gairialis* Lacépède=*Esox osseus* Linnæus).
Sarchirus Rafinesque, Jour. Acad. Nat. Sci. Phila., I, 1818, 419 (*S. vittatus* Rafinesque).
~~*Lepisosteus*~~ Agassiz; corrected spelling.
188. *Lepisosteus osseus* (Linnæus). *Long-nosed gar*; *Billfish*; *Common gar pike*; *Ohio gar*; *Ohio garfish*; *Billy gar*.
 Maryland to Florida and Louisiana, in coastwise streams; also (*huronensis*) Great Lakes and the Mississippi Valley; the most abundant of the gars.
Esox osseus Linnæus, Syst. Nat., ed. X, 1758, 313, after Artedi; based on *Acus maxima squamosa viridis*, the green gar fish of Catesby, 1738, pl. 30, Virginia.
Lepisosteus gairialis Lacépède, Hist. Nat. Poiss., V, 1803, 333, "lakes and rivers of both Indies."
Lepisosteus longirostris Rafinesque, Ichth. Ohiensis, 1820, 74, Muskingum River, Ohio; head.
Macrogathus lorreatus Gronow, Cat. Fishes, 1854, 148, after Linnæus.
Lepidosteus crassus Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, Bombay Hook, Delaware River.
Lepidosteus louisianensis Duméril, Hist. Nat. Poiss., 1870, 343, New Orleans.
Lepisosteus huronensis Richardson, Fauna Bor.-Amer., III, 1836, 237, Penitanguishene, Lake Huron.
 ?*Sarchirus vittatus* Rafinesque, loc. cit., 69, Ohio River; young; not identifiable.
 ?*Sarchirus argenteus* Rafinesque, loc. cit., 86, Ohio River; young; not identifiable.
Lepidosteus semiradiatus Agassiz, Poiss. Fossiles, II, 1836, 2, pl. 2.

- Lepisosteus bison* DeKay, New York Fauna: Fishes, 1842, 271, Buffalo, N. Y.
- Lepisosteus lineatus* Thompson, Hist. Vermont, 1842, 145, Lake Champlain.
- Lepisosteus leptorhynchus* Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 351, Devil River, Tex.
- Lepisosteus otarius* Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 86 "Platte River near Fort Riley"; but Fort Riley was on the Kansas River.
- Lepisosteus mulberti* Duméril, Hist. Nat. Poiss., 1870, 323 and 328, New York.
- Lepisosteus harlani* Duméril, loc. cit., 323 and 329, Wabash River.
- Lepisosteus smithi* Duméril, loc. cit., 323 and 330, upper Mississippi.
- Lepisosteus ayresi* Duméril, loc. cit., 323, 331, Wabash River.
- Lepisosteus copei* Duméril, loc. cit., 323, 332, northern North America.
- Lepisosteus lesueurii* Duméril, loc. cit., 323, 335, Wabash River.
- Lepisosteus elizabeth* Duméril, loc. cit., 323, 336, no locality.
- Lepisosteus troostii* Duméril, loc. cit., 323, 339, United States.
- Lepisosteus piquotianus* Duméril, loc. cit., 323, 340, Lake Erie.
- Lepisosteus thompsoni* Duméril, loc. cit., 323, 342, Upper Mississippi.
- Lepisosteus lamarr* Duméril, loc. cit., 323, 337, northern North America.
- ?*Lepisosteus clintonii* Duméril, loc. cit., 323, 338, no locality.
- Lepisosteus horatii* Duméril, loc. cit., 323, 341, northern North America.
- Genus 84. CYLINDROSTEUS Rafinesque. Short-nosed gars.**
- Cylindrosteus* Rafinesque, Ichth. Ohiensis, 1820, 72 (*Lepisosteus platostomus* Rafinesque).
- 189. *Cylindrosteus platostomus* (Rafinesque).**
- Mississippi Valley and Great Lakes; common.
- Lepisosteus platostomus* Rafinesque, Ichth. Ohiensis, 1820, 72, Ohio River.¹⁸
- Lepisosteus oxyurus* Rafinesque, loc. cit., 73, Ohio River.
- Lepisosteus albus* Rafinesque, loc. cit., 73, Ohio River.
- Lepisosteus grayi*, Agassiz, Poiss. Fossiles, II, 1836, 2, Louisiana.
- Lepisosteus (Cylindrosteus) oculatus* Winchell, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 183, Huron River, Mich.
- Cylindrosteus zadockii* Duméril, Hist. Nat. Poiss., II, 1870, 348, 353, Upper Mississippi.
- Cylindrosteus rafinesquii* Duméril, loc. cit., 348, 352, United States.
- 190. *Cylindrosteus scabriceps* Fowler.**
- Lower Missouri River.
- Cylindrosteus scabriceps* Fowler, Proc. Acad. Nat. Sci. Phila., LXII, 1910, 607, Pl. XXXVIII, figs. 8, 9, Leavenworth, Kans.; a doubtful species.
- 191. *Cylindrosteus agassizii* Duméril.**
- Missouri Basin.
- Cylindrosteus agassizii* Duméril, Hist. Nat. Poiss., II, 1870, 351, St. Louis.
- Cylindrosteus productus* Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 86, San Antonio, Tex.
- 192. *Cylindrosteus castelnaudii* Duméril.**
- Florida and Louisiana.
- Cylindrosteus castelnaudii* Duméril, Hist. Nat. Poiss., II, 1870, 355, Lake Lafayette, Florida.
- Cylindrosteus bartoni* Duméril, loc. cit., 356, New Orleans.
- Lepisosteus latirostris* Girard, Pac. R. R. Surv., X, pl. 4, Fishes, 1858, 352, Rio Pecos, Tex.
- 193. *Cylindrosteus megalops* Fowler.**
- Florida.
- Cylindrosteus megalops* Fowler, Proc. Acad. Nat. Sci. Phila., LXII, 1910, 609, Pl. XXXVIII, figs. 15, 16, Bayport, Fla.; a doubtful species.

¹⁸ Many other tributary streams mentioned.

Genus 85. *ATRACTOSTEUS* Rafinesque. *Alligator gars.*

Litholepis Rafinesque, Amer. Month. Mag., III, 1818, 447 (*L. adam-antinus* Rafinesque), based on a fancy painting by Audubon of an imaginary fish.

Atractosteus Rafinesque, Ichth. Ohiensis, 1820, 72 (*Lepisosteus ferox* Rafinesque).

194. *Atractosteus spatula* (Lacépède). *Mississippi alligator gar.*

Lower Mississippi Valley in the larger streams, reaching a length of 5 feet or more.

Lepisosteus spatula Lacépède, Hist. Nat. Poiss., V, 1803, 333, no locality.

Lepisosteus (Atractosteus) ferox Rafinesque, Ichth. Ohiensis, 1820, 73. Ohio River.

Lepisosteus (Atractosteus) berlandieri Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 353, Tamaulipas.

Atractosteus lucius Duméril, Hist. Nat. Poiss., 1870, 364, Tampico, Mexico.

195. *Atractosteus tristœchus* (Bloch and Schneider). *Cuban alligator gar; Manjuari.*

Rivers of Cuba.

Esox tristœchus Bloch and Schneider, Syst. Ichth., 1801, 395, Cuba, after Manjuari of Para.

Lepisosteus manjuari Poey, Memorias, I, 1860, 273, Cuba.

196. *Atractosteus tropicus* Gill.

Central America; Nicaragua.

Atractosteus tropicus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 172, streams near Panama.

Order M. **HALECOMORPHI**Family 36 **AMIIDÆ.** BowfinsGenus 86. *AMIA* Linnæus.

Amia Linnæus, Syst. Nat., ed. XII, 1766, 500 (*A. calva*); not of Gronow, 1763, which is a nonbinomial name for *Apogon*.

Amiatus Rafinesque, Anal. Nat., 1815, 88 (*A. calva*); substitute for *Amia* regarded as too short for symmetry.

197. *Amia calva* Linnæus. *Bowfin; John A. Grindley; Mudfish; Lawyer; Poisson de marais; Choupiquel* (Louisiana).

Great Lakes, Mississippi Valley to Virginia, Florida, and Texas, in lakes and sluggish streams.

Amia calva Linnæus, Syst. Nat., ed. XII, 1766, 500, Charleston, S. C.

Amia ocellicauda Richardson, Fauna Bor.-Amer., III, 1836, 236, Lake Huron, male.

Amia occidentalis DeKay, New York Fauna: Fishes, 1842, 269, St. Marys River, Mich.

Amia marmorata Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 412, New Orleans.

Amia ornata LeSueur, in Cuvier and Valenciennes, loc. cit., 420, Chibault, Mississippi River.

Amia viridis LeSueur, loc. cit., 421, New Orleans.

Amia canina Cuvier and Valenciennes, loc. cit., 424, Lake Erie; after Kirtland

Amia lentiginosa Cuvier and Valenciennes, loc. cit., 426; after Bonnat-terre; no locality.

Amia subcærulea Cuvier and Valenciennes, loc. cit., 427, New Orleans.

Amia cinerea Cuvier and Valenciennes, loc. cit., 430, Charleston, S. C.

Amia reticulata LeSueur, loc. cit., 431, Wabash River.

Amia thompsoni Duméril, loc. cit., II, 1870, 419, Lake Champlain.

Amia piquoti Duméril, loc. cit., 1870, 423, upper Mississippi.

Superorder TELEOSTEI. Bony fishes

Order N. ISOSPONDYLI (Physostomi; Malacopterygii). Soft-rayed fishes

Suborder ELOPOIDEA

Family 37. ELOPIDÆ. Tenpounders

Genus 87. ELOPS Linnæus.

Elops Linnæus, Syst. Nat., ed. XII, 1766, 518 (*E. saurus*).

Mugilomorus Lacépède, Hist. Nat. Poiss., V, 1803, 398 (*M. anna-carolina*).

Trichonotus Rafinesque, Anal. Nat., 1815, 88 (*Mugilomorus anna-carolina*); substitute for *Mugilomorus*, considered objectionable.

198. *Elops saurus* Linnæus. Tenpounder; John Mariggle; Bonyfish; Bonefish; Big-eyed herring; Mataquelo real; Lisa francesa.

Tropical Atlantic; West Indies, north to Long Island; common.

Elops saurus Linnæus, Syst. Nat., ed. XII, 1766, 518, Carolinas.

Argentina carolina Linnæus, loc. cit., 519, Carolinas.

Mugilomorus anna-carolina Lacépède, Hist. Nat. Poiss., V, 1803, 398, South Carolina.

Elops inermis Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 445, New York.

199. *Elops hawaiiensis* Regan. Awa.

Hawaii and Pacific coast of Mexico north to Gulf of California; Hawaii.

Elops hawaiiensis Regan, Ann. Mag. Nat. Hist., 9th ser., III, 1909, 39, Hawaii.

Family 38. MEGALOPIDÆ. Tarpons

Genus 88. TARPON Jordan and Evermann. Grande écaille.

Tarpon Jordan and Evermann, Fishes North and Mid. Amer., 1896, 409 (*Megalops atlanticus* Cuvier and Valenciennes).

200. *Tarpon atlanticus* (Cuvier and Valenciennes). Tarpon; Tarpum; Grande écaille; Silverfish; Sabalo; Savanilla.

Long Island to Brazil; abundant off Florida, breeding in the West Indies.

Megalops atlanticus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 398, Guadeloupe; San Domingo; Martinique; Porto Rico.

Megalops elongatus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 224, Long Island.

Suborder ALBULOIDEI

Family 39. ALBULIDÆ. Ladyfishes

Genus 89. ALBULA (Gronow) Bloch and Schneider. Ladyfishes.

Conorhynchus Nozeman, Act. Select., III, 1757, 382; non-binomial.

Albula Gronow, Zoophyl., 1763, 102; non-binomial

Albula Bloch and Schneider, Syst. Ichth., 1801, 432 (*A. conorhynchus* = *Esox vulpes* L.).

Butyrinus Lacépède, Hist. Nat. Poiss., V, 1803, 45 (*B. banana* = *Esox vulpes* L.).

Glossodus Cuvier, in Agassiz, Spix, Pisc. Brasil., 1829, 48 (*A. forskdli* = *Esox vulpes* L.).

201. *Albula vulpes* (Linnæus). Ladyfish; Bonefish; Macabi; Sanducha; Bananafish.

Tropical seas, ranging northward to San Diego; Long Island and southern Japan; perhaps more than one species included.

Unbarana Marcgrave, Hist. Brazil, 1648, Brazil.

Vulpes bahamensis Catesby, Nat. Hist. Carolinas, etc., 1737, Pl. II, fig. 1, Bahamas; non-binomial.

Esox vulpes Linnæus, Syst. Nat., ed. X, 1758, 313, Bahamas; based on the bonefish, *Vulpes bahamensis* of Catesby.

- Argentina glossodonta* Forskål, Descr. Anim., 1775, 68, Djidda, Arabia.
Macabi Parra, Diff. Piezas Cuba, 1787, 88, pl. 35, fig. 1, Cuba; based on *Unbarana* of Marcgrave.
Synodus argenteus Bloch and Schneider, Syst. Ichth., 1801, 398, Asia.
Clupea brasiliensis Bloch and Schneider, loc. cit., 427, Brazil.
Albula conorhynchus Bloch and Schneider, loc. cit., 432, Antilles; after Gronow and Plumier; called *Albula plumieri* on plate 86.
Amia immaculata Bloch and Schneider, loc. cit., Central America; after *Macabi* of Parra.
Butyrinus banana Lacépède, Hist. Nat. Poiss., V, 1803, 46, Ile de France.
Clupea macrocephala Lacépède, loc. cit., 426, Martinique; on a drawing by Plumier.
Glossodus forskåli Agassiz, in Spix, Pisc. Brasil., 1829, 49, Bahia; called *Engraulis sexicus* and *Engraulis bahiensis* on plates 22 and 24.
Albula parra Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 339, Martinique; Bahia; Rio de Janeiro.
Albula goreensis Cuvier and Valenciennes, loc. cit., 342, Gorea.
Albula neoguinaica Cuvier and Valenciennes, loc. cit., 350, New Guinea.
Albula seminuda Cuvier and Valenciennes, loc. cit., 351, New Guinea.
Albula erythrocheilus Cuvier and Valenciennes, loc. cit., 352, Friendly Islands.
Albula forsteri Cuvier and Valenciennes, loc. cit., 354, Tahiti.
Albula rostrata Gronow, Cat. Fishes, 1854, 189, American Ocean.

Genus 90. DIXONINA Fowler.

Dixonina Fowler, Proc. Acad. Nat. Sci. Phila., LXII, 1910, 651 (*D. nemoptera* Fowler).

202. Dixonina nemoptera Fowler.

San Domingo, West Indies.

Dixonina nemoptera Fowler, Proc. Acad. Nat. Sci. Phila., LXII, 1910, 652, San Domingo.

Suborder CLUPEOIDEI

Family 40. CHANIDÆ. Milkfishes

Genus 91. CHANOS Lacépède.

Chanos Lacépède, Hist. Nat. Poiss., V, 1803, 395 (*C. arabicus*).

Lutodeira Van Hasselt, in Ruppell, Atlas Reise Nord Africa, 1828, 17 (*L. indica* Ruppell = *Mugil chanos* L.)

Ptycholepis Gray, in Dieffenbach's Travels in New Zealand, II, about 1842, 218 (*Mugil salmoneus* Forster).

203. Chanos cyprinella Cuvier and Valenciennes. *Milkfish; Sabalo; Awa; Chani; Anged.*

Gulf of California to Panama and Hawaii; replaced in the western Pacific by *Chanos chanos* Forskål.

Chanos cyprinella Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 194, 198, Hawaiian Islands.

Family 41. HIODONTIDÆ. Mooneyes

Genus 92. HIODON LeSueur. *Mooneyes.*

Hiodon LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 334 (*H. tergus* LeSueur).

Glossodon Rafinesque, Amer. Month. Mag., II, September, 1818, 354 (*G. harengoides*).

Hyodon, corrected orthography.

204. Hiodon tergus LeSueur. *Mooneye; Toothed herring.*

Great Lakes and Mississippi Valley, north to Assiniboine River.

Hiodon tergus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 364, Ohio River.

Glossodon harengoides Rafinesque, Amer. Month. Mag., II, September, 1818, 354, Ohio River.

Glossodon heterurus Rafinesque, loc. cit., 354, Falls of Ohio River.

Hodon clodalis LeSueur, loc. cit., Ohio River at Pittsburgh.

Hyodon vernalis Rafinesque, Ichth. Ohiensis, 1820, 42, Ohio River.

205. *Hiodon selenops* Jordan and Bean. *Southern mooneye*; *Southern toothed herring*.
Tennessee, Cumberland, and Alabama Rivers.
Hyodon selenops Jordan and Bean, Bull. U. S. Nat. Mus., X, 1877, 67, Chattanooga, Tenn.
- Genus 93. AMPHIODON Rafinesque. *Goldeyes*.
Amphiodon Rafinesque, Jour. Phys., Paris, 1819, 421, Ohio River (*A. alosoides*; misprinted *alveoides*).
Elatomastius Gill and Jordan, Bull. U. S. Nat. Mus., X, 1877, 68 (*Hyodon chrysopsis* Richardson = *Amphiodon alosoides* Rafinesque).
206. *Amphiodon alosoides* Rafinesque. *Goldeye*; *La Quesche*; *Naccaysh*.
Ohio River and north to Saskatchewan; abundant and much valued in Manitoba.
Amphiodon alosoides (misprinted *alveoides*) Rafinesque, Jour. Phys., Paris, 1819, 421, Ohio River.
Hyodon amphiodon Rafinesque, Ichth. Ohiensis, 1820, 42, Ohio River near the falls.
Hyodon chrysopsis Richardson, Fauna Bor.-Amer., III, 1836, 232, Cumberland House, Saskatchewan River.
Cyprinus (*Abramis*?) *smithi* Richardson, loc. cit., 110, Richelieu River.

Family 42. CLUPEIDÆ. Herrings

- Genus 94. CLUPEA Linnæus. *Herrings*.
Clupea Linnæus, Syst. Nat., ed. X, 1758, 317 (*C. harengus* L.).
Rogenia Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 340 (*R. alba*, "the whitebait," the young of *harengus*).
207. *Clupea harengus* Linnæus. *Common herring*.
North Atlantic Ocean, south on our coast to Sandy Hook; Massachusetts; New Hampshire.
Clupea harengus Linnæus, Syst. Nat., ed. X, 1758, 317, seas of Europe.
Clupea membras Pallas, Zoogr. Rosso.-Asiat., III, 1811, 211, Baltic Sea.
Clupea halec Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 451, New York.
Clupea vittata Mitchill, loc. cit., 456, New York.
Clupea cærulea Mitchill, loc. cit., 457, New York.
Clupea elongata LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 234, Massachusetts.
Clupea leachi Yarrell, Zool. Jour., V, 1829, 277, pl. 12, England.
Clupea minima Peck, in Storer, Rept. Fish. Mass., 1839, 113, New Hampshire.
Clupea lineolata Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 256, locality unknown.
Rogenia alba Cuvier and Valenciennes, loc. cit., 340, England; the young form known as whitebait.
208. *Clupea pallasii* Cuvier and Valenciennes. *California herring*.
Pacific coast, from Kamchatka to San Diego.
Clupea harengus var. *Pallas*, Zoogr. Rosso.-Asiat., III, 1811, 209, Kamchatka.
Clupea pallasii Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 253, Kamchatka; based on Pallas's specimens.
Clupea mirabilis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 138, San Francisco.
Spratelloides bryoporus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 25, Sitka; young.
- Genus 95. POMOLOBUS Rafinesque. *Alewives*.
Pomolobus Rafinesque, Ichth. Ohiensis, 38, 1820 (*P. chrysochloris* Rafinesque).
Alausella Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 36 (*Clupea parvula* Mitchill = *Clupea medocrois* Mitchill).
- Subgenus ALAUSSELLA Gill.

209. *Pomolobus pseudoharengus* (Wilson). *Alewife*; *Branch herring*; *Gaspereau*; *Wall-eyed herring*; *Big-eyed herring*; *Sawbelly*; *Ellwife*; *Bang*. Atlantic coast of the United States; Lake Ontario and landlocked in the "Finger Lakes" of New York.
Clupea pseudoharengus Wilson, in Rees's Encycl., IX, about 1811, probably Philadelphia.
Clupea vernalis Mitchill, Rept. Fish. N. Y., 1814, 22, and Trans., Lit. Phil. Soc. N. Y., I, 1815, 454, New York.
Clupea megalops Rafinesque, Amer. Month. Mag., 1818, 206, Delaware River.
Meletta venosa Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 374, New York.
Pomolobus pseudoharengus lacustris Jordan, Man. Vert., ed. I, 1876, 265, Cayuga Lake, New York; specimens landlocked.
210. *Pomolobus mediocris* (Mitchill). *Hickory shad*; *Hickory jack*; *Fall herring*; *Tailor herring*; *Mattowacca*. Atlantic coast of United States from Maine to Florida.
Clupea mediocris Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 450, New York.
Clupea mattowacca Mitchill, loc. cit., 451, pl. 5, fig. 8, New York.
? *Clupea parvula* Mitchill, loc. cit., 452, New York.
? *Clupea pusilla* Mitchill, loc. cit., 452, New York.
Clupea fasciata LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 233, Massachusetts.
Clupea virens DeKay, New York Fauna: Fishes, 1842, 242, New York.
Alosa lineata Storer, Proc. Bost. Soc. Nat. Hist., II, 1847, 242, Massachusetts.
211. *Pomolobus æstivalis* (Mitchill). *Glut herring*; *Blueback*; *Black-belly*; *Summer herring*; *Kyack*; *Sawbelly*. Atlantic coast.
Clupea æstivalis Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 456, New York.
Alosa cyanonoton Storer, Hist. Fish. Mass., 1867, 161, Provincetown, Mass.
- Subgenus **POMOLOBUS** Rafinesque.
212. *Pomolobus chrysochloris* Rafinesque. *Skipjack*; *Blue herring*; *Nailrod*. Gulf of Mexico and Mississippi Valley; ascending streams, and introduced through the canals into Lake Erie and Lake Michigan.
Pomolobus chrysochloris Rafinesque, Ichth. Ohnensis, 1820, 38, Ohio River.
Meletta sueurii Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 375, Wabash River.
- Genus 96. **ALOSA** Linck. *Shad*. —
Alosa Linck, Mag. Neuestes Phys. Naturgesch., 1790, 35 (scarcely defined; type not specified; *Clupea alosa* L. = *Alosa* Cuvier).
Alosa Cuvier, Régne Anim., ed. II, 1829, 319 (*Clupea alosa* L.).
Alausa Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 389 (*Clupea alosa* L.).
213. *Alosa sapidissima* (Wilson). *Common shad*; *American shad*; *North River shad*; *Potomac shad*; *Connecticut River shad*. Atlantic coast of United States, from Miramachi River to Florida; from Monterey northward on the Pacific coast, as an introduced species.
Clupea sapidissima Wilson, in Rees's New Cyclopaedia, IXg., neither pagination nor date, but prior to 1812; no locality, but probably Philadelphia.
Clupea indigena Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 454, New York.
Alosa præstabilis DeKay, New York Fauna: Fishes, 1842, 255, pl. 15, fig. 4, New York.
214. *Alosa ohioensis* Evermann. *Ohio shad*. Mississippi River basin; Ohio River.
Alosa ohioensis Evermann, Rept. U. S. Fish Com., XXVII, 1901 (1902), 277, fig. 2, Ohio River at the falls.

215. *Alosa alabamæ* Jordan and Evermann. *Alabama shad*.
Gulf coast of United States; ascending streams.
Alosa alabamæ Jordan and Evermann, Rept. U. S. Fish Com., XXI,
1895 (1896), 203, Black Warrior River, Tuscaloosa, Ala.
- Genus 97. **SARDINIA** Poey. *True sardines*.
Sardinia Poey, Memorias, II, 1861, 311 (*S. pseudohispanica* Poey; not
of Jordan and Evermann, which is *Sardinella anchovia* Cuvier and
Valenciennes).
Sardina Antipa, Denkschr. Akad. Wiss. Wien, LXVIII, 1905 (*Clupea
pilchardus* L.).
216. *Sardinia cærulea* (Girard). *California sardine*.
Pacific coast, from Puget Sound to Magdalena Bay; very abundant.
Meletia cærulea Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 138,
San Francisco.
Alausa californica Gill, loc. cit., XIV, 1862, 281, California.
217. *Sardinia anchovia* (Cuvier and Valenciennes). *Sardinia de España*.
West Indies, north to Cuba.
Sardinella anchovia Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
1847, 269, Rio Janeiro; Martinique.
*Sardina*¹⁹ *pseudohispanica* Poey, Memorias, II, 1861, 311, Cuba.
- Genus 98. **SARDINELLA** Cuvier and Valenciennes. *Smooth-headed sardines*.
Sardinella Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 261,
(*S. aurita* Cuvier and Valenciennes).
218. *Sardinella aurita* Cuvier and Valenciennes. *Smooth sardine; Sardina
de España*.
Woods Hole to Florida; recorded from France; usually in rather deep
water.
Sardinella aurita Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
1847, 261, coast of France.
- Genus 99. **HARENGULA** Cuvier and Valenciennes. *Scaled sardines*.
Harengula Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 280
(*H. latula* Cuvier and Valenciennes).
219. *Harengula sardina* Poey. *Sardina de Ley*.
West Indian fauna, north to Key West.
Harengula sardina Poey, Memorias, II, 1860, 310, Cuba.
Harengula callolepis Goode, Proc. U. S. Nat. Mus., II, 1879, 152,
Bermuda.
220. *Harengula macrophthalmus* (Ranzani). *Sardina Escamuda; White-bill;
Pinceo*.
West Indies, Florida to Brazil.
Clupea macrophthalma Ranzani, Nov. Com. Acad. Sci. Bonon., V, 1842,
320, Brazil.
Harengula humeralis Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
1847, 293, Rio Janeiro; Bahia; Guadeloupe; San Domingo.
Alausa striata Cuvier and Valenciennes, loc. cit; XX, 1847, 429,
Guadeloupe; Bahia.
Alosa bishopi Müller and Troschel, in Schomburgk, Hist. Barbados,
1848, 675, Barbados.
221. *Harengula clupeiola* Cuvier and Valenciennes. *Caillou*.
West Indies.
Harengula clupeiola Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
1847, 289, Guadeloupe Island, West Indies.
222. *Harengula maculosa* Cuvier and Valenciennes.
West Indies.
Harengula maculosa Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
1847, 292, Martinique.
Alosa apicalis Müller and Troschel, in Schomburgk, Hist. Barbados,
1848, 675, Barbados.

¹⁹ Recent examination of Poey's type by Alvin Seale shows that this fish is a *Sardinia* apparently identical with *Clupea anchovia* Cuvier and Valenciennes, not the *Sardinella pseudohispanica* of Jordan and Evermann, which apparently is the same as *Sardinella aurita* Cuvier and Valenciennes.

223. *Harengula pensacolæ* Goode and Bean.
Florida.
Harengula pensacolæ Goode and Bean, Proc. U. S. Nat. Mus., II, 1879,
152, Pensacola, Fla.
224. *Harengula thrissina* (Jordan and Gilbert).
Gulf of California.
Clupea thrissina Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
353, Cape San Lucas.
- Genus 100. *LILE* Jordan and Evermann.
Lile Jordan and Evermann, Fishes North and Mid. Amer., 1896, 431
(*Clupea stolifera* Jordan and Gilbert).
225. *Lile stolifera* (Jordan and Gilbert).
Gulf of California to Panama; Mazatlan, Mexico.
Clupea stolifera Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
339, Mazatlan.
- Genus 101. *OPISTHONEMA* Gill. *Thread herring*
Opisthonema Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 37 (*Clupea*
thrissa Broussonet = *Megalops oghina* LeSueur); not of Osbeck.
226. *Opisthonema oglinum* (LeSueur). *Thread herring; Machuelo; Cailleu*
tassart; Sprat.
West Indies, ranging regularly north to Florida and Carolina; Longport,
N. J.; Newport, R. I.; Fortress Monroe.
Clupea thrissa Broussonet, Ichthyologie, fasc. I, 1782, Carolina; Jamaica;
not of Osbeck, 1757, which is a Chinese species of *Clupanodon*.
Megalops oghina LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 359,
Newport, R. I.
Megalops notata LeSueur, loc. cit., 359, Guadeloupe.
Chatoessus signifer DeKay, N. Y. Fauna: Fishes, 1842, 264, New York.
Chatoessus eumorphus Gosse, Naturalist's Sojourn in Jamaica, 1851,
290, Jamaica.
227. *Opisthonema libertate* (Günther). *Sardina machete*.
Pacific coast of Mexico and Central America.
Meletta libertate Günther, Proc. Zool. Soc. Lond., 1866, 603, Libertad,
Central America.
- Genus 102. *BREVOORTIA* Gill. *Menhadens*.
Brevoortia Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 37 (*Alosa*
menhaden Storer = *Clupea tyrannus* Latrobe).
228. *Brevoortia tyrannus* (Latrobe). *Menhaden; Mossbunker; Bonyfish;*
Whitefish; Bugfish; Fatback; Pogy.
Nova Scotia to West Indies; abundant.
Clupea tyrannus Latrobe, Trans. Amer. Phil. Soc. Phila., V, 1802, 77,
pl. 1, Chesapeake Bay.
Clupea menhaden Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 453,
New York.
Clupea neglecta Rafinesque, Amer. Month. Mag., 1818, 206, Long Island.
Clupea carolinensis Gronow, Cat. Fishes, 1854, 140, South Carolina.
229. *Brevoortia tyrannus brevipaudata* Goode. *Short-tailed menhaden*.
South Atlantic north to Connecticut.
Brevoortia tyrannus brevipaudata Goode, Proc. U. S. Nat. Mus., I, 1878,
34, Noank, Conn.
230. *Brevoortia patronus* Goode. *Gulf menhaden*.
Gulf of Mexico; coast of Texas.
Brevoortia patronus Goode, Proc. U. S. Nat. Mus., I, 1878, 39, Brazos
Santiago, Tex.
- Genus 103. *ILISHA* Gray.
Platygaster Swainson, Class. Anim., II, 1839, 294 (*Clupea africana*
Bloch); name preoccupied.
Ilisha Gray, in Richardson, Ichthy. China, in Proc. Brit. Assoc., 1845
(1846), 306 (*I. abnormis*); no description.
Pellona Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 300
(*P. orbignyana* = *Pristigaster flavipinnis* Valenciennes).

- 231. *Nisha caribbæa* Meek and Hildebrand.**
Caribbean Sea.
Nisha caribbæa Meek and Hildebrand, Marine Fishes of Panama, Pt. 1,
Field Mus. Nat. Hist., Zool., XV, Dec. 20, 1923, 191, Pl. X, fig. 1,
Porto Bello, Panama.
- Genus 104. CHIROCENTRODON** Günther.
Chirocentrodon Günther, Cat., VII, 1868, 463 (*C. tæniatus* Günther)
Medipellona Jordan and Seale, Mem. Mus. Comp. Zool., LXVII, No. 11,
May, 1926, 417 (*Pellona bleekariana* Poey).
- 232. *Chirocentrodon tæniatus* Günther.**
Jamaica.
Chirocentrodon tæniatus Günther, Cat., VII, 1868, 463, Jamaica.
- 233. *Chirocentrodon bleekariana* (Poey). *Anchoa pelada*.**
Matanzas, Cuba.
Pellona bleekariana Poey, Repertorio, II, 1867, 242, Matanzas, Cuba
- Genus 105. OPISTHOPTERUS** Gill.
Opisthopterus Gill, Proc. Acad. Nat. Sci. Phila., XII, 1861, 38 (*Pristigaster tartoon* Cuvier and Valenciennes).
- 234. *Opisthopterus dovii* (Günther).**
Panama.
Pristigaster dovii Günther, Cat., VII, 1861, 461, Panama.
- 235. *Opisthopterus macrops* (Günther).**
Panama.
Pristigaster macrops Günther, Proc. Zool. Soc. Lond., 1866, 603, Panama.
- Genus 106. PLIOSTEOSTOMA** Norman.
Pliosteostoma Norman, Ann. Mag. Nat. Hist., XI, 1923, 21 (*Pristigaster lutipinnis* Jordan and Gilbert).
- 236. *Plioosteostoma lutipinnis* (Jordan and Gilbert).**
Pacific coast of Mexico.
Pristigaster lutipinnis Jordan and Gilbert, Proc. U. S. Nat. Mus., IV,
1881, 340, Mazatlan, Mexico.
- Genus 107. ODONTOGNATHUS** Lacépède.
Odontognathus Lacépède, Hist. Nat. Poiss., II, 1799, 221 (*O. mucronatus*).
Gnathobolus Bloch and Schneider, Syst. Ichth., 1801, 556 (*Odontognathus mucronatus* Lacépède.).
- 237. *Odontognathus mucronatus* Lacépède.**
Coast of Guiana.
Odontognathus mucronatus Lacépède, Hist. Nat. Poiss., 1799, 221, pl.
7, fig. 2, Cayenne.
- 238. *Odontognathus panamensis* (Steindachner).**
Panama.
Pristigaster (Odontognathus) panamensis Steindachner, Ichth. Beitr.,
V, 1876, 24, Panama.
- Genus 108. PRISTIGASTER** Cuvier.
Pristigaster Cuvier, Règne Anim., ed. I, 1817, 176 (type not named;
later fixed as *P. cayanus* Cuvier).
- 239. *Pristigaster cayanus* Cuvier.**
Coast of Guiana and northern Brazil.
Pristigaster cayanus Cuvier, Règne Anim., ed. I, 1817, pl. 10, fig. 3,
Cayenne.
Pristigaster martin Agassiz, in Spix, Pisc. Brasil., 1829, 55, Brazil.
Pristigaster phacton Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
1847, 338, mouth of Amazon.

Family 43. DUSSUMIERIIDÆ. Round herrings

- Genus 109. JENKINSIA** Jordan and Evermann.
Jenkinsia Jordan and Evermann, Fishes North and Mid. Amer., 1896,
418 (*Dussumiera stolidifera* Jordan and Gilbert).

240. *Jenkinsia stolifera* (Jordan and Gilbert).
Gulf of Mexico from Florida to Yucatan.
Dussumiera stolifera Jordan and Gilbert, Proc. U. S. Nat. Mus., VII,
1884, 25, Key West, Fla.
241. *Jenkinsia lamprotænia* (Gosse).
Jamaica.
Clupea lamprotænia Gosse, Naturalist's Sojourn in Jamaica, 1851, 291,
pl. 1, fig. 2, Jamaica.
- Genus 110. *HALECUA* Jordan.
Halecula Jordan, Fossil Fishes of the Calif. Miocene, VIII, 1925,
(*Etrumeus acuminatus* Gilbert).
242. *Halecula acuminata* (Gilbert).
Gulf of California.
Etrumeus acuminatus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 56,
Gulf of California at Albatross station 3012 in 22 fathoms.
- Genus 111. *ETRUMEUS* Bleeker.
Etrumeus Bleeker, Verh. Bat. Gen. Japan, XXV, 1853, 58, (*Clupea micropus* Temminck and Schlegel, a Japanese species).
Perkinsia Rosa Smith Eigenmann, Amer. Nat., XXV, 1891, 153 (*P. othonops* R. S. Eigenmann).
243. *Etrumeus sadina* (Mitchill). *Round herring*.
Cape Cod to Gulf of Mexico.
Clupea sadina Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 457,
New York.
Alosa teres DeKay, New York Fauna: Fishes, 1842, 262, New York.
244. *Etrumeus micropus* Temminck and Schlegel.
Coast of Japan; Hawaii; found once off Point Loma in California.
Etrumeus micropus Temminck and Schlegel, Fauna Japonica, Poiss.,
1850, 236, pl. 107, fig. 2, Japan.
Perkinsia othonops Rosa Smith Eigenmann, Amer. Nat., XXV, 1891,
153, Point Loma, San Diego, Calif.

Family 44. DOROSOMIDÆ. Gizzard shad

- Genus 112. *DOROSOMA* Rafinesque. Gizzard shads.
Dorosoma Rafinesque, Ichth. Ohiensis, 1820, 39 (*D. notata* Rafinesque).
Chatoëssus Cuvier, Règne Anim., ed. 2, II, 1829, 320; in part (includes
Opisthonema, etc; restricted by Cuvier and Valenciennes to *Megalops cepedianus* LeSueur).
245. *Dorosoma cepedianum* (LeSueur). *Gizzard shad; Hickory shad*.
Cape Cod to Texas, entering all streams and variously landlocked.
Megalops cepedianus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818,
Delaware and Chesapeake Bays.
Clupea heterura Rafinesque, Amer. Month. Mag., 1818, 354, Ohio River.
Dorosoma notata Rafinesque, Ichth. Ohiensis, 1820, 39, Ohio River.
Chatoëssus ellipticus Kirtland, Rep. Zool. Ohio, 1839, 169, Ohio River.
Dorosoma insociabilis Abbot, Proc. Acad. Nat. Sci. Phila., XII, 1860,
365, Sturgeon Pond, Trenton, N. J.
246. *Dorosoma cepedianum exile* Jordan and Gilbert.
Streams of Texas.
Dorosoma cepedianum exile Jordan and Gilbert, Proc. U. S. Nat. Mus.,
V, 1882, 585, Galveston, Tex.
- Genus 113. *SIGNALOSA* Evermann and Kendall.
Signalosa Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897
(Feb. 9, 1898), 127 (*S. atchafalaya* Evermann and Kendall).
247. *Signalosa atchafalaya* Evermann and Kendall.
Coast of Louisiana and southward.
? *Chatoëssus mexicanus* Günther, Cat., VII, 1868, 409, Mexico.
Signalosa atchafalaya Evermann and Kendall, Bull. U. S. Fish Com.,
XVII, 1897 (Feb. 9, 1898), 127, pl. 7, fig. 4, Atchafalaya River,
Melville, La.

248. *Signalosa atchafalayæ vanhyningi* Weed.
Near Gainesville, Fla.
Signalosa atchafalayæ vanhyningi Weed, Field Mus. Nat. Hist., Zool., XII, October, 1925, 145, Prairie Creek 6 miles southeast of Gainesville, Fla.
249. *Signalosa mexicana* (Günther).
New Orleans to Central America.
Chatoessus mexicanus Günther, Cat., VII, 1868, 409, Mexico.
250. *Signalosa mexicana campi* Weed.
Streams about Brownsville, Tex.
Signalosa mexicana campi Weed, Field Mus. Nat. Hist., Zool., XII, October, 1925, 143, Resaca de la Guerra, near Brownsville, Tex.
251. *Signalosa petenensis* (Günther).
Lake Peten.
Meletia petenensis Günther, Proc. Zool. Soc. Lond., 1866, 603, Lake Peten, Yucatan.

Family 45. ENGRAULIDÆ. Anchovies

Genus 114. ENGRAULIS Cuvier. *Anchovies*.*Engraulis* Cuvier, Règne Anim., ed. I, 1817, 174 (*Chupea encrasicholus* L.).*Encrasicholus* Fleming, Brit. Anim., 1828, 183 (*Chupea encrasicholus* L.).

252. *Engraulis mordax* Girard. *California anchovy*.
Pacific coast of America, from Vancouver Island to Lower California.
Engraulis mordax Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 138, and in Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 334, Shoalwater Bay, Wash.

253. *Engraulis nanus* Girard. *Bay anchovy*.
San Francisco Bay; not known elsewhere.
Engraulis nanus Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 335, San Francisco.

Genus 115. ANCHOVIELLA²⁰ Fowler. *American anchovies*.*Anchoviella* Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1911, 211 (*Engraulis perfasciatus* Poey).*Anchoa* Jordan and Evermann, Proc. Calif. Acad. Sci., 4th Ser., XV, April 27, 1927, 502 (*Engraulis compressus* Girard).

Subgenus ANCHOVIELLA Fowler.

254. *Anchoviella perfasciata* (Poey). *Grubber broadhead*.
Florida Keys to Jamaica.
Engraulis perfasciatus Poey, Memorias, II, 1861, 313, Cuba.
255. *Anchoviella miarchus* (Jordan and Gilbert).
Pacific coast of America, from Mazatlan to Panama.
Stolephorus miarchus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 344, Mazatlan; perhaps a larval form.
256. *Anchoviella exigua* (Jordan and Gilbert).
Pacific coast of Mexico at Mazatlan.
Stolephorus exiguus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 343, Mazatlan.
257. *Anchoviella perthecata* (Goode and Bean).
Pensacola, Fla.
Stolephorus perthecatus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 434, Pensacola, Fla.

²⁰ *Anchoviella* differs from *Stolephorus* Lacépède and related Old World genera in the entire absence of ventral scales. The many species cover a variety of forms; *rostralis* and *spinifer* with its relatives being perhaps generically distinct from *perfasciatus*, *epsetus*, and related forms, having many more gillrakers; these latter species constitute the subgenus *Anchoa*, the others the subgenus *Anchoviella* Fowler.

258. *Anchoviella epsetus* (Bonnaterre). *Striped anchovy; Manjua; Anchovy fry.*
Cape Cod to Brazil.
Piquitinga Marcgrave, Hist. Brasil., 1648, 159, Brazil.
Menidia Browne, Hist. Jamaica, 1756, 441, Jamaica; nonbinomial.
Esox epsetus Bonnaterre, Tabl. Ichth., 1788, 175, Jamaica; after Browne.
Atherina brownii Gmelin, Syst. Nat., 1789, 1397, Jamaica; after Browne.
Engraulis tricolor Agassiz, in Spix, Pisc. Brasil., 1829, 51, Bahia, Pará.
Engraulis lemniscatus Cuvier, Règne Anim., ed. 2, II, 1829, 323, Brazil; after *Piquitinga* of Marcgrave.
Stolephorus hulecus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 343, Clearwater Harbor, Fla.; depth $5\frac{1}{2}$; A, 22.
259. *Anchoviella cubana* (Poey). *Bocon.*
Cuba and Porto Rico.
Engraulis cubanus Poey, Synopsis, 1868, 420, Cuba.
260. *Anchoviella ischana* (Jordan and Gilbert).
Pacific coast of America, from Mazatlan to Panama.
Stolephorus ischanus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 340, Mazatlan.
Stolephorus starksi Gilbert and Pierson, Fishes of Panama, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2813, Panama.
261. *Anchoviella cultrata* (Gilbert).
Santa Margarita Island, Lower California.
Stolephorus cultratus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 544, Santa Margarita Island, Lower California.
262. *Anchoviella delicatissima* (Girard).
San Diego Bay and southward on coast of Lower California.
Engraulis delicatissimus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 154, and Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 335, San Diego.
263. *Anchoviella helleri* Hubbs.
Gulf of California
Anchoviella helleri Hubbs, Proc. Biol. Soc., Wash., XXXIV, Mar. 31, 1921, 47, San Felipe Bay, gulf coast of Lower California.
264. *Anchoviella choerostoma* (Goode). *Hogmouth fry.*
Bermuda Islands; common in Hamilton Harbor.
Engraulis choerostomus Goode, Amer. Jour. Sci. Arts, August, 1874, 125, Bermuda.
265. *Anchoviella argyrophana* (Cuvier and Valenciennes).
Gulf Stream; Woods Hole, Mass.
Engraulis argyrophanus Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 49, equatorial Atlantic.
Stolephorus eurystole Swain and Meek, Proc. Acad. Nat. Sci. Phila., XXXVI, 1884, 34, Woods Hole, Mass.
266. *Anchoviella curta* (Jordan and Gilbert).
Mazatlan, Mexico.
Stolephorus curtus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 343, Mazatlan, Mexico.
267. *Anchoviella mitchilli* (Cuvier and Valenciennes).
Cape Cod to Texas on sandy shores, entering rivers; common.
Engraulis mitchilli Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 50, New York; Carolina; Lake Ponchartrain.
Engraulis louisiana LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 51, Lake Ponchartrain, La.
Engraulis duodecim Cope, Trans. Amer. Phil. Soc., 1866, 405, Beasley's Point, N. J. (dorsal fin said to be entirely anterior to the long and deeply concave anal, which is not true in *S. mitchilli*; the description otherwise agrees).
268. *Anchoviella lucida* (Jordan and Gilbert). *Sardina.*
Mazatlan, Mexico.
Stolephorus lucidus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 341, Mazatlan,

269. *Anchoviella naso* (Gilbert and Pierson).
Panama; common.
Stolephorus naso Gilbert and Pierson, in Jordan and Evermann, Fishes
North and Mid. Amer., 1898, 2813, Panama.
270. *Anchoviella robertsi* (Jordan and Rutter).
West Indies.
Stolephorus robertsi Jordan and Rutter, Proc. Acad. Nat. Sci. Phila.,
XLIX, 1897, 95, Jamaica.
271. *Anchoviella astilbe* (Jordan and Rutter).
West Indies.
Stolephorus astilbe Jordan and Rutter, Proc. Acad. Nat. Sci. Phila.,
XLIX, 1897, 95, Jamaica.
272. *Anchoviella scofieldi* (Jordan and Culver).
Astillero at Mazatlan; not very abundant.
Stolephorus scofieldi Jordan and Culver, in Jordan, Fishes of Sinaloa,
Proc. Calif. Acad. Sci., 3d Ser., V, 1895, 410, Mazatlan, Mexico.
273. *Anchoviella lyolepis* (Evermann and Marsh).
Porto Rico.
Stolephorus lyolepis Evermann and Marsh, Bull. U. S. Fish Com., XX,
1900 (Dec. 29), 89, fig. 13, Culebra Island, Porto Rico.
274. *Anchoviella branchiomelas* (Eigenmann).
North coast of Colombia.
Stolephorus branchiomelas Eigenmann, Proc. Am. Phil. Soc., LVI, 1917,
682, mouth of Rio Dagua, Colombia.
275. *Anchoviella arenicola* (Meek and Hildebrand).
Panama.
Anchovia arenicola Meek and Hildebrand, Marine Fishes of Panama,
in Field Mus. Nat. Hist., Zool., XV, 1923, 201, Pl. XIII, fig. 1,
Taboga Island, Panama.
276. *Anchoviella scitula* (Fowler).
Coast of California; San Diego.
Anchovia scitula Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1911, 211,
fig. 2, San Diego, Calif.
277. *Anchoviella parva* (Meek and Hildebrand).
Colon.
Anchovia parva Meek and Hildebrand, Marine Fishes of Panama, in
Field Mus. Nat. Hist., Zool., XV, 1923, 202, Pl. X, fig. 2, Porto Bello,
Panama.
278. *Anchoviella elongata* (Meek and Hildebrand).
Canal Zone.
Anchovia elongata Meek and Hildebrand, Marine Fishes of Panama, in
Field Mus. Nat. Hist., Zool., XV, 1923, 198, Pl. XII, fig. 2, Mindi
Cut, Canal Zone.
279. *Anchoviella cayorum* (Fowler).
Florida Keys.
Anchovia chærostoma cayorum Fowler, Proc. Acad. Nat. Sci. Phila.,
LVII, 1906, 85, fig. 4, Hailer's Rock, Florida Keys.
280. *Anchoviella platyargyrea* (Fowler).
West Indies.
Anchovia platyargyrea Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1911,
216, fig. 4, St. Martin's, West Indies.
281. *Anchoviella lepidentostole* (Fowler).
Coast of Surinam.
Anchovia lepidentostole Fowler, Proc. Acad. Nat. Sci. Phila., LXIII,
1911, 214, fig. 3, Surinam.
282. *Anchoviella guianensis* (Eigenmann).
Coast of Guiana.
Stolephorus guianensis Eigenmann, Mem. Carnegie Mus., V, 1912, 448,
Pl. LXII, fig. 5, Bartica Rocks, British Guiana; misspelled *Stoleferus*
in plate.

283. *Anchoviella trinitatis* (Fowler).
British West Indies.
Anchovia trinitatis Fowler, Proc. Acad. Nat. Sci. Phila., LXVII, 1915, 527, fig. 3, Port of Spain, Trinidad, British West Indies.
284. *Anchoviella filifera* (Fowler).
British West Indies.
Anchovia filifera Fowler, Proc. Acad. Nat. Sci. Phila., LXVII, 1915, 534, fig. 2, Port of Spain, Trinidad, British West Indies.
285. *Anchoviella abbotti* (Fowler).
British West Indies.
Anchovia abbotti Fowler, Proc. Acad. Nat. Sci. Phila., LXVII, 1915, 522, fig. 1, Port of Spain, Trinidad, British West Indies.
- Subgenus ANCHOA Jordan and Evermann.
286. *Anchoviella compressa* (Girard).
Point Conception to Lower California.
Engraulis compressus Girard, Pac. R. R. Surv., X, 1858, 336, San Diego.
287. *Anchoviella spinifera* (Cuvier and Valenciennes).
Coast of Guiana; recorded from Panama by Steindachner, probably an error.
Engraulis spinifer Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 39, Cayenne.
288. *Anchoviella rastralis* (Gilbert and Pierson).
Panama; abundant.
Stolephorus rastralis Gilbert and Pierson, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2811, Panama.
289. *Anchoviella mundeola* (Gilbert and Pierson).
Panama; abundant.
Stolephorus mundeolus Gilbert and Pierson, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2812, Panama.
290. *Anchoviella panamensis* (Steindachner).
Panama.
Engraulis panamensis Steindachner, Ichth. Beitr., IV, 1875, 39, Panama.
291. *Anchoviella eigenmannia* (Meek and Hildebrand).
Panama.
Anchovia eigenmannia Meek and Hildebrand, Marine Fishes of Panama, in Field Mus. Nat. Hist., Zool., XV, 1923, 20, Taboga Island, Canal Zone.
- Genus 116. ANCHOVIA Jordan and Evermann.
Anchovia Jordan and Evermann, Fishes North and Mid. Amer., 1896, 449, (*Engraulis macrolepidotus* Kner and Steindachner).
292. *Anchovia macrolepidota* (Kner and Steindachner). *Sardina bocona*.
Gulf of Mexico to Panama.
Engraulis macrolepidotus Kner and Steindachner, Abhandl. Bayer. Akad. Wiss., X, 1864, 21, Pl. III, fig. 2, Rio Bayano, Panama.
293. *Anchovia clupeioides* (Swainson). *Hechudo*; *Grubber broadhead*.
Coast of Cuba and southward; common, ascending streams.
Engraulis clupeioides Swainson, Nat. Hist. Fish., II, 1839, 388, Pernambuco.
Stolephorus surinamensis Bleeker, Nederl. Tijdschr. Dierk., 1866, 178, Surinam.
Engraulis productus Poey, Repertorio, 1866, 380, Cuba.
- Genus 117. AMPLOVA Jordan and Seale.
Amplova Jordan and Seale, Copeia, No. 141, 1925, 31 (*Stolephorus brevirostrum* Meek and Hildebrand; not *Engraulis brevirostris*, Günther, also a species of *Amplova*).
294. *Amplova balboæ* Jordan and Seale.
Panama.
Amplova balboæ Jordan and Seale, Copeia, No. 141, Apr. 30, 1925, 31, substitute for *Stolephorus brevirostrum* Meek and Hildebrand.

Anchovia brevirostrum Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, Pt. I, 193, Pl. XV, fig. 1; not *Engraulis brevirostris* Gunther from Brazil, also a species of *Amplova*.

Genus 118. CETENGRAULIS Günther.

Cetengraulis Günther, Cat., VII, 1868, 383 (*Engraulis edentulus* Cuvier).

295. *Cetengraulis edentulus* (Cuvier). Bocon; Tarpon fry.

West Indies to Brazil; common in Cuba.

Sprat Sloane, Hist. Jamaica, II, 1727, 282, Jamaica; after Sloane.

Engraulis edentulus Cuvier, Règne Anim., ed. 2, II, 1829, 323, Jamaica.

Engraulis brevis Poey, Repertorio, I, 1866, 379, Cuba.

296. *Cetengraulis gilberti* (Evermann and Marsh).

Coast of Porto Rico.

Stolephorus gilberti Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 352, Palo Seco near San Juan, Porto Rico.

297. *Cetengraulis garmani* (Evermann and Marsh). Porto Rican anchovy.

Porto Rico.

Stolephorus garmani Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899, (Dec. 19), 352, Puerto Real, Porto Rico.

298. *Cetengraulis mysticetus* (Günther).

Pacific coast of tropical America.

Engraulis mysticetus Günther, Proc. Zool. Soc. Lond., 1866, 604, Panama.

Stolephorus opercularis Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 275, Punta San Felipe, Gulf of California.

299. *Cetengraulis engymer* Gilbert and Pierson.

Panama Bay.

Cetengraulis engymer Gilbert and Pierson, in Jordan and Evermann, Fishes North and Mid. Amer. 1898, 2815, Panama Bay.

Genus 119. PTERENGRAULIS Günther.

Pterengraulis Günther, Cat., VII, 1868, 398, (*Clupea atherinoides* L.).

300. *Pterengraulis atherinoides* (Linnaeus).

Coast of Guiana and Brazil; ascending rivers.

Clupea atherinoides Linnaeus, Syst. Nat., ed. XII, 1766, 523, Surinam.

Genus 120. LYCENGRAULIS Günther.

Lycengraulis Günther, Cat., VII, 1868, 399 (*Engraulis grossidens* Cuvier).

301. *Lycengraulis grossidens* (Cuvier).

Coast of Guiana and Brazil.

Engraulis grossidens Cuvier, in Agassiz, Spix, Pisc. Brasil., 1828, 50, Rio Janeiro.

Engraulis janeiro Agassiz, loc. cit., pl. 24, fig. 1, Rio Janeiro.

Engraulis dentex Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 28, Rio Janeiro.

302. *Lycengraulis poeyi* (Kner and Steindachner).

Rio Bayano, Panama.

Engraulis poeyi Kner and Steindachner, Abhandl. Bayer. Akad. Wiss., X, 1864, 23, with plate; Rio Bayano.

Family 46. ALEPOCEPHALIDÆ

Genus 121. ALEPOCEPHALUS Risso.

Alepocephalus Risso, Mém. Acad. Nat. Sci. Turin, XXV, 1820, 270 (*A. rostratus* Risso).

303. *Alepocephalus agassizii* Goode and Bean.

Gulf Stream.

Alepocephalus agassizii Goode and Bean, Bull. Mus. Comp. Zool., X, 1883, 318, Gulf Stream.

304. *Alepocephalus productus* Gill.

Gulf Stream.

Alepocephalus productus Gill, Proc. U. S. Nat. Mus., VI, 1883, 256, Gulf Stream at Albatross station 2035 in 1,362 fathoms.

305. *Alepocephalus tenebrosus* Gilbert.
Santa Barbara Channel, California.
Alepocephalus tenebrosus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891,
545, Santa Barbara Channel.
306. *Alepocephalus asperifrons* Garman.
Off Panama in deep water.
Alepocephalus asperifrons Garman, Mem. Mus. Comp. Zool., XXIV,
1899, 291, Pl. LIX, fig. 1, Panama.
307. *Alepocephalus convexifrons* Garman
Off Pacific coast of Mexico, in deep water.
Alepocephalus convexifrons Garman, Mem. Mus. Comp. Zool., XXIV,
1899, 292, Pl. LIX, fig. 2; off Pacific coast of Mexico.
308. *Alepocephalus fundulus* Garman.
Pacific coast of Panama, in deep water.
Alepocephalus fundulus Garman, Mem. Mus. Comp. Zool., XXIV,
1899, 293, Pl. LVII, fig. 2; off Panama in deep water.
- Genus 122. MITCHILLINA Jordan and Evermann.
Mitchillina Jordan and Evermann, Fishes North and Mid. Amer.,
1896, 453, (*Alepocephalus bairdii* Goode and Bean).
309. *Mitchillina bairdii* (Goode and Bean).
Grand Banks of Newfoundland.
Alepocephalus bairdii Goode and Bean, Proc. U. S. Nat. Mus., II, 1879,
55, Grand Banks of Newfoundland.
- Genus 123. BATHYTROCTES Günther.
Bathytroctes Gunther, Ann. Mag. Nat. Hist., II, 5th ser., 1878, 249
(*B. microlepis* Günther).
310. *Bathytroctes stomias* Gilbert.
Coast of Oregon.
Bathytroctes stomias Gilbert, Proc. U. S. Nat. Mus., XII, 1890, 53,
coast of Oregon in 877 fathoms, Albatross station 3074.
311. *Bathytroctes inspector* Garman.
Pacific coast of Panama, in deep water.
Bathytroctes inspector Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
288, Pl. M., fig. 1; Panama.
312. *Bathytroctes alveatus* Garman.
Pacific coast of Panama in deep water.
Bathytroctes alveatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
Pl. LVIII, fig. 1, 287; Panama.
313. *Bathytroctes alvifrons* Garman.
Pacific coast of Panama in deep water.
Bathytroctes alvifrons Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
286, Pl. LVIII, figs. 2, 2a; Panama.
- Genus 124. XENOGNATHUS Gilbert.
Xenognathus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 311, (X.
profundorum Gilbert).
314. *Xenognathus profundorum* Gilbert.
Coast of California, in deep water, 1,350 to 2,182 fathoms.
Xenognathus profundorum Gilbert, Proc. U. S. Nat. Mus., XLVIII,
1915, 311, off Santa Catalina Island, California.
- Genus 125. NARCETES Alcock.
Narcetes Alcock, Ann. Mag. Nat. Hist., VI, 1890, 305 (*N. erimelas*
Alcock).
315. *Narcetes stomias* (Gilbert).
Off Pacific coast of America, from Oregon to southern California in
deep water.
Bathytroctes stomias Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 53,
coast of Oregon in 877 fathoms, Albatross station 3074.
316. *Narcetes pluriserialis* Garman.
Pacific coast of Panama in deep water.
Narcetes pluriserialis Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
289, Pl. LVII, fig. 3; Panama.

- Genus 126. LEPTOCHILICHTHYS** Garman.
Leptochilichthys Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 284
 (*L. agassizii* Garman).
- 317. *Leptochilichthys agassizii* Garman.**
 Pacific coast of Panama, in deep water.
Leptochilichthys agassizii Garman, Mem. Mus. Comp. Zool., XXIV,
 1899, 286, Pl. LVIII, fig. 3; off Panama.
- Genus 127. PLATYTROCTES** Günther.
Platytrectes Günther, Ann. Mag. Nat. Hist., II, 1878, 249 (*P. apus*
 Günther).
- 318. *Platytrectes apus* Günther.**
 Mid-Atlantic and the Arabian Sea.
Platytrectes apus Günther, Ann. Mag. Nat. Hist., II, 1878, 249, Mid-
 Atlantic.
- Genus 128. TALISMANIA** Goode and Bean.
Talismania Goode and Bean, Oceanic Ichth., 1895, 41 (*Bathytroctes*
homopterus Vaillant).
- 319. *Talismania antillarum* Goode and Bean.**
 Caribbean Sea.
Talismania antillarum Goode and Bean, Oceanic Ichth., 1895, 44, fig.
 49, Albatross station 2394, in Caribbean Sea, lat. 28° 38' 30" N.,
 long. 87° 2' W., 420 fathoms.
- 320. *Talismania æquatoris* Goode and Bean.**
 Coast of Ecuador.
Talismania æquatoris Goode and Bean, Oceanic Ichth., 1895, 44, fig. 50,
 off coast of Ecuador.
- Genus 129. CONOCARA** Goode and Bean.
Conocara Goode and Bean, Oceanic Ichth., 1895, 39 (*C. macdonaldi*
 Goode and Bean).
- 321. *Conocara macdonaldi* Goode and Bean.**
 Gulf of Mexico.
Conocara macdonaldi Goode and Bean, Oceanic Ichth., 1895, 39, fig. 48,
 Gulf of Mexico.
- 322. *Conocara macroptera* (Vaillant).**
 Deep seas, coast of Morocco to Soudan, banc d'Arguin, and from the
 Canaries. A specimen 8 inches in length was obtained by the
 Albatross at station 2751, lat. 16° 57' N., long. 63° 12' W., in 68
 fathoms.
Alepocephalus macropterus Vaillant, Expéd. Sci. Travailleur et Talisman,
 Poissons, 1888, 150, Pl. XI, figs 2a, 2b, 2c, coast of Morocco, Soudan,
 and Canaries.
- Genus 130. BAJACALIFORNIA** Townsend and Nichols.
Bajacalifornia Townsend and Nichols, Bull. Amer. Mus. Nat. Hist.,
 LII, 1925, 8 (*B. burragei* Townsend and Nichols).
- 323. *Bajacalifornia burragei* Townsend and Nichols.**
 West coast of Lower California in deep water.
Bajacalifornia burragei Townsend and Nichols, Bull. Am. Mus. Nat.
 Hist., LII, 1925, 8, fig. 3, off Todos Santos Bay, Lower California,
 in 590 fathoms.
- Genus 131. XENODERMICHTHYS** Günther.
Xenodermichthys Günther, Ann. Mag. Nat. Hist., II, ser. 5, 1878, 250
 (*X. nodulosus* Günther).
Aleposomus Gill, Amer. Nat., XVIII, 1884, 433 (*A. copei* Gill).
- 324. *Xenodermichthys copei* (Gill).**
 Gulf Stream.
Aleposomus copei Gill, Amer. Nat., XVIII, 1884, 433, Gulf Stream, lat.
 37° 12' 20" N., long. 69° 39' W.
- Genus 132. ERICARA** Gill and Townsend.
Ericara Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 232
 (*E. salmonea* Gill and Townsend).

325. *Ericara salmonea* Gill and Townsend.
Bering Sea, southwest of Pribilof Islands.
Ericara salmonea Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897,
232, Bering Sea.
- Genus 133. *ATOPICHTHYS* Garman.
*Atopichthys*²¹ Garman, Mem., Mus. Comp. Zool., XXIV, 1899, 325
(*A. esunculus* Garman).
326. *Atopichthys esunculus* Garman.
Pacific coast of Mexico.
Atopichthys esunculus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
327, Pl. LXV, figs. 2, 2a; off Acapulco, Mexico.

Suborder SALMONOIDEI

Family 47. SALMONIDÆ. Salmon; Trout

- Genus 134. *ONCORHYNCHUS* Suckley. *Pacific salmon*.
Oncorhynchus Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 313 (*Salmo*
scouleri Richardson=*Salmo gorbuscha* Walbaum).
Hypsifario Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Salmo*
kennerlyi Gill).
- Subgenus. *ONCORHYNCHUS* Suckley.
327. *Oncorhynchus gorbuscha* (Walbaum). *Humpback salmon; Pink salmon, Haddo; Holia; Gorbuscha*.
Pacific coast and rivers of North America and Asia, from California northward.
Salmo gorbuscha Walbaum, Artedi Pisc. 1792, 69, Kamchatka; after Gorbuscha of Pennant and Krasheninnikof.
Salmo gibber Bloch and Schneider, Syst. Ichth., 1801, 409, Kamchatka; after Krasheninnikof.
Salmo proteus Pallas, Zoogr. Rosso-Asiat., III, 1811, 376, Bering Sea.
Salmo scouleri Richardson, Fauna Bor.-Amer., III, 1836, 158; Observatory Inlet.
Salmo tschawyttschiformis Smitt, I Riksmuseum Befintliga Salmonider, 1886, 161; Port Clarence.
328. *Oncorhynchus keta* (Walbaum). *Dog salmon; Hayko; Le kai salmon; Chum; Calco salmon*.
Pacific coast of America, from San Francisco to Kamchatka.
Salmo keta vel kayko Walbaum, Artedi Pisc., 1792, 72, rivers of Kamchatka; after *keta* or *kayko* of Pennant and Krasheninnikof.
Salmo lagocephalus Pallas, Zoogr. Rosso-Asiat., III, 1811, 372; Bering Sea.
?Salmo japonicus Pallas, loc. cit., 382, Curile Islands; Amur River.
Salmo dermatinus Richardson, Voy. Herald, Zool., 1854, 167; Yukon River.
Salmo consuetus Richardson, loc. cit., 168; Yukon River.
Salmo canis Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1858, 9, and Monogr. Salmo, 101, 1861 (1874); Puget Sound.
329. *Oncorhynchus kisutch* (Walbaum). *Silver salmon; Kisutch; Medium red salmon; Skowitz; Hoopid salmon; Coho salmon; Bielaya ryba; Quisutch; Tschaviche*.
From San Francisco northward, especially in Puget Sound and the Alaskan fjords; south on the Asiatic coast to Japan.
?Salmo miltschutch Walbaum, Artedi Pisc., 1792, 70, Bering Sea; after *miltschutsch* or *miltschutsch* of Pennant and Krasheninnikof; probably the young of *kisutch*.
Salmo kisutch Walbaum, Artedi Pisc., 1792, 70, rivers and lakes of Kamchatka; after the *kisutch* of Pennant.
?Salmo striatus Bloch and Schneider, Syst. Ichth., 1801, 407, Kamchatka; after *miltschutsch* of Krasheninnikof.

²¹ This name is given not as a genus, but rather as a group name of larval fishes related to the eels or to the Alepocephalidae, commonly known as *Leptocephalus*, but *Leptocephalus morrisi* (type of *Leptocephalus*) is now known to be a young conger. A number of these larvae (*A. sicarius*, *cinctus*, *dentatus*, *falcidens*, *acutus*, *opichthys*, *cingulus*, *lychnus*, *obtusius*, and *longidens*) are described by Garman (Deep Sea Fishes), but they need not be specifically recognized here. The species called *Atopichthys esunculus* Garman, perhaps belonging to the Alepocephalidae, has been designated as the type of this group.

Salmo kisutch Bloch and Schneider, loc. cit., 407, Kamchatka; after Pennant.

Salmo sanguinolentus Pallas, Zoogr. Rosso-Asiat., III, 1811, 379, Bering Sea.

Salmo tsuppich Richardson, Fauna Bor.-Amer., III, 1836, 224, Columbia River.

Subgenus HYSIFARIO Gill.

330. *Oncorhynchus nerka* (Walbaum). *Blueback salmon*; *Redfish*; *Fraser River salmon*; *Nerka*; *Sau-qui salmon*; *Sockeye or Sukkegh salmon*; *Krashnaya ryba*.

Klamath River and Rogue River to northern Alaska, Kamchatka, and Japan.

Salmo nerka Walbaum, Artedi Pisc., 1792, 71; after the *nerka* of Pennant, *narka* of Krashenninnikof; rivers and seas of Kamchatka.

Salmo lycaodon Pallas, Zoogr. Rosso-Asiat., III, 1811, 370, Okhotsk Sea, Kamchatka.

Salmo paucidens Richardson, Fauna Bor.-Amer., III, 1836, 222, Columbia River.

Salmo tapidisma Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 365, Kamchatka; on a drawing.

Salmo arabatch Cuvier and Valenciennes, loc. cit., Kamchatka; on a drawing.

Salmo melampterus Cuvier and Valenciennes, loc. cit., Kamchatka; on a drawing.

Fario aurora Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 218; Girard, Pac. R. R. Surv., X, 1858, 308, pl. 68, Astoria, Oreg.

331. *Oncorhynchus tshawytscha* (Walbaum). *Chinook salmon*; *Tchaviche*; *King salmon*; *Columbia River salmon*; *Spring salmon*; *Sacramento River salmon*; *Tyee salmon*; *Tshawytscha*; *Quinnat salmon*.

Alaska, Oregon, and California; southward to Monterey, and to northern China.

Salmo tshawytscha Walbaum, Artedi Pisc., 1792, 71, rivers of Kamchatka; after the *tshawytscha* of Krashenninnikof, Descr. Kamchatka, 1768, 178, and the *tshawytscha* of Pennant, 1792.

Salmo orientalis Pallas, Zoogr. Rosso-Asiat., III, 1811, 367, Kamchatka.

Salmo quinnat Richardson, Fauna Bor.-Amer., III, 1836, 219, Columbia River; and of many writers.

Fario argyreus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 218, Cape Flattery; Fort Steilacoom.

Salmo confluentus Suckley, Ann. Lyc. Nat. Hist. N. Y., December, 1858, and Pacific R. R. Surv., XII, pt. 2, 1860, 334, Puyallup River near Fort Steilacoom.

Salmo cooperi Suckley, Notices New Spec. N. Amer. Salmon, New York, June, 1861, Okanogan River.

Salmo warreni Suckley, loc. cit., June, 1861, and loc. cit., 1861, 117 (1874) Fraser River, British Columbia.

Salmo richardi Suckley, loc. cit., June, 1861, and loc. cit., 1861, 117, Fraser and Skagit Rivers.

332. *Oncorhynchus kennerlyi* (Suckley).

Lakes of Oregon, Washington, Idaho, British Columbia, Alaska, and Hokkaido. A dwarf landlocked form. Probably several forms should be recognized.

Salmo kennerlyi Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 307, Lake Chiloweyuck, Washington.

Genus 135. SALMO Linnæus. Salmon and trout.

Salmo Linnæus, Syst. Nat., ed. X, 1758, 308 (*S. salar* Linnæus); after Artedi.

Trutta Linnæus, loc. cit., 308 (*trutta*, etc.: "*Truttæ corpore variegato*").

Trutta Geoffroy, Descr. de 719 Plantes, etc., 1767, 719 (*Salmo trutta* Linnæus).

Fario Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 277 (*Fario argenteus* = *Salmo fario* L.).

Salar Cuvier and Valenciennes, loc. cit., 314 (*Salmo ansonii* = *Salmo fario*).

Trutta Siebold, Süswasserfische Mitteleuropas, 1863, 280 (*Salmo trutta* L.).

Subgenus SALMO Linnæus.

- 333. *Salmo salar* Linnaeus.** *Common Atlantic salmon.*
 North Atlantic; ascending all suitable rivers in northern Europe and region north of Cape Cod; formerly abundant in the Hudson and occasional in the Delaware, its northern limit being about lat. 50° N., and southwest Greenland.
Salmo salar Linnaeus, Syst. Nat., ed. X, 1758, 308, seas of Europe.
Salmo mas Walbaum, Artedi Pisc., 1792, 58, Germany; based on "Der Hakelachs" of Bloch; probably not intended as a binomial name.
Salmo gloveri Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 86, Union River, Me.
- 334. *Salmo sebago* Girard.** *Landlocked salmon; Sebago salmon.*
 Lakes of Maine, New Hampshire, New Brunswick, and Nova Scotia.
Salmo sebago Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 380, Sebago Lake, Me.
- 335. *Salmo ouananiche* McCarthy.** *Ouananiche; Winninsh.*
 Lake St. John, Saguenay River, and northward.
Salmo salar ouananiche Eugene McCarthy, Forest and Stream, March 10, 1894, 206, and in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 487, Saguenay River, Canada.
- 336. *Salmo mykiss* Walbaum.** *Mykiss; Somka; Kamchatka salmon trout; Mukichi.*
 Kamchatka; probably not occurring in America.
Mykiss Pennant, Arctic Zool., 1792, 126, Kamchatka; after Krasheninikof, etc.
Salmo mykiss Walbaum, Artedi Pisc., 1792, 59, Kamchatka; based on *mykiss* of Pennant.
Salmo penshinensis Pallas, Zoogr. Rosso-Asiat., III, 1811, Gulf of Peshin.
Salmo purpuratus Pallas, loc. cit., 374, Bering Sea.
- 337. *Salmo regalis* Snyder.** *Royal silver trout.*
 Lake Tahoe.
Salmo regalis Snyder, Bull. U. S. Bur. Fish., XXXII, 1912 (Dec. 31), 26, Lake Tahoe near Brockway, Nev.
- Subgenus TRUTTA Linnaeus.**
- 338. *Salmo clarkii* Richardson.** *Columbia River trout; Cut-throat trout.*
 Coastwise streams, from Puget Sound south to Elk River, Humboldt County, California.
Salmo clarkii Richardson, Fauna Bor.-Amer., III, 1836, 225, Cathlamet River.
Fario stellatus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 219, Fort Steilacoom, Shoalwater Bay.
Salmo brevicauda Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 308, Puget Sound.
- 339. *Salmo lewisi* (Girard).** *Yellowstone trout; Cut-throat trout.*
 Snake River basin above the Shoshone Falls, through Two-Ocean Pass to the headwaters of the Yellowstone and other affluents of the upper Missouri; Yellowstone Lake.
Salmo lewisi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 219, Falls of Missouri River.
Salmo carinatus Cope, Hayden's Geol. Surv. Mont., 1871 (1872); locality unknown, probably Yellowstone River.
- 340. *Salmo bouvieri* (Bendire).** *Waha Lake trout.*
 Waha Lake, Washington.
Salmo purpuratus bouvieri Bendire, in Jordan and Gilbert, Synopsis, 1883, 315, Waha Lake, Wash.
- 341. *Salmo utah* Suckley.** *Cut-throat trout of Utah Lake.*
 Lakes and streams west of the Wasatch range, especially in Bear, Provo, Jordan, and Sevier Rivers and in Utah Lake.
Salmo utah Suckley, Monogr. Salmo, 1861 (1874), 136, Utah Lake; pale specimens from the lake.

342. *Salmo gibbsii* Suckley.
Middle Columbia River basin.
Fario tsuppitch Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 218,
Fort Dalles, Oreg.; not of Richardson
Salmo gibbsii Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1858, 1, Fort
Dalles; also in the Yakima, John Day, and Boise Rivers; also in the
Columbia at The Dalles, Oreg.
343. *Salmo henshawi* Gill and Jordan. *Lake Tahoe trout*; *Silver trout*; *Truckee trout*; *Pyramid Lake trout*.
Basin of the post-tertiary Lake Lahontan; Lake Tahoe, Pyramid Lake,
Wenemucca Lake, Webber Lake, Donner Lake, Independence Lake,
Truckee River, Humboldt River, Carson River, and most streams of
the eastern slope of the Sierra Nevada; also the headwaters of
Feather River, west of the Sierra Nevada; probably introduced from
Nevada.
Salmo henshawi Gill and Jordan, in Jordan, Man. Vert., ed 2, 1878,
258, Lake Tahoe.
344. *Salmo tahoensis* (Jordan and Evermann). *Great trout of Lake Tahoe*.
Deep waters of Lake Tahoe
Salmo clarkii tahoensis Jordan and Evermann, Fishes North and Mid.
Amer., 1898, 2870, Lake Tahoe.
345. *Salmo jordani* (Meek). *Jordan's trout*; *Spotted trout of Lake Southerland*.
Lake Southerland, west of Puget Sound
Salmo clarkii jordani Meek, Publ., Field Col. Mus., Zool., I, 1897
(1899), 229, Lake Southerland, Wash.
346. *Salmo declivifrons* (Meek). *Salmon trout of Lake Southerland*.
Lake Southerland, west of Puget Sound.
Salmo clarkii declivifrons Meek, Field Col. Mus., Zool., I, 1899, 230,
Lake Southerland, Wash.
347. *Salmo virginalis* (Girard). *Rio Grande trout*.
Upper Rio Grande and southward into the mountains of Chihuahua.
Salar virginalis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 220,
Utah Creek and at Sangre de Cristo Pass, Colo.
Salmo spilurus Cope, Hayden's Geol. Surv. Mont., 1871 (1872), 470;
Sangre de Cristo Pass, Colo.
348. *Salmo pleuriticus* Cope. *Colorado River trout*.
Colorado River basin.
Salmo pleuriticus Cope, Hayden's Geol. Surv. Mont., 1871 (1872),
471; headwaters of Green River, Wyo.
349. *Salmo stomias* Cope. *Greenback trout*.
Headwaters of Arkansas and Platte Rivers.
Salmo stomias Cope, Hayden's Geol. Surv. Wyo., 1870 (1871), 433;
South Platte River; locality unknown.
350. *Salmo macdonaldi* (Jordan and Evermann). *Yellowfin trout*.
Headwaters of the Arkansas River.
Salmo mykiss macdonaldi Jordan and Evermann, Proc. U. S. Nat.
Mus., XII, 1889 (January 20, 1890), 453, Twin Lakes, Colo.
351. *Salmo crescentis* (Jordan and Beardslee). *Blueback trout of Lake Crescent*.
Crescent Lake, Clallam County, Washington.
Salmo gairdneri crescentis Jordan and Beardslee, Proc. Calif. Acad.
Sci., 2d ser., VI, 1896, 207, pl. 22, Crescent Lake, Clallam County,
Wash.
352. *Salmo bathocetor* Meek. *Long-headed trout of Crescent Lake*.
Deep water of Crescent Lake, Washington.
Salmo bathocetor Meek, Field Col. Mus., Zool., I, 1899, 227, Crescent
Lake, Wash.
353. *Salmo beardslei* (Jordan and Seale). *Speckled trout of Lake Crescent*.
Crescent Lake, Clallam County, Wash.
Salmo gairdneri beardslei Jordan and Seale, Proc. Calif. Acad. Sci.,
2d Ser., VI, 1896, 209, pl. 23, Crescent Lake, Clallam County, Wash.

354. *Salmo eremogenes* Evermann and Nichols.
Streams of central Wash.
Salmo eremogenes Evermann and Nichols, Proc. Biol. Soc. Wash., XXII, June 25, 1909, 93, pl. 2, Crab Creek at Rocky Ford near Ritzville, Wash.
355. *Salmo smaragdus* Snyder. *Emerald trout*.
Truckee Basin, Nev.
Salmo smaragdus Snyder, Bull. U. S. Bur. Fish., XXXV, 1915-16 (Sept. 28, 1917), 80, fig. 9, Pyramid Lake, Nev.
356. *Salmo aquilarum* Snyder. *Eagle Lake trout*.
Lakes of northeastern California.
Salmo aquilarum Snyder, Bull. U. S. Bur. Fish., XXXV, 1915-16 (Sept. 28, 1917), 77, fig. 7, Eagle Lake near mouth of Pine Creek, Modoc Co., Calif.
357. *Salmo irideus* Gibbons. *Rainbow trout*; *Coast range trout*.
Coastwise mountain streams of the Coast Ranges from the northern coast of California southward to San Diego County, Calif.; entering the sea where it grows to 6 or 10 pounds, returning as a steelhead.
Salmo iridia Gibbons, Proc. Calif. Acad. Nat. Sci., I, 1855, 35, San Leandro Creek, Alameda County, Calif.
Salmo rivularis Ayres, loc. cit., 42, Martinez, Calif.; steelhead form.
358. *Salmo gairdnerii* Richardson. *Brook trout of western Oregon to southern Alaska*; *Steelhead*; *Salmon trout*; *Kamloops trout*; *Sht-tse*.
Southern Alaska and Puget Sound to southern Oregon; in streams of the Coast Ranges and the Cascades, descending to the sea as a "steelhead"; perhaps not separable from *Salmo irideus*. Kamloops Lake, Okanogan Lake, and other lakes tributary to Frazer River or to the upper Columbia.
Salmo gairdnerii Richardson, Fauna Bor.-Amer., III, 1836, 221, Columbia River at Fort Vancouver.
Salmo masoni Suckley, Pac. R. R. Surv., XII, part 2, 1860, 345, Cathlamet River.
Salmo truncatus Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 3, New Dungeness, straits of Fuca.
Oncorhynchus kamloops Jordan, Forest and Stream, XXXIX, No. 19, November 10, 1892, 405, Kamloops Lake, British Columbia.
359. *Salmo shasta* (Jordan). *McCloud River rainbow trout*; *Rainbow trout of fish culturists*.
Streams of the Sierra Nevada from Mount Shasta southward; known from McCloud River; the limits of its range not well known. Introduced into the Truckee River and many eastern streams, as well as in Europe and New Zealand.
Salmo irideus or rainbow trout of most writers and especially of fish-culturists; not *Salmo irideus* Gibbons, which is based on coastwise specimens.
Salmo gairdneri shasta Jordan, Thirteenth Biennial Rept. Calif. Fish Com., 1894, 142, with plate; McCloud River at Baird, Shasta County, Calif.
360. *Salmo stonei* (Jordan). *Nissuee trout*; *No-shee trout*.
McCloud River and upper Sacramento River.
Salmo gairdneri stonei Jordan, Thirteenth Biennial Rept. Calif. Fish Com., 1894, 142, with plate; McCloud River at Baird, Calif.; perhaps a variant of *Salmo shasta*.
361. *Salmo newberrii* Girard.
Klamath River.
Fario or *Salmo newberrii* Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 225; new name for fish misidentified and described as "*Salmo gairdneri* Richardson" in Pacific R. R. Survey, X, 1859, 313, Pl. LXXI, figs. 1-4, Klamath River, Oreg.
362. *Salmo gilberti* (Jordan). *Kern River trout*.
Kern River, Calif.
Salmo gairdneri gilberti Jordan, Thirteenth Biennial Rept. Calif. Fish Com., 1894, 143, with plate; Kern River at Soda Springs, Calif.

363. *Salmo rosei* Jordan and McGregor. *Rose trout*.
Certain headwaters of Kern River, Calif.
Salmo rosei Jordan, Proc. Acad. Nat. Sci. Phila., LXXVI, 1924, 19,
Pl. II, Lake Culver, Calif., and neighboring lakes.
364. *Salmo agua-bonita* (Jordan). *Golden trout of south fork of Kern River*.
South fork of Kern River, Calif.; introduced into Cottonwood Lakes,
east of Mount Whitney.
Salmo mykiss agua-bonita Jordan, Proc. U. S. Nat. Mus., XV, 1892,
481, "Volcano or Whitney Creek, California," by error; the original
type came from the Cottonwood Lakes on the east side of Mount
Whitney, into which the species had been introduced from the south
fork of Kern River. This species does not occur in Volcano Creek
nor in Whitney Creek.
365. *Salmo whitei* Evermann. *Golden trout of western tributaries of Kern
River; Stewart White trout*.
Western tributaries of Kern River, Calif.
Salmo whitei Evermann, Bull. U. S. Bur. Fish., XXV, 1905 (May 19,
1906), 20, Pl. XVI; south fork of Kaweah River, Calif., into which
the species had been transplanted from Coyote Creek, a western
tributary of Kern River.
366. *Salmo roosevelti* Evermann. *Golden trout of Volcano Creek; Roosevelt
trout; Volcano Creek trout*.
Native only to Volcano Creek, an eastern tributary of Kern River,
Calif.
Salmo roosevelti Evermann, Bull. U. S. Bur. Fish., XXV, 1905 (May 19,
1906), 26, Pl. I, Volcano Creek, Kern River Basin, Calif.
367. *Salmo evermanni* Jordan and Grinnell. *San Bernardino trout; Evermann's
trout*.
Headwaters of Santa Ana River, Calif., above the falls.
Salmo evermanni Jordan and Grinnell, Proc. Biol. Soc. Wash., XXI,
1908, 31, Pl. II, south fork of the Santa Ana River, Calif., at 8,200
feet altitude.
368. *Salmo nelsoni* Evermann. *Lower California trout; Nelson's trout*.
Mountain streams of northern Lower California.
Salmo nelsoni Evermann, Proc. Biol. Soc. Wash., XXI, 1908, 26, Pl. I,
San Ramon River, San Pedro Martir Mountains, Lower California.
- Genus 136. CRISTIVOMER Gill and Jordan. *Great Lakes trout*.
Cristivomer Gill and Jordan, in Jordan, Man. Vert., ed. 2, 1878, 356.
(*Salmo namaycush* Walbaum).
369. *Cristivomer namaycush* (Walbaum). *Great Lakes trout; Mackinaw
trout; Longue (Vermont); Togue (Maine); Namaycush; Masamacush*.
Great Lakes region and lakes of northern New York, New Hampshire,
and Maine; headwaters of Columbia and Fraser Rivers; streams of
Vancouver Island, and north to the Arctic Circle.
Namaycush salmon Pennant, Arctic Zool., 1792, 191, Hudson Bay.
Salmo namaycush Walbaum, Artdi Pisc., 1792, 68, Hudson Bay; based
on the *Namaycush salmon* of Pennant.
Salmo pallidus Rafinesque, Amer. Month. Mag., December, 1817,
120, Lake Champlain.
Salmo amethystinus Mitchell, Jour. Acad. Nat. Sci. Phila., I, 1818, 410,
Sault Ste. Marie.
Salmo confinis DeKay, N. Y. Fauna: Fishes, 1842, 238, Louis Lake,
Hamilton County, N. Y.; Silver Lake, Pa.
Salmo symmetricus Prescott, Am. Jour. Sci. Arts, XI, 1851, 340, Lake
Winnepiseogee, N. H.
Salmo toma Hamlin, The Togue, in Holmes, 2d Ann. Rept., Maine
Fish Com., 1862, 109; lakes of Maine.
Salmo adirondacus Norris, Amer. Angler's Book, 1865, 255; Adirondack
lakes.

370. *Crystivomer siscowet* (Agassiz). *Siscowet*.

Lake Superior, in deep water, where it feeds on the fat ciscoes; a deep-water variant of *C. namaycush*.

Salmo siscowet Agassiz, Lake Superior, 1850, 333, Lake Superior.

Salmo siskawitz Agassiz, in Herbert Frank Forester's Fish and Fishing, 1850, 112, with plate; Lake Superior.

Salmo ursinus Barnston, Rept. Fish. Canada, Lake Superior; reference uncertain.

Genus 137. *SALVELINUS* Richardson. *Charrs*.

Salvelini Nilsson, Prodr. Ichth. Scand., 1832, 7 (*alpinus*); group name.

Salvelinus Richardson, Fauna Bor.-Amer., III, 1836, 169 (*Salmo salvelinus* L.); after Nilsson.

Baione DeKay, N. Y. Fauna: Fishes, 1842, 244 (*B. fontinalis* DeKay).

Umbla Rapp, Fische Bodensees, 1854, 32 (*Salmo umbla*=*alpinus* L.).

371. *Salvelinus fontinalis* (Mitchill). *Brook trout*; *Speckled trout*; *Eastern brook trout*.

Maine to the Saskatchewan and northward to Labrador, southward in the Alleghenies to headwaters of the Savannah, Chattahoochee, Catawba, and French Broad Rivers; largely introduced into western streams but not native west of the Mississippi except in Minnesota and Iowa. Some unnamed variants included.

Salmo fontinalis Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 435, near New York City.

Salmo alleghaniensis Rafinesque, Ichth. Ohiensis, 1820, 44, brooks falling into the Allegheny and Monongahela Rivers.

Salmo nigrescens Rafinesque, Ichth. Ohiensis, 1820, 45, near the Laurel Hills, Pa.

Salmo canadensis Hamilton Smith, in Griffith's Cuvier, X, 1834, 474, Canada; dots blood red, each "in white circular spot"

Salmo hudsonicus Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 310, Hudson Bay and tributaries; Labrador; Newfoundland.

372. *Salvelinus agassizii* Garman. *Dublin Pond trout*.

Dublin Pond, N. H.

Salmo agassizii Garman, Nineteenth Rept., Mass. Fish Com., 1885, 20, Dublin Pond (Lake Monadnock), N. H.; Center Pond, N. H.

373. *Salvelinus timagamensi* Henn and Rinkenbach. *Aurora trout*.

White Pine Lake, Clearwater Lake, Timigami region, Ontario, of the St. Lawrence Basin.

Salvelinus timagamensis Henn and Rinkenbach, Ann. Carnegie Mus., XVI, 1925, 131, figs. 1, 2, and 3; White Pine Lake.

374. *Salvelinus aureolus* Bean. *Sunapee trout*; *American saibling*; *Golden trout of Sunapee Lake*.

Sunapee Lake, N. H.; Dan Hole Pond, Carroll County, N. H., tributary to Saco River; and Floods Pond, Otis, Me., tributary to Union River; also in Averill Pond, Vt.

Salvelinus aureolus Bean, Proc. U. S. Nat. Mus., X, 1887, 628, Sunapee Lake, N. H.

375. *Salvelinus rossii* (Richardson). *Ross's arctic salmon*.

Lakes of Greenland and Boothia Felix.

?*Salmo hearneri* Richardson, Franklin's First Voy. 1823, 706, and in Fauna Bor.-Amer., III, 1836, 167, Bloody Fall, Coppermine River, Lat. 67°; not certainly recognizable, description imperfect.

Salmo rossii Richardson, App. Ross's Voy., LVI, 1835; and in Fauna Bor.-Amer., III, 1836, 163, Regent's Inlet, Boothia Felix.

Salmo alipes Richardson, loc. cit., LVII, 1835, and in Fauna Bor.-Amer., III, 1836, 169, Lakes in Regent's Inlet, Boothia Felix.

Salmo nitidus Richardson, loc. cit., LVII, 1835, and Fauna Bor.-Amer., 1836, 171; Lake of Boothia Felix.

376. *Salvelinus stagnalis* (Fabricius). *Greenland charr*.

Waters of Greenland, Boothia, and neighboring regions.

Salmo stagnalis Fabricius, Fauna Groenlandica, 1780, 175; Alpine ponds of Greenland; not migratory.

Salmo rivalis Fabricius, loc. cit., 176, Alpine brooks of Greenland; not entering the sea.

Salmo immaculatus H. R. Storer, Jour. Bost. Soc. Nat. Hist., VI, 1850, 364, lower St. Lawrence; Canadian "salmon trout." Name preoccupied. Sea-run forms.

377. *Salvelinus naresi* (Günther). Lakes of Arctic America, Discovery Bay, and Cumberland Gulf.
Salmo naresi Gunther, Proc. Zool. Soc. Lond., 1877, 476, plate, fresh-water lakes near Discovery Bay.
378. *Salvelinus arcturus* (Günther). *Arctic charr*.
Victoria Lake and Flocberg Beaches of Arctic America, latitude 82° 34'.
Salmo arcturus Gunther, Proc. Zool. Soc. Lond., 1877, 294, Pl. XXXII, Victoria Lake, Flocberg Beach
379. *Salvelinus malma* (Walbaum). *Malma trout*.
Waters of Kamchatka and Alaska, everywhere abundant; descending to the sea; apparently intergrading southward with *Salvelinus spectabilis*.
Salmo malma Walbaum, Artdi Pisc., 1792, 66, Kamchatka; based on *Malma* of Pennant.
Salmo curilus Pallas, Zoogr. Rosso-Asiat., III, 1811, 251, Curile Islands.
Salmo callaris Pallas, loc. cit., 353, Bering Sea.
Salmo lævigatus Pallas, loc. cit., 385, Curile Islands.
Salmo nummifer Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 365, Kamchatka; on a drawing by Mertens.
Salmo erythrorhynchus Cuvier and Valenciennes, loc. cit., 367, Kamchatka.
Salmo tudes Cope, Proc. Amer. Phil. Soc. Phila., XIII, 1873, 24, Captains Harbor, Unalaska.
Salmo pluvius Hilgendorf, Mitth., Deutsch. Ges. Ostasiens, XI, 1876, 25; clear streams throughout Japan.
380. *Salvelinus malma spectabilis* (Girard). *Dolly Varden trout*; *Oregon charr*; *Bull trout*; *Red-spotted trout*; *Golet*.
Streams west of the Cascade Range, from the upper Sacramento to Montana and Alaska; in the Columbia basin as far up as Montana and Idaho.
Salmo spectabilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 218; Fort Dalles, Oreg.
Salmo bairdii Suckley, loc. cit., 309, tributary of Flathead River, Mont.
Salmo parkei Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 309, Kootenay River.
Salmo campbelli Suckley, loc. cit., 313, Fort Dalles, Oreg.; substitute for *spectabilis*, preoccupied in *Fario*, not in *Salmo* or *Salvelinus*.
Salmo lordii Gunther, Cat., VI, 1866, 148, Skagit River; dwarf specimens.
381. *Salvelinus oquassa* (Girard). *Oquassa trout*; *Blueback trout*; *Quasky*.
Rangeley Lakes, western Maine.
Salmo oquassa Girard, Proc. Bost. Soc. Nat. Hist., IV, 1854, 262, Oquassa or Oquossoc Lake, Me.
382. *Salvelinus marstoni* (Garman). *Lac de Marbre trout*; *Marston trout*.
Salmo marstoni Garman, Science, July 14, 1893, Lac de Marbre, Ottawa County, Quebec.

Family 48. COREGONIDÆ. Whitefishes

Genus 138. STENODUS Richardson. *Inconnu*.

Stenodus Richardson, in Back's Narrative Arctic Land Exped., back append., 1836, 521 (*Salmo mackenzii* Richardson).

383. *Stenodus mackenzii* (Richardson). *Inconnu*.

Delta of MacKenzie River.

Salmo mackenzii Richardson, Franklin's Jour., 1823, 707, MacKenzie River.

Genus 139. LEUCICHTHYS Dybowski. *Lake herrings and ciscoes*.

Argyrosomus Agassiz, Lake Superior, 1850, 330 ("*clupeiformis*" of DeKay, not of Mitchill=*Salmo harengus* Rich); not of M. de la Pylaie, which is *Sciæna aquila* Cuvier.

Leucichthys Dybowski, Fische Baikal-Wassersysteme, Verk. K.K. Zool.-Bot. Gesell. Wien, Bd. XXIV, 1874, 300 (*Salmo omul* Pallas).
Allosomus Jordan, Man. Vert., ed. 2, 1878, 361 (*Coregonus tullibee* Richardson).

Thrissomimus Gill, in Jordan and Evermann, Bull. U. S. Bur. Fish., XXIX, 1909 (1911), 3 (*Coregonus artedi* LeSueur).
Cisco Jordan and Evermann, Bull. U. S. Bur. Fish., XXIX, 1909 (1911), 3 (*Argyrosomus nigripinnis* Gill).

Subgenus **THRISOMIMUS** Gill.

384. *Leucichthys harengus* (Richardson). *Saginaw Bay herring*; *Georgian Bay herring*.
 Shallow bays of Lakes Huron and Michigan; Georgian Bay, Saginaw Bay, Green Bay, etc.
Salmo (*Coregonus*) *harengus* Richardson, Fauna Bor.-Amer., III, 1836, 210, Pl. XC, fig. 2; Georgian Bay at Penetanguishene, Ontario.
Coregonus clupeiformis Agassiz, Lake Superior, 1850, 339, the Pic; Shallow Bays, Michipicoten Island; not *clupeiformis* of Mitchill.
Coregonus albus Agassiz, loc. cit., p. 342, the Pic; not of LeSueur.
385. *Leucichthys osmeriformis* (Smith). *Seneca Lake herring*; *Seneca Lake smelt*.
 Lakes of central New York, tributary to Lake Ontario.
Coregonus hoyi Bean, Proc. U. S. Nat. Mus., V, 1882, 658; not of Gill,
Coregonus osmeriformis Smith, Bull. U. S. Fish. Com., XIV, 1894, Pl. I, II, Seneca Lake, Skaneateles Lake, N. Y.
386. *Leucichthys sisco* (Jordan). *Cisco of Lake Tippecanoe*.
 Small lakes of Wisconsin and northern Indiana.
Argyrosomus sisco Jordan, Amer. Nat., 1875, 135, Lake Tippecanoe near Warsaw, Ind.
387. *Leucichthys sisco huronius* (Jordan and Evermann). *Lake Huron herring*; *Blueback or Michigan herring*.
 Lakes Huron and Michigan. The common form from which *L. sisco* is derived.
Argyrosomus huronius Jordan and Evermann, Proc. U. S. Nat. Mus., XXXVI, March 3, 1909, 167, fig. 2, Port Stanley, Ontario.
388. *Leucichthys ontariensis* Jordan and Evermann.
 Lake Ontario and Cayuga Lake, N. Y.
Coregonus clupeiformis DeKay, New York Fauna: Fishes, 1842, 248, pl. 60, fig. 198, Lake Ontario; not *clupeiformis* of Mitchill.
Leucichthys ontariensis Jordan and Evermann, Bull. U. S. Bur. Fish., XXIX, 1909 (1911), 13, Lake Ontario and Cayuga Lake, N. Y.
389. *Leucichthys lucidus* (Richardson). *Great Bear Lake herring*.
 Mackenzie River and tributaries; Great Bear Lake and River.
Salmo (*Coregonus*) *lucidus* Richardson, Fauna Bor.-Amer., III, 1836, 207, Pl. XC, fig. 1, with figure, Great Bear Lake.
390. *Leucichthys laurettae* (Bean). *Lauretta whitefish*.
 Lakes and streams of northern and western Alaska.
Coregonus laurettae Bean, Proc. U. S. Nat. Mus., IV, 1881, 156, Point Barrow, Alaska.
391. *Leucichthys alascanus* (Scotfield). *Arctic lake herring*.
Argyrosomus alascanus Scotfield, in Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2817, northern Alaska near Bering Straits.
392. *Leucichthys pusillus* (Bean). *Least lake herring*.
 Lakes of Yukon basin and Alaska generally.
Coregonus pusillus Bean, Proc. U. S. Nat. Mus., XI, 1888, 526, Kobuk River, Alaska.
393. *Leucichthys athabascæ* Harper and Nichols. *Athabasca herring*.
 Saskatchewan River.
Leucichthys athabascæ Harper and Nichols, Bull. Amer. Mus. Nat. Hist., XLI, 1919, 269, Saskatchewan River.
394. *Leucichthys macrognathus* Harper and Nichols. *Mackenzie River herring*.
 Mackenzie River.
Leucichthys macrognathus Harper and Nichols, Bull. Amer. Mus. Nat. Hist., XLI, 1919, 269, Mackenzie River.

395. *Leucichthys entomophagus* Harper and Nichols.
Mackenzie River.
Leucichthys entomophagus Harper and Nichols, Bull. Amer. Mus. Nat. Hist., XLI, 1919, 267, Mackenzie River.
396. *Leucichthys gemmifer* Snyder. *Cisco of Lake Bonneville*.
Bear River Basin.
Leucichthys gemmifer Snyder, Bull. U. S. Bur. Fish., XXXVI, 1817-18 (May 7, 1919), 4, Bear Lake, near Fish Haven, Idaho.
397. *Leucichthys artedi* (LeSueur). *Lake herring; Erie herring; Common lake herring; Grayback*.
Lake Erie and Lake St. Clair; Lake Huron at Port Huron.
Coregonus artedi LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 231, Lake Erie (at Buffalo) and Niagara River (Lewiston).
398. *Leucichthys artedi bisselli* (Bollman). *Rawson Lake herring; Bissell's herring*.
Glacial lakes of southern Michigan once tributary to Lake Erie.
Coregonus tullibee bisselli Bollman, Bull. U. S. Fish Com., VIII, 1888, 223, Rawson Lake and Howard Lake, Mich.
399. *Leucichthys eriensis* (Jordan and Evermann). *Jumbo herring; Erie great herring*.
Lake Erie and Lake Huron.
Argyrosomus eriensis Jordan and Evermann, Proc. U. S. Nat. Mus., XXXVI, March 3, 1909, 165, fig. 1, Lake Erie at Port Stanley.
400. *Leucichthys reighardi* Koelz.
Lake Michigan.
Leucichthys reighardi Koelz, Occ. Papers, Mus. Zool. Univ. Mich., No. 146, March 22, 1924, 5; Lake Michigan 18 miles off Michigan City, Ind.
401. *Leucichthys alpenae* Koelz.
Lake Michigan and Lake Huron, including Georgian Bay.
Leucichthys alpenae Koelz, Occ. Papers, Mus. Zool. Univ. Mich., No. 146, I, March 22, 1924; Lake Michigan off Isle aux Galets.
Subgenus CISCO Jordan and Evermann.
402. *Leucichthys supernas* Jordan and Evermann. *Cisco of Lake Superior*.
Deep waters of Lake Superior.
Leucichthys supernas Jordan and Evermann, Bull. U. S. Bur. Fish., XXIX, 1909 (1911), 22, Lake Superior.
403. *Leucichthys prognathus* (H. M. Smith). *Cisco of Lake Ontario; Ontario longjaw; Bloater; Chub*.
Deep waters of Lake Ontario, in depths of 60 fathoms and more
Coregonus prognathus Smith, Bull. U. S. Fish Com., XIV, 1894, 4, pl. 1, fig. 3, Lake Ontario at Wilson, N. Y.
404. *Leucichthys johannae* (Wagner). *Lake Michigan cisco; Bloater of Lake Michigan*.
Northwestern part of Lake Huron in deep water, and also for the whole length of Lake Michigan.
Argyrosomus johannae Wagner, Science, N. S., XXXI, No. 807, June 17, 1910, 957-958; Lake Michigan in about 25 fathoms, some 18 miles off Racine, Wis.
405. *Leucichthys nigripinnis* (Gill). *Blackfin of Lake Michigan*.
Deep waters of Lake Michigan and certain small lakes in Wisconsin.
Argyrosomus nigripinnis Gill, in Hoy, Trans. Wis. Acad. Sci., I, 1872, 100, Lake Michigan off Racine; name only.
Coregonus nigripinnis Jordan, Man. Vert., ed. 2, 1878, 275; description.
406. *Leucichthys cyanopterus* Jordan and Evermann.
Deep waters of Lake Superior.
Leucichthys cyanopterus Jordan and Evermann, Bull. U. S. Bur. Fish., XXIX, 1909 (1911), 27, fig. 13, Lake Superior, off Marquette, Mich.

407. *Leucichthys hoyi* (Gill). *Cisco of Lake Michigan; Kiyi; Chub; Mooneye cisco*.
Lake Michigan in deep water.
Argyrosomus hoyi Gill, in Hoy, Trans. Wis. Acad. Sci., I, 1872, 100,
Lake Michigan off Racine; no description.
Argyrosomus hoyi Gill, in Jordan, Amer. Nat., March, 1875, 135, Lake
Michigan off Racine, Wis.; description same specimens.
408. *Leucichthys kiyi* Koelz. *Kiyi; Mooneye; Waterbelly*.
Lake Michigan.
Leucichthys kiyi Koelz, Occ. Papers, Mus. Zool. Univ. Mich., No. 104,
Dec. 31, 1921; Lake Michigan, 12 miles E. by S. of the mouth of
Sturgeon Bay ship channel in about 60 fathoms.
409. *Leucichthys birgei* Wagner. *Cisco of Green Lake*.
Green Lake, Wisconsin.
Leucichthys birgei Wagner, Bull. Wis. Nat. Hist. Soc., IX, 1911, 73,
Green Lake, Wis., 28 meters below the surface.
410. *Leucichthys zenithicus* (Jordan and Evermann). *Longjaw of Lake Superior*.
Lake Superior, in deep water; possibly in other lakes.
Argyrosomus hoyi Milner, Rept. U. S. Fish Com., II, 1872-73 (1874),
86, Lake Superior at Outer Island, Wis.; not of Gill, Hoy, or Jordan,
and not original type.
Argyrosomus zenithicus Jordan and Evermann, Proc. U. S. Nat.
Mus., XXXVI, March 3, 1909, 169, fig. 3, Lake Superior between
Duluth and Isle Royale.

Subgenus ALLOSOMUS Jordan.

411. *Leucichthys manitoulinus* Jordan and Evermann. *Manitoulin tullibee*.
North channel of Lake Huron, and probably lakes of Minnesota.
Argyrosomus tullibee, Evermann and Smith, Rept. U. S. Fish Com., XX,
1894, 320, pl. 28: in part, the plate being tullibee.
Leucichthys manitoulinus Jordan and Evermann, Bull. U. S. Bur. Fish.,
XXIX, 1909, 31, fig. 16, Blind River, North channel, Lake Huron.
412. *Leucichthys tullibee* (Richardson). *Tullibee; Tulipi*.
Lake Winnipeg Basin, perhaps entering Lake Superior Basin.
Salmo (Coregonus) tullibee Richardson, Fauna Bor.-Amer., III, 1836,
201, Cumberland House, Pine Island Lake, near Lake Winnipeg.
Leucichthys nipigon Koelz, Occ. Papers Mus. Zool. Univ. Mich., No.
158, April 6, 1925; Lake Nipigon, Lake Winnipeg.
- Genus 140. COREGONUS Linnæus. *Whitefishes; Lavarets*.
Coregonus Artedi, Genera Pisc., 1738, 9.
Coregoni Linnæus, Syst. Nat., Ed. X, 1758, 310 (*Salmo lavaretus* L.).
Tripteronotus Lacépède, Hist. Nat. Poiss., V, 1803, 48 (*T. havutin*
Lacépède = *Salmo lavaretus* L.).
Coregonus Lacépède, loc. cit., X, 263 (*Salmo lavaretus* L.).
Coregonus Cuvier, Règne Anim. Ed. I, 1817, 162 (*thymallus, lavaretus*,
etc.).
413. *Coregonus clupeaformis* (Mitchill). *Whitefish; Labrador whitefish; Lake Superior whitefish; Manitoba whitefish; Musquaw River whitefish; Whiting of Lake Winnepesaukee; Shad of Lake Champlain*.
Great Lakes, excepting Lake Erie; and neighboring waters.
Salmo clupeaformis Mitchill, Amer. Month. Mag., II, 1818, 321, Falls
of St. Mary, northern extremity of Lake Huron.
Salmo otsego Clinton, account of the *Salmo otsego* or the Otsego basse,
1822, 1, with plate, Otsego Lake.
Coregonus labradoricus Richardson, Fauna Bor.-Amer., III, 1836, 206,
Musquaw River, Labrador.
Salmo (Coregonus) sapidissimus Agassiz, Lake Superior, 1850, 344,
Lake Superior.
Salmo labor Agassiz, loc. cit., 348, Lake Superior.
Coregonus neohantoni Prescott, Amer. Jour. Sci. Arts, XI, 1851, 343,
Lake Winnepesaukee, N. H.
? *Coregonus richardsonii* Günther, Cat., VI, 1866, 185, Arctic America.

414. *Coregonus clupeaformis albus* (LeSueur). *Lake Erie whitefish; Common whitefish.*
Lake Erie and Lake St. Clair; Lake Ontario and various lakes to western New York and New Hampshire. New York; probably not separable from *Coregonus clupeaformis*, although local differences appear.
Coregonus albus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 232, Lake Erie.
415. *Coregonus nelsonii* Bean. *Humpback whitefish, Nelson's whitefish.*
Alaska, from Bristol Bay northward.
Coregonus nelsonii Bean, Proc. U. S. Nat. Mus., VII, 1884, 48, Nulato, Alaska.
- Genus 141. PROSOPIUM Milner.
Prosopium Milner, in Jordan, Man Vert., ed. 2, 1878, 361 (*Coregonus quadrilaterals* Richardson).
416. *Prosopium quadrilaterale* (Richardson). *Menominee whitefish; Pilotfish; Round whitefish; Shadwarter.*
Alaska and upper Great Lakes to New England, in lakes.
Coregonus quadrilaterals Richardson, Franklin's Narrative, 1823, 714, Pl. XXV, fig. 2, Fort Enterprise, British America.
Coregonus novæ-angliæ Prescott, Amer. Jour. Sci. Arts, XI, 1851, 342, Lake Winnepesaukee, N. H.
417. *Prosopium kennicotti* (Milner). *Kennicott's whitefish.*
Mackenzie River, Canada; Yukon River and other streams of the Alaska region; Lake Bennett, Alaska.
Coregonus kennicotti Milner, in Jordan and Gilbert, Synopsis, 1883, 298, Fort Good Hope, British America.
418. *Prosopium preblei* (Harper and Nichols).
Mackenzie River, Canada.
Coregonus preblei Harper and Nichols, Bull. Amer. Mus. Nat. Hist., XLI, 1919, 266, Pl. XV, fig. 2, Mackenzie River.
419. *Prosopium stanleyi* (Kendall). *Stanley's whitefish.*
Lakes of northern Maine.
Coregonus stanleyi Kendall, Bull. U. S. Fish Com., XXII, 1902 (1904), 366, with figure, thoroughfare between Mud and Cross Lakes, Aroostook County, Me.
420. *Prosopium williamsoni* (Girard). *Rocky Mountain whitefish.*
Rivers of Sierra Nevada and west slope of the Rocky Mountains, from the Fraser and the Columbia to the Truckee and other streams of the Lahontan Basin of Nevada; abundant, especially in lakes of northern Idaho, western Montana, Washington, and Utah.
Coregonus willamsoni Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 136, Des Chutes River, Oreg.
421. *Prosopium williamsoni cismontanum* (Jordan). *Yellowstone whitefish.*
Streams of the Rocky Mountain region tributary to the upper Missouri.
Coregonus willamsoni cismontanum Jordan, Bull. U. S. Fish Com., IX, 1889, 49, pl. 9, figs. 8, 9, Horsethief Creek, Madison River, Mont.
Hardly distinct from *P. willamsoni*.
422. *Prosopium coulteri* (Eigenmann and Eigenmann). *Coulter's whitefish.*
Headwaters of the Columbia.
Coregonus coulteri Eigenmann and Eigenmann, Amer. Nat., Nov., 1892, 961, Kicking Horse River at Field, British Columbia.
423. *Prosopium couesi* (Milner).
Headwaters of Saskatchewan River.
Coregonus couesi Milner, Rept U. S. Fish Com., II, 1872-73 (1874), 88, Chief Mountain Lake, Mont.
424. *Prosopium spilonotus* (Snyder). *Bonneville whitefish.*
Bear River Basin.
Coregonus spilonotus Snyder, Bull. U. S. Bur. Fish., XXXVI, 1917-18 (May 7, 1919), 6, Bear Lake, near Fish Haven, Idaho.

425. *Prosopium abyssicola* (Snyder). *Bear Lake whitefish*.
 Bear Lake, Idaho.
Coregonus abyssicola Snyder, Bull. U. S. Bur. Fish., XXXVI, 1917-18
 (May 7, 1919), 8, Bear Lake, near Fish Haven, Idaho.
- Genus 142. **IRILLION** Jordan.
Irillion Jordan, Proc. Acad. Nat. Sci. Phila., LXX, 1918, 342 (*Coregonus oregonius* Jordan and Snyder).
426. *Irillion oregonius* (Jordan and Snyder). *Chiselmouth jack*.
 Mountain streams of Oregon tributary to the Columbia.
Coregonus oregonus Jordan and Snyder, Proc. U. S. Nat. Mus., XXXVI,
 1911, 425, with fig., McKenzie River, Ore.

Family 49. THYMALLIDÆ. Graylings

- Genus 143. **THYMALLUS** Cuvier. *Graylings*.
Thymallus Cuvier, Règne Anim., ed. 2, II, 1829, 306 (*Salmo thymallus* Linnaeus); not *Thymalus* Latreille, 1802, a genus of *Coleoptera*.
Choregon Minding, Lehrbuch Naturgesch. Fische, 1832, 119 (*Salmo thymallus* L.) substitute for *Thymallus*, regarded as preoccupied.
Orthocolus Gistel, Thierreich, 1848, XI, 1838 (*Salmo thymallus* L.); substitute for *Thymallus*, regarded as preoccupied.
427. *Thymallus signifer* (Richardson). *Arctic grayling; Poisson bleu*.
 Mackenzie River to Alaska and the Arctic Ocean.
Coregonus signifer Richardson, Franklin's Jour., 1823, 711; Winter Lake near Fort Enterprise.
Coregonus thymalloides Richardson, Fauna Bor.-Amer., III, 1836, 714, Winter River.
428. *Thymallus tricolor* Cope. *Michigan grayling*.
 Streams of northern Michigan; formerly abundant in Au Sable River, Jordan River, and other streams in the southern peninsula, and Otter Creek, near Keweenaw, in the northern peninsula.
 ?*Thymallus ontariensis* Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 452; said to have been brought by Milbert from Lake Ontario; most likely from Lake Geneva, Switzerland.
Thymallus tricolor Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 80, Au Sable River, Mich.
429. *Thymallus montanus* Milner. *Montana grayling*. A hardy and attractive aquarium fish.
 Madison and Gallatin Rivers, Montana.
Thymallus montanus Milner, Rept. U. S. Fish Com., II, 1872-73 (1874), 741, tributary of Missouri River at Camp Baker.

Family 50. OSMERIDÆ. Smelts

- Genus 144. **MALLOTUS** Cuvier. *Capelins*.
Mallotus Cuvier, Règne Anim., ed. 2, II, 1829, 305 (*Clupea villosa* Muller).
430. *Mallotus villosus* (Müller). *Capelin; Lodde*.
 Arctic America, south to Cape Cod and southern Alaska; Kamchatka.
Clupea villosa Muller, Prodr. Zool. Dan., 1777, 245, Greenland.
Salmo arcticus Fabricius, Fauna Grœnlandica, 1780, 177, Greenland.
Salmo grœnlandicus Bloch, Ichthyologie, VIII, 1794, pl. 99, Greenland.
Salmo socialis Pallas, Zoogr. Ross-Asiat., III, 389, islands between Asia and America.
Osmerus macrodon Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 385, no locality.
- Genus 145. **THALEICHTHYS** Girard. *Eulachon*.
Thaleichthys Girard, Pac. R. R. Surv., X, 1858, Fishes, 325 (*T. stevensi* Girard=*Salmo* (*Mallotus*) *pacificus* Richardson).
431. *Thaleichthys pacificus* (Richardson). *Eulachon; Candlefish; Oolachan*.
 Oregon to Alaska.
Salmo (*Mallotus*) *pacificus* Richardson, Fauna Bor.-Amer., III, 1836, 226, Columbia River.
Thaleichthys stevensi Girard, Pac. R. R. Surv., X, 1858, 325, pl. 75, figs. 1-4, Puget Sound.

432. *Thaleichthys albatrossis* (Jordan and Gilbert). *Kodiak smelt*.
Off Kodiak Island, Alaska.
Osmerus albatrossis Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2823, Kodiak Island; probably not distinct from *Thaleichthys pacificus*.
- Genus 146. *SPIRINCHUS* Jordan and Evermann.
Spirinchus Jordan and Evermann, Fishes North and Mid. Amer., I, 1896, 522 (*Osmerus thaleichthys* Ayres).
433. *Spirinchus thaleichthys* (Ayres).
Pacific coast of America from San Francisco northward to Bristol Bay, Alaska.
Osmerus thaleichthys Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 62, San Francisco.
- Genus 147. *ALLOSMERUS* Hubbs.
Allosmerus Hubbs, Proc. Biol. Soc. Wash., XXXVIII, 1925, 53 (*Osmerus attenuatus* Lockington).
434. *Allosmerus attenuatus* (Lockington).
Coast of California, from San Francisco northward.
Osmerus attenuatus Lockington, Proc. U. S. Nat. Mus., III, 1880, 66, San Francisco.
- Genus 148. *OSMERUS* Lacépède. *Smelts*.
Osmerus Lacépède, Hist. Nat. Poiss., V, 1803, 229 (*Salmo eperlanus* L.).
Eperlanus Gaimard, Voy. Island and Greenland, 1851 (*Salmo eperlanus* L.).
435. *Osmerus dentex* Steindachner. *Rainbow herring*.
Coast of Alaska and south on the Asiatic coast to northern China.
Osmerus dentex Steindachner, Sitzungsber. Kais. Akad. Wiss. Wien, LXI, 1870, 429, northern China.
436. *Osmerus mordax* (Mitchill). *American smelt*; *Icefish*.
Atlantic coast of United States, from Virginia northward to Gulf of St. Lawrence; Lakes Champlain and Memphremagog.
Atherina mordax Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 446, New York.
Osmerus viridescens LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 230, coast of Maine.
437. *Osmerus mordax spectrum* (Cope). *Wilton smelt*.
Wilton Pond, Kennebec County, Me.
Osmerus spectrum Cope, Proc. Amer. Phil. Soc., XI, 1870, 490, Wilton Pond, Kennebec County, Me.
438. *Osmerus mordax abbotti* (Cope). *Cobessicentic smelt*.
Landlocked in Cobessicentic Lake, Kennebec County, Me.
Osmerus abbotti Cope, Proc. Amer. Phil. Soc., XI, 1870, 490, Cobessicentic Lake, Me.
439. *Osmerus sergeanti* ²² Norris.
New York Bay and southward.
Osmerus sergeanti Norris, Proc. Acad. Nat. Sci. Phila., XX, 1868, 93, Schuylkill River, N. J.
- Genus 149. *HYPOMESUS* Gill. *Surf smelts*.
Mesopus ²³ Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 14 (*Argentina pretiosa* Girard).
Hypomesus Gill, loc. cit., 15 (*Argentina pretiosa* Girard).
440. *Hypomesus pretiosus* (Girard). *Surf smelt*.
Coast of California and Oregon, from Monterey northward.
Argentina pretiosa Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 155, San Francisco.
Osmerus elongatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 17, San Francisco.

²² The fish is described, but not named, in Proc. Acad. Nat. Sci. Phila., XIII, 1861, 58. It is named in Proc. Acad. Nat. Sci. Phila., XX, 1868, 93.

²³ *Mesopus*, a name left in type through failure to correct proof sheets. *Hypomesus* of same date, a page lower, chosen by author as name.

441. *Hypomesus olidus* (Pallas). *Pond smelt*.

Alaska and Kamchatka.

Salmo (*Osmerus*) *olidus* Pallas, Zoogr. Rosso.-Asiat., III, 1811, 391, lakes and rivers of Kamchatka.? *Osmerus oligodon* Kner, Denkschr. Akad. Wiss. Wien, 1865, 185, tab. 4, fig. 1.**Genus 150. *THEROBROMUS* Lucas. *Ghost smelts*.***Therobromus* Lucas, in Jordan and Gilbert, Fishes of Bering Sea, 1898, 440 (*T. callorhini* Lucas).**442. *Therobromus callorhini* Lucas. *Ghost smelt*.**

Bering Sea.

Therobromus callorhini Lucas, in Jordan and Gilbert, Fishes of Bering Sea, in Jordan, Fur Seals and Fur-Seal Islands, Pt. III, 1898, 440, Bering Sea, in southern portion; known only from skeletons and bones found in the stomachs of fur seals.**Family 51. ARGENTINIDÆ. Argentines****Genus 151. ARGENTINA Linnæus. *Argentines*.***Argentina* (Artedi) Linnæus, Syst. Nat., Ed. X, 1758, 315 (*A. sphyxæna* L.).*Silus* Reinhardt, Bemærk., Skandinav. Ichthy., 1833, 11 (*Argentina silus* Ascanius).*Acantholepis* Krøyer, Danmarks Fiske, III, 1846-49, 98 (*Argentina silus* Ascanius).**443. *Argentina silus* Ascanius.**

Northern Europe; occasionally taken on the Grand Banks and off the coast of Maine.

Argentina silus Ascanius, Icon. Rev. Nat., 1763, Pl. III, 3, 24. Norway.*Silus ascanii* Reinhardt, Bemærk., Skandinav. Ichth., 1833, 11, Norway.*Argentina syrtensium* Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 261, off Sable Island in 200 fathoms; from stomach of *Phycis tenuis*.**444. *Argentina striata* Goode and Bean.**

Gulf of Mexico.

Argentina striata Goode and Bean, Oceanic Ichth., 1895, 52, fig. 62, Gulf Stream.**445. *Argentina sialis* Gilbert.**

Coast of California.

Argentina sialis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 56, Albatross station 3017 in 58 fathoms.**Genus 152. *LEUROGLOSSUS* Gilbert.***Leuroglossus* Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 57 (*L. stilbius* Gilbert).**446. *Leuroglossus stilbius* Gilbert.**

Coast of California, in deep water.

Leuroglossus stilbius Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 57, from Albatross stations 2997 and 2998 in 221 and 40 fathoms.**Family 52. MICROSTOMIDÆ****Genus 153. *NANSENIA* Jordan and Evermann.***Nansenia* Jordan and Evermann, Fishes North and Mid. Amer., 1896, 528 (*Microstomus grænlandica* Reinhardt).**447. *Nansenia grænlandica* (Reinhardt).**

Greenland.

Microstomus grænlandica Reinhardt, Vidensk. Selsk. Natury. Math. Afhandl., VIII, 1841, LXXIV, Greenland.**Family 53. BATHYLAGIDÆ****Genus 154. *BATHYLAGUS* Günther.***Bathylagus* Günther, Ann. Mag. Nat. Hist., II, 1878, 248 (*B. antarcticus* Günther).

448. *Bathylagus benedicti* Goode and Bean.
Gulf Stream.
Bathylagus benedicti Goode and Bean, Oceanic Ichth., 1895, 55, fig. 64,
Gulf Stream.
449. *Bathylagus euryops* Goode and Bean.
Gulf Stream.
Bathylagus euryops Goode and Bean, Oceanic Ichth., 1895, 55, fig. 63,
Gulf Stream, in about lat. 40° N. and long. 70° W.
450. *Bathylagus pacificus* Gilbert.
Coast of Washington, in deep water.
Bathylagus pacificus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 55,
Albatross stations 3071 and 3074, off coast of Washington.
451. *Bathylagus borealis* Gilbert.
Coast of Alaska, in deep water.
Bathylagus borealis Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1896),
402, north of Unalaska Island at *Albatross* station 3327 in 322
fathoms.
452. *Bathylagus milleri* Jordan and Gilbert.
Off coast of southern California in deep water.
Bathylagus milleri Jordan and Gilbert, in Jordan and Evermann, Fishes
North and Mid. Amer., 1898, 2825, Cortez Banks, off San Diego, Calif.
- Genus 155. *XENOPHTHALMICHTHYS* Regan.
Xenophthalmichthys Regan, Ann. Mag. Nat. Hist., ser. 9, XV, 1925, 59
(*X. danæ* Regan).
453. *Xenophthalmichthys danæ* Regan.
Depths of the Caribbean Sea.
Xenophthalmichthys danæ Regan, Ann. Mag. Nat. Hist., ser. 9, XV,
1925, 59, off Santa Lucia in deep water.

Suborder STOMIATIOIDEI

Family 54. STOMIATIDÆ

- Genus 156. *STOMIAS* Cuvier.
Stomias Cuvier, Règne Anim., Ed. I, 1817, 184 (*Esox boa* Risso).
454. *Stomias ferox* Reinhardt.
Greenland and southward; Gulf Stream, from Bahama Channel to the
Grand Banks.
Stomias ferox Reinhardt, Vidensk. Selsk. Naturv. Math. Afhandl., X,
1842, 77, Greenland.
455. *Stomias affinis* Günther.
Sombrero Island, West Indies.
Stomias affinis Günther, *Challenger* Rept., XXII, 1887, 205, Pl. LIV,
Fig. A; Sombrero Island, West Indies.
456. *Stomias colubrinus* Garman.
Off Panama in deep water.
Stomias colubrinus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
275, Pl. LVII, fig. 7, station 3360, lat. 6° 17' N., long. 82° 5' W. in
1,672 fathoms.
457. *Stomias hexagonatus* Garman.
Eastern tropical Pacific in deep water.
Stomias hexagonatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
Pl. LVI, fig. 5; various stations in the Pacific.
458. *Stomias atriventer* Garman.
Eastern Pacific in deep water.
Stomias atriventer Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 277,
Pl. LVI, fig. 4, *Albatross* station 3433, lat. 25° 76' 15" N., long. 109°
48' W., in 1,218 fathoms.
- Genus 157. *DACTYLOSTOMIAS* Garman.
Dactylostomias Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 279
(*D. filifer* Garman).

459. *Dactylostomias filifer* Garman.
Off Panama in deep water.
Dactylostomias filifer Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 279, Pl LVI, fig 6, *Albatross* stations 3413, 3414, and 3418
Genus 158. *ZASTOMIAS* Gilbert.
Zastomias Gilbert, Proc., U. S. Nat. Mus., XLVIII, 1915, 322 (*Z. scintillans* Gilbert).
460. *Zastomias scintillans* Gilbert.
Coast of California; Monterey Bay in deep water.
Zastomias scintillans Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 322, pl. 15, off Monterey Bay, Calif.
Genus 159. *ECHIOSTOMA* Lowe.
Echiostoma Lowe, Proc. Zool. Soc. Lond., 1843, 87 (*E. barbatum* Lowe).
Hyperchoristus Gill, Proc. U. S. Nat. Mus., VI, 1883, 256 (*H. tanneri* = *Echiostoma barbatum*)
461. *Echiostoma barbatum* Lowe.
Madeira; Gloucester, Mass.; and southward in the Gulf Stream to the old Bahama Channel.
Echiostoma barbatum Lowe, Proc. Zool. Soc. Lond., 1843, 88, Madeira.
Hyperchoristus tanneri Gill, Proc. U. S. Nat. Mus., VI, 1883, 256, Gulf Stream at *Albatross* station 2083, at 40° 26' 40" N., 60° 58' W., in 956 fathoms.
462. *Echiostoma margaritum* Goode and Bean.
Middle of the Gulf of Mexico.
Echiostoma margarita Goode and Bean, Oceanic Ichth., 1895, 109, fig. 131, Gulf of Mexico.
Genus 160. *GRAMMATOSTOMIAS* Goode and Bean.
Grammatostomias Goode and Bean, Oceanic Ichth., 1895, 110 (*G. dentatus* Goode and Bean).
463. *Grammatostomias dentatus* Goode and Bean.
Gulf Stream.
Grammatostomias dentatus Goode and Bean, Oceanic Ichth., 1895, 110, fig. 133, Gulf Stream.
Genus 161. *PHOTONECTES* Günther.
Lucifer Döderlein, Weigmann's Archiv, 1882, 26 (*L. albipinnis* Döderlein); name preoccupied in Crustacea
Photonectes Günther, *Challenger* Rept., XXII, 1887, 212 (*L. albipinnis* Döderlein).
464. *Photonectes gracilis* Goode and Bean.
Martinique.
Photonectes gracilis Goode and Bean, Oceanic Ichth., 1895, 112, fig. 137, off Martinique at *Blake* station XL in 472 fathoms.

Family 55. IDIACANTHIDÆ.

- Genus 162. *IDIACANTHUS* Peters.
Idiacanthus Peters, Monatsber. Akad. Wiss. Berlin, 1876, 846 (*I. fasciola* Peters).
Bathyopsis Gunther, Ann. Mag. Nat. Hist., ser. 5, II, 1878, 181 (*B. ferox* Gunther).
465. *Idiacanthus ferox* (Günther).
North Atlantic, nearly midway, in 2,750 fathoms.
Bathyopsis ferox Gunther, Ann. Mag. Nat. Hist., ser. 5, II, 1878, 181, middle of North Atlantic in 2,750 fathoms.
466. *Idiacanthus antrostomus* Gilbert.
Coast of southern California in deep water.
Idiacanthus antrostomus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 54, off southern California, at *Albatross* station 2980 in 603 fathoms.

Family 56. MALACOSTEIDÆ

Genus 163. MALACOSTEUS Ayres.

Malacosteus Ayres, Jour. Bost. Soc. Nat. Hist., 1849, 53 (*M. niger* Ayres).

467. *Malacosteus niger* Ayres.

Open sea; Gulf Stream and southward to Barbados.

Malacosteus niger Ayres, Jour. Bost. Soc. Nat. Hist., 1849, 53, Gulf Stream, 42° N., 60° W.

Family 57. ASTRONESTHIDÆ

Genus 164. ASTRONESTHES Richardson.

Astronesthes Richardson, Ichth. Voy. *Sulphur*, 1845, 97 (*A. nigra*, Richardson).

Phænodon Lowe, Proc. Zool. Soc. Lond., XVIII, 1850, 250 (*P. ringens* Lowe).

468. *Astronesthes richardsoni* (Poey).

Deep waters of West Indies; Indian and Pacific Oceans.

Chauliodus richardsoni Poey, Memorias, I, 1852, 176, Cuba.

469. *Astronesthes gemmifer* Goode and Bean.

Gulf Stream.

Astronesthes gemmifer Goode and Bean, Oceanic Ichth., 1895, 105, fig. 124, Grand Banks, lat. 44° 25' N., long 53° 12' W., in 300 fathoms, in halibut stomach.

470. *Astronesthes niger* Richardson.

Deep waters of all seas.

Astronesthes niger Richardson, Ichth. Voy. *Sulphur*, 1845, 97, Atlantic.

Family 58. STYLOPHTHALMIDÆ

Genus 165. STYLOPHTHALMUS Brauer.

Stylophthalmus Brauer, Zool. Anzeiger, XXV, 1902, 298 (*S. paradoxus* Brauer).

471. *Stylophthalmus paradoxus* Brauer.

Open sea, north to southern California (San Pedro, W. F. Thompson).

Stylophthalmus paradoxus Brauer, Tiefseefische *Valdivia* Exped. (Zool. Anzeiger, XXV), 1902, 298; deep sea off the Cape of Good Hope and the Indian Ocean.

Family 59. CHAULIODONTIDÆ

Genus 166. CHAULIODUS Bloch and Schneider.

Chauliodus Bloch and Schneider, Syst. Ichth., 1801, 430 (*C. sloani* Bloch and Schneider).

Leptodes Swainson, Nat. Hist. Anim., II, 298 (*Chauliodes sloani* Bloch and Schneider).

472. *Chauliodus sloani* Bloch and Schneider.

Mediterranean and deep waters of the Atlantic; Georges Banks.

Chauliodus sloani Bloch and Schneider, Syst. Ichth., 1801, 430, Atlantic; after *Vipera marina* of Catesby.

473. *Chauliodus macouni* Bean.

Coast of California to British Columbia and Queen Charlotte Islands.

Chauliodus macouni Bean, Proc. U. S. Nat. Mus., XIII, 1890, 44, off Cape St. James, Queen Charlotte Islands.

474. *Chauliodus barbatus* Garman.

Off Pacific coast of Panama.

Chauliodus barbatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 271, Pl. K, figs. 2, 2a; off Panama.

Family 60. GONOSTOMIDÆ

Genus 167. GONOSTOMA Rafinesque.

Gonostoma Rafinesque, Indice, 1810, 64 (*G. denudatum* Rafinesque).

475. *Gonostoma denudatum* Rafinesque.

Gulf of Mexico at Albatross station 2665; off Morocco and Cape Verdes; Mediterranean Sea.

Gonostoma denudatum Rafinesque, Indice, 1810, 65, Palermo.

Genus 168. ZAPHOTIAS Goode and Bean.

Bonapartia Goode and Bean, Oceanic Ichth., 1895, 102 (*B. pedaliota* Goode and Bean); name preoccupied.

Zaphotias Goode and Bean, in Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2826 (*Bonapartia pedaliota* Goode and Bean).

476. *Zaphotias pedaliotus* (Goode and Bean).

Gulf Stream.

Bonapartia pedaliota Goode and Bean, Oceanic Ichth., 1895, 102, fig. 120, Albatross station 2642, in 25° 20' 30'' N. Lat., 79° 58' W. Long., depth 217 fathoms.

Genus 169. CYCLOTHONE Goode and Bean.

Cyclothone Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 221 (*C. lusca* Goode and Bean).

Sigmops Gill, Proc. U. S. Nat. Mus., VI, 1883, 256 (*S. stigmaticus* Gill).
Neostoma Filhol, La Nature, 1884, 184 (*N. bathyphilum* Filhol).

Subgenus CYCLOTHONE Goode and Bean.

477. *Cyclothone microdon* (Günther).

Bermuda; Atlantic, Pacific, and Antarctic Oceans; Pacific coast from Oregon to Panama.

Gonostoma microdon Gunther, Ann. Mag. Nat. Hist., II, 1878, 188, near Bermuda.

Cyclothone lusca Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 221, Gulf Stream.

478. *Cyclothone bathyphila* Vaillant.

Gulf of Gascony and off the Azores; western Atlantic.

Neostoma bathyphilum Filhol, La Nature, 1884, 184; name and rough figure only.

Cyclothone bathyphila Vaillant, Expéd. Sci. Travailleur et Talisman, Poissons, 1888, 96, pl. 8, fgs. 1, 1a; Gulf of Gascony, Azores.

479. *Cyclothone acclimdens* Garman.

Off California in deep water.

Cyclothone acclimdens Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 247, Pl. T, fig. 4; numerous stations of Albatross off California coast.

480. *Cyclothone signata* Garman.

Off Panama in deep water.

Cyclothone signata Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 246, Pl. T, fig. 3; Albatross station 3382, off Panama.

Subgenus SIGMOPS Gill.

481. *Cyclothone elongata* (Günther).

New Guinea; Banda; American coast; Gulf Stream; Indian Ocean.

Gonostoma elongatum Günther, Ann. Mag. Nat. Hist., II, 1878, 187, New Guinea; Banda.

Sigmops stigmaticus Gill, Proc. U. S. Nat. Mus., VI, 1883, 256, Gulf Stream, in 38° 19' 26'' N., 68° 20' 20'' W.

Genus 170. YARRELLA Goode and Bean.

Yarrella Goode and Bean, Oceanic Ichth., 1895, 103 (*Y. blackfordi* Goode and Bean).

482. *Yarrella blackfordi* Goode and Bean.

Gulf Stream.

Yarrella blackfordi Goode and Bean, Oceanic Ichth., 1895, 103, fig. 121, Gulf Stream at Albatross station 2376, in 324 fathoms.

483. *Lychnopoles argenteolus* Garman.
Off Panama in deep water.
Lychnopoles argenteolus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 244, Pl. LIII, figs. 4, 4a; off Panama.

Family 61. MAUROLICIDÆ

Genus 172. MAUROLICUS Cocco.

Maurolicus Cocco, Nuovi Ann. Sci. Nat., II, 1838, 32 (*M. amethystino punctatus* Cocco).

484. *Maurolicus pennanti* (Walbaum). *Pearlsides*.
Open seas, widely distributed; New England coast at Nahant, Provincetown, Woods Hole, etc.
Argentina pennanti Walbaum, Artedi Pisc., 1792, 47, England; after "sheppy Argentine" of Pennant.

485. *Maurolicus oculatus* Garman.
Deep water off Pacific coast of Mexico.
Maurolicus oculatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 241, Pl. LIII, fig. 3; off Pacific coast of Mexico.

Genus 173. VINCIGUERRIA Jordan and Evermann.

Vinciguerria Jordan and Evermann, in Goode and Bean, Oceanic Ichth., 1895, 513 (*V. attenuata* Cocco).
Zalarges Jordan and Williams, in Jordan and Starks, Proc. Calif. Acad. Sci., ser. 2, V, 1895, 793 (*Z. nimbarius* Jordan and Williams).

486. *Vinciguerria attenuata* (Cocco).
Open Atlantic, west to the Bahama regions.
Maurolicus attenuatus Cocco, Nuovi Ann. Sci. Nat., II, 1838, 33, coast of Italy.

487. *Vinciguerria lucetia* (Garman).
Eastern Pacific, in deep water.
Maurolicus lucetius Garman, Deep Sea Fishes, in Mem. Mus. Comp. Zool., XXIV, 1899, 242, Pl. T, fig. 2; various Albatross stations, eastern Pacific.

488. *Vinciguerria nimbaria* (Jordan and Williams).
Open sea between Oregon and Hawaii.
Zalarges nimbarius Jordan and Williams, Proc. Calif. Acad. Sci., ser. 2, V, 1895 (1896), 793, open Pacific.

Genus 174. VALENCIENNELLUS Jordan and Evermann.

Valenciennellus Jordan and Evermann, Fishes North and Mid. Amer., 1896, 577 (*Maurolicus tripunctulatus* Esmark).

489. *Valenciennellus tripunctulatus* (Lutken).
Madagascar; Denmark Straits between Greenland and Iceland.
Maurolicus tripunctulatus Esmark, Christiania Vid. Selsk. Forh., 1870, 488, Madagascar.

490. *Valenciennellus stellatus* Garman.
Deep water off California.
Valenciennellus stellatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 239, Pl. LIII, fig. 2; off California coast, lat. 30° 31' 35" N., long. 140° 5' 30" W. in 300 fathoms.

Family 62. STERNOPTICHIDÆ

Genus 175. STERNOPTIX Herrmann.

Sternoptix Herrmann, Naturforscher, 2 Stück, XVI, 1781, 8 (*S. diaphana* Herrmann).
Sternoptyx, authors' amended spelling.

491. *Sternoptix obscura* (Garman).
Eastern Pacific in deep water.
Sternoptyx obscura Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 232, Pl. LIII, fig. 1; numerous stations in eastern Pacific in depths ranging from 421 to 1,823 fathoms.

492. *Sternoptix diaphana* Herrmann.
Atlantic; Gulf Stream, from Santa Cruz Island to the Grand Banks.
Sternoptix diaphana Herrmann, Naturforscher, XVI, 1781, 8, Jamaica.
- Genus 176. ARGYROPELECUS Cocco.
Argyrolepecus Cocco, Arch. del R. Acad. Peloritano, 1829, 146 (*A. hemigymnus* Cocco).
Pleurothyris Lowe, Hist. Fish. Madeira, 1843, 64 (*Sternoptyx olfersi* Cuvier).
493. *Argyrolepecus hemigymnus* Cocco.
Atlantic and Mediterranean; Gulf Stream off Southern New England.
Argyrolepecus hemigymnus Cocco, Arch. del R. Acad. Peloritano, 1829, 146, coast of Italy.
Sternoptyx mediterraneus Cocco, Giorn. Faro de Messina, IV, 1838, 7, coast of Italy.
494. *Argyrolepecus olfersi* (Cuvier).
Open Atlantic; coast of Norway to Brazil and Cape of Good Hope; Grand Banks southward in the Gulf Stream.
Sternoptyx olfersi Cuvier, Règne Anim., ed. 2, II, 1829, 316, near Cape of Good Hope.
Argyrolepecus durvillii Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849, 405, open Atlantic.
495. *Argyrolepecus lychnus* Garman.
Western Pacific in deep water.
Argyrolepecus lychnus Garman, Deep Sea Fishes, in Mem. Mus. Comp. Zool., XXIV, 1899, 234, Pl. T, fig. 1, 16; many stations in eastern Pacific.
496. *Argyrolepecus affinis* Garman.
Tropical Atlantic in deep water.
Argyrolepecus affinis Garman, Deep Sea Fishes, in Mem., Mus. Comp. Zool., XXIV, 1899, 237; sta. 2117, N. lat. 15° 24' 40'', W. long. 63° 31' 30'', in 683 fathoms.
- Genus 177. POLYIPNUS Günther.
Polyipnus Gunther, Challenger Rept., 1887, 170 (*P. spinosus* Günther).
497. *Polyipnus laternatus* Garman.
Off Barbados.
Polyipnus laternatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 238; Blake station 280, off Barbados in 221 fathoms.
498. *Polyipnus spinosus* Günther.
Gulf of Mexico; between the Philippines and Borneo.
Polyipnus spinosus Günther, Challenger Rept., XXII, 1887, 170, from the surface near the Philippine Islands.

Order O. LYOPOMI

Family 63. HALOSAURIDÆ

- Genus 178. HALOSAURUS Johnson.
Halosaurus Johnson, Ann. Mag. Nat. Hist., XIV, 1863, 74 (*H. oweni* (oweni) Johnson).
Halosaurichthys Alcock, loc. cit., 1889, 454, (*H. carinicauda* Alcock).
499. *Halosaurus güntheri* Goode and Bean.
Gulf Stream.
Halosaurus güntheri Goode and Bean, Oceanic Ichth., 1895, 131, Gulf Stream, in lat. 39° 13' N., long. 72° 1' W., Albatross station 2722.
500. *Halosaurus radiatus* Garman.
Off Panama in deep water.
Halosaurus radiatus Garman, Deep Sea Fishes, in Mem. Mus. Comp. Zool., XXIV, 1899, 298, Pl. LX, figs. 2, 2a, Pl. LXXXIV, figs. 3-6; several stations off Panama in depths from 259 to 511 fathoms.

501. *Halosaurus attenuatus* Garman.
Off Panama in deep water.
Halosaurus attenuatus Garman, Deep Sea Fishes, in Mem. Mus. Comp. Zool., XXIV, 1899, 296, Pl. LX, figs 1, 1a, Albatross station 3413, lat. 2° 34' N., long 92° 06' W., in 1,360 fathoms.
- Genus 179. *ALDROVANDIA* Goode and Bean.
Aldrovandia Goode and Bean, Oceanic Ichth., 1895, 132 (*Halosaurus rostrata* Günther).
Halosauropsis Collett, Poiss. Prov. Comp. l'*Hirondelle*, Fasc. *X., Monaco, 1896, 143 (*Halosaurus macrochir*).
502. *Aldrovandia rostrata* (Günther).
Mid-Atlantic.
Halosaurus rostratus Günther, Ann. Mag. Nat. Hist., II, 1878, 252, Mid-Atlantic.
503. *Aldrovandia macrochir* (Günther).
Marion Island; Blake stations 308 and 325.
Halosaurus macrochir Günther, Ann. Mag. Nat. Hist., II, 1878, 251, between Cape of Good Hope and Kerguelen Island.
504. *Aldrovandia goodei* (Gill).
Gulf Stream, at Albatross stations 2037 in 1,731 fathoms; 2051 in 1,106 fathoms; 2035 in 1,362 fathoms, and 2052 in 1,098 fathoms.
Halosaurus goodei Gill, Proc. U. S. Nat. Mus., VI, 1883, 257, Gulf Stream off South Carolina.
505. *Aldrovandia gracilis* Goode and Bean.
Gulf Stream and Gulf of Mexico
Aldrovandia gracilis Goode and Bean, Oceanic Ichth., 1895, 134, fig. 157, off Guadeloupe and in Gulf Stream at Albatross stations 2380 and 2381.
506. *Aldrovandia pallida* Goode and Bean.
Gulf of Mexico, in deep water.
Aldrovandia pallida Goode and Bean, Oceanic Ichth., 1895, 135, fig. 158, Gulf of Mexico, at Albatross station 173.

Order P. HETEROMI. Spiny eels

Family 64. LIPOGENYIDÆ

- Genus 180. *LIPOGENYS* Goode and Bean.
Lipogenys Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894 (1895), 469 (*L. gillii* Goode and Bean).
507. *Lipogenys gillii* Goode and Bean
Gulf Stream.
Lipogenys gillii Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894 (1895), 469, pl. 18, fig. 3, and in Oceanic Ichth., 173, 1895, station 2742, Gulf Stream.

Family 65. NOTACANTHIDÆ

- Genus 181. *NOTACANTHUS* Bloch.
Notacanthus Bloch, Abhandl., Bohmischen Gesellsch. Wiss., I, 1787, 278 (*N. chemnitzii* Bloch).
Acanthonotus Bloch, Ichthyologia, XII, 1797, 113 (*A. nasus* Bloch = *N. chemnitzii* Bloch).
Campylodon (Fabricius, 1798) Reinhardt, Vidensk. Selsk. Naturv. Math. Afhandl., 1838, 120 (*C. fabricii* Reinhardt).
508. *Notacanthus chemnitzii* Bloch.
Greenland; West Indies; Iceland; Scandinavia; South Greenland.
Notacanthus chemnitzii Bloch, Abhandl., Bohmischen Gesellsch. Wiss., I, 1787, 278, northern seas.
Notacanthus nasus Bloch, Ichthyologia, XII, 1795, 113, pl. 431, "India."
Campylodon fabricii Reinhardt, Vidensk. Selsk. Naturv. Math. Afhandl., 1838, 120, Greenland.

509. *Notacanthus analis* Gill.
Gulf Stream.
Notacanthus analis Gill, Proc. U. S. Nat. Mus., VI, 1883, 255, lat. 40° N., long. 69° W., Albatross stations 2677 in 478 fathoms and 2676 in 407 fathoms.
510. *Notacanthus phasganorus* Goode.
Grand Banks, Newfoundland.
Notacanthus phasganorus Goode, Proc. U. S. Nat. Mus., III, 1880, 535, Grand Banks of Newfoundland.
511. *Notacanthus spinosus* Garman.
Off Panama.
Notacanthus spinosus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 301, Pl. L, figs. 4, 4a, 4b; Albatross stations 3384 and 3354, 322 and 458 fathoms.
- Genus 182. *POLYACANTHONOTUS* Bleeker.
Polyacanthonotus Bleeker, Versl. Akad. Amsterdam (2), VIII, 368 (*Notacanthus rissoanus* Filippi and Verany).
Zanotacanthus Gill, Johnson's Cyclopædia, III, 1876, 883 (*Notacanthus rissoanus* Filippi and Verany).
Macdonaldis Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894 (1895), 467 (*Notacanthus rostratus* Collett).
512. *Polyacanthonotus challenger* (Vaillant).
North Pacific and Bering Sea.
Notacanthus rissoanus Gunther, Fishes, Challenger, XXII, Part LVII, Pl. LXI, fig. 13 (not of Filippi and Verany).
Notacanthus challenger Vaillant, Expéd. Sci. Travailleur et Talisman, 1888, south of Yeddo; based on Gunther's description.
513. *Polyacanthonotus rostratus* (Collett).
Gulf Stream.
Notacanthus rostratus Collett, Bull. Soc. Zool. France, 1889, 307, off Newfoundland, at Albatross station 2216 in 963 fathoms, and station 2553 in 551 fathoms.

Order Q. SYMBRANCHIA. Symbranchoid eels

Suborder HOLOSTOMI

Family 66. SYNBRANCHIDÆ

Genus 183. SYNBRANCHUS Bloch.

- Synbranchus* Bloch, Ichthyologia, IX, 1795, 87 (*Synbranchus marmoratus* Bloch).
Umbbranchaperta Lacépède, Hist. Nat. Poiss., V, 1803, 658 (*Synbranchus marmoratus*).
Ophisternon McClelland, Calcutta Jour. Nat. Hist., V, 1845, 197 (*O. bengalensis*).
Tetrabanchus Bleeker, Nat. Tijds. Ned. Ind., II, 1862, 69 (*T. micropthalmus* McClelland).
Synbranchus of authors generally; corrected spelling.

514. *Synbranchus marmoratus* Bloch.

- Tropical America, in streams from the Amazon northward to southern Mexico and St. Lucia; generally abundant; recorded northward from Vera Cruz, Trinidad, St. Lucia, Lake Peton, Huamuchal, Rio Chisoy, etc.
Synbranchus marmoratus Bloch, Ichthyologia, IX, 1795, 87, pl. 418, tropical America.
Synbranchus immaculatus Bloch, loc. cit., pl. 419, fig. 1.
Synbranchus transversalis Bloch and Schneider, loc. cit., 1801, 524, Guinea; after Gronow.
Umbbranchaperta grisea, *marmorata*, *immaculata*, and *lineata* Lacépède, Hist. Nat. Poiss., V, 1803, 658, Surinam.
Synbranchus fuliginosus Ranzani, Nov. Com. Acad. Sci. Bonon, IV, 1840, 75, pl. 11, fig. 1, Brazil.
Muraena humbricus Gronow, Cat. Fish., 1854, 18, Sea of Guinea.
Symbranchus vittatus Castelnau, Anim. Amer. Sud. Poiss., 84. 1855.

Order R. CARENCELI. Long-necked eels

Family 67. DERICHTHYIDÆ

Genus 184. DERICHTHYS Gill.

Derichthys Gill, Amer. Nat., XVIII, 1884, 433 (*D. serpentinus* Gill).

515. *Derichthys serpentinus* Gill.

Gulf Stream.

Derichthys serpentinus Gill, Am. Nat., XVIII, 1884, 433, Gulf Stream at Albatross station 2094 in 1,022 fathoms.

Genus 185. GORGASIA Meek and Hildebrand.

Gorgasia Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 133 (*G. punctata* Meek and Hildebrand).

516. *Gorgasia punctata* Meek and Hildebrand.

Panama.

Gorgasia punctata Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 133, Chame Point, Panama.

Order S. APODES

Suborder ENCHELYCEPHALI

Series ANGUILLOIDEI

Family 68. ANGUILLIDÆ. True eels

Genus 186. ANGUILLA Shaw. Eels.

Anguilla Shaw, Gen. Zool., IV, 1803, 15 (*A. anguilla*).

Muraena Bleeker, Poey, etc. (taking as type *Muraena anguilla*, the first species mentioned by Artedi under *Muraena*).

Tribranchnus (Peters) Muller, Abhandl., Akad. Wiss. Berlin, 1844 (*T. anguillaris* Peters = *Anguilla australis* Rich).

517. *Anguilla rostrata* (LeSueur). American eel; Fresh-water eel.

Atlantic coast of the United States; very abundant from Maine to Mexico, ascending all rivers south of Canada and east of the Rocky Mountains, individuals sometimes resident throughout the Mississippi Valley; common in the West Indies.

Muraena rostrata LeSueur, Jour. Acad. Nat. Sci. Phila., 1817, 81, Cayuga Lake.

Muraena bostonensis LeSueur, loc. cit., 81, Boston.

Muraena serpentina LeSueur, loc. cit., 82, Newport, R. I.

Muraena macrocephala LeSueur, loc. cit., 82, Saratoga, N. Y.

Muraena argentea LeSueur, loc. cit., 82, Boston Bay.

Anguilla chrysypa Rafinesque, Amer. Month. Mag. and Crit. Rev., 1817, 120, Lake George; Hudson River; Lake Champlain.

Anguilla blephura Rafinesque, loc. cit., 120, Long Island.

Anguilla laticauda Rafinesque, loc. cit., 445, Ohio River.

Anguilla aterrima Rafinesque, Ichth. Ohiensis, 1820, 78, Tennessee and Cumberland Rivers.

Anguilla xanthomelas Rafinesque, loc. cit., 78, Ohio River.

Anguilla lutea Rafinesque, loc. cit., 78, Ohio River.

Anguilla tenuirostris DeKay, N. Y. Fauna Fishes, 1842, 310, New York.

Anguilla novæorleansis Kaup, Apodal Fishes, 1856, 43, fig. 33, New Orleans, La.

Anguilla punctatissima Kaup, loc. cit., 44, Niagara River.

Anguilla cubana Kaup, loc. cit., 44, Cuba.

Anguilla novæterræ Kaup, loc. cit., 45, fig. 35, Newfoundland.

Anguilla texana Kaup, loc. cit., 45, fig. 36, Texas.

Anguilla wabashensis Kaup, loc. cit., 46, Wabash River.

Anguilla tyrannus Girard, U. S. Mex. Bound. Surv., Ichth., 1859, 75, Rio Grande.

Family 69. SIMENCHELYIDÆ. Snub-nosed eels

Genus 187. SIMENCHELYS Gill.

Simenchelys Gill, in Goode and Bean, Bull. Essex Inst., 1879, 27 (*S. parasiticus* Gill).

Conchognathus Collett, Bull. Soc. Zool. France, 1889, 123 (*C. grimaldii* Collett).

518. *Simenchelys parasiticus* Gill.

Offshore banks, in deep water, south of Newfoundland; also recorded from the Azores.

Simenchelys parasiticus Gill, in Goode and Bean, Bull. Essex Inst., 1879, 27, Newfoundland banks.

Conchognathus grimaldii Collett, Bull. Soc. Zool. France, 1889, 122, Azores.

Family 70. SYNAPHOBANCHIDÆ

Genus 188. SYNAPHOBANCHUS Johnson.

Synphobanchus Johnson, Proc. Zool. Soc. Lond., 1862, 169 (*S. kaupii* Johnson = *Muraena pinnata* Gronow).

519. *Synphobanchus pinnatus* (Gronow).

North Atlantic and western Pacific; common about the Madeiras, Canaries, etc., and also about the banks of Newfoundland.

Muraena pinnata Gronow, Cat. Fish., 1854, 19, locality unknown.

Synphobanchus kaupii Johnson, Proc. Zool. Soc. Lond., 1862, 169, Madeira.

Genus 189. HISTIOBRANCHUS Gill.

Histiobranchus Gill, Proc. U. S. Nat. Mus., VI, 1883, 255 (*H. infernalis* Gill).

520. *Histiobranchus bathybius* (Günther).

Northern and western Pacific in deep water; Bering Strait; off Japan and off Cape of Good Hope; one specimen obtained by Doctor Gilbert in Bering Sea in 1890.

Synphobanchus bathybius Günther, Ann. Mag. Nat. Hist., XX, 1877, 445, and in Voy. Challenger, 254, Pl. LXII, fig. 6, 1887, off Yeddo; North Pacific; Kerguelen Island.

521. *Histiobranchus infernalis* Gill.

Gulf Stream.

Histiobranchus infernalis Gill, Proc. U. S. Nat. Mus., VI, 1883, 255, Gulf Stream, at Albatross station 2037, 38° 30' N., 69° W., in 1,731 fathoms.

Family 71. ILYOPHIDÆ. Ooze eels

Genus 190. ILYOPHIS Gilbert.

Ilyophis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 351 (*I. brunneus* Gilbert).

522. *Ilyophis brunneus* Gilbert.

One specimen from Chatham Island, Galapagos Archipelago, in 634 fathoms.

Ilyophis brunneus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 352, Chatham Island, Galapagos Archipelago, at Albatross station 2808 in 654 fathoms.

Family 72. MURÆNESOCIDÆ

Genus 191. MURÆNESOX McClelland.

Muraenox McClelland, Calcutta Jour. Nat. Hist., V, 1844, 408 (*M. tricuspidata* McClelland).

Branchyconger Bleeker, Nederl. Tijdschr. Dierk., II, 1865, 236 (*Muraena savanna* Cuvier).

523. *Muraenox savanna* (Cuvier).

Cuba to Rio Janeiro, not common; occasional in the Mediterranean Sea.

Muraena savanna Cuvier, Règne Anim., ed. 2, II, 1829, 350, Martinique.

Conger brasiliensis Ranzani, De Novis Spec. Pisc. Diss. Prima., IV, 1838, 17, pl. 13, fig. 1, Brazil.

Congrus curvidens Richardson, Voy. Erebus and Terror, 1844, 111, no locality.

Conger limbatus Castelnau, Anim. Amér. Sud, 1855, 83, pl. 43, fig. 3, Rio Janeiro.

524. *Muraenesox coniceps* Jordan and Gilbert.
Cape San Lucas to the coast of Colombia; generally common.
Muraenesox coniceps Jordan and Gilbert, Proc. U. S. Nat. Mus., IV,
1881, 348, Mazatlan, Mexico.

Family 73. CONGRIDÆ (LEPTOCEPHALIDÆ). Conger eels

Genus 192. CONGER Cuvier. Conger eels.

- a. Adult forms (Conger):
Conger Cuvier, Oken, in Règne Anim., Ed. I, 1817, 231 ("Les Congres"
Cuvier, Oken, 1182) (*Muraena conger* Linnæus).
Congrus Richardson, Voy. *Erebus* and *Terror*, 1844, 107 (*Muraena conger* Linnæus).
b. Larval forms (*Leptocephali*):
Leptocephalus Scopoli, Int. Hist. Nat., 1777, 453 (*L. morrissi* Gmelin).
Oxyurus Rafinesque, Caratteri, 1810, 19 (*O. vermiformis* Lacépède).
Helmictus Rafinesque, Indice, 1810, 62.
Helmichthys Costa, Fauna Napoli, Pesci, tavoli 31, 1854 (*H. diaphanus* Costa).
? *Leptocephalichthys* Bleeker, Act. Soc. Sci. Indo-Neder., I, 69 (*L. hypselosoma* Bleeker, a larval form allied to *Conger*).
? *Diaphanichthys* Peters, Sitzungsber., Akad. Wiss. Berlin, 1864, 399 (*D. brevicaudus* Peters); larva allied to *Conger*.

525. *Conger conger* (Linnæus). Conger eel.
Atlantic Ocean; generally common on both coasts, from Cape Cod to Brazil, also on coasts of Asia and Africa.

- a. Adult forms:
Muraena supremo marginæ pinnæ dorsalis nigro Artedi, Synon., 40, 1738, 2, Mediterranean.
Muraena conger Linnæus, Syst. Nat., ed. X, 1758, 245; based on Artedi.
Muraena nigra Risso, Ichth. Nice, 1810, 93 (black variety), Nice.
Anguilla oceanica Mitchell, Jour. Acad. Nat. Sci. Phila., 1818, 407, off New York.
Conger verus Risso, Europ. Mérid., III, 1826, 201, Nice.
Conger vulgaris Cuvier, Règne Anim., ed. 2, II, 1829, 350, France.
Conger rubescens Ranzani, De Novis Spec. Pisc. Diss. Prima, 1838, 19, Pl. V, fig. 5, Mediterranean.
Ophisoma obtusa Swainson, Nat. Hist. Fish., Reptiles, and Amphib., II, 1839, 395, Sicily.
Conger orbignyanus Valenciennes, D'Orbigny, Voy. Am. Merid., Poiss., 1839, pl. 12, 1, South America.
Conger verreauxi Kaup, Apodal Fishes, 1856, 115, no locality.
Conger esculentus Poey, Memorias, II, 1861, 346, Cuba.

- b. Larval forms:
Leptocephalus morrissi Gmelin, Syst. Nat., 1150, 1877, Holyhead, England.
Ophidium pellucidum Couch, Lond. Mag. Nat. Hist., V, 1832, 3113, 742, England.
Leptocephalus gracilis Storer, Mem. Amer. Acad., II, 1839, 524, Massachusetts.
Leptocephalus spallanzani, *candidissimus*, etc., of European writers.

526. *Conger cardilimbatus* (Poey).
Tropical parts of Atlantic; Pensacola to Cuba and Madeira.
Echelus caudilimbatus Poey, Repertorio, II, 1867, 249, Cuba.
Conger macrops Günther, Cat., VIII, 1870, 40, Madeira, Bahamas.
Conger caudicula Bean, Proc. U. S. Nat. Mus., V, 1882, 435, Pensacola.

Genus 193. ARIOSOMA Swainson.

- Ariosoma* Swainson, Nat. Hist. Fish., I, 1838, 220; II, 1839, 195 (no type named, logotype as restricted by Bleeker and by Swainson).
Ophisoma acuta Swainson = *Muraena balearica* De la Roche).
Ophisoma Swainson, loc. cit., 1839; *Ophisoma acuta* Swainson; substitute for *Ariosoma*; the name preoccupied.
Congermuraena Kaup, Apodal Fishes, 1856, 71 (108) (*Muraena balearica* De la Roche, as restricted by Bleeker).
Congrellus Ogilby, Proc. Linn. Soc. New South Wales, XXIII, 1898, 288 (*Muraena balearica* De la Roche).

527. *Ariosoma opisthophthalma* (Ranzani).
West Indies to Brazil.
Conger opisthophthalmus Ranzani, De Novis Spec. Pisc. Diss. Prima, 1838, 16, Pl. V, fig. 1, Bahia.
Conger microstomus Castelnau, Anim. Amér. Sud, 1855, 83, pl. 42, fig. 4, Rio Janeiro.
Conger analis Poey, Memorias, II, 1861, 318, Habana.
Conger impressus Poey, loc. cit., 318, Cuba.
528. *Ariosoma gilberti* (Ogilby).
Panama; probably also at the Galapagos and Cape San Lucas.
Ophisoma balearicum Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 349, Albatross station 2797, Panama Bay, 33 fathoms; not of De la Roche.
Congrellus gilberti Ogilby, Proc. Linn. Soc. New South Wales, 1898, 288, after Gilbert.
529. *Ariosoma caudale* (Garman).
Off Panama in deep water.
Congermuræna caudalis Garman, Mus. Comp. Zool., XXIV, 1899, 305; Albatross station 3354, lat. 7° 9' 45'' N., long. 80° 50' W., in 322 fathoms.
- Genus 194. ANAGO²⁴ Jordan and Hubbs.
Anago Jordan and Hubbs, Mem. Carnegie Mus., X, 1925, 191 (*Conger anago* Schlegel).
530. *Anago nitens* (Jordan and Bollman).
One specimen dredged at Albatross station 2801, off Panama, 8° 47' N., 79° 29' 30'' W., in 14 fathoms.
Ophisoma nitens Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 153, Panama.
531. *Anago macrurus* (Gilbert).
Gulf of California.
Ophisoma macrurum Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 351, Gulf of California, at Albatross station 3105.
- Genus 195. HILDEBRANDIA Jordan and Evermann.
Hildebrandia Jordan and Evermann, Proc. Calif. Acad. Sci., 4th Ser., XVI, Apr. 27, 1927, 502 (*Congermuræna flava* Goode and Bean).
532. *Hildebrandia flava* (Goode and Bean).
Albatross stations 2121 and 2122 in 31 and 34 fathoms; station 2402 in 111 fathoms; also Blake station 264 in 84 fathoms; Gulf Stream.
Congermuræna flava Goode and Bean, Oceanic Ichth., 1895, 138, fig. 159, Gulf Stream.
- Genus 196. GNATHOPHIS Kaup.
Gnathophis Kaup, Abhandl., Natur. Verein Hamburg, 1859 (1860), IV, Abth. 2, 7 (*G. heterognathus* Kaup); teeth blunt.
533. *Gnathophis guppii* (Norman).
West Indies.
Congermuræna guppii Norman, Ann. Mag. Nat. Hist., ser. 9, XV, 1925, 313, Tobago.
534. *Gnathophis proriger* (Gilbert).
Panama to Ecuador.
Ophisoma prorigerum Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 350, coast of Ecuador, at Albatross station 2792 in 401 fathoms; also at station 2799, near Panama.
- Genus 197. CONGROSOMA Garman.
Congrosoma Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 308 (*Congrosoma evermanni* Garman).
535. *Congrosoma evermanni* Garman.
Off Panama.
Congrosoma evermanni Garman, Mem. Mus. Comp. Zool., XXIV, 1899, Pl. LXII, fig. 1, Albatross station 3355, lat. 7° 12' 20'' N., long. 80° 55' W. in 182 fathoms.

²⁴ This shore species named below may not, any of them, belong to Anago or, in fact, to the same genus. They seem nearer the Japanese form *Anago* than to *Ariosoma*.

Genus 198. UROCONGER Kaup.*Uroconger* Kaup, Apodal Fishes, 1856, 110 (*U. lepturus* Kaup).**536. *Uroconger vicinus* Vaillant.**Deep waters off the coast of northern Africa; a young individual taken by the *Albatross* at station 2161 in 146 fathoms.*Uroconger vicinus* Vaillant, Expéd. Sci. Travailleur et Talsman, 1883, 86, Pl. VI, fig. 1, Banc d'Arguin, off Soudan, off Cape Verde Islands.**537. *Uroconger varidens* Garman.**

Off Panama in deep water.

Uroconger varidens Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 304, Pl. LXI, fig. 1, *Albatross* stations 3396 and 3357 in 259 and 782 fathoms.**Genus 199. MICROCONGER Fowler.***Microconger* Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 9 (*Leptocephalus caudalis* Fowler)**538. *Microconger caudalis* (Fowler).**

Coast of Lower California.

Leptocephalus (Microconger) caudalis Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 9; off Lower California.**Genus 200. MAYERINA Silvester.***Mayerina* Silvester, Carnegie Inst. Year Book, XIV, 1915, 214 (*Mayerina mayeri* Silvester).**539. *Mayerina mayeri* Silvester.**

Porto Rico.

Mayerina mayeri Silvester, Carnegie Inst. Year Book, XIV, 1915, 214, Porto Rico.**Genus 201. XENOMYSTAX Gilbert.***Xenomystax* Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 348 (*X. atrarius* Gilbert).**540. *Xenomystax atrarius* Gilbert.**

Coast of Ecuador.

Xenomystax atrarius Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 348, 1° S., 81° W., at *Albatross* station 2792 in 401 fathoms.**541. *Xenomystax rictus* Garman.**

Off Panama.

Xenomystax rictus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 315, Pl. N, several *Albatross* stations (3384, 3394, 3354, 3404, 3417) in from 322 to 511 fathoms.**Genus 202. HOPLUNNIS Kaup.***Hoplunnus* Kaup, Aalen. Fische Hamburg Mus., 1860, 19 (*H. schmidtii* Kaup).**542. *Hoplunnus schmidtii* Kaup.**

Caribbean Sea at Puerto Cabello, near Aspinwall; Colon.

Hoplunnus schmidtii Kaup, Aalen. Fische Hamburg Mus., 1859, 19, pl. 2, fig. 4, Puerto Cabello, near Aspinwall, Colon.**543. *Hoplunnus diomedianus* Goode and Bean.**A single individual obtained by the *Albatross* in the Gulf of Mexico*Hoplunnus diomedianus* Goode and Bean, Oceanic Ichth., 146, 1895, fig. 163, Gulf of Mexico, *Albatross* station 2402, 28° 36' N., 86° 50' W., 111 fathoms.**Genus 203. NEOCONGER Girard.***Neoconger* Girard, Proc. Acad. Nat. Sci. Phila., X, 1858 (1859), 171 (*N. mucronatus* Girard).**544. *Neoconger mucronatus* Girard.**

Coast of Texas.

Neoconger mucronatus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858 (1859), 171, St. Joseph Island, Texas.**545. *Neoconger vermiformis* Gilbert.**

Lower California to Panama

Neoconger vermiformis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 57, off Lower California, at *Albatross* station 3035 in about 30 fathoms.

Genus 204. **LEPTOCONGER** Poey.

Leptoconger Poey, Anal. Soc. Española Hist. Nat., Madrid, 1880, 250
(*L. perlongus* Poey).

546. **Leptoconger perlongus** (Poey).

Matanzas, Cuba.

Neoconger perlongus Poey, Ann. Lyc. Nat. Hist. N. Y., XI, 1874, 67,
pl. 9, figs. 3 and 4, Matanzas, Cuba.

Family 74. **NETTASTOMIDÆ**. Duckbill eels; SorcerersGenus 205. **CHLOPSIS** Rafinesque.

Chlopsis Rafinesque, Indice, 1810, 58 (*C. bicolor* Rafinesque).

Saurenehelys Peters, Sitzungsber. Akad. Wiss. Berlin, 1864, 397 (*S. can-*
crowa Peters)

547. **Chlopsis equatorialis** Gilbert.

Coast of Ecuador.

Chlopsis equatorialis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 347,
off coast of Ecuador, 1° S., 81° W., at Albatross station 2792 in 401
fathoms.

548. **Chlopsis gilberti** Garman.

Off Panama and San Diego.

Chlopsis gilberti Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
Pl. LXII, fig. 2, various stations of the Albatross, from 322 to 511
fathoms.

Genus 206. **VENEFICA** Jordan and Davis.

Venefica Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 651 (*Net-*
tastoma procerum Goode and Bean).

549. **Venefica procera** (Goode and Bean).

Gulf Stream; also taken off San Pedro, Calif.

Nettastoma procerum Goode and Bean, Bull. Mus. Comp. Zool., X,
1883, 224, Gulf Stream, at Albatross station 325.

550. **Venefica ocella** Garman.

Off Panama in deep water.

Venefica ocella Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 318,
Pl. LXI, fig. 2, Albatross station 3366, lat. 5° 30' N., long. 86° 45' W.,
in 1,067 fathoms.

551. **Venefica tentaculata** Garman.

Off Panama in deep water; San Diego.

Venefica tentaculata Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
319, Pl. M, fig. 2, various Albatross stations at depths ranging from
770 to 978 fathoms.

Genus 207. **TÆNIOCONGER** Herre.

Tænioconger Herre, Philippine Jour. Sci., XXIII, No. 2, August, 1923,
152 (*T. chapmani* Herre).

552. **Tænioconger digueti** Pellegrin.

Gulf of California.

Tænioconger digueti Pellegrin, Compt. Rend. Acad. Sci. Paris, CLXXVII,
1923, 789, near Island of Espiritu Santo, Gulf of California.

Family 75. **ECHELIDÆ**. Worm eelsGenus 208. **AHLIA** Jordan and Davis.

Ahlia Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 639 (*Myro-*
phis egmontis Jordan).

553. **Ahlia egmontis** (Jordan).

Florida Keys.

Myrophis egmontis Jordan, Proc. Acad. Nat. Sci. Phila., XXXVI, 1884,
44, Egmont Key, Fla.

Genus 209. **MYROPHIS** Lütken.

Myrophis Lütken, Vid. Med. Naturh. Foren. Kjøbenhavn, 1, 1851
(*M. punctatus* Lütken).

554. *Myrophis punctatus* Lütken.

West Indies; coast of Texas to Surinam; common along our Gulf coast.
Myrophis punctatus Lütken, Vid. Med. Naturh. Foren. Kjøbenhavn, 1, 1851, 9, West Indies.

Myrophis microstigmus Poey, Repertorio, II, 1867, 250, Cuba.

Myrophis lumbricus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 261, Galveston, Tex.

555. *Myrophis vafer* Jordan and Gilbert.

Pacific coast of tropical America, from Guaymas to Panama.

Myrophis vafer Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 645, Panama.

Genus 210. *CHILORHINUS* Lütken.

Chilorhinus Lütken, Vid. Med. Naturh. Foren. Kjøbenhavn, 1, 1851 (C. *suensonii* Lütken).

556. *Chilorhinus suensonii* Lütken.

St. Croix, West Indies.

Chilorhinus suensonii Lütken, Vid. Med. Naturh. Foren. Kjøbenhavn, 1, 1851, St. Croix.

Family 76. NEMICHTHYIDÆ. Thread eels; Snipe eels**Genus 211. *NEMICHTHYS* Richardson.**

Nemichthys Richardson, Voy. *Samarang*, 1848, 25 (*N. scolopaceus* Richardson).

Leptorhynchus Lowe, Ann. Mag. Nat. Hist., X, 1852, 54 (*L. leuchtenbergi* Lowe); preoccupied.

Belonopsis Brandt, Mem. Acad. Sci. St. Petersb., Savans Etrangères, 1854, 174 (*Leptorhynchus leuchtenbergi* Lowe).

557. *Nemichthys scolopaceus* Richardson. *Snipe eel*.

Atlantic Ocean, in deep water; many specimens taken off the New England coast and off the Grand Banks; common about Madeira.

Nemichthys scolopacea Richardson, Voy. *Samarang*, 1848, 25, South Atlantic.

Leptorhynchus leuchtenbergii Lowe, Mem. Acad. Sci. St. Petersb., Savans Etrangères 1854, 171, Madeira.

Belonopsis leuchtenbergii Brandt, loc. cit., 1854, 174 (with good plate).

558. *Nemichthys avocetta* Jordan and Gilbert.

Puget Sound near Seattle; only the type known.

Nemichthys avocetta Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 409, harbor of Port Gamble, near Seattle.

559. *Nemichthys fronto* Garman

Off Pacific coast of Mexico, in deep water.

Nemichthys fronto Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 324, Pl. LXV, fig. 1, *Albatross* stations 3384 and 3434, in 458 and 1,588 fathoms.

Genus 212. *AVOCETTINA* Jordan and Davis.

Avocettina Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 655 (*Nemichthys infans* Gunther).

560. *Avocettina infans* (Gunther).

Known from West Indies, and middle Atlantic off Pernambuco.

Nemichthys infans Gunther, Ann. Mag. Nat. Hist., 5th Ser., II, 1878, 251, and in *Challenger* Rept., XXII, 1887, 264, mid. Atlantic in 2,500 fathoms.

561. *Avocettina gilli* (Bean).

Coast of southeastern Alaska in deep water.

Labichthys gilli Bean, Proc. U. S. Nat. Mus., XIII, 1890, 45, east of Prince of Wales Island, Alaska.

562. *Avocettina elongata* (Gill and Ryder).

Gulf Stream.

Labichthys elongatus Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 262, Gulf Stream at *Albatross* station 2100, 39° 22' N., 68° 34' W.

Genus 213. **SERRIVOMER** Gill and Ryder.

Serrivomer Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 260 (*S. beanii* Gill and Ryder).

563. **Serrivomer beanii** Gill and Ryder.

Gulf Stream; also taken in the Gulf of California.

Serrivomer beanii Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 261, Gulf Stream, lat. 41° 40' 30'', long. 65° 28' 30'' in 855 fathoms.

564. **Serrivomer sector** Garman.

Gulf of California to Panama in deep water.

Serrivomer sector Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 320, Pl. LXIII, numerous stations of the *Albatross* and off Guaymas, 50 miles south, surface to 700 fathoms.

Genus 214. **SPINIVOMER** Gill and Ryder.

Spinivomer Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 261 (*S. goodei* Gill and Ryder).

565. **Spinivomer goodei** Gill and Ryder.

Gulf Stream.

Spinivomer goodei Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 261, Gulf Stream, at lat. 38° 19' 26'', long. 68° 20' 20'' in 2,361 fathoms.

Genus 215. **LABICHTHYS** Gill and Ryder.

Labichthys Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 261 (*L. carinatus* Gill and Ryder).

566. **Labichthys carinatus** Gill and Ryder.

Gulf Stream.

Labichthys carinatus Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 253, 255, 261, Gulf Stream, 41° 13' N., 65° 33' W.

567. **Labichthys bowersi** Garman.

Off Panama in deep water.

Labichthys bowersi Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 823, Pl. XIV, fig. 1, *Albatross* stations 3414, 3388, and 3361, in 1,168 to 2,232 fathoms.

Family 77. **OPHICHTHYIDÆ**. Snake eelsGenus 216. **SPHAGEBRANCHUS** Bloch.

Sphagebranchus Bloch, Ichthyologia, IX, 1795, 88, pl. 419 (*S. rostratus* Bloch).

Cæcilia Lacépède, Hist. Nat. Poiss., II, 1800, 135 (*C. branderiana* Lacépède = *Muræna cæca* Linnæus); not *Cæcilia* L. a genus of *Batrachia*.

Apterichthys Duméril, Zool. Anal., 112, 1806, 331 (*Muræna cæca* Linnæus).

Branderius Rafinesque, Anal. Nat., 1815, 93; substitute for *Cæcilia* Lacépède.

Ichthyapus Brisout de Barneville, Revue Zool., 1847, 219 (*I. acutirostris*).

Ophisuraphis Kaup, Apodal Fishes, 1856, 29 (*O. gracilis* Kaup).

568. **Sphagebranchus anguiformis** (Peters).

Open Atlantic, near the West Indies.

Ophichthys (Sphagebranchus) anguiformis Peters, Monatsber. Akad. Wiss. Berlin, 1876, 849, Atlantic Ocean, 15° 40' N., 23° 5' W.

569. **Sphagebranchus selachops** (Jordan and Gilbert).

Rocks about Cape San Lucas.

Apterichthys selachops Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 356, Cape San Lucas.

570. **Sphagebranchus ophioneus** Evermann and Marsh.

Porto Rico.

Sphagebranchus ophioneus Evermann and Marsh, Fish. Porto Rico, Bull. U. S. Fish Com., XX, 1900, 73, fig. 7, off Mayaguez, Porto Rico.

Genus 217. **VERMA** Jordan and Evermann.

Verma Jordan and Evermann, Fishes North and Mid. Amer., 1896, 374 (*Sphagebranchus kendalli* Gilbert).

571. **Verma kendalli** (Gilbert).
Coast of Florida, in rather deep water.
Sphagebranchus kendalli Gilbert, Bull. U. S. Fish Com., IX, 1889 (1891),
310, off west coast of Florida, 25° 34' N., 82° 50' W., in 25 fathoms.
- Genus 218. **CRYPTOPTERENCHELYS** Fowler.
Cryptopterus Kaup, Aalen. Fische Hamburg Mus., II, 1860 (*C. puncticeps*
Kaup); name preoccupied.
Cryptopterenchelys Fowler, Amer. Mus. Novitates, CLXII, March 31,
1925 (*Cryptopterus puncticeps* Kaup); substitute for *Cryptopterus*
Kaup, preoccupied.
572. **Cryptopterenchelys puncticeps** (Kaup).
Caribbean Sea.
Cryptopterus puncticeps Kaup, Aalen. Fische Hamburg Mus., II, 1860,
pl. 1, fig. 2, Puerto Cabello.
573. **Cryptopterenchelys frontalis** (Garman).
Off Panama.
Ophichthys (*Cryptopterus*) *frontalis* Garman, Mem. Mus. Comp. Zool.,
XXIV, 1899, 309, Albatross stations 3386, 3389, 3391.
- Genus 219. **MYRICHTHYS** Girard.
Pisodonophis Kaup, Apodal Fishes, 1856, 15 (in part; not type, as
restricted by Bleeker, which is *P. cancrivorus*).
Myrichthys Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 58 (*M.*
tigrinus Girard).
Ophisurus Swainson, Bleeker, etc. (not of Lacépède, nor of Risso, nor
Kaup, who restrict the name to *O. serpens*).
574. **Myrichthys tigrinus** Girard.
Pacific coast of Mexico; rather common about Mazatlan, and occa-
sionally ranging northward.
Myrichthys tigrinus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 58,
Adair Bay, Oregon; doubtless an error.
Ophisurus xysturus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
346, Mazatlan.
575. **Myrichthys xysturus** (Jordan and Gilbert).
Pacific coast of Mexico; common among the rocks about Mazatlan.
Ophichthys xysturus Jordan and Gilbert, Proc. U. S. Nat. Mus., IX,
1881, 346, Mazatlan, Mexico.
576. **Myrichthys oculatus** (Kaup).
Tropical Atlantic; Cuba to Surinam and Cape Verde Islands.
Pisodonophis oculatus Kaup, Apodal Fishes, 1856, 22, Curaçao.
Ophisurus latimaculatus Poey, Repertorio, II, 1867, 252, pl. 3, fig. 1,
Cuba.
577. **Myrichthys acuminatus** (Gronow).
West Indies, occasionally northward to Florida Keys.
Muraena acuminata Gronow, Fish. Brit. Mus., 1854, 21, Insula Div.,
Eustachii.
Pisodonophis guttulatus Kaup, Apodal Fishes, 1856, 21, fig. 10, Mar-
tinique.
Ophisurus longus Poey, Repertorio, II, 1867, 254, Cuba.
Ophichthys psavarius Poey, Anal. Soc. Española Hist. Nat., 1875, 196,
Cuba.
578. **Myrichthys pantostigmus** Jordan and McGregor.
Revillagigedo Islands.
Myrichthys pantostigmus Jordan and McGregor, Rept. U. S. Fish Com.,
XXIV, 1898 (1899), 274, pl. 4, Clarion Island.
- Genus 220. **LETHARCHUS** Goode and Bean.
Letharchus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 436 (*L.*
velifer Goode and Bean).
579. **Letharchus pacificus** Osburn and Nichols.
Gulf of California.
Letharchus pacificus Osburn and Nichols, Bull. Amer. Mus. Nat. Hist.,
XXXV, 1916, 146, fig. 3, Cape San Lucas.

580. *Letharchus velifer* Goode and Bean.
Coast of Florida, in rather deep water; known only from the Snapper Banks off Pensacola and Tampa.
Letharchus velifer Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 437, West Florida.
- Genus 221. **PISODONOPHIS** Kaup.
Pisodonophis Kaup, Apodal Fishes, 1856, 17 (*P. cancrivorus* Kaup as restricted by Bleeker; later written *Pisoodonophis* by Kaup and since spelled *Pisodontophis* by purists).
581. *Pisodonophis daspilotos* Gilbert.
Panama.
Pisodonophis daspilotos Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2803, Panama.
- Genus 222. **OMOCHELYS** Fowler.
Omoichelys Fowler, Proc. Acad. Nat. Sci. Phila., LXX, 1918, 3 (*Pisodonophis cruentifer* Goode and Bean).
Styletor Jordan, loc. cit., LXX, 1918 (1919), 343 (*Pisoodonophis cruentifer* Goode and Bean); name preoccupied.
582. *Omoichelys cruentifer* (Goode and Bean).
Two specimens at station 1035 of U. S. Fish Commission steamer *Fish Hawk*, 39° 57' N., 69° 28' W., in 120 fathoms; four others at nearly same region in 245 fathoms.
Pisoodonophis cruentifer Goode and Bean, Oceanic Ichth., 1895, 147, fig. 166, Gulf Stream.
- Genus 223. **CALLECHELYS** Kaup.
Callechelys Kaup, Apodal Fishes, 1856, 28 (*C. guichenoti* Kaup).
583. *Callechelys muræna* Jordan and Evermann.
Snapper Banks off Pensacola.
Callechelys muræna Jordan and Evermann, Proc. U. S. Nat. Mus., IX, 1886, 466, Snapper Banks.
- Genus 224. **BASCANICHTHYS** Jordan and Davis.
Bascanichthys Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 621 (*Cæcula bascanium* Jordan).
584. *Bascanichthys bascanium* (Jordan).
Egmont Key, Florida.
Cæcula bascanium Jordan, Proc. Acad. Nat. Sci. Phila., XXXVI, 1884, 48, Egmont Key, Fla.
585. *Bascanichthys scuticaris* (Goode and Bean).
West coast of Florida.
Sphagebranchus scuticaris Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 343, Cedar Key.
Sphagebranchus teres Goode and Bean, loc. cit., V, 1882, 436, west Florida.
586. *Bascanichthys cylindricus* Meek and Hildebrand.
Panama.
Bascanichthys cylindricus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 152, Pl. VIII, Chame Point, Panama.
587. *Bascanichthys panamensis* Meek and Hildebrand.
Panama.
Bascanichthys panamensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, Pl. VII, 151, Chame Point, Panama.
588. *Bascanichthys bascanoides* Osburn and Nichols.
Coast of Lower California.
Bascanichthys bascanoides Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 147, fig. 4, San Cristobal, from the stomach of *Mycteroperea venadorum*.
589. *Bascanichthys peninsulæ* (Gilbert).
La Paz Bay, Gulf of California.
Callechelys peninsulæ Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 548, La Paz Bay, Lower California.

Genus 225. QUASSIREMUS Jordan and Davis.

Quassiremum Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 622
(*Ophichthys evionthas* Jordan and Bollman).

590. *Quassiremum nothochir* (Gilbert).

San Josef Island, Gulf of California.

Ophichthys nothochir Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 58,
San Josef Island, Gulf of California.

591. *Quassiremum evionthas* (Jordan and Bollman).

Hood Island, Galapagos Archipelago.

Ophichthys evionthas Jordan and Bollman, Proc. U. S. Nat. Mus., XII,
1889, 154, Hood Island, Galapagos Archipelago.

Genus 226. OPHICHTHUS Thunberg and Ahl.

Ophichthys Ahl, De Muræna et Ophichtho, 1789 (*Muræna ophis* L.).

Cogrus Rafinesque, Caratteri, 1810, 62 (*C. maculatus* Rafinesque).

Centruropus Kaup, Apodal Fishes, 1856, 2 (*C. spadiceus* Kaup).

Pæcilocephalus Kaup, loc. cit., 5 (*P. bonapartei* Kaup).

Microdontophis Kaup, loc. cit., 6 (*M. altipinnus* Kaup).

Cæcilophis Kaup, loc. cit., 6 (*C. compar* Kaup).

Herpetorhynchus Kaup, loc. cit., 7 (*H. ornatissimus* Kaup).

Elapsopsis Kaup, loc. cit., 9 (*E. versicolor* Kaup).

Murænopsis Kaup, loc. cit., 11 (*Murænophis ocellatus* LeSueur); the
name wrongly credited to LeSueur.

Scytalophis Kaup, loc. cit., 13 (*S. magnoculus* Kaup).

Leptorhinophis Kaup, loc. cit., 14 (*Ophisurus gomesi* Castelnau).

Uranichthys Poey, Repertorio, II, 1867, 256 (*Muræna havannensis*
Bloch and Schneider).

Oxyodontichthys Poey, Anales Soc. Española Hist. Nat., 1880, 254
(*macrurus*).

Ophichthys Bleeker, Günther, and of recent authors generally.

592. *Ophichthys ophis* (Linnæus).

West Indies; apparently not common.

? *Serpens marinus maculosus* Willoughby, Hist. Pisc., 1686, Tab. G., 9,
no locality.

? *Muræna ophis* Linnæus, Syst. Nat., ed. X, 1758, 244; after Willoughby;
may be identical with *Ophichthys regius*, a St. Helena species likewise
spotted with black.

Innomnado Parra, Dif. Piezas Hist. Nat., 1787, pl. 37, fig. 2, Habana.

Muræna havannensis Bloch and Schneider, Syst. Ichth., 1801, 491;
after Parra.

? *Ophisurus guttatus* Cuvier, Règne Anim., ed. 1, 1817, 232, Surinam;
after Bloch, pl. 154.

? *Muræna maculosa* Cuvier, loc. cit., Surinam; after *Ophisurus ophis*
Lacépède, which is based on Bloch, pl. 154.

Herpetorhynchus sulcatus Kaup, Apodal Fishes, 1856, 8, fig. 5, locality
unknown.

Uranichthys brachycephalus Poey, Repertorio, II, 1867, 257, Cuba.

593. *Ophichthys retropinnis* Eigenmann.

Snapper Banks off Pensacola, Fla.

Ophichthys retropinnis Eigenmann, Proc. U. S. Nat. Mus., X, 1887, 116,
Snapper Banks off Pensacola.

594. *Ophichthys guttifer* (Bean and Dresel).

Snapper Banks off Pensacola, Fla.

Ophichthys guttifer Bean and Dresel, Proc. Biol. Soc. Wash., II, 1884,
100, Snapper Banks.

595. *Ophichthys ocellatus* (LeSueur).

West Indies, south to Brazil and north to Pensacola.

Murænophis ocellatus LeSueur, Jour. Acad. Nat. Sci. Phila., V, 1825,
108, pl. 4, fig. 3, South America.

596. *Ophichthus triserialis* (Kaup).
Pacific coast of tropical America, rather common from Lower California to the Galapagos.
Muraenopsis triserialis Kaup, Apodal Fishes, 1856, 12, Pacific.
Herpetoichthys callisoma Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 475, locality unknown.
Ophisurus californiensis Garrett, Proc. Calif. Acad. Nat. Sci., III, 1863, 66, Margarita Bay, Lower California.
Ophichthus rugifer Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 155, Charles Island, Galapagos; young with longer pectorals.
597. *Ophichthus biserialis* (Garman).
Galapagos Islands.
Ophichthys biserialis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 311, Chatham Island.
598. *Ophichthus chamensis* Meek and Hildebrand.
Panama.
Ophichthys chamensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 155, Pl. VIII, fig. 2, Chame Point, Panama.
- Subgenus *SCYTALOPHIS* Kaup.
599. *Ophichthus gomesii* (Castelnau). *Sea serpent; Sea eel; Whipsnake eel.*
South Carolina to Rio Janeiro; generally common, especially about the Florida Keys and Cuba.
Ophisurus gomesii Castelnau, Anim. Amér. Sud, 1855, 84, pl. 44, fig. 2; Rio Janeiro.
Ophisurus chrysops Poey, Memorias, II, 1861, 321, Habana.
Oxydontichthys brachyurus Poey, Synopsis, 1868, 426; and Poey, Enumeratio, II, 1875, 155, Habana.
Oxydontichthys macrurus Poey, Anal. Soc. Española Hist. Nat., 1880, 254, Habana.
Oxydontichthys limbatus Poey, loc. cit., 254, Habana; name a substitute for *brachyurus*.
600. *Ophichthus zophochir* (Jordan and Gilbert).
Pacific Coast of Mexico, Guaymas to Acapulco.
Ophichthys zophochir Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 347.
601. *Ophichthus magniocus* (Kaup).
West Indies to Brazil; Aspinwall.
Scytalophis magniocus Kaup, Apodal Fishes, 1856, 13, fig. 7, St. Croix; Brazil.
602. *Ophichthus parilis* (Richardson).
Cuba to Brazil.
Ophisurus parilis Richardson, Voy. *Erebus* and *Terror*, 1844, 105, West Indies.
Ophichthys pauciporus Poey, Repertorio, II, 1868, 255, pl. 3, fig. 5.
- Genus 227. *MYSTRIOPHIS* Kaup.
Mystriophis Kaup, Apodal Fishes, 1856, 10 (*Ophisurus rostellatus* Richardson).
Crotolopsis Kaup, Abhandl. Natur. Verein Hamburg, IV, 1860, 12 (*C. quinctifer* Kaup).
Echiopsis Kaup, loc. cit., 13, (*Ophisurus intertinctus* Richardson).
Macrodonophis Poey, Repertorio, II, 1867, 251 (*M. mordax ophiurus* Poey).
- Subgenus *ECHIOPSIS* Kaup
603. *Mystriophis intertinctus* (Richardson).
West Indies, north to Pensacola, Fla.
Ophisurus intertinctus Richardson, Voy. *Erebus* and *Terror*, Fishes, 1844, 102, West Indies.
Ophisurus sugillatus Richardson, loc. cit., 103, probably West Indies.
Crotolopsis punctifer Kaup, Abhandl. Natur. Verein Hamburg, IV, 1860, 2, 12, pl. 1, fig. 3, Puerto Cabello.
Conger mordax Poey, Memorias, II, 1861, 319, Cuba.
Ophichthys schneideri Steindachner, Ich. Beitr., VIII, 1879, 66, Brazil.

Genus 228. SCYTALICHTHYS Jordan and Davis.

Scytalichthys Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 635 (*Ophichthys miurus* Jordan and Gilbert).

604. Scytalichthys miurus (Jordan and Gilbert).

Cape San Lucas, Lower California.

Ophichthys miurus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 357, Cape San Lucas.

Genus 229. BRACHYSOMOPHIS Kaup.

Brachysomophis Kaup, Apodal Fishes, 1856, 9 (*B. horridus* Kaup).

?*Achrophichthys* Bleeker, Poisson Ined. Murènes, Ned Tijdschr. Dierk., II, 242, 1865, (*A. typus*), young.

605. Brachysomophis crocodilinus (Bennett).

East Indies; a specimen recorded by Gunther from the Galapagos.

Ophisurus crocodilinus Bennett, Proc. Zool. Soc. Lond., 1833, 32, Mauritius.

Brachysomophis horridus Kaup, Apodal Fishes, 1856, 9, fig. 6, Otaheite [Tahiti].

?*Achrophichthys typus* Bleeker, Nederl. Tijdschr. Dierk., II, 1865, 242, Celebes.

Family 78. ACANTHENCHELYIDÆ

Genus 230. ACANTHENCHELYS Regan.

Acanthenchelys Regan, Ann. Mag. Nat. Hist., X, 1922, 296 (*A. spinicauda* Regan).

606. Acanthenchelys spinicauda Regan.

West Indies.

Acanthenchelys spinicauda Regan, Ann. Mag. Nat. Hist., X, 1922, 296, Island of Tobago.

Family 79. MORINGUIDÆ

Genus 231. MORINGUA Gray.

Moringua Gray, Zool. Misc., 1831, 9 (*M. lateralis* Gray).

Rataboura Gray, Illust. Ind. Zool., 1831, 95 (*Muræna ratibouri* Hamilton = *Rataboura hardwickei* Gray).

Pachyrurus Swainson, Nat. Hist. Fish, Reptiles and Amphib., II, 1839, 335 (*Moringua linearis* Gray; not of Agassiz, 1829).

Pseudomoringua Bleeker, Nederl. Tijdschr. Dierk., II, 1865, 114 (*Moringua lumbricoides* Richardson).

607. Moringua boekei Metzelaar.

Caribbean Sea.

Moringua boekei Metzelaar, Trop. Atl. Visschen, 1919, 17, Curaçao.

Genus 232. APHTHALMICHTHYS Kaup.

Aphthalmichthys Kaup, Apodal Fishes, 1856, 68 (*A. javanicus* Kaup).

608. Aphthalmichthys caribbeus Gill and Smith.

Porto Rico.

Aphthalmichthys caribbeus Gill and Smith, Science, XI, No. 286, 1900, 974, San Geronimo, near San Juan, Porto Rico.

Genus 233. STILBISCUS Jordan and Bollman.

Stilbiscus Jordan and Bollman, Proc. U. S. Nat. Mus., XI, 1888, 549 (*S. edwardsi* Jordan and Bollman).

609. Stilbiscus edwardsi Jordan and Bollman.

Green Turtle Cay, one of the Bahamas.

Stilbiscus edwardsi Jordan and Bollman, Proc. U. S. Nat. Mus., XI, 1888, 549, Green Turtle Cay, Bahamas.

Genus 234. GORDIICHTHYS Jordan and Davis.

Gordichthys Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 644 (*G. irretitus* Jordan and Davis).

610. Gordichthys irretitus Jordan and Davis.

Snapper Banks off west Florida in rather deep water.

Gordichthys irretitus Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 644, off Pensacola, Fla.

Order T. COLOCEPHALI

Family 80. MURÆNIDÆ. Morays

Genus 235. ENCHELYCORE Kaup.

Enchelycore Kaup, Apodal Fishes, 1856, 72 (*Muræna nigricans* Bonnaterre).

611. *Enchelycore nigricans* (Bonnaterre).

West Indies; rather common.

Muræna unicolor maxillus elongatus teretiusculus, inferiore longiore, etc., Gronow, Zoophy., 1, 1763, 163, South America.

Muræna nigricans Bonnaterre, Encycl. Meth. Ichth., 1788, 34; after Gronow.

Muræna anguina Gronow, Cat. Fish., 1854, 18, South America.

Enchelycore euryrhina Kaup, Apodal Fishes, 1856, 73, no locality.

612. *Enchelycore nigrocastaneus* (Cope).

West Indies.

Gymnothorax nigrocastaneus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 483, St. Martin.

?*Gymnothorax umbrosus* Poey, Ann. Lyc. Nat. Hist. N. Y., XI, 1874, 67, Habana, Cuba.

Genus 236. MURÆNA Linnæus. Morays.

Muræna Artedi, Gen. Pisc., 1738, 23, in part; includes all eels.

Muræna Klein, Hist. Pisc., 1742, 28, in part; includes all eels without pectoral fins.

Muræna Linnæus, Syst. Nat., ed. X, 1758, 244 (*helenæ*, etc.); includes all eels.

Muræna Thunberg and Ahl, De *Muræna* et *Ophichthus*, 1789, 6 (restricted to *helenæ*, etc.), includes species without pectoral fins.

Muræna Gunther and of authors generally; not of Bleeker.

Murænophis Lacépède, Hist. Nat. Poiss., V, 1803, 630 (*helenæ*, etc.).

Limamuræna Kaup, Apodal Fishes, 1856, 95 (*L. guttata* Kaup).

613. *Muræna melanotis* (Kaup).

Tropical Atlanta, from Africa to South America.

Limamuræna melanotis Kaup, Aalen. Fische Hamburg Mus., 1859, 27, pl. 4, fig. 3.

614. *Muræna lentiginosa* Jenyns. *Morena pinta*.

Pacific coast of America, from Gulf of California to Galapagos.

Muræna lentiginosa Jenyns, Voy. *Beagle*, Zool., 1842, 143, Galapagos Islands.

Muræna pinta Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 345, Mazatlan.

615. *Muræna retifera* Goode and Bean.

Coast of South Carolina, in rather deep water; one taken at Tucker-nuck Island, Mass.

Muræna retifera Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 435, off Charleston.

616. *Muræna argus* (Steindachner).

Gulf of California.

Gymnothorax (Limamuræna) argus Steindachner, Ichth. Notizen, X, 1870, 17, Pl. IV, Altata, Gulf of California.

617. *Muræna insularum* Jordan and Davis.

Galapagos Islands.

Muræna insularum Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 609, Chatham Island.

618. *Muræna myrialeucosticta* Fowler.

West Indies.

Muræna myrialeucosticta Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 18, St. Martin.

619. *Muræna clepsydra* Gilbert.

Panama.

Muræna clepsydra Gilbert, in Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2805, Panama.

Genus 237. PYTHONICHTHYS Poey.

Pythonichthys Poey, Repertorio, II, 1867, 265 (*P. sanguineus* Poey).

620. *Pythonichthys sanguineus* Poey.
Coast of Cuba, in rather deep water.
Pythonichthys sanguineus Poey, Repertorio, II, 1867, 265, pl. 2, fig. 7, Cuba.
- Genus 238. *RABULA* Jordan and Davis.
Rabula Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 589
(*R. aquæ-dulcis* Jordan and Davis=*Rabula davis* Fowler, not *Muraena aquæ-dulcis* Cope).
621. *Rabula davis* Fowler.
Said to be from San Diego, which is doubtful.
Rabula davis Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 21;
after *Rabula aquæ-dulcis* Jordan and Davis; not of Cope.
622. *Rabula marmorea* (Valenciennes).
Galapagos Islands.
Muraenophis marmoreus Valenciennes, Voy. *Vénus*, Zool., 1855, 347, pl. 10, fig. 1, Galapagos.
623. *Rabula panamensis* (Steindachner).
Pacific coast of Central America.
Muraena panamensis Steindachner, Ichth. Beitr., V, 1876, 19, Panama.
624. *Rabula longicauda* (Peters).
Tropical Atlantic, off the West Indies.
Muraena longicauda Peters, Monatsberichte, 1876, 850, open Atlantic,
with *Sphagebranchus anguiformis*.
- Genus 239. *GYMNOTHORAX* Bloch. *Morays*.
Gymnothorax Bloch, Naturgesch. Ausl. Fische, IX, 1795, 83, restricted
to allies of *M. afra* by Gunther (*G. muraena* Bloch).
Lycondontis McClelland, Calcutta Jour. Nat. Hist., V, No XVIII,
1844, 173 (*L. literata*=*tule*).
Throndontis McClelland, loc. cit., V, No. XVIII, 1844, 174 (*T. reticulata*=*tesselata*).
Eurymyctera Kaup, Apodal Fishes, 1856, 72, (*E. crudelis* Kaup).
Polyuranodon Kaup, loc. cit., 96 (*Muraena polyuranodon* Bleeker).
Tæniophis Kaup, Aalen. Fische Hamburg Mus., Nachträge, 1859, 10
(*T. westphali* Kaup).
Priodontophis Kaup, loc. cit., 1859, 22 (*Gymnothorax ocellatus* Agassiz).
Neomuraena Girard, U. S. Mex. Bound. Surv., Fishes, 1859, 76 (*N. nigromarginata* Girard=*ocellatus*).
Pseudomuraena Johnson, Proc., Zool. Soc. Lond., 1860, 167 (*P. maderirensis* Johnson).
625. *Gymnothorax mordax* (Ayres). *Conger eel of California*.
Point Conception to Cedros Island; abundant about Santa Barbara Islands.
Muraena mordax Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 30, Cerros [Cedros] Island.
626. *Gymnothorax moringa* (Cuvier). *Common spotted moray; Hamlet; Fel*.
Gulf of Mexico, Pensacola, and West Indies to Rio Janeiro and St. Helena.
Muraena maculata nigra; the black moray, Catesby, Nat. Hist. Carolinas, etc., 1738, pl. 21, Bahamas.
? *Gymnothorax afer* Bloch, Syst. Ichth., 1795, pl. 417, Africa.
Muraena moringa Cuvier, Règne Anim., ed. 2, II, 1829, 352; after Catesby.
Gymnothorax rostratus Agassiz, Spix, Pisc. Brasil., 1830, 91, pl. 50a, Brazil.
Muraena moringua Richardson, Voy. *Erebus* and *Terror*, Fishes, 1844, 89, Jamaica.
Muraena punctata Gronow, Cat. Fish., 1854, 18, North America.
Muraenophis curvilineata Castelnau, Anim. Amér. Sud, Poiss., 1855, 81, pl. 42, fig. 2, Rio Janeiro.
Muraenophis caramuru Castelnau, loc. cit., 85, pl. 43, Bahia.
Gymnothorax flavoscriptus Poey, Enumeratio, 1875, 158, Cuba.
Gymnothorax picturatus Poey, Anal. Soc. Española Hist. Nat., 1880, 257, Cuba.

627. *Gymnothorax funebris* Ranzani. *Black moray; Morena verde*.
Tropical America; Rio Janeiro; West Indies.
Muraena maculata nigra et viridis (the moray) Catesby, Nat. Hist. Carolinas, etc., 1738, pl. 20, Bahamas.
Gymnothorax funebris Ranzani, Nov. Com. Acad. Sci. Inst. Bonon., IV, 1840, 76, Brazil.
Muraena lineopinnus Richardson, Voy. *Erebus and Terror*, Fishes, 1844, 89, Puerto Cabello.
Muraena infernalis Poey, Memorias, II, 1861, 347, 354, Cuba.
Thyrsoidea concolor Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 479, Vera Cruz.
Muraena erebus Poey, Memorias, II, 1861, 426, Cuba.
628. *Gymnothorax castaneus* (Jordan and Gilbert).
Pacific coast of Mexico; common about Mazatlan; representing *G. funebris* of the West Indies
Sidera castanea Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 647, Mazatlan.
629. *Gymnothorax sanctæ-helenæ* (Günther).
Tropical Atlantic; recorded from St. Helena and the Bermudas.
Muraena sanctæ-helenæ Gunther, Cat., VIII, 1870, 115, St. Helena.
630. *Gymnothorax vicinus* (Castelnau).
Tropical Atlantic, Cuba to Africa and Brazil.
Muraenophis vicina Castelnau, Anim. Amér. Sud, Poiss., 1855, 81, pl. 42, fig. 4, Bahia.
Gymnothorax versipunctatus Poey, Enumeratio, 1875, 156, Cuba.
Thyrsoidea maculipinnus Kaup, Apodal Fishes, 1856, 86, Gold Coast.
631. *Gymnothorax punctarenæ* Fowler.
Puntarenas, Costa Rica, one specimen known.
Gymnothorax punctarenæ Fowler, Copeia, No. 39, Jan. 24, 1917, 3, Puntarenas, Costa Rica.
632. *Gymnothorax brunneus* Nichols.
Bermudas.
Gymnothorax brunneus Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 59, Bermuda.
633. *Gymnothorax virescens* Poey.
Cuba.
Gymnothorax virescens Poey, Enumeratio, 1875, 156, Cuba.
634. *Gymnothorax polygonius* Poey.
Cuba.
Gymnothorax polygonius Poey, Ann. Lyc. Nat. Hist. N. Y., XI, 1870, 68, Habana.
635. *Gymnothorax dovii* (Günther). *Morena pintita*.
Gulf of California to the Galapagos Archipelago.
Muraena dovii Gunther, Cat., VIII, 1870, 103, Panama.
Muraena pintita Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 346, Mazatlan.
636. *Gymnothorax conspersus* Poey.
Cuba to Rio Janeiro.
Muraenophis punctata Castelnau, Anim. Amér. Sud, 1855, 82, Rio Janeiro; not *Gymnothorax punctatus* Bloch.
Gymnothorax conspersus Poey, Repertorio, II, 1868, 259, Cuba.
637. *Gymnothorax jordani* (Evermann and Marsh).
Porto Rico.
Lycondonitis jordani Evermann and Marsh, Fish. Porto Rico, Bull. U. S. Fish Com., XX, 1900 (1902), pt. 1, 78, pl. 2; off Mayaguez, Porto Rico.
638. *Gymnothorax miliaris* (Kaup).
West Indies.
Thyrsoidea miliaris Kaup, Apodal Fishes, 1856, 90, Martinique.
Muraena multocellata Poey, Memorias, II, 1861, 324, Cuba; based on specimens with smaller spots than usual.
Gymnothorax scriptus Poey, Repertorio, II, 1868, 261, Cuba.
Muraena myrialeucostictus Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 18, St. Martins, West Indies.

639. *Gymnothorax elaboratus* (Poey).
Cuba.
Muraena elaborata Poey, *Memorias*, II, 1861, 323, Cuba.
640. *Gymnothorax obscuratus* Poey.
Cuba.
Gymnothorax obscuratus Poey, *Ann. Lyc. Nat. Hist. N. Y.*, IX, 1870, 320, Cuba.
641. *Gymnothorax chlevastes* (Jordan and Gilbert).
Galapagos Islands.
Sidera chlevastes Jordan and Gilbert, *Proc. U. S. Nat. Mus.*, VI, 1883, 208, Galapagos.
642. *Gymnothorax albimentis* (Evermann and Marsh).
Porto Rico.
Lycodontis albimentis Evermann and Marsh, *Fish. Porto Rico*, *Bull. U. S. Fish Com.*, XX, 1900 (1902), pt. 1, 78, fig. 9; off Culebra Island.
643. *Gymnothorax concolor* (Abbott).
West Indies.
Thyrsoidea concolor Abbott, *Proc. Acad. Nat. Sci. Phila.*, XII, 1860, 479, Vera Cruz.
? *Muraena erebus* Poey, *Memorias*, II, 1861, 426, Cuba.
644. *Gymnothorax aquædulcis* (Cope).
Rio Grande, Costa Rica.
Muraena aquædulcis Cope, *Rept.*, U. S. Geol. Surv., 1871 (1872), 474; said to be from Rio Grande, Costa Rica. It is evidently not a fresh-water fish. According to Fowler, it is not a *Rabula*, the dorsal beginning well before the gill opening.
645. *Gymnothorax carcinognathus* Fowler.
West Indies.
Gymnothorax carcinognathus Fowler, *Proc. Acad. Nat. Sci. Phila.*, LXIV, 1912, 22, St. Martins, West Indies.
646. *Gymnothorax verilli* (Jordan and Gilbert).
Panama; one specimen known.
Sidera verrilli Jordan and Gilbert, *Proc. U. S. Nat. Mus.*, V, 1882, 623, Panama.
647. *Gymnothorax ocellatus* Agassiz. *Spotted moray*; *Moray eel*.
Pensacola to Rio Janeiro.
Gymnothorax ocellatus Agassiz, *Pisc. Brasil.*, 1828, 91, pl. 50b, Brazil.
648. *Gymnothorax ocellatus saxicola* Jordan and Davis.
Snapper Banks off Pensacola and deep water off Cuba.
Muraena meleagris Quoy and Gaimard, *Voy. Freycinet*, *Zool.*, 1824, 245, pl. 52, fig. 2; not of Shaw.
Gymnothorax ocellatus saxicola Jordan and Davis, *Rept. U. S. Fish Com.*, XVI, 1888, 606, Snapper Banks off Pensacola.
649. *Gymnothorax ocellatus nigromarginatus* (Girard).
Coast of west Florida and Texas.
Neomuraena nigromarginata Girard, *U. S. Mex. Bound. Surv.*, 1859, 76, pl. 41, St. Joseph Island, Tex.
- Genus 240. *SIDEEA* Kaup.
Siderea Kaup, *Uebersicht der Aale*, 1856, 59 (*Muraena pfeifferi* Bleeker = *Muraena picta* Ahl); name later written *Sidera*.
650. *Siderea picta* (Ahl).
South Seas; East Indies; Hawaii; Revillagigedos; taken at Clarion Island.
Muraena picta Ahl, *De Muraena et Ophichtho*, VI, 1787, 6, tab. 2.
Muraenopsis pantherina Lacépède, *Hist. Nat. Poiss.*, V, 1803, 628.
Muraena variegata Quoy and Gaimard, *Voy. Uranie*, 1824, 246, pl. 52.
Muraena lita Richardson, *Voy. Erebus and Terror*, 1844, 84, Moluccas.
Muraena siderea Richardson, *loc. cit.* 85, pl. 48, Australia.
Muraena pfeifferi Bleeker, *Nat. Tijds. Ned. Ind.*, V, 173, Celebes.

Genus 241. AHYNNODONTOPHIS Fowler.

Ahynnodontophis Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 25 (*A. stigmatotus* Fowler).

651. Ahynnodontophis stigmatotus Fowler.

West Indies (?).

Ahynnodontophis stigmatotus Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 25; supposed to be from the West Indies.

Genus 242. ECHIDNA Forster.

Echidna Forster, Bibliotheca Banksia, 1777, 181 (*E. variegata* Forster).
Gymnomuraena Lacépède, Hist. Nat. Poiss., V, 1803, 648 (*G. dolata* Lacépède).

Gymnopsis Rafinesque, Anal. Nat., 1815, 93 (substitute for *Gymnomuraena*).

Megaderia Rafinesque, loc. cit., 93 (substitute for *Echidna* Forster).

Molarii Richardson, Voy. Erebus and Terror, 1844, 79 (*Muraena ophis* = *Echidna nebulosa*).

Pacilophis Kaup, Apodal Fishes, 1856, 98 (*Gymnothorax catenatus* Bloch).

652. Echidna catenata (Bloch).

West Indies, from Bermuda to Surinam; our specimens from Santa Lucia.

Muraena seu conger brasiliensis Seba, Thesaurus, II, 1738, 72, pl. 69, figs. 4, 5, Brazil.

Gymnothorax catenatus Bloch, Naturgesch. Ausl. Fische, XII, 1795, 74, pl. 415, fig. 1, Coromandel; an error.

Muraena sordida Cuvier, Règne Anim., ed. 1, 1817, 233, after Seba.

Muraenophis catenula Lacépède, Hist. Nat. Poiss., V, 1803, 628 and 641, Palmerston Island; after Bloch.

Echidna fuscomaculata Poey, Repertorio, II, 1868, 263, Cuba.

Echidna flavoscripta Poey, loc. cit., 264, Cuba.

653. Echidna nocturna (Cope).

Pacific coast of Mexico, the two known specimens from the Rio Grande, in Costa Rica and from Cape San Lucas.

Pacilophis nocturnus Cope, Rept. U. S. Geol. Surv. Mont., 1871 (1872), 474, Rio Grande at San José, Costa Rica.

654. Echidna scabra Garman

Cocos Island.

Echidna scabra Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 314, Cocos Island.

655. Echidna cocosa Garman.

Cocos Island.

Echidna cocosa Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 313, Cocos Island.

656. Echidna chionostigma Fowler.

Gulf of California (?).

Echidna chionostigma Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 28; supposed to be from the Gulf of California.

Genus 243. UROPTERYGIUS Rüppell.

Gymnomuraena Lacépède, Hist. Nat. Poiss., V, 1803, 648 (*G. dolata* Lacépède); restricted first by Kaup, in 1856, to *dolata*, which is a species of *Echidna*.

Ichthyophis Lesson, Voy. Coquille, II, 1830, 120 (*I. pantherinus* Lesson); not of Fitzinger, 1829, a genus of reptiles.

Uropterygius Rüppell, Neue Wirbelthiere, Fische, 1835, 83 (*Nebrius concolor* Rüppell).

Muraenoblenna Kaup, Apodal Fishes, 1856, 67, (*Ichthyophis tigrina* Lesson); not of Lacépède, 1803, which is *Myxine*.

Scutica Jordan and Evermann, Fish. North and Mid. Amer., 1896, 403, (*Uropterygius neclurus* Jordan and Gilbert).

Subgenus SCUTICA Jordan and Evermann.

657. *Uropterygius necturus* (Jordan and Gilbert).

Gulf of California.

Gymnomuraena nectura Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 356, Cape San Lucas.

Genus 244. CHANNOMURÆNA Richardson.

Channo-Muræna Richardson, Voy. *Erebus* and *Terror*, 1844, 96 (*Ichthyophis vittata* Richardson).658. *Channomuræna vittata* (Richardson).

Coast of Cuba

Raro Parra, Dif. Piezas Hist. Nat., 1780, 66, pl. 30, fig. 3, Habana.*Ichthyophis vittatus* Richardson, Voy. *Sulphur*, Fishes, 1844, 114, pl. 53, figs. 7-9; locality uncertain.*Channomuræna cubensis* Poey, Repertorio, II, 1867, 266, pl. 3, fig. 6, Cuba.

Order U. LYOMERI

Family 81. SACCOPHARYNGIDÆ. Gulpers

Genus 245. SACCOPHARYNX Mitchill.

Saccopharynx Mitchill, Ann. Lyc. Nat. Hist. N. Y., 1824, 86 (type afterwards called *S. flagellum* Mitchill).*Ophiognathus* Harwood, Philos. Trans., 1827, 277 (*O. ampullaceus* Harwood).659. *Saccopharynx ampullaceus* (Harwood).

Atlantic.

Ophiognathus ampullaceus Harwood, Philos. Trans., 1827, 52, Atlantic.*Saccopharynx flagellum* Cuvier, Règne Anim., ed. 2, 1829, 355, Atlantic; after Mitchill.*Saccopharynx chordatus* Storer, Synopsis, 1846, 237, open Atlantic; after Mitchill.

Family 82. EURYPHARYNGIDÆ. Swallowers

Genus 246. GASTROSTOMUS Gill and Ryder.

Gastrostomus Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 271 (*G. bairdii* Gill and Ryder).660. *Gastrostomus bairdii* Gill and Ryder.

Deep water off Newfoundland Banks and Davis Strait.

Gastrostomus bairdii Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 271, off the Grand Banks of Newfoundland.661. *Gastrostomus pacificus* Bean.

Pacific Ocean in deep water.

Gastrostomus pacificus Bean, Smiths. Misc. Coll., XLV, 1904, 254, fig. 31; halfway between Midway Island and Guam.

Series OSTARIOPHYSI

Order V. HETEROGNATHI

Family 83. CHARACINIDÆ. Characins

Genus 247. PHENAGONIATES Eigenmann and Wilson.

Phenagoniates Eigenmann and Wilson, Indiana Univ. Studies, XIX, 1914, 2 (*P. wilsoni* Eigenmann = *Raboides macrolepis* Meek and Hildebrand).662. *Phenagoniates macrolepis* (Meek and Hildebrand).

Atlantic slope of Panama.

Raboides macrolepis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 84, Rio Cupe, Boca de Cupe, Tuyra River Basin.

Genus 248. COMPSURA Eigenmann.

Compsura Eigenmann, Mem. Carnegie Mus., VII, 1915, 60 (*Compsura heterura* Eigenmann).

663. *Compsura gorgonæ* (Evermann and Goldsborough).

Both slopes of Panama.

Cheirodon gorgonæ Evermann and Goldsborough, Proc. Biol. Soc. Wash., XXII, 1909, 99, figs. 1, 3, Gorgona, Atlantic slope, Panama Canal Zone.**Genus 249. PSEUDOCHEIRODON Meek and Hildebrand.***Pseudocheirodon* Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 275 (*P. affinis* Meek and Hildebrand).**664. *Pseudocheirodon affinis* Meek and Hildebrand.**

Both slopes of Panama.

Cheirodon insignis Evermann and Goldsborough, Proc. Biol. Soc. Wash., XXII, 1909, 98, Tabernilla, Atlantic slope; not of Steindachner.*Pseudocheirodon affinis* Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 275, Pl. XVIII, Rio Gatun, Monte Liria, Canal Zone, Panama.**Genus 250. GEPHYROCHARAX Eigenmann.***Gephyrocharax* Eigenmann, Indiana Univ. Studies, No. 16, 1912, 23 (*G. chocoensis* Eigenmann).**665. *Gephyrocharax atricaudatus* (Meek and Hildebrand).**

Both slopes, Canal Zone, Panama.

Deuterodon atricaudata Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1912, 68, Rio Frijoles, Canal Zone.*Gephyrocharax atricaudata* Eigenmann, Indiana Univ. Studies, No. 20, 1914, 41, Panama Canal Zone.**666. *Gephyrocharax intermedius* Meek and Hildebrand.**

Rio Chama, Panama.

Gephyrocharax intermedius Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 278, Rio Chama.**Genus 251. ASTYANAX Baird and Girard.***Astyanax* Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 26 (*A. argentatus* Baird and Girard=*Tetragonopterus mexicanus* Filippi).*Pacilurichthys* Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 417 (*P. brevoorti* Gill).All the species mentioned below belong to the subgenus or genus *Astyanax*, distinguished from *Tetragonaplenes* Cuvier by the absence of teeth on the maxillary and by the shorter anal of less than 40 rays.**667. *Astyanax mexicanus* (Filippi).**

Rio Balsas and Rio Tehuacan to tributaries of the Rio Grande in Texas, the northernmost species of the multitude of Characins.

Tetragonopterus mexicanus Filippi, Guerin's Rev. Mag. Zool., 1853, 166, Mexico.*Astyanax argentatus* Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 27, Comanche Spring and Brownsville, Tex.*Tetragonopterus fulgens* Bocourt, Ann. Sci. Nat. Zool., IX, 62, 1868.*Tetragonopterus nitidus* Vaillant and Pellegrin, Bull. Mus. Hist. Nat. Paris, 1904, 324.*Tetragonopterus brevimanus* Günther, Cat., V, 1864, 325, Yzabel; Rio San Geronimo, Guatemala.**668. *Astyanax æneus* (Günther).**

Oaxaca, Mexico, and Rio Chagres, and other streams about Panama, south to Brazil.

Tetragonopterus æneus Günther, Proc. Zool. Soc. Lond., 1860, 319, Oaxaca, Mexico.*Tetragonopterus microphthalmus* Günther, Cat., V, 1864, 324, Lake Amatitlan, Guatemala; Rio Rimag.*Tetragonopterus panamensis* Gunther, loc. cit., 324, Pacific coast of Panama; Yzabel.*Tetragonopterus humilis* Günther, loc. cit., 327, Lake Amatitlan, Guatemala.*Tetragonopterus ærstedii* Krøyer, in Lütken, Ict. Bidrag., III, 1874, 229, Rio San Juan, Nicaragua.

669. *Astyanax petenensis* (Gunther).
Nicaragua to western Ecuador; southeast to Argentine Republic.
Tetragonopterus petenensis Gunther, Cat., V, 1864, 326, Lake Peten, Nicaragua.
670. *Astyanax scabripinnis* (Jenyns).
Jamapa, Mexico, to Rio Janeiro.
Tetragonopterus scabripinnis Jenyns, Voy. *Beagle*, Zool., Fishes, 1842, 125, Rio Janeiro, Brazil.
671. *Astyanax macrophthalmus* (Regan).
Streams of eastern Mexico.
Tetragonopterus macrophthalmus Regan, Biol. Centrali-Amer., 1915, 171, eastern Mexico; Monzorangi, Vera Cruz.
672. *Astyanax nasutus* Meek.
Astyanax nasutus Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 108, Lake Managua, Nicaragua.
673. *Astyanax fasciatus* (Cuvier).
Coasts of Brazil, from Para to Napo and on the Pacific side from Panama to Ecuador.
Tetragonopterus fischeri Steindachner, Flussfische Südamerikas, I, 1879, 166, Mamo River at Chepo, Colombia.
674. *Astyanax ruberrimus* Eigenmann.
Panama, both slopes.
Astyanax ruberrimus Eigenmann, Indiana Univ. Studies, XVIII, 1913, 25, Isthmeria, Atlantic slope.
675. *Astyanax brevoortii* (Gill).
Island of Trinidad.
Pacilurichthys brevoortii Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 417, Trinidad.
Pacilurichthys taenirurus Gill, loc. cit., 418, Trinidad.
Pacilurichthys pulcher Gill, loc. cit., 419, Trinidad.
- Genus 252. **BRYCONAMERICUS** Eigenmann.
Bryconamericus Eigenmann, Ann. Carnegie Mus., IV, 1907, 139, (*B. exodon* Eigenmann).
676. *Bryconamericus emperador* (Eigenmann and Ogle).
Streams of both slopes, Panama.
Astyanax emperador Eigenmann and Ogle, Proc. U. S. Nat. Mus., XXXIII, 1907, 26, Empire Station, Panama.
677. *Bryconamericus cascajalensis* Meek and Hildebrand.
Panama.
Bryconamericus cascajalensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 284, Rio Cascajal, Puerto Bello, Panama.
- Genus 253. **HEMIBRYCON** Günther.
Hemibrycon Günther, Cat., V, 1864, 318 (*Tetragonopterus (Hemibrycon) polyodon* Günther).
678. *Hemibrycon dariensis* Meek and Hildebrand.
Basin of Rio Tuyra, Panama.
Hemibrycon dariensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 285, Pl. XX, mouth of Rio Yape, Panama.
- Genus 254. **HYPHESOBRYCON** Durbin.
Hyphessobrycon Durbin, in Eigenmann, Bull. Mus. Comp. Zool., LII, 1908, 100, (*Hemigrammus compressus* Meek).
Dermatocheir Durbin, Ann. Carnegie Mus., 1909, 55 (*D. calablepta* Durbin).
679. *Hyphessobrycon panamensis* Durbin.
Rio Chagres Basin, Panama.
Hyphessobrycon panamensis Durbin, in Eigenmann, Bull. Mus. Comp. Zool., LII, 1908, 101, Rio Boqueron, Atlantic slope, Panama.
Hemigrammus minutus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1912, 67, Agua Clara, Panama, Atlantic side of Isthmus of Panama.

Genus 255. *HEMIGRAMMUS* Gill.

Hemigrammus Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 420 (*H. unilineatus* Gill).

680. *Hemigrammus unilineatus* (Gill).

Streams of Trinidad.

Pecilurichthys unilineatus Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1854, 420, Trinidad.

Genus 256. *CREAGRUTUS* Günther.

Creagrutus Günther, Cat., V, 1864, 339 (*Leporinus mulleri* Günther).

681. *Creagrutus notropoides* Meek and Hildebrand.

Rio Chagres, Panama.

Creagrutus notropoides Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1912, 68, Rio Indio, Chagres basin.

682. *Creagrutus simus* Meek and Hildebrand.

Rio Tuyra Basin, Panama.

Creagrutus simus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 85, Rio Cupe, Cituro, Panama.

Genus 257. *RÆBOIDES* Günther.

Ræboides Günther, Cat., V, 1864, 347 (*Epicyrthus microlepis* Reinhardt).

683. *Ræboides guatemalensis* (Günther).

Rio Chagres and streams of Guatemala.

Anacyrtus guatemalensis Günther, Cat., V, 1864, 347, Rio Chagres; Huamuchal.

684. *Ræboides salvadoris* Hildebrand. *Plateado; Sardina; Alma seca; Umina.*

Lakes Guija and Metapan and Rio Sucio and Lempa, El Salvador.

Ræboides salvadoris Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925, 246, fig. 9, Rio Sucio, Sitio del Niño.

685. *Ræboides occidentalis* Meek and Hildebrand.

Pacific slope of Panama and southward.

Ræboides occidentalis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 293, Pl. XXIII, Rio Cardenas, Corozal, Panama.

Genus 258. *BRYCON* Muller and Troschel.

Brycon Müller and Troschel, Horæ Ichth., I, 1845, 15 (*B. falcatus* Müller and Troschel).

Chalcinopsis Kner, Sitzungsab. Akad. Wiss. München, 1863, 226 (*C. striatulus* Kner).

Megalobrycon Günther, Proc. Zool. Soc. Lond., 1869, 423 (*Brycon cephalus* Günther).

686. *Brycon striatulus* (Kner).

Rio Chagres and streams about Panama.

Chalcinopsis striatulus Kner, Sitzungsab. Akad. Wiss. München, 1863, 226, Panama.

Chalcinopsis chagresensis Kner, loc. cit., 1863, 226, Rio Chagres.

687. *Brycon guatemalensis* Regan.

Brycon guatemalensis Regan, Biol. Centrali-Amer., 1908, 168, lakes and streams of Guatemala and Nicaragua.

688. *Brycon petrosus* Meek and Hildebrand. *Pipon.*

Chagres River basin, Atlantic slope of Panama.

Brycon petrosus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 84, Pl. XXIV, 1916; upper Chagres, Panama.

689. *Brycon dentex* Günther.

Yucatan to Ecuador.

Brycon dentex Günther, Proc. Zool. Soc. Lond., 1860, 240, Esmeraldas, Ecuador.

690. *Brycon argenteus* Meek and Hildebrand.

Rio Charrera, and in Rio Bayano and Rio Tuyra basins, Pacific slope of Panama.

Brycon argenteus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 84, Pl. XXV, 298, 1916; Rio Aruza, Aruza, Panama.

691. *Brycon oligolepis* Regan.
Streams of Panama
Brycon oligolepis Regan, Ann. Mag. Nat. Hist., IX, 1913, 462, Rio Condoto, Panama.
- Genus 259. **BRAMOCHARAX** Gill.
Bramocharax Gill, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 189 (*B. transfordi* Gill).
692. *Bramocharax transfordi* Gill.
Lake Nicaragua.
Bramocharax transfordi Gill, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 190, Lake Nicaragua.
693. *Bramocharax elongatus* Meek. *Sabalito*.
Bramocharax elongatus Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 110, Lake Managua, Nicaragua.
- Genus 260. **PIABUCINA** Cuvier and Valenciennes.
Piabucina Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849, 122 (161) (*P. erythrinoides* Cuvier and Valenciennes).
694. *Piabucina panamensis* Gill.
Tripoli, Atlantic slope of the Isthmus of Panama.
Piabucina panamensis Gill, Proc. Acad. Nat. Sci. Phila., XXVIII, 1876, 336, Rio Tripoli, Isthmus of Panama.
695. *Piabucina festæ* Boulenger.
Panama.
Piabucina festæ Boulenger, Bull. Mus. Torino, XIV, 1899, 346, I, Rio Sabina, Panama.
- Genus 261. **HOPLIAS** Gill.
Macrodon Müller, in Wiegmann's Archiv, 1843 163, (*Macrodon trahira* = *Synodus malabaricus* Schneider); name preoccupied in *Sciænidæ*.
Hoplias Gill, Proc. U. S. Nat. Mus., XXVI, 1903, 1016 (*Synodus malabaricus* Schneider).
696. *Hoplias malabaricus* (Bloch).
Rio Tuyra Basin; both slopes of Colombia; Ecuador, and south to Brazil.
Esox malabaricus Bloch, Naturgesch. Ausl. Fische, pt. 8, 1794, 149, pl. 392, Tranquebar; locality wrongly given.
Macrodon tereira Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 508, Bahia; San Francisco; Amazon; Maracaibo.
Erythrinus trahira Agassiz, in Spix, Pisc. Brasil., 1829, pl. 18, Bahia; San Francisco; Amazon; Maracaibo.
Erythrinus macrodon Agassiz, loc. cit., 43, pl. 18, Alma, Prov. Bahia, San Francisco.
Erythrinus macrocephalus Agassiz, loc. cit., 44, San Francisco.
Erythrinus brasiliensis Agassiz, loc. cit., 45, pl. 20, Peruaguacu.
Macrodon auritus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 519, Montevideo.
Macrodon teres Cuvier and Valenciennes, loc. cit., 521, Lake Maracaibo.
Macrodon patana Cuvier and Valenciennes, loc. cit., 522, Cayenne.
Macrodon amara Cuvier and Valenciennes, loc. cit., 586, Cayenne.
Macrodon ferox Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 51, Trinidad.
697. *Hoplias microlepis* (Günther).
Panama to Ecuador.
Macrodon microlepis Günther, Cat., V, 1864, 282, western Ecuador, Guatemala.

Family 84. **GASTEROPELICIDÆ**

- Genus 262. **GASTEROPELECUS** Gronow.
Gasteropelecus Gronow, Mus. Ichth., II, 1763, 7 (type given in Gen. Fish. as *Clupea sternica* L.).

698. *Gasteropelecus maculatus* Steindachner. *Peche-peche*.
Rio Mamoni, near Panama.
Gasteropelecus maculatus Steindachner, Flussfische Südamerikas, 1879, 168, Rio Mamoni, at Chepo, Panama.

699. *Gasteropelecus sternicla* Pallas.

Surinam; Brazil; Essequibo.

Gasteropelecus sternicla Pallas, Spic. Zool., 1780, fasc. 8, p. 50, tab. 3, fig. 4.

Family 85. XIPHOSTOMIDÆ

Genus 263. CTENOLUCIUS Gill.

Ctenolucius Gill, Suppl., Proc. Acad. Nat. Sci. Phila., XIII, '1861, 8 (*Luciocharax insculptus* Steindachner).*Luciocharax* Steindachner, Denkschr. Akad. Wiss. Wien, XXXIX, 1878 (1879), 67 (*L. insculptus* Steindachner).700. *Ctenolucius insculptus* (Steindachner).

Rio Magdalena and Rio Momoni near Panama.

Luciocharax insculptus Steindachner, Denkschr. Akad. Wiss. Wien, XXXIV, 1878 (1879), 67 Rio Magdalena, and Rio Mamoni near Panama.701. *Ctenolucius beani* (Fowler).

Pacific coast of Panama; both slopes of Colombia.

Beloncharax beani Fowler, Proc. Acad. Nat. Sci. Phila., LVIII, 1906, 464, Rio Truando, a tributary of Rio Atrato, Panama.*Luciocharax striatus* Boulenger, Ann. Mag. Nat. Hist., ser. 8, VII, 1911, 312, Boca de Calmia, Choco.

Family 86. ANOSTOMIDÆ. (Curimatidæ)

Genus 264-265. CURIMATA Walbaum.

Curimata Walbaum, Artdi Pisc., 1792, 80 (*Salmo marcgravi* Walbaum, based on *Charax maxilla superiore longiore* Gronow, which is *Salmo cyprinoides* Linnaeus).*Les Curimates* Cuvier, Règne Anim., ed. 1, II, 1817, 165 (*Salmo edentulus* Bloch).*Curimatus* of authors generally; altered spelling.*Curimatella* Eigenmann and Eigenmann, Ann. N. Y. Acad. Sci., IV, 1889, 415 (*C. lepidurus* Eigenmann and Eigenmann).*Semilapicus* Eigenmann and Eigenmann, loc. cit., 415 (*Curimatus spilurus* Günther).701a. *Curimata magdalene* Steindachner. *Sardina blanca*.

Rio Magdalena and Rio Mamoni, Colombia.

Curimatus magdalene Steindachner, Denkschr. Akad. Wiss. Wien, XXXIX, 1878 (1879), 34, Rio Magdalena, Colombia.

Family 87. HEMIODONTIDÆ

Genus 266. CHARACIDIUM Reinhardt.

Characidium Reinhardt, Overs. Dansk. Forh. Kjøbenhavn, 1866, 56, (*C. fasciatum* Reinhardt).702. *Characidium marshi* Breder.

Tuyra Basin, Pacific slope of Panama.

Characidium marshi Breder (C. M.), Amer. Mus. Novitates, CLXXX, July 5, 1925, 5, fig. 5, Rio Sucubti, Darien, Panama.

Genus 267. APAREIODON Eigenmann.

Apareiodon Eigenmann, Ann. Carnegie Mus., X, 1916, 71 (*Parodon piracicabæ* Eigenmann).703. *Apareiodon compressus* Breder.

Tuyra Basin, Pacific slope of Panama.

Apareiodon compressus C. M. Breder, jr., Amer. Mus. Novitates, CLXXX, July 5, 1925, 4, fig. 3, 4, Rio Chucunaque, Panama.704. *Apareiodon dariensis* (Meek and Hildebrand).

Rio Cupe, Panama.

Parodon dariensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 83, Pl. XVII, 1916, Rio Cupe, Pacific slope of Panama.

Order W. GYMNONOTI

Family 83. GYMNOTIDÆ

Genus 268. GYMNOTUS Linnæus.

Gymnotus Linnæus, Syst. Nat., Ed. X, 1758, 246 (*G. carapo* L.).

Carapus Rafinesque, Indice, 37, 1810, 57 (*Gymnotus acus* Linnæus, prior to *Pterasfer* Cuvier, not *Carapus* Cuvier=*Giton* Kaup).

Carapus Müller and Troschel, Horæ Zool., III, 1849, 13 (*Gymnotus fasciatus* Pallas=*Gymnotus carapo* L.).

705. *Gymnotus carapo* Linnæus.

Carapo Marcgrave, Hist. Pisc. Brasil., 1648, 170, Brazil.

Gymnotus carapo Linnæus, Syst. Nat., Ed. X, 1758, 246.

Gymnotus fasciatus Pallas, Spic. Zool., VII, 1769, 35, fresh waters of America; anal rays about 180.

Gymnotus albus Pallas, loc. cit., 36, Surinam.

Gymnotus putao Lacépède, Hist. Nat. Poiss., II, 1801, 176, Brazil.

Genus 269. APTERONOTUS Lacépède.

Apteronotus Lacépède, Hist. Nat. Poiss., II, 1800, 208 (*A. passan* Lacépède=*Gymnotus albifrons* L.).

Sternarchus Schneider, Syst. Ichth., Bloch, 1801, 497 (*Gymnotus albifrons* L.).

Memarchus Kaup in Duméril, Ichth. Anal., 1856, 201 (*Gymnotus albifrons* L.).

706. *Apteronotus rostratus* (Meek and Hildebrand).

Streams of Panama, in Atlantic drainage, and of Colombia (Rio Cauca).

Sternarchus rostratus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 85, Rio Grande near Cana, Panama.

Genus 270. STERNOPTYGUS Müller and Troschel.

Sternopygus Müller and Troschel, Horæ Zool., III, 1849, 13 (*Gymnotus macrurus* Bloch and Schneider).

Carapus Cuvier, Règne Anim., Ed. I, 1817, 237 (*Gymnotus macrourus* Bloch and Schneider); not *Carapus* Rafinesque, 1810.

Carapo Oken, Isis, 1817, 1183; after Cuvier.

Giton Kaup, in Duméril, Ichth. Anal., 1856, 201 (*Gymnotus fasciatus* Pallas).

707. *Sternopygus dariensis* Meek and Hildebrand.

Streams of Panama, Atlantic slope.

Sternopygus dariensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 309, Rio Tuyra at Marrigante.

Genus 271. HYPOPOMUS Gill.

Hypopomus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 152 (*Rhamphichthys mulleri* Kaup).

Brachyrhamphichthys Günther, Cat., VIII, 1890, 6 (*Rhamphichthys artemis* Kaup).

708. *Hypopomus brevirostris* (Steindachner).

Panama, on both slopes to Brazil and Rio de la Plata.

Rhamphichthys brevirostris Steindachner, Sitzungs., Akad. Wiss. Wien, LVIII, 1868, 254, fig. 2, Garapore, Amazon, at Santarem.

Hypopomus occidentalis Regan, Ann. Mag. Nat. Hist., XIV, 1914, (8), 32, Condoto River, Colombia.

Genus 272. EIGENMANNIA Jordan and Evermann.

Cryptops Eigenmann, Ann. N. Y. Acad. Sci., VII, 1894, 626 (*Sternopygus humboldti* Steindachner); name preoccupied.

Eigenmannia Jordan and Evermann, Fish. North and Mid. Amer., 1896, 341 (*Sternopygus humboldti* Steindachner=*Sternarchus virescens* Valenciennes).

709. *Eigenmannia virescens* (Valenciennes).

Brazil, north in Atlantic drainage to Panama.

Sternarchus virescens Valenciennes, D'Orbigny Voyage, V, 1847, 11.

Sternopygus lineatus Müller and Troschel, Schomburgk's Reise, III, 1848, 640, British Guiana.

Sternopygus tumifrons Müller and Troschel, Horæ Zool., III, 1849, 14.

Sternopygus humboldti Steindachner, Denkschr. Akad. Wiss. Wien, 1878, 55, Rio Magdalena.

Order X. EVENTOGNATHI. Carps

Family 89. CATOSTOMIDÆ. Suckers and buffalofishes

Genus 273. CYCLEPTUS Rafinesque. *Blackhorse*.

Cycleptus Rafinesque, Jour. Phys., LXXXVIII, 1819, 421 (*C. nigrescens* Rafinesque = *Catostomus elongatus* LeSueur).

Rhytidostomus Heckel, Fisch. Syriens, in Russegger's Reisen, 1842, 1023 (*Catostomus elongatus* LeSueur).

710. *Cycleptus elongatus* (LeSueur). *Blackhorse*; *Gourdseed sucker*; *Missouri sucker*; *Suckerel*.

Mississippi valley; rather common in large streams.

Catostomus elongatus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 103, Ohio River.

Cycleptus nigrescens Rafinesque, Jour. Phys., LXXXVIII, 1819, 421, Ohio River.

Genus 274. MEGASTOMATOBUS Fowler. *Big-mouthed buffalofish*.

Megastomatobus Fowler, Proc., Acad. Nat. Sci. Phila., LXV, 1913, 45 (*Sclerognathus cyprinella* Cuvier and Valenciennes, the chief de filo of these authors for *Sclerognathus* being *Sclerognathus cyprinus*).

Ichthyobus Agassiz, Amer. Jour. Sci. Arts, 1855, 192 (*I. bubalus* Agassiz; not *Catostomus bubalus* Rafinesque; revised spelling of *Ictiobus*).

711. *Megastomatobus cyprinella* (Cuvier and Valenciennes). *Redmouth buffalofish*; *Common buffalofish*; *Bigmouth buffalo*; *Gourdseed buffalo*. Mississippi Valley to Red River of the North; generally abundant in larger streams.

Sclerognathus cyprinella Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 355 (477), Lake Pontchartrain, La.

Genus 275. ICTIOBUS Rafinesque. *Buffalofishes*.

Ictiobus Rafinesque, Ichth. Ohiensis, 1820, 55 (*Ambiodon bubalus* Rafinesque).

Bubalichthys Agassiz, Amer. Jour. Sci. Arts, XVIII, 1855, 192 (*Carpiodes urus* Agassiz).

Subgenus BUBALICHTHYS Agassiz.

712. *Ictiobus urus* (Agassiz). *Mongrel buffalo*; *Round buffalo*; *Black buffalo*, *Chopper*.

Mississippi Valley; less common than the others in the large streams; the largest and least common of the buffalo fish.

?*Carpiodes niger* Rafinesque, Ichth. Ohiensis, 1820, 56, Ohio River; may be *M. cyprinella*.

Carpiodes urus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 355, Tennessee River, Ala.

Bubalichthys urus Agassiz, loc. cit., Ohio River.

?*Bubalichthys niger* Agassiz, loc. cit., 193, Ohio River.

?*Bubalichthys bonasus* Agassiz, loc. cit., 195, Osage River, Mo.

Subgenus ICTIOBUS Rafinesque.

713. *Ictiobus bubalus* (Rafinesque). *Smallmouth buffalo*; *Razor-backed buffalo*; *Suckermouth buffalo*; *Quillback buffalo*; *Brown buffalo*; *Rooter*. Mississippi Valley and southward; generally abundant.

?*Ambiodon bubalus* Rafinesque, Jour. Phys., LXXXVIII, 1819, 421, Ohio River.

?*Carpiodes taurus* Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 355, Mobile River.

?*Carpiodes vitulus* Agassiz, loc. cit., 356, Wabash River at New Harmony.

Ichthyobus rauchi Agassiz, loc. cit., XIX, 1855, 192.

Ichthyobus cyanellus Nelson, Bull., Ill. Lab. Nat. Hist., I, 1876, 49, Illinois River.

Bubalichthys altus Nelson, in Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 73, Cairo, Ill.

Bubalichthys bubalinus Jordan, Bull. U. S. Nat. Mus., IX, 1877, 50, Ohio River.

- Genus 276. CARPIODES Rafinesque.** *Carp suckers.*
Carpiodes Rafinesque, Ichth. Ohnensis, 1820, 56 (*Catostomus cyprinus* LeSueur).
Sclerognathus Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 472 (*Catostomus cyprinus* and *Sclerognathus cyprinella* Cuvier and Valenciennes; restricted to the former by Agassiz, 1854, to the latter by Gunther, 1868).
- 714. *Carpiodes cyprinus* (LeSueur).** *Carp sucker; American carp; Carp; Quillback.*
Streams about Chesapeake Bay; common in the Chesapeake and Potomac, rare in the Delaware.
Catostomus cyprinus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 91, Elk River and other tributaries of Chesapeake Bay.
Carpiodes vacca Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 356, Susquehanna River.
- 715. *Carpiodes carpio* (Rafinesque).** *Carp sucker; Common river carp; Sucker.*
Ohio Valley and southwest to central Texas; generally abundant and variable.
Catostomus carpio Rafinesque, Ichth. Ohnensis, 1820, 56, falls of the Ohio River.
Carpiodes nummifer Cope, Proc. Amer. Phil. Soc., XI, 1870, 484, Wabash River.
- 716. *Carpiodes cutisanterinus* Cope.**
Allegheny Basin.
Carpiodes cutisanterinus Cope, Proc., Amer. Phil. Soc. Phila., XI, 1870, 481, Kiskiminitas River, Pennsylvania (Allegheny Basin).
Carpiodes selene Cope, loc. cit., 1870, 41; supposed to be from Root River, Mich.
- 717. *Carpiodes difformis* Cope.** *Blunt-nosed river carp; Sucker.*
Ohio valley and westward; generally common.
Carpiodes difformis Cope, Proc., Amer. Phil. Soc., XI, 1870, 480, Wabash River.
- 718. *Carpiodes bison* Agassiz.**
Osage River.
Carpiodes bison Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 356, Osage River, Mo.
- 719. *Carpiodes thompsoni* Agassiz.** *Lake carp; Drum; Shad* (Sandusky, Ohio).
Great Lakes region from Lake Champlain to Lake of the Woods, chiefly northward.
Carpiodes thompsoni Agassiz, Amer. Jour. Sci. Arts, XVIII, 1855, 191, Lake Champlain.
- 720. *Carpiodes velifer* (Rafinesque).** *Silver carp; Quillback; Spearfish; Sailfish; Skimback; Sailing sucker; Sailorfish.*
Mississippi Valley and westward to the upper Missouri; more common northward.
Catostomus velifer Rafinesque, Ichth. Ohnensis, 1820, 56, Ohio.
Carpiodes damalis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 170, Milk River, Mont.
Carpiodes grayi Cope, Proc., Amer. Phil. Soc., XI, 1870, 482, Western States.
- 721. *Carpiodes tumidus* Baird and Girard.** *Metalote; Buffalo.*
Lowland streams and lagoons, from the Rio Grande to the Rio Panuco and Tampico
Carpiodes tumidus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 28, Rio Grande at Fort Brown, Tex.; perhaps not separable from *C. velifer*.
- 722. *Carpiodes labiosus* Meek.**
Streams of eastern Mexico.
Carpiodes labiosus Meek, Field Col. Mus., Zool., V, 1904, 29, fig. 6, Valles, San Luis Potosi, Mexico.

- 723. *Carpiodes elongatus* Meek.**
Streams of the east coast of Mexico.
Carpiodes elongatus Meek, Field Col. Mus., Zool., V, 1904, 28, fig. 5,
Rio San Juan, Rio Conchos, and Rio Sarto Maruta in Tamaulipas,
Linares, Nueva Leon.
- 724. *Carpiodes microstomus* Meek.**
Tributaries of the Rio Grande, on the Mexican plateau.
Carpiodes microstomus Meek, Field Col. Mus., Zool., V, 1904, 27, fig.
4, Rio Conchos in Chihuahua.
- 725. *Carpiodes meridionalis* (Günther).**
Southern Mexico and Guatemala.
Sclerognathus meridionalis Günther, Cat., VII, 1868, 23, Rio Usumacinta, Guatemala.
- Genus 277. NOTOLEPIDOMYZON Fowler.**
Notolepidomyzon Fowler, Proc. Acad. Nat. Sci. Phila., LXV, 1913, 47
(*Pantosteus arizonæ* Gilbert=*Catostomus clarki* Baird and Girard).
- 726. *Notolepidomyzon plebeius* (Baird and Girard).**
Basin of the Rio Grande, Colorado to Chihuahua; Rio Piedras, Rio Casas Grandes; very common.
Catostomus plebeius Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 28, Rio Mimbres, a tributary of Lake Guzman, Chihuahua, Mexico.
Catostomus (Acomus) guzmaniensis Girard, loc. cit., VIII, 1856, 173, Rio Janos, tributary to Lago de Guzman, Chihuahua, Mexico.
Catostomus nebuliferus Garman, Bull. Mus. Comp. Zool., VIII, 1881, 89, Rio Nazas, Coahuila, Mexico.
- 727. *Notolepidomyzon clarkii* (Baird and Girard).**
Rio Gila, Salt River, and their tributaries in Arizona.
Catostomus clarkii Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 27, Rio Santa Cruz, Gila Basin, Ariz.
Pantosteus arizonæ Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 170, Salt River, Tempe, Ariz.
- 728. *Notolepidomyzon santæ-anæ* (Snyder).**
Streams of southern California.
Pantosteus santæ-anæ Snyder, Proc. U. S. Nat. Mus., XXXIV, 1908, 33, Santa Ana River, at Riverside, Calif.
- 729. *Notolepidomyzon generosus* (Girard). *Mountain sucker*.**
Colorado Basin.
Catostomus (Acomus) generosus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 174, Cottonwood Creek; not in Utah, "no doubt from Colorado" (Snyder).
- Genus 278. PANTOSTEUS Cope. *Mountain suckers*.**
Pantosteus Cope, Wheeler's Surv., V, 1875, 673 (*Minomus platyrhynchus* Cope).
- 730. *Pantosteus delphinus* (Cope). *Bluehead sucker*.**
Mountain streams tributary to the Colorado River in western Colorado and Wyoming; very abundant.
Minomus delphinus Cope, Hayden's Geol. Surv. Wyo., 1870, 435; probably Henry Fork of Green River, Wyo.
? *Minomus bardus* Cope, loc. cit., 436; probably Green River, Wyo.
- 731. *Pantosteus platyrhynchus* (Cope).**
Streams of Utah Basin (Provo and Sevier Rivers).
Minomus platyrhynchus Cope, Proc. Am. Phil. Soc., XIV, 1874, 134, Utah Lake at Provo.
? *Minomus jarrovi* Cope, loc. cit., 135; "near Provo"; the type lost.
- 732. *Pantosteus jordani* Evermann.**
Upper Missouri and Columbia River Basins.
Pantosteus jordani Evermann, Bull. U. S. Fish Com., XII, 1892 (Jan. 27, 1893), 51, fig. on p. 53; Whitewood Creek and other streams of the Black Hills, S. Dak.

- 733. *Pantosteus columbianus* Eigenmann.**
Columbia Basin.
Pantosteus columbianus Eigenmann, Amer. Nat., Feb. 4, 1893, 151,
Columbia River; perhaps identical with *Pantosteus jordani*.
- 734. *Pantosteus virescens* Cope.**
Tributaries of Lake Bonneville basin of Utah; abundant in Weber and
Bear Rivers; largest of the hard-mouthed or mountain suckers.
Pantosteus virescens Cope, Wheeler's Surv., Zool., 1875, 675; by error
ascribed to Arkansas River at Pueblo; must have come from the
basin of Utah.
- 735. *Pantosteus lahontan* Rutter.**
Streams of western Nevada; Lahontan Basin.
Pantosteus lahontan Rutter, Bull. U. S. Fish Com., XXII, 1902, 146,
with figure; Susan River, a tributary of Honey Lake, Little Truckee
River, Prosser Creek; misprint for *lahontan*.
- Genus 279. CATOSTOMUS LeSueur. Fine-scaled sucker.**
Catostomus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 89 (*Cyprinus*
catostomus Forster).
Decactylus Rafinesque, Ichth. Ohiensis, 1820, 60 (*Catostomus bostoniensis*
Rafinesque = *Cyprinus commersoni* Lacépède).
Stomocatus Bonaparte, Fauna Italica, 1840, Fasc. 129 (*Cyprinus catosto-*
mus Forster).
Minomus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 173 (*Catos-*
tomus insignis Baird and Girard).
Acomus Girard, loc. cit., 173 (*Catostomus forsterianus* Richardson =
Cyprinus catostomus Forster).
- Subgenus DECACTYLUS Rafinesque.**
- 736. *Catostomus commersonii* (Lacépède).** *Common sucker; White sucker;*
Brook sucker; Fine-scaled sucker, Black mullet (Sandusky, Ohio); Fine-
scaled mullet; Bay mare.
Quebec and the Great Lakes to Montana, Colorado, southward to
Missouri and Georgia; the commonest of the suckers, excessively
abundant from Massachusetts west to Kansas
Cyprinus commersonii Lacépède, Hist. Nat. Poiss., V, 1803, 503, locality
unknown; wrongly accredited to Commerson, the description and
figure drawn from an old example long in alcohol
Cyprinus teres Mitchell, Trans. Lit. Phil. Soc. N. Y., 1815, 458, New
York.
Catostomus communis LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817,
95, 106, Delaware River.
Catostomus bostoniensis LeSueur, loc. cit., 106, Boston.
Catostomus flexuosus Rafinesque, Ichth. Ohiensis, 1820, 59, Ohio River.
Catostomus reticulatus Richardson, Fauna Bor.-Amer., Fishes, 1836,
303, Albany River.
Catostomus gracilis Kirtland, Rept. Zool. Ohio, 1838, 169, Cleveland,
Ohio
Catostomus pallidus DeKay, New York Fauna: Fishes, 1842, 200, New
York.
Catostomus sucklii Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 175,
Milk River, Mont.
Catostomus chloropteron Abbott, loc. cit., XII, 1860, 473, Kansas.
? *Catostomus alticola* Cope, Proc. Amer. Phil. Soc., XIV, 1874, 138,
Twin Lake, Colo.
Moxostoma trisignatum Cope, Wheeler's Surv., V, Zool., 1875, 679,
Arkansas River.
? *Catostomus utawana* Mather, Twelfth Rept., N. Y. Fish Com., Surv.
of Adirondacks, 1884, 36, Blue Mountain Lakes, N. Y.; dwarf form.
- 737. *Catostomus ardens* Jordan and Gilbert.** *Mullet of Utah Lake.*
Great basin of Utah and upper Snake River; Idaho; locally abundant.
Catostomus ardens Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880,
464, Utah Lake, Provo, Utah.

738. *Catostomus occidentalis* Ayres. *Sacramento sucker*.
Streams of northern and central California and southern Oregon.
Catostomus occidentalis Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 18,
San Francisco.
Catostomus labiatus Ayres, loc. cit., 1855, 32, San Joaquin River, Stock-
ton, Calif.
Catostomus aræopus Jordan, Bull. U. S. Nat. Mus., XII, 1878, 173,
South Fork of Kern River, Calif.
739. *Catostomus tsilcoosensis* Evermann and Meek.
Coastal streams of Oregon.
Catostomus tsilcoosensis Evermann and Meek, Bull. U. S. Fish Com.,
XVII, 1897, 66, fig. 1, Tsilcoos Lake, Lane Co., Ore.
740. *Catostomus snyderi* Gilbert.
Klamath Basin, California and Oregon.
Catostomus snyderi Gilbert, Bull. U. S. Fish Com., XVII, 1897 (Jan. 6,
1898), 3, Upper Klamath Lake, Ore.
741. *Catostomus microps* Rutter.
Streams of the lava beds of California.
Catostomus microps Rutter, Bull. U. S. Bur. Fish., XXVII, 1907 (1908),
120, fig. 1; Rush Creek, a tributary of Ash Creek, Aden, Modoc Co.,
Calif.
742. *Catostomus macrocheilus* Girard. *Columbia River sucker*; *Bighorn sucker*.
Columbia River Basin, as far up as Flathead Lake, Montana; Upper
Salmon Falls and the headwaters of Salmon River, Idaho.
Catostomus macrocheilus Girard, Proc. Acad. Nat. Sci. Phila., VIII,
1856, 175, Astoria, Ore.
743. *Catostomus insignis* Baird and Girard.
Gila Basin, Ariz.
Catostomus insignis Baird and Girard, Proc. Acad. Nat. Sci. Phila.,
VII, 1854, 28, Rio San Pedro, Ariz.
Catostomus gila Kirsch, Proc. U. S. Nat. Mus., XI, 1888, 555, Rio Gila,
Fort Thomas, Ariz.
744. *Catostomus bernardini* Girard.
San Bernardino Creek, a tributary of Rio Yaqui, on the boundary of
Arizona and Sonora.
Catostomus bernardini Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
175, San Bernardino Creek, Ariz.
745. *Catostomus sonoriensis* Meek.
Rivers of Sonora on the Pacific slope.
Catostomus sonoriensis Meek, Field Col. Mus., Zool., Ser. III, No. 6,
May, 1902, 75, Pl. XVI, Rio Paphigochic, Miñaca, Chihuahua,
Mexico.
- Subgenus CATOSTOMUS LeSueur.
746. *Catostomus catostomus* (Forster). *Long-nosed sucker*; *Northern sucker*;
Red sucker; *Black sucker* (Sandusky, Ohio); *Sturgeon sucker*.
Great Lakes; upper Missouri River; Ohio Basin; West Virginia;
upper Columbia; eastward to Maine and northwestward to Alaska;
very abundant northward.
Cyprinus catostomus Forster, Philos. Trans., LXIII, 1773, 155, tab. 6,
streams about Hudson Bay.
Catostomus longirostrum and *hudsonius* LeSueur, Jour. Acad. Nat. Sci.
Phila., I, 1817, 102, 107, Vermont; after Forster.
Catostomus forsterianus Richardson, Franklin's Jour., 1823, 720, Lake
Huron; Great Slave Lake.
Catostomus aurora Agassiz, Lake Superior, 1850, 360, Lake Superior.
Catostomus nanomyzon Mather, Twelfth Rep., N. Y. Fish Com., Surv.
of Adirondacks, 1884, 36, Big Moose Lake, northern New York;
dwarf form.
747. *Catostomus richardsoni* Harper and Nichols. *Gray sucker*.
Streams of northwestern Canada.
Catostomus richardsoni Harper and Nichols, Bull. Amer. Mus. Nat.
Hist., XLI, 1919, 263, Pl. XV, fig. 1, type locality not designated.

748. *Catostomus griseus* Girard.
Platte River and upper Missouri region.
Catostomus (Acomus) griseus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 174, Sweetwater Fork of Platte River.
? *Catostomus (Acomus) lactarius* Girard, loc. cit., 174, Milk River, Mont.
749. *Catostomus retropinnis* Jordan.
Northern Montana.
Catostomus retropinnis Jordan, Bull. U. S. Nat. Mus., XII, 1878, 178, Milk River, Mont.
750. *Catostomus pocatello* Gilbert and Evermann. *Moo-gad-ee of the Fort Hall Indians.*
Upper Snake River.
Catostomus pocatello Gilbert and Evermann, Bull., U. S. Fish Com., XIV, 1894, 189, pl. 21, fig. 1, Ross fork of Snake River, near Pocatello, Idaho.
751. *Catostomus tahoensis* Gill and Jordan. *Tahoe sucker.*
Lake Tahoe and Humboldt River, Nev.
Catostomus tahoensis Gill and Jordan, in Jordan, Bull., U. S. Nat. Mus., XII, 1878, 173, Lake Tahoe, Calif.
752. *Catostomus discobolus* Cope.
Upper Colorado Basin.
Catostomus discobolus Cope, Hayden's Geol. Surv. Wyo., 1870 (1872), 435, Green River, Wyo.
753. *Catostomus chamberlaini* (Rutter).
Chasmistes chamberlaini Rutter, Bull. U. S. Fish Com., XXII, 1902 (1904), 147, Eagle Lake (a lake without outlet in the Lahontan Basin); apparently a variant of *C. tahoensis*.
754. *Catostomus latipinnis* Baird and Girard. *Flannel-mouthed sucker.*
Rio Colorado and Rio Gila and their tributaries.
Catostomus latipinnis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 388, Rio San Pedro, Gila Basin, Ariz.
755. *Catostomus conchos* Meek.
Catostomus conchos Meek, Field Col. Mus., Zool., III, 1902, 75, Rio Conchos, Jimenez, Chihuahua, Mexico.
- Genus 280. **HYPENTELIUM** Rafinesque.
Hypentelium Rafinesque, Jour. Acad. Nat. Sci. Phila., II, 1818, 421 (*H. macroplectum* Rafinesque = *Catostomus nigricans* LeSueur).
Eurystomus Rafinesque, Ichth. Ohiensis, 1820, 59 (*Catostomus megastomus* Rafinesque), based on an incorrect drawing of *Hypentelium nigricans*.
Hylomyzon Agassiz, Amer. Jour. Sci. Arts, 1855, 205 (*Catostomus nigricans* LeSueur).
756. *Hypentelium nigricans* (LeSueur). *Black sucker; Mud sucker; Hog sucker; Stoneroller; Toter; Crawl-a-bottom; Hammerhead; Stonelugger; Hogmolly; Bootleg sucker; Spotted sucker; Niggerhead.*
New York to Minnesota, Kansas, Arkansas, and the Carolinas; abundant in swift or rocky streams.
Catostomus nigricans LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 102, Lake Erie.
Catostomus maculosus LeSueur, loc. cit., 103, Pipe Creek, Md.
Hypentelium macropteryum Rafinesque, loc. cit., I, 1817, 420, Ohio River.
Catostomus xanthopus Rafinesque, Ichth. Ohiensis, 1820, 57, Ohio River.
Catostomus nigricans etowanus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876 (1877), 345, Etowah River, Rome, Ga.
Catostomus megastomus Rafinesque, Ichth. Ohiensis, 1820, 59; based on incorrect drawing.
Catostomus planiceps Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 334 (450), 1844, Wabash River.
- Genus 281. **THOBUERNIA** Jordan and Snyder.
Thoburnia Jordan and Snyder, Copeia, October 4, 1917, 88 (*Catostomus rhothæcus* Thoburn).

757. *Thoburnia rathœca* (Thoburn).
Headwaters of Tennessee River.
Catostomus rathœcus Thoburn, in Jordan and Evermann, Fishes North and Mid Amer., 1896, 181, upper tributaries of Tennessee River; supposed to be from the French Broad River at Wolf Creek, Tenn.
- Genus 282. CHASMISTES Jordan.
Chasmistes Jordan, Bull., Hayden's Geol. Surv. Terr., IV, No. 2, 1878, 417 (*C. fecundus* Jordan; not *Catostomus fecundus* Cope and Yarrow = *Chasmistes lorus* Jordan).
Lipomyzon Cope, Amer. Nat., XV, 1881, 59 (*Chasmistes brevirostris* Cope).
Pithecomyzon Fowler, Proc. Acad. Nat. Sci. Phila., LXV, 1913, 54 (*Chasmistes cujus* Cope).
- Subgenus LIPOMYZON Cope.
758. *Chasmistes fecundus* (Cope and Yarrow). *Webug sucker*.
Utah Lake.
Catostomus fecundus Cope and Yarrow, Wheeler's Surv., Zool., V, 1875, 678, Utah Lake.
759. *Chasmistes brevirostris* Cope.
Klamath Lake, Oreg.
Chasmistes brevirostris Cope, Amer. Nat., XIII, 1879, 785, Klamath Lake, Oreg.
760. *Chasmistes stomias* Gilbert.
Upper Klamath Lake, Oreg.
Chasmistes stomias Gilbert, Bull., U. S. Fish Com., XVII, 1897 (1898), 5, text figure, Upper Klamath Lake, Oreg.
761. *Chasmistes lorus* Jordan. *June sucker*.
Utah Lake.
Chasmistes lorus Jordan, Bull., U. S. Nat. Mus., XII, 1878, 249, Utah Lake at Provo, Utah.
762. *Chasmistes cujus* Cope. *Cousa*; *Kweewee*; *Quirwee*.
Pyramid Lake, Nevada; locally abundant.
Chasmistes cujus Cope, Proc. Acad. Nat. Sci. Phila., XXXV, 1883, 149, Pyramid Lake, Nev.
763. *Chasmistes copei* Evermann and Meek. *Tswam*.
Klamath Lakes, Oreg.
Chasmistes copei Evermann and Meek, Bull., U. S. Fish Com., XVII, 1894 (1898), 70, fig. 3; Pelican Bay, Upper Klamath Lake, Oreg.
- Genus 283. DELTISTES Seale.
Deltistes Seale, Proc. Calif. Acad. Sci., ser. 2, VI, 1896, 269 (*Chasmistes luxatus* Cope).
764. *Deltistes luxatus* (Cope). *Tswam*; *Lost River sucker*.
Klamath Lake and River, Oreg.; abundant.
Chasmistes luxatus Cope, Amer. Nat., XIII, 1879, 784, Klamath Lake, Oreg.
Catostomus rex Eigenmann, loc. cit., XXV, July, 1891, 667, Lost River, Oreg.
- Genus 284. XYRAUCHEN Eigenmann and Kirsch. *Razorback suckers*.
Xyrauchen Eigenmann and Kirsch, Proc. U. S. Nat. Mus., XI, 1888, 556 (*Catostomus cypho* Lockington = *Catostomus texanus* Abbott).
765. *Xyrauchen texanus* (Abbott). *Razorback sucker*; *Humpback sucker*.
Basin of the Colorado and Gila Rivers (not in Texas).
Catostomus texanus Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 473, Colorado and New Rivers, Ariz.
Catostomus cypho Lockington, loc. cit., 1880, 237, Colorado River at mouth of the Gila, Ariz.
Xyrauchen uncomphagre Jordan and Evermann, Bull. U. S. Fish Com., 1889, 26, Uncomphagre River delta, Colo.; young example, probably the same.
- Genus 285. ERIMYZON Jordan. *Chub suckers*.
Erimyzon Jordan, Bull. Buffalo Soc. Nat. Hist., III, 1876, 95 (*Cyprinus oblongus* Mitchill).

- 766. *Erimyzon sucetta* (Lacépède).** *Chub; Chub sucker; Creekfish; Creek sucker.*
Coastwise streams, Pennsylvania and south; very abundant southward.
Cyprinus sucetta Lacépède, Hist. Nat. Poiss., V, 1803, 606, South Carolina.
Moxostoma kennerlyi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 171, Dry Creek, Victoria, Tex.
Moxostoma campbelli Girard, loc. cit., 172, Live Oak Creek and Devil River, Tex.
Moxostoma tenue Agassiz, Amer. Jour. Sci. Arts, 1855, 203, Mobile, Ala.
Erimyzon goodii Jordan, Bull. U. S. Nat. Mus., XII, 1878, 148, St. Johns River, Fla.
- 767. *Erimyzon oblongus* (Mitchill).** *Chub sucker; Sweet sucker; Mullet; Creek fish; Pin minnow.*
Great Lakes region to Maine and the Dakotas, south to Virginia and Oklahoma; very abundant; replacing the deeper-bodied *sucetta* northward; the two probably intergrading.
Cyprinus oblongus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 459, New York.
Catostomus gibbosus LeSueur, Connecticut River, Northampton; *tuberculatus*, Germantown, Pa.; and *vittatus*, Wissahickon River, Pa.; Jour. Acad. Nat. Sci. Phila., I, 1817, 92, 93, 104.
Catostomus fasciolaris Rafinesque, Ichth. Ohiensis, 1820, 58, Ohio River.
Labeo elegans, New York; *esopus*, New York; and *elongatus*, Mohawk River; DeKay, N. Y. Fauna: Fishes, 1842, 192, 195, 394.
Moxostoma claviformis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 171, Coal Creek, Canadian River, Indian Territory [Okla.]
- Genus 286. MINYTREMA** Jordan. *Spotted suckers.*
Mynitrema Jordan, Man. Vert., ed. 2, 1878, 318 (*Catostomus melanops* Rafinesque).
- 768. *Minytrema melanops* (Rafinesque).** *Spotted sucker; Striped sucker; Winter sucker; Sand sucker; Black winter sucker.*
Great Lakes region to North Carolina and west to Texas and Montana; rather common westward.
Catostomus melanops Rafinesque, Ichth. Ohiensis, 1820, 57, Ohio River.
Catostomus fasciatus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 333 (449), Mississippi River.
Moxostoma victorix Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 171, Dry Creek, Victoria, Tex.
Ptychostomus haydeni Girard, loc. cit., 172, Yellowstone River; Missouri River at Fort Pierre, Neb.
- Genus 287. MOXOSTOMA** Rafinesque. *Redhorse.*
Moxostoma Rafinesque, Ichth. Ohiensis, 1820, 54 (*Catostomus anisurus* Rafinesque).
Teretulus Rafinesque, loc. cit., 57 (*Catostomus aureolus* LeSueur).
Ptychostomus Agassiz, Amer. Jour. Sci. Arts., XIX, 1855, 203 (*Catostomus aureolus* LeSueur).
Myxostoma Jordan; corrected orthography.
- 769. *Moxostoma papillosum* (Cope).** *White mullet.*
Lowland streams, from the Dismal Swamp in Virginia to the Ocmulgee River, Ga.
Ptychostomus papillosus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 470, Catawba and Yadkin Rivers, N. C.
- 770. *Moxostoma anisurum* (Rafinesque).** *White-nosed sucker.*
Ohio River, Great Lakes region and Lake of the Woods; not very common, but widely distributed.
Catostomus anisurus Rafinesque, Ichth. Ohiensis, 1820, 54, Ohio River.
Catostomus carpio Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 457, Lake Ontario, not of Rafinesque.
Ptychostomus velatus Cope, Proc. Amer. Phil. Soc., XI, 1870, 471, Youghiogheny River, Pa.
Moxostoma valenciennesi Jordan, Proc. U. S. Nat. Mus., VIII, 1885, 73; substitute for *Catostomus carpio* Cuvier and Valenciennes; preoccupied.

771. *Moxostoma bucco* (Cope).
Missouri River at St. Joseph.
Ptychostomus bucco Cope, Hayden's Geol. Surv. Wyo., 1870, 437, St. Joseph, Mo.; a doubtful species allied to *M. anisurum*.
772. *Moxostoma pidiense* (Cope).
Great Pedee Basin, the Carolinas.
Ptychostomus pidiensis Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 471, Yadkin River, N. C.
773. *Moxostoma collapsum* (Cope).
Lowland streams of North Carolina.
Ptychostomus collapsus Cope, Proc. Amer. Phil. Soc., XI, 1870, 471, Neuse, Yadkin, and Catawba Rivers, N. C.
774. *Moxostoma coregonus* (Cope). *Blue mullet*.
Catawba and Yadkin Rivers, N. C.
Ptychostomus coregonus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 472, Catawba and Yadkin Rivers
775. *Moxostoma album* (Cope).
Rivers of North Carolina.
Ptychostomus albus Cope, Proc., Amer. Phil. Soc. Phila., XI, 1870, 472, Catawba River, N. C.
776. *Moxostoma thalassinum* (Cope).
Yadkin River, N. C.
Ptychostomus thalassinus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 472, Yadkin River, N. C.
777. *Moxostoma congestum* (Baird and Girard). *Texas redhorse*.
Streams of Texas.
Catostomus congestus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 27, Rio Salado, Tex.
Ptychostomus albidus Girard, loc. cit., VIII, 1856, 172, Rio San Juan, Monterey, Nuevo Leon, Mexico.
778. *Moxostoma austrinum* (Bean).
Rio de Santiago, Mexico; known from Michoacan, Mexico (west of Sierra Madre); Rio Lerna at Salamanca.
Myxostoma austrina Bean, Proc. U. S. Nat. Mus., II, 1879, 302, Piedad in Morelia, Michoacan, Mexico.
779. *Moxostoma mascotæ* Regan.
Streams of Jalisco (Pacific drainage), Mexico.
Moxostoma mascotæ Regan, Biol. Centrali-Amer., 1907, 147, Tab. XXIV, fig. 2, Rio Mascota, Mexico.
780. *Moxostoma parvidens* Regan.
Rio Grande Basin.
Moxostoma parvidens Regan, Biol. Centrali-Amer., 1907, 147, Tab. XXIV, fig. 3, Rio Grande at Juarez, Mexico.
781. *Moxostoma aureolum* (LeSueur). *Common redhorse; Mullet; White sucker; Large-scaled sucker; Sucker*.
Lake Ontario and Lake Michigan to the Missouri River and Lake of the Woods, south to Arkansas and Georgia; very abundant west of Alleghenies.
Catostomus aureolus LeSueur, Jour., Acad. Nat. Sci. Phila., I, 1817, 95, Lake Erie near Buffalo.
Catostomus duquesnii LeSueur, loc. cit., I, 105, 1817, Ohio River at Pittsburgh, Pa.
Catostomus erythurus Rafinesque, Amer. Month. Mag., III, 1818, 354, 355, Ohio River.
Catostomus oneida DeKay, N. Y. Fauna: Fishes, 1842, 198, Oneida Lake.
Rutilus melanurus Rafinesque, Ichth. Ohiensis, 1820, 51, Ohio River.
Myxostoma euryops Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 348, Oostanaula River, Rome, Ga.; probably a monstrosity of this species.

782. *Moxostoma lesueurii* (Richardson). *Picconou*.
Tributaries of Hudson Bay and westward.
Catostomus lesueurii Richardson, Franklin's Jour., 1823, 1772, Pine Island Lake, British America.
783. *Moxostoma macrolepidotum* (LeSueur). *Mullet*; *Red mullet*; *Eastern redhorse*; *River sucker*.
Streams about Chesapeake and Delaware Bays and southward to North Carolina; east of the Alleghenies.
Catostomus macrolepidotus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 94, Delaware River.
Ptychostomus lachrymans Cope, Proc., Amer. Phil. Soc., XI, 1870, 474, Neuse River at Newbern, N. C.
784. *Moxostoma robustum* (Cope).
Yadkin River, N. C.
Ptychostomus robustus Cope, Proc. Amer. Phil. Soc., XI, 1870, 473, Yadkin River, N. C.
785. *Moxostoma crassilabre* (Cope).
Streams of eastern North Carolina.
Ptychostomus crassilabris Cope, Proc. Amer. Phil. Soc., XI, 1870, 477, Neuse River near Raleigh, N. C.
786. *Moxostoma conus* (Cope).
Neuse and Yadkin Rivers, N. C.
Ptychostomus conus Cope, Proc. Amer. Phil. Soc., XI, 1870, 478, Yadkin River, N. C.
787. *Moxostoma breviceps* (Cope). *Short-headed redhorse*.
Ohio Valley and Great Lakes region; abundant in Lake Erie.
Ptychostomus breviceps Cope, Proc. Amer. Phil. Soc., XI, 1870, 478, Youghiogheny River, Pa.
788. *Moxostoma pæcilurum* (Jordan).
Southern Mississippi to eastern Texas; Pearl, Tangipahoa, and Sabine Rivers.
Myxostoma pæcilura Jordan, Bull. U. S. Nat. Mus., X, 1877, 66, Tangipahoa River, La.
789. *Moxostoma alleghaniensis* Nichols.
Mountain streams of North Carolina.
Moxostoma alleghaniensis Nichols, Bull. Amer. Mus. Nat. Hist., XXX, 1911, 275, Pl. XI, fig. 1, Marshall, N. C.
- Genus 288. SCARTOMYZON Fowler.
Scartomyzon Fowler, Proc. Acad. Nat. Sci. Phila., LXV, 1913, 59 (*Ptychostomus cervinus* Cope).
790. *Scartomyzon rupiscartes* (Jordan and Jenkins). *Jumprocks*.
Rivers of the Carolinas and Georgia, from the Catawba to the Chattahoochee; locally abundant.
Moxostoma rupiscartes Jordan and Jenkins, Proc. U. S. Nat. Mus., XI, 1888, 353, Catawba River at Morgantown, and various other streams, representing *Scartomyzon cervinus* southward
791. *Scartomyzon cervinus* (Cope). *Jumping mullet*; *Crawl-a-bottom*.
Rivers of the South Atlantic States, from the James to the Neuse.
Teretulus cervinus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 236, headwaters of the Roanoke and James Rivers, Va.
- Genus 289. PLACOPHARYNX Cope.
Placopharynx Cope, Proc. Amer. Phil. Soc., XI, 1870, 467 (*P. carinatus* Cope).
792. *Placopharynx carinatus* Cope. *White sucker*; *Big-toothed redhorse*.
Michigan to Tennessee, Georgia, and Arkansas; abundant in large streams, especially in the Ozarks.
Placopharynx carinatus Cope, Proc. Amer. Phil. Soc., XI, 1870, 467, Wabash River.

Genus 290. LAGOCHILA Jordan and Brayton. *Rabbitmouth suckers*.
Lagochila Jordan and Brayton, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 280, (*L. lacera* Jordan and Brayton); not *Lagocheilus*, an earlier genus of mollusks.
Quassilabra Jordan and Brayton, Man. Vert., Ed. II, 1878, 406 (*Lagochila lacera* Jordan and Brayton); substitute for *Lagochila*, regarded as preoccupied.

- 793. Lagochila lacera** Jordan and Brayton. *Harelip sucker; Cut-lips; Split-mouth sucker; May sucker; Rabbitmouth sucker; Pealip sucker*.
 Mississippi Valley; Ozark Mountains; Scioto, Tippecanoe, Clinch, Cumberland, and Chickamauga Rivers, and the White River of Arkansas; nowhere very common.
Lagochila lacera Jordan and Brayton, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 280, Chickamauga River, Ringgold, Ga.

Family 90. CYPRINIDÆ. Daces and minnows

In arranging the genera of this by far the largest natural family of fishes, we have begun the series with the forms having closest affinity with Asiatic and then with European types—these having largely five teeth in the main row on the pharyngeals. Later follow the characteristically American forms, more or less degenerate as compared with the possible European ancestors, the series closing with aberrant forms from the same stock.

Genus 291. ORTHODON Girard.
Orthodon Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 182 (*Gila microlepidota* Ayres).

- 794. Orthodon microlepidotus** (Ayres). *Blackfish*.
 Larger rivers and lakes of central and northern California; generally abundant.
Gila microlepidota Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 21, Sacramento and San Joaquin Rivers, Calif.

Genus 292. ACROCHEILUS Agassiz. *Chiselmouths*.
Acrocheilus Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 96 (*A. alutaceus* Agassiz and Pickering).

- 795. Acrocheilus alutaceus** Agassiz and Pickering. *Chiselmouth; Square-mouth; Harpmouth*.
 Lakes and rivers of southeastern Oregon; lower Columbia and tributaries as far up as Spokane and Shoshone Falls.
Acrocheilus alutaceus Agassiz and Pickering, Amer. Jour. Sci. Arts, XIX, 1855, 99, Willamette Falls and Walla Walla River, Oreg.

Genus 293. MYLOCHEILUS Agassiz.
Mylocheilus Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 229 (*M. lateralis* Agassiz and Pickering).

- 796. Mylocheilus lateralis** Agassiz and Pickering. *Columbia River chub*.
 Columbia and Frazer River Basins; ascending to Flathead Lake and Shoshone Falls.
Mylocheilus lateralis Agassiz and Pickering, Amer. Jour. Sci. Arts, 1855, 231, Columbia River at Fort Vancouver; confused by recent writers with *Leuciscus caurinus* Richardson, which is a near relative of *Richardsonius balteatus*.

- 797. Mylocheilus fraterculus** Girard.
Mylocheilus fraterculus Girard, Proc., Acad. Nat. Sci. Phila., VIII, 1856, 169, "Monterey, Calif."; a doubtful record, as no *Mylocheilus* has been yet found in California; scales about 85; 77 in *Mylocheilus lateralis*.

Genus 294. MYLOPHARODON Ayres.
Mylopharodon Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 33 (*M. robustus* Ayres=*Mylopharodon conocephalus* Girard).

798. *Mylopharodon conocephalus* (Baird and Girard). *Kaweah chub*.
Sacramento and San Joaquin Rivers, Calif.; southeastern Oregon, Pit River and Big Valley.
Gila conocephala Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 135, San Joaquin River, Calif.
Mylopharodon robustus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 33, San Joaquin River, Calif.
- Genus 295. **STYPODON** Garman.
Stypodon Garman, Bull. Mus. Comp. Zool., VIII, 1881, 90 (*S. signifer* Garman).
799. *Stypodon signifer* Garman.
Lago de Parras, an isolated lagoon in Coahuila, Mexico.
Stypodon signifer Garman, Bull. Mus. Comp. Zool., VIII, 1881, 90, Lago de Parras, Coahuila, Mexico.
- Genus 296. **CHROSOMUS** Rafinesque.
Chrosomus Rafinesque, Ichth. Ohiensis, 1820, 47 (*Luxilus erythrogaster* Rafinesque).
800. *Chrosomus erythrogaster* Rafinesque. *Red-bellied dace*.
New Brunswick to Ohio, Wisconsin, Colorado, and northern Alabama; St. Lawrence River; chiefly or entirely west of the Alleghenies, very abundant northward.
Luxilus or *Chrosomus erythrogaster* Rafinesque, Ichth. Ohiensis, 1820, 47, Ohio River.
Chrosomus pyrrhogaster Jordan, Man. Vert., Ed. I, 1876, 284, White River, Ind.
801. *Chrosomus eos* Cope.
Streams east of the Alleghenies, north to the Susquehanna; scarce.
Chrosomus eos Cope, Proc., Acad. Nat. Sci. Phila., XIII, 1861, 523, Meshoppen Creek, Susquehanna River, Pa.; perhaps a variation of *Chrosomus erythrogaster*.
802. *Chrosomus oreas* Cope.
Headwaters of Roanoke River, and clear tributaries of the Tennessee River.
Chrosomus oreas Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 233, head of Roanoke River, Montgomery County, Va.
803. *Chrosomus dakotensis* Evermann and Cox.
Chrosomus dakotensis Evermann and Cox, Rept. U. S. Fish Com., XIX, 1895, Crow Creek, Chamberlain, S. Dak.
- Genus 297. **LAVINIA** Girard.
Lavinia Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 137 (*L. exilicauda* Baird and Girard).
Luxilinus Jordan, Cat. Fishes N. Amer., 1885, 33 (*Larucosomus occidentalis* Baird and Girard); young.
804. *Lavinia exilicauda* Baird and Girard. *Hitch; Chigh*.
Streams of the coast ranges about San Francisco and Monterey; locally common as far north as Clear Lake, Calif.
Lavinia exilicauda Baird and Girard, Proc., Acad. Nat. Sci. Phila., VII, 1854, 137, Sacramento River, Calif.
Leucosomus occidentalis Baird and Girard, loc. cit., VII, 1854, 137, Poso Creek; Four Creeks, Calif.; young of *L. exilicauda*.
Lavinia harengus Girard, loc. cit., VIII, 1856, 184, Monterey, Calif.
- Genus 298. **OXYGENEUM** Forbes.
Oxygeneum Forbes, Bull. Ill. Lab. Nat. Hist., II, 1885, 136 (*O. pulverulentum* Forbes).
805. *Oxygeneum pulverulentum* Forbes.
Illinois River, Ill.
Oxygeneum pulverulentum Forbes, Bull. Ill. Lab. Nat. Hist., II, 1885, 136, Illinois River, Ill.
- Genus 299. **PTYCHOCEILUS** Agassiz. *Squawfish*.
Ptychocheilus Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 227 (*P. gracilis* Agassiz=*Cyprinus* (*Leuciscus*) *oregonensis* Richardson).

806. *Ptychocheilus oregonensis* (Richardson). *Squawfish*; *Chappaul*; *Whitefish*.
Rivers of Oregon and Washington, in the Columbia Basin to Montana and Idaho, and northward to British Columbia.
Cyprinus (Leuciscus) oregonensis Richardson, Fauna Bor.-Amer., III, 1836, 305, Columbia River.
Ptychocheilus gracilis Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 229, Willamette River.
807. *Ptychocheilus grandis* (Ayres). *Sacramento pike*; *Sacramento squawfish*.
Rivers of California and southeastern Oregon; Pit River.
Gila grandis Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 18, San Francisco.
Ptychocheilus major Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 229, San Francisco.
Ptychocheilus harfordi Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 72, Sacramento River; scales 90 to 95, a variant with smaller scales.
808. *Ptychocheilus lucius* Girard. *White salmon of the Colorado*; *Colorado River squawfish*.
Colorado River Basin, in all larger streams from Rio Uncompahgre to the Gulf of California; largest of American *Cyprinidæ*, reaching 80 pounds.
Ptychocheilus lucius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 209, Rio Colorado.
809. *Ptychocheilus umpquæ* Snyder.
Coastal streams of Oregon.
Ptychocheilus umpquæ Snyder, Bull. U. S. Bur. Fish., XXVII, 1907 (1908), 170, fig. 2, in part, Callopoia Creek, Oakland, Ore.
- Genus 300. *GILA* Baird and Girard.
Gila Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 368 (*G. robusta* Baird and Girard).
810. *Gila elegans* Baird and Girard. *Bony-tail*; *Gila trout*; *Verde trout*.
Colorado and Gila Rivers; abundant in river channels.
Gila elegans Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 369, Zuñi, Colorado, and Gila Rivers.
Gila emoryi Baird and Girard, loc. cit., 388, Gila River.
811. *Gila robusta* Baird and Girard. *Roundtail*.
Tributaries of the Rio Colorado and Rio Gila; very common up to the foot of the mountains in Colorado.
Gila robusta Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 368, Zuñi River.
Gila gracilis Baird and Girard, loc. cit., 369, Zuñi River.
Gila grahami Baird and Girard, loc. cit., 389, Rio San Pedro, tributary to Rio Gila.
Ptychocheilus vorax Girard, loc. cit., 1856, 209, locality unknown.
Leuciscus zunniensis Gunther, Cat., VII, 1868, 241, Zuñi River; substitute for *gracilis*, preoccupied in *Leuciscus*.
Gila affinis Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 474, probably erroneously ascribed to Kansas River.
Gila nacreæ Cope, Hayden's Geol. Surv. Wyo., 1870 (1871), 441, tributary of Green River, Wyo.; young, perhaps of *G. elegans*.
812. *Gila seminuda* Cope and Yarrow.
Rio Virgen, in Utah and Nevada.
Gila seminuda Cope and Yarrow, Wheeler's Surv., Zool., V, 1875, 666, Rio Virgen, Nev.
813. *Gila minacæ* Meek.
Streams of Chihuahua, Mexico.
Gila minacæ Meek, Field Col. Mus., Zool., III, May, 1902, 80, Rio Paphogochic, Miñaca, Chihuahua, Mex.
- Genus 301. *NOTEMIGONUS* Rafinesque.
Notemigonus Rafinesque, Jour. Phys., LXXXVIII, 1819, 420, (*N. auratus* Rafinesque = *Cyprinus crysoleucas* Mitchell).
Hemiphus Rafinesque, Anal. Nat., I, 1820, 6 (*H. lacustris* Rafinesque).

814. *Notemigonus crysoleucas* (Mitchill). *Roach; Sunfish; Dace; Butter-head; Shiner; Chub; Bream; Gudgeon; Young shad; Windfish.*
Nova Scotia and Maryland to the Dakotas.
Cyprinus crysoleucas Mitchill, Rept., Fish. N. Y., 1814, 23, New York.
Cyprinus hemiplus Rafinesque, Amer. Month. Mag., II, 1817, 121, Lake George, Saratoga, N. Y.
- *Notemigonus auratus* Rafinesque, Jour. Phys., LXXXVIII, 1819, 421, Ohio River
Hemiplus lacustris Rafinesque, Anal. Nat., I, 1820, 6, Lake George.
Abramis versicolor DeKay, N. Y. Fauna: Fishes, 1842, 191, Connecticut and Hudson Rivers; Peekskill.
815. *Notemigonus bosci* (Cuvier and Valenciennes).
Rivers of the South Atlantic States; abundant from the James to the Altamaha.
Leuciscus bosci Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 232 (313), Carolina, Pennsylvania, and New York.
Cyprinus americanus Linnæus, Syst. Nat., ed. XII, 1766, 530, Charlestown; name preoccupied.
Notemigonus ischanus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 364, South Fork Ocmulgee River, Flat Shoals, Ga.
Luxilus leptosomus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 203, Dry Creek, Victoria, Tex.
Luxilus seco Girard, loc. cit., 203, Rio Seco, tributary of Rio Nueces, Tex.
816. *Notemigonus gardoneus* (Cuvier and Valenciennes).
South Carolina.
Leuciscus gardoneus Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 235 (316), Charleston, S. C.; perhaps a hybrid.
- Genus 302. OPSOPŒODUS Hay.
Opsopæodus Hay, Proc. U. S. Nat. Mus., III, 1880, 507 (*O. emiliæ* Hay).
Trychærodon Forbes, in Jordan and Gilbert, Bull. U. S. Nat. Mus., XVI, 1882, 247 (*T. megalops* Forbes).
817. *Opsopæodus emiliæ* Hay.
Lake Erie and southern Indiana and Illinois to Georgia and Mississippi.
Opsopæodus emiliæ Hay, Proc. U. S. Nat. Mus., III, 1880, 507, Artesia, Macon, and Enterprise, Miss.
818. *Opsopæodus osculus* Evermann.
Rio Neches and other streams about Palestine and Houston, Tex.
Opsopæodus osculus Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 82, Neches River near Palestine, Tex.
819. *Opsopæodus borealis* Harper and Nichols. *Athabaska minnow.*
Streams of Alberta.
Opsopæodus borealis Harper and Nichols, Bull. Amer. Mus. Nat. Hist., XLI, 266, Lake Athabaska at Fort Chipewyan, Alberta, Canada.
820. *Opsopæodus megalops* (Forbes).
Western Ohio to Illinois.
Trychærodon megalops Forbes, in Jordan and Gilbert, Synopsis, 1883, 247, Illinois River at Pekin and Peoria, Ill.
- Genus 303. OPSOPŒA Jordan and Evermann.
Opsopæa Jordan and Evermann, Fishes North and Mid. Amer., 1896, 247 (*Opsopæodus bollmani* Gilbert).
821. *Opsopæa bollmani* (Gilbert).
Lowland swamps of the Southern States; known from Satilla River, Waycross, Ga., and Obion River, Cypress, Tenn.
Opsopæodus bollmani Gilbert, Bull. U. S. Fish Com., VIII, 1888 (1890), 226, Buckland Creek, Millen, Ga.
- Genus 304. ALGANSEA Girard.
Algansea Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 182 (*Leuciscus tincella* Cuvier and Valenciennes).
822. *Algansea tincella* (Cuvier and Valenciennes).
Lakes about the City of Mexico and in streams of central Mexico.
Leuciscus tincella Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 240 (323), City of Mexico.

823. *Algansea dugesi* Bean.
Central Mexico, in streams tributary to Rio de Lerma, Lake Yuriria in Guanajuato.
Algansea dugesi Bean, Proc. U. S. Nat. Mus., XV, 1892, 283, Lake Yuriria, Guanajuato, Mex.
? *Algansea tarascorum* Steindachner, Einige Fischarten Mexicos, 1895, 11, pl. 3, figs. 2-2c, Lake Patzcuaro, Mex.
824. *Algansea lacustris* Steindachner.
Lake Patzcuaro, Jalisco, Mexico.
Algansea lacustris Steindachner, Einige Fischarten Mexicos, 1895, 10, pl. 3, figs. 1 and 1b, Lake Patzcuaro.
825. *Algansea sallæi* (Günther).
Streams of central Mexico.
Ceratichthys sallæi Günther, Cat., VII, 1868, 484, Cuernavaca, Mex.
Zophendum australe Jordan, Proc. U. S. Nat. Mus., II, 1879, 300, Lake Tupatario, Mex., Rio Lerma.
826. *Algansea rubescens* Meek.
Lake Patzcuaro, Michoacan.
Algansea rubescens Meek, Field Col. Mus., Zool., V, 1905, 46, fig. 9, Lake Patzcuaro.
827. *Algansea stigmatura* Regan.
Northern Mexico
Algansea stigmatura Regan, Biol. Centrali-Amer., 1907, 152, Tab. XXV, fig. 2, Rio Grande de Santiago, Jalisco.
828. *Algansea affinis* Regan
Upper tributaries of Rio Lerma and Rio Panuco, Mexico.
Algansea affinis Regan, Biol. Centrali-Amer., 1907, 151, Tab. XXV, fig. 1, Aguas Calientes, Salamanca, and San Juan del Rio, Mexico.
- Genus 305. *POGONICHTHYS* Girard.
Pogonichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 136 (*P. inaequilobus* Baird and Girard = *Leuciscus macrolepidotus* Ayres).
Symmetrurus Jordan, Hayden's Geol. Surv. Terr., IV, 1878, 788, (*Pogonichthys argyreus* Girard); young.
829. *Pogonichthys macrolepidotus* (Ayres). *Splittail*.
Streams of central California; very common in the Sacramento and San Joaquin Rivers.
Leuciscus macrolepidotus Ayres, Proc. Calif. Acad. Nat. Sci., I, in Placer Times and Transcript, May 30, 1854, San Francisco, Calif.
Pogonichthys inaequilobus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 136, San Joaquin Valley, Calif.
Pogonichthys argyreus Girard, loc. cit., 153, Presidio, Calif.
- Genus 306. *MARGARISCUS* Cockerell.
Margariscus Cockerell, Proc. Biol. Soc. Wash., XXII, 1909, 217 (*Clinostomus margarita* Cope).
As a small barbel is normally present in this genus and placed above the tip of the maxillary, as in *Semotilus*, the genus must be placed near *Leucosomus*.
830. *Margariscus margarita* (Cope). *Minnow; Goldthread shiner; Pearl minnow*.
Lake Erie, south to James River and headwaters of the Kanawha.
Clinostomus margarita Cope, Trans. Amer. Phil. Soc., XIII, 1866, 377, Conestoga River, Lancaster, Pa.
831. *Margariscus nachtriebi*²⁵ (Cox).
North Dakota to New Brunswick.
Leuciscus nachtriebi U. O. Cox, Rept. U. S. Fish Com., XX, 1894 (1896), 615, Minnesota, Mille Lacs Lake.
Couesius plumbeus rubrilateralis Philip Cox, Bull. Nat. Hist., 2, 1901, 42, Miramichi, Canada.
Leuciscus carletoni Kendall, Bull. U. S. Fish Com., XXII, 1902, 357, fig. 1, Penobscot River, Me.

²⁵ "With much doubt we leave this species (*M. nachtriebi*) with *Margariscus*. The actual relation of this and other related forms is yet to be determined." (Jordan.)

Genus 307. LEUCOSOMUS Heckel. *Fallfishes.*

Leucosomus Heckel, Russeger's Reisen, 1, 1843, 1042 (*L. chrysoleucus* Heckel = *Cyprinus corporalis* Mitchell); not *Leucosoma* Gray, a genus of Salangidae.

Cheilonemus Baird, in Storer, Hist. Fish. Mass., 1855, 285 (*Leuciscus pulchellus* Storer = *Cyprinus corporalis* Mitchell).

832. *Leucosomus corporalis* (Mitchill). *Chub; Dace; Fallfish; Silver chub; Windfish; Corporal; Chwin.*

Eastern United States, from the St. Lawrence to James River, Virginia; largest of our eastern Cyprinidae; common northward, east of the Alleghenies.

Cyprinus corporalis Mitchell, Amer. Month. Mag., I, 1817, 289, Wallkill River, N. Y.

Cyprinus bullaris Rafinesque, loc. cit., II, Dec., 1817, 120, Hudson River.

Leuciscus argenteus Storer, Hist. Fish. Mass., 1839, 90, Worcester Co., Mass.

Leuciscus pulchellus Storer, loc. cit., 90, Walpole, Mass.

Leuciscus nitidus DeKay, N. Y. Fauna: Fishes, 1842, 209, Lake Champlain.

Leuciscus chrysopterus DeKay, loc. cit., 211, N. Y. Harbor.

Leuciscus rotengulus Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 236 (318), Carolinas.

Leucosomus cataractus Baird, Iconog. Encyl., II, 216, and in Cope, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 523, Susquehanna River.

Leucosomus rhotheus Cope, loc. cit., XIII, 1861, 523, Brandywine Creek and Rancocas Creek, Pa.

Squalius hyalope Cope, loc. cit., XVI, 1864, 280, Conestoga River, Pa.

Genus 308. SEMOTILUS Rafinesque. *Horned dace.*

Semotilus Rafinesque, Ichth. Ohiensis, 1820, 49 (*S. dorsalis* Rafinesque = *Cyprinus atromaculatus* Mitchell).

833. *Semotilus atromaculatus* (Mitchill). *Horned dace; Creek chub; Chub; Silver chub; Mud chub.*

Maine to New Jersey; westward to Kansas and Wyoming.

Cyprinus atromaculatus Mitchell, Amer. Month. Mag., II, 1818, 324, Wallkill River, N. Y.

Semotilus dorsalis Rafinesque, Ichth. Ohiensis, 1820, 49, Kentucky River.

Semotilus cephalus Rafinesque, loc. cit., 49, Ohio River.

Semotilus diplema Rafinesque, loc. cit., 50, Ohio River.

Leuciscus irris Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 190, (255), New York and Carolina.

Leuciscus storeri Cuvier and Valenciennes, loc. cit., 237 (255), New York.

Leuciscus pulchelloides Ayres, Proc. Bost. Soc. Nat. Hist., III, 1849, 157, Connecticut.

Leucosomus pallidus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 190, Antelope Creek, Ark.

Leucosomus macrassatus Girard, loc. cit., 1856, 190, Choctaw Agency.

Semotilus macrocephalus Girard, loc. cit., 1856, 204, Fort Pierre, Nebr.

Semotilus speciosus Girard, loc. cit., 1856, 204, Sweetwater River, Nebr.

Semotilus hammondi Abbott, loc. cit., XII, 1860, 474, Kansas.

834. *Semotilus atromaculatus thoreauanus* (Jordan).

Flint River, Ga.

Semotilus thoreauanus Jordan, Bull., U. S. Nat. Mus., X, 1877, 63, Flint River, Ga.

Genus 309. RICHARDSONIUS ²⁵ Girard.

Richardsonius Girard, Proc., Acad. Nat. Sci. Phila., VIII, 1856, 201 (*Cyprinus (Abramis) baileatus* Richardson).

²⁵ The various American allies of *Leuciscus* Cuvier (type *Cyprinus leuciscus* L. of Europe, *Squalius*), we now divide among several genera, these corresponding to the subgenera in the previous check list. No species belonging to *Leuciscus*, as strictly defined, occurs in America or Asia. Many of the species, both in the groups allied to *Leuciscus* and *Rutilus* as well as in those related to *Hybopsis* and *Notropis* need comparison and verification.

Richardsonius is here restricted to the single species, *baileatus*, distinguished, with its variant, by the very long anal fin. The genus *Cheonida* is provisionally recognized for species more or less similar to *Richardsonius* but with the anal fin shorter. This group seems to merge into *Tygoma*, species with the anal fin short and the scales rather loosely imbricated. One species, *caurina*, forms another intermediate type with short anal and very small scales. Any arrangement of these fishes must be still wholly provisional.

835. *Richardsonius siuslawi* (Evermann and Meek).
Streams of western Oregon.
Leuciscus siuslawi Evermann and Meek, Bull. U. S. Fish Com., XVII,
1897 (1898), 72, Siuslaw River, Mapleton, Oreg.
836. *Richardsonius lateralis* Girard.
Streams of Washington and Montana, a doubtful species, nominally
distinguished by the smaller number of anal rays.
Richardsonius lateralis Girard, Proc. Acad. Nat. Sci. Phila., VIII,
1856, 202, Fort Steilacoom, Wash.
Leuciscus gilli Evermann, Bull., U. S. Fish Com., XI, 1891, 44, Browns
Gulch Creek, Silver Bow, Mont.
837. *Richardsonius balteatus* (Richardson). *Red-sided bream*.
Columbia and Fraser Rivers, and streams about Puget Sound.
Cyprinus (Abramis) balteatus Richardson, Fauna Bor.-Amer., III,
1836, 301, Columbia River.
838. *Richardsonius thermophilus* Evermann and Cockerell.
Warm Springs, Silver Creek, and Silvies River, Oreg.
Richardsonius thermophilus Evermann and Cockerell, Proc. Biol. Soc.
Wash., XXII, December 8, 1909, 185, Warm Springs, Harney Co.,
Oreg.
- Genus 310. *CLARKINA* Jordan and Evermann.
Cyprinus (Leuciscus) caurinus Richardson is an ally of the genus
Richardsonius, differing in the short anal fin of 9 rays only. The
body is elongate and the scales (86) very small, well imbricated,
differing from the type of *Tigoma*.
Clarkina Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI,
1927 (April 27), 502 (*Cyprinus caurinus* Richardson).
839. *Clarkina caurina* (Richardson).
Cyprinus (Leuciscus) caurinus Richardson, Fauna Bor.-Amer., III,
1836, 304, Columbia River at Fort Vancouver.
Leuciscus caurinus Snyder, Bull. U. S. Fish Com., XXVII, 1907, 173;
not *Mylocheilus caurinus* Agassiz; Willamette River near Corvallis,
Oreg.
- Genus 311. *CHEONDA* Girard.
Cheonda Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 207 (*C.*
cooperi Girard); scales moderate; anal rays 8 to 11, usually 9.
840. *Cheonda cooperi* Girard.
Lower Columbia River Basin.
Cheonda cooperi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 207,
Columbia River, Fort Vancouver, Wash.
841. *Cheonda microdon* (Snyder).
Richardsonius microdon Snyder, Bull. U. S. Bur. Fish., XXXV, 1915-16
(1918), 58; scales 55; anal rays 8 or 9.
842. *Cheonda egregia* (Girard).
Nevada to northern California; known from the Humboldt, Truckee,
and Sacramento-San Joaquin Basins and Napa Valley, Calif.
Tigoma egregia Girard, Pac. R. R. Surv., X, 1858, 291, locality Hum-
boldt River at Imlay, Nev.
Gila ardesiaca Cope, Wheeler Surv., Zool., V, 1875, 660, Pl. XXX,
figs. 1, 1a, locality unknown, probably Nevada.
Squalius galtii Cope, Proc. Acad. Nat. Sci. Phila., XXXV, 1883, 148,
Pyramid Lake, Nev.
Phoxinus clevelandi C. H. and R. S. Eigenmann, West Amer. Scientist,
1889, 149, Etna Springs, Napa County, Calif. (locality probably
erroneous).
843. *Cheonda evermanni* (Juday).
South Platte basin.
Leuciscus evermanni Juday, Bull. U. S. Bur. Fish., XXIV, 1904 (1905),
226, with plate, Boulder Creek, Boulder, Colo.; A. 8; scales 47; teeth
2, 5-5, 1.

844. *Cheonda hydrophlox* (Cope). *Silverside minnow*; *Po-hs-wa*.
Salt Lake basin and upper Snake River.
Clinostomus hydrophlox Cope, Hayden's Geol. Surv. Mont., 1871, 475,
Blackfoot Creek, Idaho.
? *Clinostomus tzenia* Cope, Proc., Amer. Phil. Soc., XIV, 1874, 133,
Utah Lake, Provo, Utah.
845. *Cheonda humboldti* (Girard).
Locality doubtful.
Tigoma humboldti Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 206,
Humboldt River, Nev.; may be identical with *C. hydrophlox*, but this
seems unlikely.
846. *Cheonda montana* (Cope).
Clinostomus montanus Cope, Hayden's Geol. Surv. Mont., 1871, 476,
Grass Creek, Idaho.
? *Clinostomus tzenia* Cope, Proc., Amer. Phil. Soc., XIV, 1874, 133,
Utah Lake, Provo, Utah.
May prove to be the same as *C. hydrophlox*.
847. *Cheonda copei* (Jordan and Gilbert). *Leather-sided minnow*.
Great Basin of Utah and Wood River, Idaho.
? *Tigoma gracilis* Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 206,
locality unknown; species unidentifiable, name preoccupied in
Leuciscus.
Squalius copei Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 461,
Bear River, Evanston; not *Leuciscus copei* Gunther, a *Notropis*.
Squalius aliciae Jouy, loc. cit., IV, 1881, 19, Provo River near Utah
Lake, Utah.
- Genus 312. **TIGOMA** Girard.
Tigoma Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 205 (*Gila
pulchella* Baird and Girard).
Protoporus Cope, Hayden's Geol. Surv. Mont., 1871 (1872), 473 (*P.
domninus* Cope, young of *Tigoma atraria* Girard).
Myloleucus Cope, loc. cit., 475 (*M. pulverentulus* Cope); the teeth of
Siphateles accidentally placed in the pharynx of a *Tigoma*.
Temeculina Cockerell, Proc. Biol. Soc. Wash., XXII, 1909, 216 (*Rich-
ardsonius orcutti* Eigenmann and Eigenmann).
848. *Tigoma atraria* (Girard).
Great Basin of Utah, and Snake River Basin above Shoshone Falls.
? *Tigoma lineata* Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 206,
locality unknown; not identifiable.
Siboma atraria Girard, loc. cit., 208, Fish Springs, Toole Co., Utah,
near Desert.
Tigoma squamata Gill, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 42,
Great Salt Lake Basin.
Protoporus domninus Cope, Hayden's Geol. Surv. Mont., 1871, 473,
Snake River, Fort Hall, Idaho.
Hybopsis bivittatus Cope, loc. cit., 474, Warm Springs, Utah.
Hybopsis timpanogensis Cope, Proc. Amer. Phil. Soc., XIV, 1874, 134,
Timpanogus, Utah; young.
Squalius cruoreus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880,
460, Provo River, Utah.
Squalius rhomaleus Jordan and Gilbert, loc. cit., 461, Utah Lake, Utah.
849. *Tigoma gibbosa* (Baird and Girard).
Rio Gila, Ariz.
Gila gibbosa Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854,
28, Rio Santa Cruz, Ariz.
Tigoma intermedia Girard, loc. cit., VIII, 1856, 206, Rio San Pedro of
the Gila, Ariz.
Gila nigra Cope, Wheeler's Surv., Zool., V, 1875, 663, Ash Creek and
San Carlos, Ariz.
Squalius lemmoni Rosa Smith, Bull. Calif. Acad. Sci., I, 1884, 3, Rillitto
Creek, near Tucson, Ariz.
850. *Tigoma purpurea* Girard.
San Bernardino Creek in southern Arizona, a tributary of Rio Yaqui.
Tigoma purpurea Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 206,
San Bernardino Creek, Ariz.

- 851. *Tigoma pulchella* (Girard).** *Pescadito; Chub of the Rio Grande.*
 Rio Grande and other Mexican rivers.
Gila pulchella Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 29, Rio Mimbres, Chihuahua; not *Leuciscus pulchella* Storer, a *Cheilonemus*.
Tigoma nigrescens Girard, loc. cit., VIII, 1856, 207, Boca Grande and Rio Janos, Chihuahua, Mexico.
Tigoma pulchra Girard, loc. cit., 207, Boca Grande and Rio Janos; Chihuahua River, Mexico.
Clinostomus pandora Cope, Hayden's Geol. Surv. Mont., 1871, 475, tributaries Rio Grande, Sangre de Cristo Pass.
Gila gila Cope, Wheeler's Surv., Zool., V, 1875, 661, Rio de Acama and near Fort Wingate, N. Mex.
Cheonda modesta Garman, Bull. Mus. Comp. Zool., VIII, 1881, 92, Rio Salinas, Saltillo, Coahuila.
Gila conspersa Garman, loc. cit., VIII, 1881, 91, Rio Nazas, Coahuila, Mexico.
- 852. *Tigoma bicolor* Girard.**
 Klamath Lake, Oreg.
Tigoma bicolor Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 206, Klamath Lake, Oreg.
Cheonda cærulea Girard, loc. cit., 207, Lost River, Oreg.
- 853. *Tigoma pulverulenta* (Cope).**
Myloleucus pulverulentus Cope, Hayden's Geol. Surv. Mont., 1871, 475, Warm Springs, Utah.
Myloleucus is a synonym of *Tigoma*. In his study of these species, according to Fowler, Professor Cope by accident replaced the single-rowed teeth of a *Siphateles* in his type example of *Myloleucus pulverulentus*, a species having teeth in two rows.
- Subgenus **TEMECULINA** Cockerell.
- 854. *Tigoma orcutti* (Eigenmann and Eigenmann).**
 Temecula, San Luis Rey, San Jacinto, and Santa Ana Rivers, California.
Phoxinus orcutti Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2nd ser., III, 1890, 2, Temecula River, Riverside County, Calif.
- Genus 313. SIBOMA** Girard.
Siboma Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 209 (*S. crassicauda* Girard).
- 855. *Siboma crassicauda* (Baird and Girard).** *Sacramento chub.*
 Rivers of California; generally abundant in the Sacramento and San Joaquin Rivers.
Lavinia crassicauda Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 137, San Joaquin River.
Tigoma crassa Girard, loc. cit., VIII, 1856, 207, Sacramento River.
- 856. *Siboma conformis* (Baird and Girard).**
 Tulare Valley and Sacramento-San Joaquin Basin, Calif.
Lavinia conformis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 137; Poso Creek, Tulare County, Calif.
- Genus 314. CLINOSTOMUS** Girard.
Clinostomus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 211 (*Luxilus elongatus* Kirtland).
- 857. *Clinostomus elongatus* (Kirtland).** *Red-sided shiner.*
 Great Lakes, upper Mississippi Valley, and New York.
Luxilus elongatus Kirtland, Rept. Zool. Ohio, 1836, 169, 193, Mahoning River, Ohio, and Lake Erie near Cleveland.
Leuciscus productus Storer, Synopsis, 1846, 416, Wabash River.
 ?*Leuciscus longirostris* Kirtland, The Family Visitor, Mar. 14, 1850, Cleveland, Ohio, and Hudson River.
Squalius proriger Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 280, Michigan.

- 858. *Clinostomus vandoisulus*** (Cuvier and Valenciennes). *Minnow; Long-mouthed minnow; Rosy dace; Pike shiner.*
Leuciscus vandoisulus Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 236 (317), South Carolina.
Clinostomus affinis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 212, James River, Va.
Clinostomus funduloides (Potomac River at Washington) and *carolinus* (Salem, N. C.) Girard, loc. cit., 212.
Gila estor Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 66, Elk River, Estill Springs, Tenn.
- Genus 315. HEMITREMIA** Cope.
Hemitremia Cope, Proc. Amer. Phil. Soc., XI, 1870, 462 (*H. vittata* Cope=*Phoxinus flammeus* Jordan and Gilbert).
- 859. *Hemitremia vittata*** Cope.
 Tributaries of the Tennessee River; common in clear streams in northern Alabama.
Hemitremia vittata Cope, Proc. Amer. Phil. Soc., XI, 1870, 462, Holston River, Knoxville, Tenn.; *vittatus* is preoccupied in *Leuciscus*.
Phoxinus flammeus Jordan and Gilbert, in Jordan, Man. Vert., ed. 2, 1878, 303, Elk River, Tenn.
- 860. *Hemitremia milneriana*** (Cope).
 Upper Missouri River Basin.
Phoxinus milnerianus Cope, Amer. Nat., July, 1879, 440, upper Missouri River Basin; probably Battle Creek, Mont.
- Genus 316. PFRILLE** Jordan.
Pfrille Jordan, Copeia, No. 32, 1924, 71 (*Phoxinus neogæus* Cope).
- 861. *Pfrille neogæa*** (Cope). *Bronze minnow.*
 Upper Mississippi Valley and neighboring waters; also Maine and Connecticut lakes.
Phoxinus neogæus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 375, New Hudson, Mich.
 ?*Leuciscus burtonianus* LeSueur, in Vaillant, Bull. Soc. Philom., VIII, 1896, 28, with plate, Burton Mine, Mo.; perhaps a distinct species.
- Genus 317. IOTICHTHYS** Jordan and Evermann.
Iotichthys Jordan and Evermann, Fishes North and Mid. Amer., 1896, 228 (*Clinostomus phlegethontis* Cope).
- 862. *Iotichthys phlegethontis*** (Cope).
 Tributaries of Great Salt Lake and Sevier Lake; basin of Lake Bonneville.
Clinostomus phlegethontis Cope, Proc. Amer. Phil. Soc., XIV, 1874, 137, Beaver River, Utah.
- Genus 318. HESPEROLEUCUS** Snyder.
Hesperoleucus Snyder, Bull. U. S. Bur. Fish., XXXII, 1912, 63 (*Pogonichthys symmetricus* Girard).
Myloleucus Cope, Hayden's Geol. Surv. Mont., 1871, 475 (*M. pulverulentus* Cope=*Pogonichthys symmetricus* Baird and Girard, in part, not type, the teeth having been misplaced).
 This genus and others related, having the teeth 4-5 or 5-5, were placed in the former check list and the Fishes of North and Middle America, in the European genus *Rutilus* Rafinesque; but no American or Asiatic dace is congeneric with *Rutilus rutilus* L.
- 863. *Hesperoleucus symmetricus*** (Baird and Girard).
 Rivers of California and Nevada and waters of Lake County, Oreg.
Pogonichthys symmetricus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 136, Fort Miller, San Joaquin Valley, Calif.
Myloleucus parovanus Cope, Proc. Amer. Phil. Soc., XIV, 1874, 131, Beaver River, Utah.
- 864. *Hesperoleucus navarroensis*** Snyder. *Navarro roach.*
 Navarro Basin on the western coast of northern California.
Hesperoleucus navarroensis Snyder, Bull. U. S. Bur. Fish., XXXII, 1912 (1914), 65, Navarro River near Philo, Calif.

865. *Hesperoleucus parvipinnis* Snyder. *Short-finned roach*.
Gualala River Basin, Calif.
Hesperoleucus parvipinnis Snyder, Bull. U. S. Bur. Fish., XXXII, 1912 (1914), 66, Gualala River, Sonoma County, Calif.
866. *Hesperoleucus venustus* Snyder. *Venus roach*.
Russian River and the streams entering San Pablo, Suisun, and San Francisco Bays.
Hesperoleucus venustus Snyder, Bull. U. S. Bur. Fish., XXXII, 1912 (1914), 64, fig. 2, Coyote Creek near Gilroy Hot Springs, Calif.
867. *Hesperoleucus mitrulus* Snyder. *Northern roach*.
Streams tributary to the north end of Goose Lake, Oreg.
Hesperoleucus mitrulus Snyder, Bull. U. S. Bur. Fish., XXXII, 1912, (1914), 67, Drew Creek, Lake County, Oreg.
868. *Hesperoleucus subditus* Snyder. *Monterey roach*.
Pajaro River system in the region of foothills and lower mountain ranges of coast ranges.
Hesperoleucus subditus Snyder, Bull. U. S. Bur. Fish., XXXII, 1902 (1904), 67, fig. 3, Uvas Creek, Pajaro River Basin, Santa Clara County, Calif.
869. *Hesperoleucus mearnsi* (Snyder).
Rio Yaqui.
Notropis mearnsi Snyder, Proc., U. S. Nat. Mus., XLIX, 1916 (Dec. 23, 1915), 582, fig., Rio San Bernardino, Yaqui Basin, Sonora, Mexico.
870. *Hesperoleucus anticus* (Cope).
Texas.
Algansea antica Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 282, Texas.
- Genus 319. SIPHATELES Cope.
Siphateles Cope, Proc. Acad. Nat. Sci. Phila., XXXV, 1883, 146 (*S. vittatus* Cope); young example.
871. *Siphateles olivaceus* (Cope).
Lake Tahoe; Pyramid Lake; Eagle Lake and Willow Creek, Calif.
Leucus olivaceus Cope, Proc. Acad. Nat. Sci. Phila., XXXV, 1883, 145, Pyramid Lake, Nev.
872. *Siphateles obesus* (Girard).
Lakes and streams of Lahontan Basin.
Algansea obesa Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 183, Humboldt River, Nev.
Leucus dimidiatus Cope, loc. cit., XXXV, 1883, 145, Pyramid Lake, Nev.
Siphateles vittatus Cope, loc. cit., 146, Pyramid Lake, Nev.
873. *Siphateles bicolor* (Girard).
Lakes of southeastern Oregon (Klamath, Warner, Goose, etc.); Sacramento-San Joaquin Basin; Pit River.
Algansea bicolor Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 183, Klamath Lake.
874. *Siphateles formosus* (Girard).
Algansea formosa Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 183, Merced and Mohave Rivers, Calif.
Myloleucus thalassinus Cope, loc. cit., XXXV, 1883, 144, Goose Lake, Oreg.
875. *Siphateles columbianus* (Snyder).
Lakes of southeastern Oregon.
Rutilus columbianus Snyder, Bull. U. S. Bur. Fish., XXVII, 1907, 92, fig. 4, Warm Springs near Harney Lake, Harney County, Oreg.
876. *Siphateles oregonensis* (Snyder).
Lakes of southeastern Oregon.
Rutilus oregonensis Snyder, Bull. U. S. Bur. Fish., XXVII, 1907, 87, fig. 3, XL, Spring, Abert Lake, Oreg.

877. *Siphateles mohavensis* Snyder.
Mohave River, Calif.
Siphateles mohavensis Snyder, Proc. U. S. Nat. Mus., LIV, 1919 (Mar. 15, 1918), 298, figure, Mohave River near Victor, Calif.
- Genus 320. *LEUCIDIUS* Snyder.
Leucidrus Snyder, Bull. U. S. Bur. Fish., XXXV, 1917, 64 (*L. pectinifer* Snyder).
878. *Leucidius pectinifer* Snyder. *Lake minnow*.
Lahontan system, Nevada and northeastern California.
Leucidrus pectinifer Snyder, Bull. U. S. Bur. Fish., XXXV, 1916-17 (Sept. 28, 1917), 64, figs. 5 and 6, Pyramid Lake, Nev.
- Genus 321. *NOTROPIS* Rafinesque.
*Notropis*²⁷ Rafinesque, Amer. Month. Mag., II, 1818, 204 (*Notropis atherinoides* Rafinesque = ?*Alburnus rubellus* Agassiz).
Minnilus Rafinesque, Ichth. Ohiensis, 1820, 45 (*M. dinemus* Rafinesque).
Alburnellus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856 (*A. dilectus* Girard).
Photogenis Cope, Trans. Amer. Phil. Soc., XIII, 1867, 378 (*Squalius photogenis* Cope = *Photogenis leucops* Cope).
879. *Notropis atherinoides* Rafinesque. *Shiner*; *Emerald minnow*; *Rosy minnow*.
Great Lakes region and Ohio and Mississippi Valleys, and north to Winnipeg.
Notropis atherinoides Rafinesque, Amer. Month. Mag., II, 1818, 204, Lake Erie.
Minnilus dinemus Rafinesque, Ichth. Ohiensis, 1820, 46, Ohio River.
Alburnus rubellus Agassiz, Lake Superior, 1850, 364, pl. 3, figs 1-3, Sault Ste. Marie and the Pic, Lake Superior.
Alburnus nitidus Kirtland, Cleveland Ann. Sci., 1854, 44, tributaries of Lake Erie.
Alburnellus jaculus Cope, Trans., Amer. Phil. Soc., XIII, 1866 (1869). 387, St. Joseph River and Dowagiac River, Mich.
Leuciscus copii Günther, Cat., VII, 1868, 255, Virginia and Michigan; *l. b. titulus jaculus*, preoccupied.
880. *Notropis photogenis* (Cope). *Silvery minnows*.
Allegheny region.
? *Ruhulus compressus* Rafinesque, Ichth. Ohiensis, 1820, 51, Monongahela River, not certainly identifiable; may be *Luzilis cornutus*.
Squalius photogenis Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 280, Youghiogheny River, Pa.
Photogenis leucops Cope, loc. cit., XIX, 1867, 164, Sinking Creek, and near Austinville, Va.
Photogenis leucops engraulinus Cope, loc. cit., 164, tributaries of Kanawha River, Austinville, Va.
881. *Notropis amœnus* (Abbott).
Clear streams of the Alleghenies from the Raritan to the Neuse.
Alburnellus amœnus Abbott, Amer. Nat., VIII, 1874, 334, Raritan River, N. J.
882. *Notropis scepticus* (Jordan and Gilbert).
Rivers of the Carolinas from Cape Fear to the Santee.
Notropis photogenis ("pale variety"), Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 23.
Minnilus scepticus Jordan and Gilbert, Synopsis, 1883, 200, Saluda River, S. C.

²⁷ The name *Notropis atherinoides* was given by Rafinesque to a small and very slender minnow from Lake Erie. It was apparently a semi-dried example having therefore a ridge on the back as implied in the name *Notropis*. It had a long anal fin of 11 rays, and the dorsal inserted behind the ventrals. Among the small minnows of the Great Lakes it could hardly be anything else than *rubellus*, yet the original description is very incomplete and perhaps not identifiable. As this is the oldest name for any of the small American minnows with four teeth in the main row of the pharyngeals, and as distinctions of genera among these is especially difficult, the writers in 1893 extended the name to include them all. But this assemblage is hardly a natural one, as the species diverge in various directions. Hence, in this catalogue the former subgenera are given generic rank.

883. *Notropis telescopus* (Cope).
Tennessee River Basin; very abundant in cold mountain streams.
Photogenis telescopus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867
165, Holston River, Va.
884. *Notropis arcansanus* (Meek).
Ozark region
Notropis telecopus arcansanus Meek, Bull. U. S. Fish Com., IX, 1889
(1891), 133, White River, Eureka Springs, Ark.
885. *Notropis universitatis* Evermann and Cockerell.
Boulder Creek, Boulder, Colo.
Notropis notatus var., Univ. of Colo. Stud., Apr., 1908, 170.
Notropis universitatis Evermann and Cockerell, Proc. Biol. Soc. Wash.,
XXII, Dec., 1909, 187, Boulder Creek, Boulder, Colo.
886. *Notropis welaka* Evermann and Kendall.
St. Johns River, Fla.
Notropis welaka Evermann and Kendall, Bull. U. S. Fish Com., XVII,
1897 (Feb. 9, 1898), 126, pl 6, fig. 2, St. Johns River near Welaka,
Fla.
887. *Notropis leuciodus* (Cope).
Tennessee River Basin, in mountain streams.
Photogenis leuciodus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867,
165, Holston River, Va.
888. *Notropis micropteryx* (Cope).
Headwaters of Cumberland and Tennessee Rivers; also in Ozark region.
Alburnellus micropteryx Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868,
233, Holston River, Va.
889. *Notropis metallicus* Jordan and Meek.
Swamp streams in Georgia and Florida, from the Suwanee Basin to
the Escambia.
Notropis metallicus Jordan and Meek, Proc. U. S. Nat. Mus., VII,
1884, 475, Allapaha River, Nashville, Ga.
890. *Notropis louisianæ* Evermann.
Atchafalaya River, La.
Notropis louisianæ Evermann, in Jordan and Evermann, Fishes North
and Mid. Amer., 1893, 2801, Atchafalaya River, Melville, La.
891. *Notropis socius* (Girard).
Live Oak Creek, southwestern Texas.
Alburnus socius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 193,
Live Oak Creek, Tex.
892. *Notropis amabilis* (Girard).
Rio Leona, a tributary of Rio Nueces, Tex.
Alburnus amabilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
193, Rio Leona, Tex.
893. *Notropis arge* (Cope).
Upper Mississippi Valley.
Alburnellus arge Cope, Trans. Amer. Phil. Soc., XIII, 1866, 388, Detroit
River or St. Joseph River, Mich.
894. *Notropis pilsbryi* Fowler.
Notropis pilsbryi Fowler, Proc. Acad. Nat. Sci. Phila., LVI, 1904, 245,
figure, Rogers, White River Basin, Ark.
895. *Notropis rubrifrons* (Cope). *Rosy-faced minnow*.
New York and western Pennsylvania to southern Michigan, Kansas,
and Kentucky.
Alburnus rubrifrons Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 85,
Kiskiminitas River, a tributary of Allegheny River.
Alburnellus percobromus Cope, Hayden's Geol. Surv. Wyo., 1870, 440,
Missouri River at St. Joseph, Mo.

896. *Notropis dilectus* (Girard).
Lower Ohio to the Rio Grande, abundant in Arkansas and eastern Texas in sandy streams
Alburnus dilectus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 193,
Arkansas River, Fort Smith, Ark.
Alburnus lepidulus Girard, loc. cit., 212, Black Warrior River, Ala
Alburnus oligaspiis Cope, loc. cit., XVI, 1864, 282, Kansas.
Alburnellus jemezianus Cope, Wheeler Surv., Zool., V, 1875, 650, Rio Grande, San Ildefonso, Mexico.
897. *Notropis stilbius* (Jordan).
Alabama River and tributaries.
Nototropis stilbius Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 343,
Etowah and Oostanaula Rivers, Rome, Ga.
898. *Notropis notemigonoides* Evermann
Rio Neches near Palestine, Tex., and streams about Houston, Tex
Notropis notemigonoides Evermann, Bull. U. S. Fish Com., XI, 1891
(1892), 81, Neches River, Palestine, Tex., and Sims Bayou, Houston,
Tex.
899. *Notropis fumeus* Evermann.
Hunter Creek, Houston, Tex.
Notropis fumeus Evermann, Bull. U. S. Fish Com., XI, 1891 (1892),
81, Hunter Creek, Houston, Tex.
- Genus 322. *PARANOTROPIS* Fowler.
Episema Cope and Jordan, Bull. U. S. Nat. Mus., X, 1877, 64 (*Photogenis*
scabriceps Cope, name preoccupied)
Paranotropis Fowler, Proc. Acad. Nat. Sci. Phila., LVI, 1904, 245
(*Photogenis scabriceps* Cope).
900. *Paranotropis scabriceps* (Cope).
Kanawha River, W. Va.
Photogenis scabriceps Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867,
166, Sinking and Walker Creeks, tributaries of Kanawha River.
901. *Paranotropis ariommus* (Cope).
Ohio and Tennessee Valleys.
Photogenus ariommus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 378,
no locality given.
902. *Paranotropis jejuna* (Forbes)
Western Pennsylvania to Kansas and north to Winnipeg.
Episema jejuna Forbes, Bull. Ill. Lab. Nat. Hist., I, 1878, 60, Illinois
River
903. *Paranotropis megalops* (Girard).
Rivers of Texas from the Colorado westward.
Alburnus megalops Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
193, San Felipe Creek, Tex.; the name preoccupied in *Notropis*
(*Luxilus*) but not in *Paranotropis*.
- Genus 323. *LYTHRURUS* Jordan.
Lythrurus Jordan, Man. Vert., ed. 1, 1876, 272 (*Notropis lythrurus*
Jordan = *Semotilus diplæmius* of authors, not of Rafinesque).
904. *Lythrurus umbratilis* (Girard).
Arkansas River and streams of Kansas and southwestern Iowa.
Alburnus umbratilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
193, Sugar Loaf Creek, tributary of Poteau River, Ark
Luxilus lucidus Girard, loc. cit., 203, Coal Creek, tributary of South Fork
of Canadian River.
Minnilus nigripinnis Gilbert, Bull., Washburn Lab. Nat. Hist., I, 1884,
14, Shunganunga Creek near Topeka, Kans.
? *Notropis macrolepidotus* Forbes, Bull. Ill. Lab. Nat. Hist., II, 1885, 138,
Illinois.

905. *Lythrurus cyanocephalus* Copeland.
Rivers of southern Wisconsin to Ohio and Kentucky and Indiana.
Lythrurus cyanocephalus Copeland, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 70, Racine River, Wis.
Rutilus ruber Rafinesque, Ichth. Ohiensis 1820, 52, Elkhorn and Kentucky Rivers; named but not described.
Notropis lythrurus Jordan, Proc., U. S. Nat. Mus., VII, 1884, 476, White River, Indianapolis; a rather more slender southern form.
906. *Lythrurus atripes* Jordan. *Blackfin; Compressed redfin*.
Southern Illinois and Iowa.
Lythrurus atripes Jordan, Bull. Ill. Lab. Nat. Hist., No. 2, 1878, 59, streams of Union and Johnson Counties, Ill.
907. *Lythrurus ardens* (Cope).
Roanoke River, Va.
Hypsilepis ardens Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 163, headwaters of Roanoke River, Montgomery County, Va.
908. *Lythrurus fasciolaris* (Gilbert).
Southern bend of Tennessee River, in limestone streams.
Notropis umbratilis fasciolaris Gilbert, Bull. U. S. Fish Com., IX, 1889 (1891), 148, streams about Florence, Ala.
909. *Lythrurus matutinus* (Cope).
Neuse and Pamlico Rivers; common in sandy brooks.
Alburnellus matutinus Cope, Proc. Amer. Phil. Soc., XI, 1870, 465, Neuse River, Wake County, N. C.
910. *Lythrurus punctulatus* (Hay).
Tributaries of Big Hatchee River, northern Mississippi.
Minnilus punctulatus Hay, Proc. U. S. Nat. Mus., III, 1880, 508, Tuscumbia River, a tributary of the Big Hatchee, near Corinth, Miss.
911. *Lythrurus bellus* (Hay).
Tombigbee River, Miss.
Minnilus bellus Hay, Proc. U. S. Nat. Mus., III, 1880, 510, Tombigbee River, Artesia and Macon, Miss.
912. *Lythrurus roseipinnis* (Hay).
Sandy streams of Gulf States, from Escambia River to the Chickasawha.
Notropis roseipinnis Hay, in Jordan, Cat. Fishes N. Amer., 1885, 27, Chickasawha River, Enterprise, Miss; substitute for *rubripinnis*, preoccupied.
913. *Lythrurus lirus* (Jordan).
Alabama River Basin; abundant in sandy streams.
Nototropis lirus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 342, Etowah River, Rome, Ga.
Notropis alabamæ Jordan and Meek, Proc. U. S. Nat. Mus., VII, 1884, 476, Alabama River, Montgomery, Ala.
- Genus 324. **ORCELLA** Jordan and Evermann
Orcella Jordan and Evermann, Fish. North and Mid. Amer., 1896, 254 (*N. orca* Woolman); not "*Orcella*," a genus of Cetacea.
Orcula Jordan and Evermann, loc. cit., 1900, 3140, substitute for *Orcella*, regarded as preoccupied by *Orcella*.
914. *Orcella nazas* (Meek).
Headwaters of the Rio Nazas in Durango, Mexico
Nototropis nazas Meek, Field Col. Mus., Zool., V, 1904, 70, fig. 19, Santiago Papasquiaro, Durango, Mexico.
915. *Orcella orca* (Woolman).
Rio Grande at El Paso, Tex.
Notropis orca Woolman, Bull. U. S. Fish Com., XIV, 1894, 56, Rio Grande, El Paso, Tex.
- Genus 325. **HYDROPHLOX** Jordan and Brayton
Hydrophlox Jordan and Brayton, Bull. U. S. Nat. Mus., XIII, 1878, 32 (*Hybopsis rubricroceus* Cope).

916. *Hydrophlox rubricroceus* (Cope). *Red fallfish*.
Headwaters of the Tennessee and Savannah Rivers.
Hybopsis rubricroceus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 231,
Tumbling Creek, tributary of the North Fork of Holston River, Va.
917. *Hydrophlox chlorocephalus* (Cope).
Santee Basin, in North and South Carolina
Hybopsis chlorocephalus Cope, Proc. Amer. Phil. Soc., XI, 1870, 461,
Catawba River, N. C.
918. *Hydrophlox lutipinnis* Jordan and Brayton.
Santee and Oconee Rivers, Ga.
Hydrophlox lutipinnis Jordan and Brayton, Bull. U. S. Nat. Mus.,
XII, 1878, 36, Oconee River, Halls Springs, Ga.
919. *Hydrophlox chiliticus* (Cope).
Basin of Great Pedee River, N. C.; common in upland streams.
Hybopsis chiliticus Cope, Proc. Amer. Phil. Soc., XI, 1870, 462, Yadkin
River, Rowan County, N. C.
920. *Hydrophlox altipinnis* (Cope).
Yadkin, Great Pedee, and Cape Fear Rivers, N. C.
Alburnellus altipinnis Cope, Proc. Amer. Phil. Soc., XI, 1870, 464,
Yadkin River, Rowan County, N. C.
921. *Hydrophlox roseus* (Jordan).
Lowland streams of Gulf States from Ogeechee River to the Mississippi.
Luzilus roseus Jordan, Bull. U. S. Nat. Mus., X, 1877, 61, Notalbany
River near Tikfaw, La.
922. *Hydrophlox stonei* (Fowler).
Streams of South Carolina.
Notropis stonei Fowler, Proc. Acad. Nat. Sci. Phila., LXXII, 1920, 391,
figure, Pocatigo River, Manning, S. C.
923. *Hydrophlox chalybæus* (Cope).
Delaware River to Ogeechee River.
Hybopsis chalybæus Cope, Trans., Amer. Phil. Soc., XIII, 1866, 383,
Schuylkill River, Pa.
924. *Hydrophlox abbotti* (Fowler).
Notropis chalybæus abbotti Fowler, Proc. Acad. Nat. Sci. Phila., LVI,
1904, 239, Pl. XVII, upper figure, Batsto River, N. J.
925. *Hydrophlox chrosomus* (Jordan).
Alabama Basin; very abundant in clear streams and outlets of springs.
Hybopsis chrosomus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 333,
Etowah River, etc., Rome, Ga.
926. *Hydrophlox xænocephalus* (Jordan).
Georgia to Mississippi; common in streams of the pine woods, descending
to brackish water; abundant in Perdido Bay, Pensacola, Fla.
Hybopsis xænocephalus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876,
334, Etowah River, etc., Rome, Ga.
- Genus 326. COCCOTIS Jordan
Coccotis Jordan, Rept. Zool. Ohio, 1882, 852 (*Hypsilepis coccogenis*
Cope).
Coccogenia Cockerell, Proc. Biol. Soc. Wash., XXII, 1909, 190 (*Hypsi-*
lepis coccogenis Cope).
927. *Coccotis coccogenis* (Cope).
Cumberland, Tennessee, and Savannah Rivers.
Hypsilepis coccogenis Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867,
160, Holston River, Va.
928. *Coccotis brimleyi* (B. A. Bean).
Notropis brimleyi B. A. Bean, Proc. U. S. Nat. Mus., XXVI, 1903, 913,
Cane River, N. C.
929. *Coccotis macdonaldi* (Jordan and Jenkins).
Mountain streams of Virginia tributary to the James and Shenandoah.
Notropis macdonaldi Jordan and Jenkins, Proc. U. S. Nat. Mus., XI,
1888, 354, plate 44, fig. 4, Shenandoah River, Luray, Va.

930. *Coccotis zonatus* (Agassiz).
Mountain streams in the Ozark region of Missouri and Arkansas.
Alburnus zonatus Agassiz, Bull. Mus. Comp. Zool., I, 1863, 9, Osage River, Mo.
931. *Coccotis zonistius* (Jordan).
Chattahoochee River
Luxilus zonistius Jordan, Proc. U. S. Nat. Mus., II, 1879, 239, Suwannee Creek, tributary of Chattahoochee River in northern Georgia.
- Genus 327. *LUXILUS* Rafinesque.
Luxilus Rafinesque, Ichth. Ohiensis, 1820, 47 (*L. chrysocephalus* Rafinesque = *Cyprinus cornutus* Mitchill).
Hypsoplepis Baird, Amer. Jour. Sci. Arts, XVII, 1854, 359 (*Cyprinus cornutus* Mitchill); later written *Hypsoplepis*.
932. *Luxilus cornutus* (Mitchill). *Shiner*; *Dace*; *Redfin*.
Entire region east of the Rocky Mountains excepting the south Atlantic States and Texas; very abundant; also Potomac River and tributaries; not found south of the James.
Cyprinus cornutus Mitchill, Amer. Month. Mag., I, July, 1817, 289, Wallkill River, N. Y.; preliminary notice, and II, 1818, 324, Westchester County, N. Y.
Cyprinus megalops Rafinesque, loc. cit., II; December, 1817, 121, Hudson River, above the falls.
Cyprinus melanurus Rafinesque, loc. cit., 121, Hudson River, above the falls.
? *Luxilus interruptus* Rafinesque, loc. cit., 49, Ohio River; may be *Notropis rubrifrons*.
Rutilus plargyrus Rafinesque, Ichth. Ohiensis, 1820, 50, Ohio River in Kentucky.
Rutilus compressus Rafinesque, loc. cit., 51, Monongahela River, etc., not certainly identifiable; may be *Notropis photogenus*.
Leuciscus vittatus DeKay, N. Y. Fauna: Fishes, 1842, 212, Chittenonga Creek, tributary of Mohawk River.
Aigyreus rubripinnis Heckel, Russeger's Reisen, 1843, 1040.
Leuciscus spinlingulus Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 239 (321), New Jersey; New Harmony, Ind.
Leuciscus obesus Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 48, Florence, Ala.
Plargyrus typicus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 196; after Rafinesque.
Plargyrus bowmani Girard, loc. cit., 196, Sweetwater River, Nebr.
Plargyrus argentatus Girard, loc. cit., 212, James River, Va.
Alburnops plumbeolus Cope, loc. cit., XVI, 1864, 282, Saginaw River, Flint, Mich.
Hypsoplepis cornutus cyaneus Cope, loc. cit., XIX, 1867, 160, Montreal River, Keweenaw, Mich.
Leuciscus frontalis Agassiz, Lake Superior, 1850, 368, Montreal River, eastern shore of Lake Superior.
Leuciscus gracilis Agassiz, loc. cit., 370, Lake Huron.
Hypsoplepis cornutus gibbus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 158, Monroe County, Mich.
933. *Luxilus cornutus isolepis* (Hubbs and Brown).
Oklahoma and Arkansas.
Notropis cornutus isolepis Hubbs and Brown, Proc. Okla. Acad. Sci., Pt. I, VI, Jan. 1, 1927, 129, Mountain Fork River, McCurtain County Okla.
934. *Luxilus cornutus chrysocephalus* (Rafinesque).
Ohio River and southward; the southern form of *cornutus*.
Luxilus chrysocephalus Rafinesque, Amer. Month. Mag., 1820, 47, Ohio River.
935. *Luxilus cerasinus* (Cope).
Roanoke River Basin, in mountain brooks.
Hypsoplepis cornutus cerasinus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 159, headwaters of Roanoke River, Va.

936. *Luxilus albeolus* (Jordan).
Roanoke, Tar, and Neuse Rivers.
Notropis megalops albeolus Jordan, Bull. U. S. Fish Com., VIII, 1888
(1891), 123, Roanoke River near Roanoke, Va
937. *Luxilus lacertosus* (Cope).
Headwaters of Tennessee River.
Hybopsis lacertosus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 232,
Bear Creek, tributary of the Middle Fork of Holston River
- Genus 328. *EROGALA* Jordan and Brayton
Erogala Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1898, 20
(*Photogenis stigmaturus* Jordan).
938. *Erogala pyrrhomelas* (Cope).
North and South Carolina in the Santee Basin.
Photogenis pyrrhomelas Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870,
463, Catawba River, N. C.
939. *Erogala formosa* (Putnam).
Southern Georgia and Alabama.
Alburnus formosus Putnam, Bull. Mus. Comp. Zool., I, 1863, 9, Mobile;
not *Monana formosa* Girard
Leuciscus hypselopterus Gunther, Cat., VII, 1868, 255, Mobile; substitute
for *formosus*, preoccupied in *Leuciscus* but not in *Erogala*.
Photogenis grandipinnis Jordan, Bull. U. S. Nat. Mus., X, 1877, 62,
Flint River, Ga.
940. *Erogala whipplii* (Girard). *Silverfin; Lemonfin; Satinfin; Steel-colored minnow.*
Central and northern New York to Minnesota, northern Alabama, and
Arkansas, in clear streams.
Cyprinella whippli Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
198, Sugarloaf Creek, tributary Poteau River, Ark.
Photogenis spilopterus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 378, St.
Joseph River, southwestern Michigan.
Hybopsis fretensis Cope, loc. cit., 382, Detroit, Mich.; young of *C.*
whippli, fide Hubbs.
941. *Erogala analostana* (Girard). *Silverfin, Satinfin, Lacefin; Shiner.*
Streams about Delaware and Chesapeake Bays; representing *Erogala*
whippli east of the Alleghenies.
Cyprinella analostana Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859,
59, at Analostan Island, Washington, D. C.
942. *Erogala galactura* (Cope).
Ozark region to Cumberland, Tennessee, and Savannah Rivers; abundant
in mountain streams.
Hypsilepis galacturus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 160,
Holston River, Va.
943. *Erogala stigmatura* (Jordan). *Spotted-tail minnow.*
Alabama Basin; very abundant.
Photogenis stigmaturus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876,
337, Etowah River, Rome, Ga.
Cyprinella callura Jordan, Bull. U. S. Nat. Mus., X, 1877, 61, Black
Warrior River, Ala.
944. *Erogala camura* (Jordan and Meek).
Arkansas River and tributaries
Chiola camura Jordan and Meek, Proc. U. S. Nat. Mus., VII, 1884, 474,
Arkansas River, Fort Lyon, Colo.
945. *Erogala nivea* (Cope).
Southern Virginia to South Carolina.
Hybopsis nveus Cope, Proc. Amer. Phil. Soc., XI, 1870, 460, Catawba
River, N. C.
946. *Erogala chloristia* (Jordan and Brayton).
Santee River and tributaries.
Codoma chloristia Jordan and Brayton, Bull. U. S. Nat. Mus., XII,
1878, 21, tributary of Saluda River, Greenville, S. C.

947. *Erogala trichroistia* (Jordan and Gilbert).
Alabama River Basin.
Codoma trichroistia Jordan and Gilbert, Bull. U. S. Nat. Mus., XII, 1878, 50, Etowah River, etc., Rome, Ga.
948. *Erogala callistia* (Jordan).
Alabama River Basin
Photogenis callistius Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 337, Etowah River, etc., Rome, Ga.
949. *Erogala cærulea* (Jordan).
Alabama River Basin.
Photogenis cæruleus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 338, Etowah River, Rome, Ga.
950. *Erogala eurystoma* (Jordan).
Chattahoochee River.
Photogenis eurystomus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 356, Nancys Creek, tributary to Chattahoochee River, near Atlanta, exclusive of the smaller specimens mentioned, which are *Notropis zonistis*; not *Codoma eurystoma* Jordan.
Photogenis leucopus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 41, Chattahoochee River, Gainesville, Ga.
951. *Erogala cercostigma* (Cope).
Sandy streams tributary to the Gulf of Mexico, from Pearl River and Red River to Rio Nueces; abundant.
Cyprinella cercostigma Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 157, Pearl River, Monticello, Miss.
Luxilus chickasawensis Hay, Proc. U. S. Nat. Mus., III, 1880, 506, Chickasawha River, Enterprise, Miss.
952. *Erogala xænura* (Jordan).
Altamaha River Basin, Ga.
Mimulus xænurus Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 79, Ocmulgee River, Flat Shoals, Ga.
953. *Erogala notata* (Girard).
Streams of Texas from Austin westward.
Cyprinella notata Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 198, Rio Seco, Tex.
954. *Erogala venusta* (Girard).
Streams of Texas.
Cyprinella venusta Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 198, Rio Sabinal, Tex.
955. *Erogala texana* (Girard).
Rivers of Texas from Trinity River to Rio Salado
Cyprinella texana Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 198, Rio Salado and Turkey Creek, Tex.
- Genus 329. *CYPRINELLA* Girard.
Cyprinella Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 196 (*Leuciscus bubalinus* Baird and Girard).
956. *Cyprinella bubalina* (Baird and Girard).
Streams of Arkansas and westward; not common.
Leuciscus bubalinus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 391, Otter Creek, Ark.
Cyprinella umbrosa Girard, loc. cit., VIII, 1856, 197, Coal Creek, tributary of South Fork Canadian River.
Cyprinella gunnisoni Girard, loc. cit., 197, Cottonwood Creek of the Rio Grande, Colo.
Cyprinella beckwithi Girard, loc. cit., 197, Arkansas River, near Fort Makee.
Cyprinella lepida Girard, loc. cit., 197, Rio Frio, Tex.
957. *Cyprinella ludibunda* Girard.
Rio Grande Basin.
Cyprinella ludibunda Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 199, locality now known to be Cottonwood Creek, Colo.

958. *Cyprinella macrostoma* Girard.
 Rio Grande region, northward perhaps to Kansas.
Cyprinella macrostoma Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 198, Devils River, Tex.
 ?*Cyprinella luxloldes* Girard, loc. cit., 198, San Pedro Creek, Tex.
 ?*Cyprinella lugubris* Girard, loc. cit., 199, Cottonwood Creek, Colo.
 ?*Notropis umbrifer* Hay, Proc. U. S. Nat. Mus., X, 1887, 245, Solomon River at Beloit and Saline River near Wakeeney, Kans.
959. *Cyprinella forlonensis* (Meek).
 Streams of Tamaulipas, Mexico.
Nototropis forlonensis Meek, Field Col. Mus., Zool., V, 1904, 70, fig. 20, Rio Forlon, Tamaulipas, Mexico.
960. *Cyprinella rubripinna* Garman.
 Tributaries of Lago del Muerto, Coahuila, Mexico.
Cyprinella rubripinna Garman, Bull. Mus. Comp. Zool., VIII, 1881, 91, Lago del Muerto; preoccupied.
Notropis garmani Jordan, Cat. Fish. N. Amer., 1885, 813; substitute for *rubripinna*; preoccupied in *Luxilus* but not in *Cyprinella*.
- Genus 330. *MONIANA* Girard.
Moniana Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 199 (*Leuciscus lutrensis* Baird and Girard); probably inseparable from *Cyprinella*.
961. *Moniana lutrensis* (Baird and Girard). *Redfin*.
 Southern Illinois to South Dakota, Kansas, and the Rio Grande; very abundant in clear brooks southward.
Leuciscus lutrensis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 391, Otter Creek, tributary of North Fork of Red River, Ark.
Cyprinella suavis Girard, loc. cit., VIII, 1856, 197, San Antonio, Tex.
Moniana leonina Girard, loc. cit., 199, Leon River, tributary to Rio San Antonio, Tex.
Moniana complanata Girard, loc. cit., 200, Brownsville, Tex.
Moniana latibilis Girard, loc. cit., 200, Hurah Creek, tributary of Rio Pecos.
Moniana pulchella Girard, loc. cit., 200, Sugar Loaf Creek, tributary to the Poteau River.
Moniana couchi Girard, loc. cit., 201, China, Nuevo Leon, Mexico.
Moniana rutila Girard, loc. cit., 201, Cadereita, Nuevo Leon, Mexico.
Moniana gracilis Girard, loc. cit., 201, Monterrey, Nuevo Leon, Mexico.
Moniana gibbosa Girard, loc. cit., 201, Brownsville, Tex.
Cyprinella billingsiana Cope, Hayden's Geol. Surv. Wyo., 1870, 439, St. Joseph, Mo.
Moniana jugalis Cope, loc. cit., 439, St. Joseph, Mo.
Hypsilepis iris Cope, Wheeler Surv., Zool., V, 1875, 653, Rio Grande, San Ildefonso, N. Mex.
Cyprinella forbesi Jordan, Bull. Ill. State Lab. Nat. Hist., I, 1878, 57, clear streams of southern Illinois.
Chiola montiregis Cope, Proc. Amer. Phil. Soc., XII, 1885, 168 (read Dec. 7, 1884), Monterey, Nuevo Leon, Mexico.
962. *Moniana formosa* Girard.
 Rio Mimbres, Chihuahua, Mexico.
Moniana formosa Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 201, Rio Mimbres, Chihuahua, Mexico.
963. *Moniana frigida* Girard.
 Rio Frio and tributaries of Rio San Antonio, Tex., and Lake Santa Maria, Chihuahua, Mexico.
Moniana frigida Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 200, Rio Frio, Tex.
964. *Moniana santamarie* (Evermann and Goldsborough).
 Rio Grande Basin in Mexico.
Notropis santamarie Evermann and Goldsborough, Bull., U. S. Fish Com., XXI, 1901 (1902), 147, fig. 2, Lake Santa Maria in Chihuahua, Mexico.

965. *Moniana proserpina* Girard.
Rio Grande region
Moniana proserpina Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 200, Devils River, Tex.
Moniana aurata Girard, loc. cit., 200, Piedra Pointe, N. Mex.
? *Moniana tristis* Girard, loc. cit., 1856, 201, locality unknown.
966. *Moniana boucardi* (Gunther). *Salmichi*.
Central Mexico.
Leuciscus boucardi Gunther, Cat., VII, 1868, 485, Cuernavaca, Mexico.
Graodius nigrotæniatus Gunther, loc. cit., 485, Atlusco, Mexico.
967. *Moniana robusta* (Meek).
Santa Rosalia and Jimenez, Mexico.
Notropis robustus Meek, Field Col. Mus., Zool., III, 1902, 82, Rio Conchos, Jimenez, Chihuahua, Mexico, figured as *Nototropis robustus*, in Field Col. Mus., Zool., V, 1904, 66.
968. *Moniana chihuahua* (Woolman).
Streams of Chihuahua, Mexico.
Notropis chihuahua Woolman, Amer. Nat., March, 1892, 260, Rio de los Conchos, Chihuahua, Mexico.
969. *Moniana callisema* (Jordan).
Ocmulgee River, Ga.
Episema callisema Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 363, South Fork of Ocmulgee River, Flat Shoals, Ga.
- Genus 331. HUDSONIUS Girard.
Hudsonius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 210 (*Clupea hudsonia* Clinton)
970. *Hudsonius chamberlaini* (Evermann).
Atchafalaya River, La.
Notropis chamberlaini Evermann, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2800, Atchafalaya River, Melville, La.
971. *Hudsonius hudsonius* (De Witt Clinton). *Spawneater*; *Spot-tailed minnow*; *Shiner*; *Smelt*.
The Dakotas and Lake Superior to New York, and southward; abundant in the Great Lakes.
Clupea hudsonia De Witt Clinton, Ann. Lyc. Nat. Hist. N. Y., I, 1824, 49, Hudson River.
Hudsonius fluviatilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 210, Hudson River.
Luxilus selene Jordan, Bull. U. S. Nat. Mus., X, 1877, 60, Lake Superior, Bayfield, Wis.
Notropis scopiferus Eigenmann and Eigenmann, Amer. Nat., 1893, 153.
Winnipeg, Brandon, Fort Qu'Appelle and Medicine Hat, Assiniboia.
972. *Hudsonius amarus* Girard. *Gudgeon*; *Spawneater*; *Silverfin*; *Shiner*; *Smelt shiner*; *Silvery minnow*; *Blue-lined minnow*.
Delaware and Potomac Rivers.
Hudsonius amarus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 210, Potomac River at Washington.
Hybopsis phaenna Cope, loc. cit., 1864, 279, Delaware River, Trenton, N. J.
973. *Hudsonius saludanus* (Jordan and Brayton).
Coastwise streams from the James to the Ocmulgee.
Alburnops saludanus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 16, a tributary of Saluda River, Greenville, S. C.
Hudsonius euryopa Bean, Proc. U. S. Nat. Mus., II, 1879, 285, McBean Creek, tributary of Savannah River, Ga.
974. *Hudsonius aletes* Jordan and Evermann.
Southern Indiana.
Hudsonius aletes Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, 1927 (Apr. 27), 502, Switz City Swamp, Green County, Ind.

Genus 332. HYBOPSIS Agassiz.

Hybopsis Agassiz, Amer. Jour. Sci. Arts, XVII, 1854 (*H. gracilis* Agassiz = *Rutilus amblops* Rafinesque?).

Alburnops Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 194 (*Hybopsis gracilis* Agassiz = *A. blennius* Girard).

Graodus Gunther, Cat., VII, 1868, 485 (*Graodus nigrotæniatus* Girard, specimen with pharyngeal teeth lost = *Alburnops blennius* Girard).

Chrope Jordan, Hayden's Geol. Surv. Terr., IV, 1878 (*Hybopsis bifrenatus* Cope).

Minellus Jordan, Man. Vert., ed. 5, 1888, 56 (*Hybopsis procne* Cope).

975. *Hybopsis gracilis* Agassiz.

Tennessee River.

Hybopsis gracilis Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 358, Huntsville, Ala.

976. *Hybopsis blennius* (Girard). *Straw-colored minnow*.

Ohio, Michigan, and New York to Tennessee, Dakota, and Kansas, thence southwestward to Texas.

?*Minnilus macrostomus* Rafinesque, Ichth. Ohiensis, 1820, 47, Kentucky River; not certainly identifiable.

?*Rutilus amblops* Rafinesque, loc. cit., 51; not certainly identifiable.

Alburnops blennius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 194, Arkansas River, near Fort Smith, Ark.

Hybognathus stramineus Cope, loc. cit., XVI, 1864, 283, Detroit River, Grosse Isle.

Hybopsis missouriensis Cope, Hayden's Geol. Surv. Wyo., 1870, 437, Missouri River, St. Joseph, Mo.

Notropis reticulatus Eigenmann and Eigenmann, Amer. Nat., 1893, 152, Fort Qu'Appelle, Assiniboia.

977. *Hybopsis winchelli* Girard.

Black Warrior River, Ala.

Hybopsis winchelli Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 211, Black Warrior River, Ala.

This species and the preceding may prove to be identical with *Hybopsis gracilis*. The synonymy of these little fishes is still unsettled.

978. *Hybopsis procne* (Cope). *Minnow; Minnie; Swallowminnow shiner*.

Delaware River and Cayuga Lake Basin southward in coastwise streams as far as the Neuse River, N. C.

Hybognathus procne Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 283, Conestoga River, a tributary of the Susquehanna.

Hybopsis longiceps Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 231, headwaters of James and Roanoke Rivers, Va.

979. *Hybopsis cummingsi* (Myers).

Streams near Wilmington, N. C.

Notropis cummingsi Myers, Amer. Mus. Novitates, No. 168, April 23, 1925, 1, fig. 1, upper Burnt Mill Creek, Wilmington, N. C.

980. *Hybopsis cayuga* (Meek).

Connecticut River, northern New York, and upper Mississippi Valley.

Notropis cayuga Meek, Ann. Acad. Sci. N. Y., IV, 1888, 305, Cayuga Lake, N. Y.

981. *Hybopsis atrocaudalis* (Evermann).

Rio Neches, Palestine, Tex., also in Rio Comal at New Braunfels, Tex.

Notropis cayuga atrocaudalis Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 76, Neches River, Palestine, Tex.

982. *Hybopsis keimi* (Fowler).

Allegheny River, western Pennsylvania.

Notropis keimi Fowler, Proc. Acad. Nat. Sci. Phila., LX, 1908, 533, Plate XXVII, tributary of Allegheny River, Port Allegheny, Pa.

983. *Hybopsis volucella* (Cope).

Michigan, Wisconsin, and northern Indiana.

Hybognathus volucellus Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 283, Grosse Isle, Detroit River, Mich.

984. *Hybopsis phenacobius* (Forbes).
Streams of central Illinois.
Notropis phenacobius Forbes, Bull. Ill. Lab. Nat. Hist., II, 1885, 2, 137, Peoria Ill.
985. *Hybopsis spectrunculus* Cope.
Headwaters of Tennessee River in Tennessee, Virginia, and North Carolina.
Hybopsis spectrunculus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 231, tributary of Middle Fork of Holston River, Va.
986. *Hybopsis shumardi* (Girard).
Missouri region and southwestward.
Alburnops shumardi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 194, Arkansas River, Fort Smith, Ark.
Alburnops illecebrosus Girard, loc. cit., 194, Arkansas River, Fort Smith, Ark.
987. *Hybopsis boops* (Gilbert). *Bigeye minnow*.
Maumee River, west to Iowa and Missouri.
Notropis boops Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 201, Salt Creek, Brown County, Ind., and Flat Rock Creek, Decatur County, Ind.
988. *Hybopsis kanawha* (Jordan and Jenkins).
Tributaries of Kanawha River, southwestern Virginia.
Notropis kanawha Jordan and Jenkins, Proc. U. S. Nat. Mus., XI, 1888, 354, Plate 44, fig. 5, Reed Creek, Wytheville, Va.
989. *Hybopsis piptolepis* (Cope).
North Platte River.
Photogenus piptolepis Cope, Hayden's Geol. Surv. Wyo., 1870, 438, Red Cloud Creek, a tributary of North Platte River.
990. *Hybopsis gilberti* (Jordan and Meek).
Illinois to eastern Colorado.
Notropis gilberti Jordan and Meek, Proc. U. S. Nat. Mus., VIII, 1885, 4, Des Moines River and Village Creek, Ottumwa, Iowa; perhaps not distinct from *Hybopsis piptolepis*.
991. *Hybopsis horatii* (Cockerell).
Platte Basin.
Notropis horatii Cockerell, Science, XXXIV, No. 879, Nov. 3, 1911, 614, Julesburg, Colo.
992. *Hybopsis deliciosa* (Girard). *Straw-colored minnow*.
Michigan, and southwest to Arkansas.
? *Minnilus macrostomus* Rafinesque, Ichth. Ohiensis, 1820, 47, Ohio River; not certainly identifiable.
? *Hybopsis dorsalis* Agassiz, Amer. Jour. Sci. Arts., XVII, 1854, 358, Burlington, Iowa; unidentifiable.
Moniana deliciosa Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 199, Leon River, a tributary of the San Antonio River, Tex.
993. *Hybopsis deliciosa straminea* Cope. *Northern straw-colored minnow*.
Great Lakes region, mostly in small brooks, the northern form of *H. deliciosus*.
Hybopsis stramineus Cope, Trans. Amer. Phil. Soc., XIII, 1886, 381, Grosse Isle, Detroit River.
994. *Hybopsis callientis* (Jordan and Snyder).
Streams of central Mexico.
Notropis callientis Jordan and Snyder, Bull. U. S. Fish. Com., XIX, 1899 (1900), 122, fig. 4, Rio Verde at Aguas Calientes, Mexico.
995. *Hybopsis buchanani* (Meek).
Poteau River, Okla.
Notropis buchanani Meek, Bull. U. S. Fish. Com., XV, 1895 (1896), 342, Poteau River near Poteau, Ind. Ter.
996. *Hybopsis heterolepis* (Eigenmann and Eigenmann).
Notropis heterolepis Eigenmann and Eigenmann, Amer. Nat., 1893, 152, Cayuga Lake.
Notropis muskoka Meek, Field Col. Mus., Zool., I, 1898, 308, Muskoka Lake near Georgian Bay, Ontario.

997. *Hybopsis kendalli* (Evermann and Cockerell).
Cross Lake Thoroughfare, Me.
Notropis kendalli Evermann and Cockerell, Proc. Biol. Soc. Wash., XXII,
Dec. 8, 1909, Cross Lake Thoroughfare, Me.
998. *Hybopsis ozarcanus* (Meek).
White River, Ark.
Notropis ozarcanus Meek, Bull., U. S. Fish Com., IX, 1889 (1891), 129,
North Fork of White River, Ark.
999. *Hybopsis scylla* Cope.
Illinois River to western Kansas and Montana, abundant in the shallow
sandy streams of the plains.
Alburnus lineolatus Agassiz, in Putnam, Bull., Mus. Comp. Zool., I,
1863, 9, Osage River, Mo.; unidentifiable.
Hybopsis scylla Cope, Hayden's Geol. Surv. Wyo., 1870, 438, Red
Cloud Creek, tributary of Platte River.
Chola chlora Jordan, Bull. Hayden's Geol. Surv. Terr., IV, 1878, 791,
upper Missouri River.
1000. *Hybopsis longirostris* (Hay).
Chickasawha River, Miss., to Escambia River, Fla.
Alburnops longirostris Hay, Proc., U. S. Nat. Mus., III, 1880, 504,
Chickasawha River at Enterprise, Miss.
1001. *Hybopsis nocomis* (Evermann).
Trinity, San Marcos, and Comal Rivers, Tex.
Notropis nocomis Evermann, Bull. U. S. Fish Com., XI, 1891 (1892),
78, Trinity River, Palestine, Tex., and San Marcos River, San
Marcos, Tex.
1002. *Hybopsis ortenburgeri* (Hubbs)
Mountain Fork River, Okla.
Notropis ortenburgeri Hubbs, Proc. Okla. Acad. Sci., pt. 1, VI, Jan. 1,
1927, 127, Mountain Fork River, McCurtain County, Okla.
1003. *Hybopsis nux* (Evermann).
Neches and Trinity Rivers near Palestine, Tex.
Notropis nux Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 77,
Neches River, Palestine, Tex.
1004. *Hybopsis simus* (Cope).
Rio Grande at San Ildefonso, N. Mex.
Alburnellus simus Cope, Wheeler Surv., V, Zool., 1875, 649, San Ilde-
fonso, N. Mex.
1005. *Hybopsis sabineæ* (Jordan and Gilbert).
Sabine River, Tex.
Notropis sabineæ Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1886,
15, Sabine River at Longview, Tex.
1006. *Hybopsis nitida* (Girard).
Tributaries of Rio San Juan at Cadereita, Nuevo Leon, Mexico.
Moniana nitida Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 201,
Cadereita, Mexico; not *Alburnus nitidus* Kirtland, also a *Notropis*
but not a *Hybopsis*.
Notropis braytoni Jordan and Evermann, Fishes North and Mid.
Amer., 1896, 264, Cadereita, Mexico; substitute for *M. nitida*
Girard, preoccupied.
- Subgenus **CHRIOPE** Jordan.
1007. *Hybopsis bifrenata* Cope. *Bridled minnow*; *Minnæ*.
Maine to Maryland.
Hybopsis bifrenatus Cope, Trans., Amer. Phil. Soc., XIII, 1867, Schuyl-
kill River, Conshohocken, Pa.
1008. *Hybopsis maculata* (Hay).
Chickasawha River, Miss.
Hemitremia maculata Hay, Proc. U. S. Nat. Mus., III, 1880, 1505,
Chickasawha River, Enterprise, Miss.

1009. *Hybopsis anogenus* (Forbes).
Western New York (Cayuga Lake) to northern Illinois.
Notropis anogenus Forbes, Bull. Ill. Lab. Nat. Hist., II, 1885, Art. 2, 138, Fox River, McHenry, Ill.
1010. *Hybopsis heterodon* (Cope).
New York to Michigan and Kansas; common.
Alburnops heterodon Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 281, Lansing, Grosse Isle, Mich.
Notropis germanus Hay, Proc. U. S. Nat. Mus., X, 1887, 252, Smoky Hill River, Wallace, Kans.
1011. *Hybopsis heterodon richardsoni* (Hubbs and Greene).
Fox River System, tributary to Green Bay, Wis.
Notropis heterodon richardsoni Hubbs and Greene, Univ. Mich. Mus. Zool., Misc. Pub. 15, 1926, 39, Pl. III, fig. 1, Fox River, opposite Lock 25, Columbia County, Wis.
1012. *Hybopsis jordani* (Eigenmann and Eigenmann).
South Saskatchewan River, Medicine Hat, Assiniboia.
Notropis albeolus Eigenmann and Eigenmann, Amer. Nat., 1893, 152, Medicine Hat, Assiniboia; name preoccupied.
Notropis jordani Eigenmann and Eigenmann, loc. cit., 592, substitute for *Chriope albeolus*, preoccupied in *Notropis* but not in *Chriope*.
- Genus 333. **PLATYGOBIO** Gill.
Platygobio Gill, Trans. Amer. Phil. Soc., V, 1863, 178 (*Pogonichthys communis* Girard = *Cyprinus* (*Leuciscus*) *gracilis* Richardson).
1013. *Platygobio gracilis* (Richardson). *Flat-headed chub*.
East slope of the Rocky Mountains, from the Missouri and Yellowstone Rivers to the Saskatchewan; abundant in river channels as far south as Kansas City.
Cyprinus (*Leuciscus*) *gracilis* Richardson, Fauna Bor.-Amer., Fishes, 1836, 120, Saskatchewan River at Carleton House.
Coregonus angusticeps Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 389-390 (534), Saskatchewan River.
Pogonichthys communis Girard, Proc., Acad. Nat. Sci. Phila., VIII, 1856, 188, Yellowstone River, Fort Pierre, Nebr., etc.
Pogonichthys (*Platygobio*) *gulonellus* Cope, loc. cit., XVI, 1864, 277, near Bridgers Pass, Wyo.
1014. *Platygobio physignathus* (Cope).
Upper waters of Arkansas River; very abundant in the channels; the most common fish at Pueblo, Colo.
Ceratichthys physignathus Cope, Wheeler Surv., V, Zool., 1875 (1876), 651, Arkansas River, Pueblo, Colo.
1015. *Platygobio pallidus* Forbes.
One specimen from Ohio River at Cairo, Ill.
Platygobio pallidus Forbes, in Jordan and Gilbert, Synopsis, 1883, 220, Ohio River at Cairo, Ill.
- Genus 334. **COUESIUS** Jordan.
Couesius Jordan, Hayden's Geol. Surv. Terr., IV, 1878, 785 (*Nocomis milneri* Jordan = *Gobio plumbeus* Agassiz).
1016. *Couesius squamilentus* (Cope).
Henry Fork of Green River in southwestern Wyoming.
Ceratichthys squamilentus Cope, Hayden's Geol. Surv. Wyo., 1870, 442, Henry Fork of Green River, Wyo., Colorado Basin.
1017. *Couesius plumbeus* (Agassiz). *Chub minnow*; *Lake chub*; *Plumbeus minnow*; *Moose Lake minnow*.
Streams and lakes from Lake Superior east to the Adirondack region and New Brunswick; not very common except northward.
Gobio plumbeus Agassiz, Lake Superior, 1850, 366, Lake Superior.
Ceratichthys prosthemi Cope, Trans. Amer. Phil. Soc., XIII, 1867, Montreal River, Keweenaw Point, Lake Superior.
Nocomis milneri Jordan, Bull. U. S. Nat. Mus., X, 1877, 64, Lake Superior.

- 1018. *Couesius greeni* Jordan.**
 Stuart Lake, headwaters of Fraser River, British Columbia, and Lake
 Pend d'Oreille, Idaho.
Couesius greeni Jordan, Proc. U. S. Nat. Mus., XVI, 1893, 313, Stuarts
 Lake, British Columbia.
- 1019. *Couesius dissimilis* (Girard).**
 Upper Missouri and Black Hills region.
Leucosomus dissimilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
 189, Milk River and Little Muddy River, Mont.
- 1020. *Couesius adustus* Woolman.**
 Rio de los Conchos, Chihuahua, Mexico
Couesius adustus Woolman, Bull. U. S. Fish Com., XIV, 1894, 57, Rio
 de los Conchos, Chihuahua, México.
- Genus 335. NOCOMIS Girard.**
Nocomis Girard, Proc. Acad. Nat. Sci. Phila., 1856, 190 (*N. nebrascensis*
 Girard = *Luxilus kentuckiensis* Rafinesque)
Ceratichthys Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853,
 391 (*C. nigralis* Baird and Girard; not of Baird and Girard, 1856); no
 description.
- 1021. *Nocomis kentuckiensis* (Rafinesque).** *Chub; Hornyhead; River chub;*
Jerker; Indian chub.
 Pennsylvania to Wyoming and Alabama, on both sides of the Alle-
 ghenies; abundant in the larger streams
Luxilus kentuckiensis Rafinesque, Ichth. Ohioensis, 1820, 48, Ohio River.
 ?*Catostomus melanotus* Rafinesque, loc. cit., 58, Ohio River at the falls.
Semotilus biguttatus Kirtland, Bost. Jour. Nat. Hist., III, 1840, 344, Pl. V,
 Yellow Creek, Ohio.
Nocomis nebrascensis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
 190, Sweetwater River, Nebr.
Ceratichthys leptocephalus Girard, loc. cit., 213, Salem, N. C.
Nocomis bellicus Girard, loc. cit., 213, Black Warrior River.
Ceratichthys cyclois Cope, loc. cit., XVI, 1864, 277, Grosse Isle, Water-
 ford; Clinton River and Bruce, Mich.
Ceratichthys stigmaticus Cope, loc. cit., 1864, 278, Michigan.
- 1022. *Nocomis micropogon* (Cope).**
 Streams of Pennsylvania.
Ceratichthys micropogon Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864,
 277, Conestoga River near Lancaster, Pa
- Genus 336. YURIRIA Jordan and Evermann**
Yuriria Jordan and Evermann, Fishes North and Mid Amer., 1896,
 321 (*Hudsonius altus* Jordan).
- 1023. *Yuriria alta* (Jordan).** *Pesca blanca.*
 Lakes and streams of Guanajuato, tributary to Rio Lerma, Mexico, in
 Pacific drainage.
Hudsonius altus Jordan, Proc. U. S. Nat. Mus., II, 1879, 301, Lake
 Tupatato, Guanajuato, Mexico.
- Genus 337. ERINEMUS Jordan**
Erinemus Jordan, Man. Vert., 1876, 279 (*Ceratichthys hyalinus* Cope).
- 1024. *Erinemus storerianus* (Kirtland).** *Storer's chub; Lake minnow.*
 Great Lakes and westward to Wyoming, Tennessee, and Arkansas;
 abundant in the larger streams and lakes, especially in Iowa
Rutilus storerianus Kirtland, Proc. Bost. Soc. Nat. Hist., I, 1842, 71,
 Lake Erie (name only); description given as *Leuciscus storerianus* in
 Bost. Jour. Nat. Hist., V, 1845, 30, Pl. I, fig. 2,
Gobio vernalis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 189,
 Arkansas River, Fort Smith, Ark.
Ceratichthys lucens Jordan, Proc. U. S. Nat. Mus., II, 1879, 238, Falls
 of Ohio River near New Albany, Ind.
- 1025. *Erinemus hyalinus* (Cope).** *Silver chub.*
Ceratichthys hyalinus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 226,
 tributary of Holston River.

1026. *Erinemus labrosus* (Cope).
Basin of the Santee in North and South Carolina, common.
Ceraticthys labrosus Cope, Proc. Amer. Phil. Soc., XI, 1870, 458,
tributaries of Catawba River, McDowell and Burke Counties, N. C.
Ceraticthys zanemus Jordan and Brayton, Bull. U. S. Nat. Mus., XII,
1878, 24, Saluda River, Greenville, N. C.
1027. *Erinemus hypsinotus* (Cope).
Santee Basin, in North and South Carolina
Ceraticthys hypsinotus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870,
458, tributaries of Catawba River, McDowell County, N. C.
1028. *Erinemus rubrifrons* (Jordan).
Basin of the Altamaha, Ga.
Nocomis rubrifrons Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876,
330, Ocmulgee River, Flat Rock, Ga.
- Genus 338. **ERIMYSTAX** Jordan.
Erimystax Jordan, Rept. Geol. Surv. Ohio, IV, 1882, 858 (*Luxilus dis-*
similis Kirtland).
1029. *Erimystax dissimilis* (Kirtland). *Spotted shiner*.
Lake Erie to the headwaters of the Tennessee and west to Arkansas
and Iowa.
Luxilus dissimilis Kirtland, Bost. Jour. Nat. Hist., III, 1840, 341, pl.
4, fig. 2, Mahoning River, Ohio, and Lake Erie near Cleveland,
Ohio.
1030. *Erimystax watauga* (Jordan and Evermann).
Holston River, Va.; Watauga River, Tenn.; White River, Ark.;
Big Barren River, Ky.; and Tippecanoe River, Ind.
Hybopsis watauga Jordan and Evermann, Proc. U. S. Nat. Mus., XI,
1888, 355, Watauga River, Elizabethtown, Tenn.; North Fork of
Holston River, Saltville, Va.
- Genus 339. **ERIMONAX** Jordan.
Erimonax Jordan, Copeia, 1924, 52 (*Ceraticthys monacus* Cope).
1031. *Erimonax monacus* (Cope).
Tennessee Basin in the river channels.
Ceraticthys monacus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 227,
Holston River, Washington County, Va.
- Genus 340. **OREGONICHTHYS** Hubbs.
Oregonichthys Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., 1927
(*Hybopsis crameri* Snyder).
1032. *Oregonichthys crameri* (Snyder).
Hybopsis crameri Snyder, Bull. U. S. Bur. Fish., XXVII, 1907, 181,
fig. 5, Willamette River at Oregon City, Oreg.
1033. *Oregonichthys cumingii* (Günther).
California; only the type known.
Ceraticthys cumingii Günther, Cat., VII, 1868, 177, California; type
locality uncertain.
- Genus 341. **MACRHYBOPSIS** Cockerell and Alliston.
Macrhybopsis Cockerell and Alliston, Proc. Biol. Soc. Wash., XXII,
1909, 162 (*Gobio gelidus* Girard).
1034. *Macrhybopsis æstivalis* (Girard).
Arkansas River to the Rio Grande; abundant in sandy river channels,
not in small brooks.
Gobio æstivalis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 189,
Rio San Juan, near Cadereita, Nuevo Leon.
1035. *Macrhybopsis marconis* (Jordan and Gilbert).
Abundant in Rio San Marcos, San Marcos, Tex.; also known from the
Guadalupe River near San Marcos and the Rio Comal at New
Braunfels, Tex.
Hybopsis æstivalis marconis Jordan and Gilbert, Proc. U. S. Nat. Mus.,
IX, 1886, 22, Rio San Marcos, San Marcos, Tex.

1036. *Macrhybopsis sterletus* (Cope).
Rio Grande.
Ceraticthys sterletus Cope, Wheeler Surv., V, Zool., 1875 (1876), 652,
Rio Grande, at San Ildefonso, New Mexico.
1037. *Macrhybopsis gelida* (Girard).
Middle Missouri River Basin, from Wyoming to eastern Nebraska.
Gobio gelidus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 188,
Milk River, Mont.
1038. *Macrhybopsis montana* (Meek).
Type supposed to be from the upper Missouri.
Hybopsis montanus Meek, Proc. U. S. Nat. Mus., VII, 1884, 526,
locality unknown but collected by Dr. F. V. Hayden, probably in
the upper Missouri Basin.
1039. *Macrhybopsis meeki* (Jordan and Evermann).
Missouri River at St. Joseph and elsewhere in the river channel.
Hybopsis meeki Jordan and Evermann, Fish. North and Mid. Amer.,
1896, 317, Missouri River at St. Joseph, Mo.
1040. *Macrhybopsis hyostoma* (Gilbert).
Indiana to Iowa, and south to the Alabama River.
Nocomis hyostomus Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 203,
White River, Bedford, Ind.
- Genus 342. *EXTRARIUS* Jordan.
Extrarvus Jordan, Proc. Acad. Nat. Sci. Phila., LXX, 1918, 342 (*Hy-*
bopsis tetranemus Gilbert).
1041. *Extrarius tetranemus* (Gilbert).
Tributaries of Arkansas River in Kansas and Arkansas.
Hybopsis tetranemus Gilbert, Bull. Washburn Lab. Nat. Hist., 1886,
208, Elm and Spring Creeks, Medicine Lodge, Kans.
- Genus 343. *RHINICHTHYS* Agassiz.
Argyreus Heckel, Russeger's Reisen, 1842 (*Cyprinus atronasus* Mitchell;
name preoccupied).
Rhinichthys Agassiz, Lake Superior, 1850, 353 (*Cyprinus atronasus*
Mitchell).
1042. *Rhinichthys cataractæ* (Cuvier and Valenciennes). *Long-nosed dace*;
Black-nosed dace; *Black minnow*; *Niagara gudgeon*.
New England to Virginia and Wisconsin
Gobio cataractæ Cuvier and Valenciennes, Hist. Nat. Poiss., XVI, 1842,
240 (315), pl. 483, Niagara Falls.
Leuciscus nasutus Ayres, Jour. Bost. Soc. Nat. Hist., IV, 1843, 290,
West Hartford, Conn.
Rhinichthys marmoratus Agassiz, Lake Superior, 1850, 354, Pl. II, figs.
1 and 2, Sault Ste. Marie, Lake Superior.
1043. *Rhinichthys bowersi* Goldsborough and Clark.
Rhinichthys bowersi Goldsborough and Clark, Bull. U. S. Bur. Fish.,
XXVII, 1907, 36, Dry Fork, Herman, W. Va.
1044. *Rhinichthys dulcis* (Girard).
Upper Missouri, Platte, Arkansas, Rio Grande, and Colorado Rivers;
Columbia River; Utah Basin; and coastwise streams of Washington
and Oregon.
Argyreus dulcis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 185,
Sweetwater River, Nebr.
Rhinichthys maxillosus Cope, loc. cit., XVI, 1864, 278, Kansas
Rhinichthys transmontanus Cope, Amer. Nat., 1879, 441, tributaries of
Rio Grande in Colorado and New Mexico.
Rhinichthys ocella Garman, Science Observer, III, 1881, 59, northeastern
Wyoming and Montana.
Rhinichthys luteus Garman, Bull. Mus. Comp. Zool., VIII, 1881, 87,
Bear River, Ogden, Utah.

1045. *Rhinichthys atronasmus* (Mitchill) *Dace; Black-nosed dace; Brook minnow; Striped dace*
New England to Minnesota, northern Alabama, and Virginia, very abundant in clear brooks and mountain streams.
Cyprinus atronasmus Mitchill, Trans Lit. Phil. Soc. N. Y., I, 1815, 460, Wallkill River, brooks of New York.
Cyprinus vitatus Rafinesque, Amer. Month. Mag., II, 1817, 121, upper Hudson River.
1046. *Rhinichthys croceus* (Storer)
Tennessee Basin; abundant in clear brooks.
Leuciscus croceus Storer, Proc. Bost. Soc. Nat. Hist., 1845, 48, Florence, Ala.
Rhinichthys obtusus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 357, Huntsville, Ala.
Rhinichthys badius Garman, Science Observer, III, 1881, 60, Clinch River, Tenn.
1047. *Rhinichthys lunatus* Cope.
Lakes and brooks of Michigan, Indiana, Wisconsin, and Minnesota, and in Rainy River.
Rhinichthys lunatus Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 278, Grosse Isle, Mich.
Rhinichthys arenatus Garman, Science Observer, 1881, 62, Sand Hill River, Minn.
1048. *Rhinichthys meleagris* Agassiz.
Illinois and Iowa.
Rhinichthys meleagris Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 357, Burlington, Iowa.
1049. *Rhinichthys simus* Garman.
Coahuila, Mexico.
Rhinichthys simus Garman, Science Observer, III, 1881, 61, Coahuila, Mexico.
- Genus 344. *AGOSIA* Girard.
Agosia Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 186 (*Agosia chrysogaster* Girard)
Zophendum Jordan and Gilbert, Synopsis, 1883, 155 (*Hybognathus siderius* Cope = *Agosia chrysogaster* Girard).
1050. *Agosia chrysogaster* Girard.
Tributaries of the Gila River; common in the mountain brooks.
Agosia chrysogaster Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 187, Rio Santa Cruz, Sonora.
Agosia metallica Girard, loc. cit., 1856, 187, Rio San Pedro, Ariz.
Hyborhynchus (*Hybognathus*) *siderius* Cope, Wheeler Surv., V, Zool., 1875, 670, Pl. XXXI, figs. 6 and 6a, Camp Lowell, Ariz.
- Genus 345. *APOCOPE* Cope.
Apocope Cope, Hayden's Geol. Surv. Mont., etc., 1871 (1872), 472 (*A. carringtoni* Cope).
Eritrema Cope, Wheeler's Surv., V, Zool., 1876, 645 (*Apocope henshawi* Cope).
1051. *Apocope nubila* (Girard).
Basin of the Columbia River from western Idaho, below Shoshone Falls of Snake River, to the coast, and in coastwise streams from Washington southward into Oregon.
Argyreus nubilus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 186, Fort Steilacoom, Puget Sound.
1052. *Apocope robusta* (Rutter).
Upper Sacramento Basin.
Agosia robusta Rutter, Bull. U. S. Fish Com., XXII, 1902, 148, Prosser Creek, Calif.

1053. *Apocope carringtonii* Cope.
Upper Snake River Basin to Heart Lake in Yellowstone Park, thence extending southward in the Great Basin and the Coast Ranges; southeastern Oregon.
Apocope carringtoni Cope, Hayden's Geol. Surv. Mont., 1871, 472, Warm Springs, Utah.
Apocope vulnerata Cope, loc. cit., 473, Logan, Utah.
Tigoma rhinichthyoides Cope, loc. cit., 473, Logan, Utah.
Rhinichthys henshawi Cope, Proc. Amer. Phil. Soc., XIV, 1874, 133, Provo, Utah.
Agosia novemradiata Cope, Proc. Acad. Nat. Sci. Phila., XXV, 1883, 141, Weber River at Echo, Utah.
1054. *Apocope oscula* (Girard).
Lower Colorado and Gila Rivers, not rare.
Argyreus osculus Girard, Proc. Acad. Nat. Sci. Phila., 1856, 186, Babocomori Creek, a tributary of the Rio San Pedro, Ariz.
Argyreus notabilis Girard, loc. cit., 1856, 186, Rio Santa Cruz, Sonora, a tributary of the Rio San Pedro, Ariz.
Ceratichthys ventricosus Cope, Proc. Amer. Phil. Soc., XXV, 1874, 136, Ariz.
1055. *Apocope couesii* Yarrow.
Colorado River Basin.
Apocope couesii Yarrow, Field and Forest, 1876, and Wheeler Surv., V, Zool., 1875, 648, Camp Apache, Ariz.
1056. *Apocope yarrowi* (Jordan and Evermann).
Streams of Colorado in the Colorado River Basin.
Agosia yarrowi Jordan and Evermann, Bull. U. S. Fish Com., IX, 1889 (1891), 28, Tomichi Creek and Gunnison River, Gunnison, Colo.
1057. *Apocope adobe* (Jordan and Evermann).
Sevier River, Utah; locally abundant.
Agosia adobe Jordan and Evermann, Bull. U. S. Fish Com., IX, 1889 (1891), Sevier River, Juab, Utah.
1058. *Apocope klamathensis* (Evermann and Meek).
Agosia klamathensis Evermann and Meek, Bull. U. S. Fish Com., XVII, 1897 (1898), 75, 9, fig. 5, Pelican Bay, Upper Klamath Lake, Oreg.
1059. *Apocope nevadensis* (Gilbert).
Warm springs in the deserts of southwestern Nevada; Ash Meadows, Indian Creek, and Vegas Creek, Nev.
Rhinichthys (Apocope) nevadensis Gilbert, Death Valley Exped., 1893, 230, pl. 6, fig. 1, Ash Meadows, Amargosa Desert, Nev.
1060. *Apocope velifera* (Gilbert).
Pahranagat Valley, southwestern Nevada.
Rhinichthys (Apocope) velifer Gilbert, Death Valley Exped., 1893, 229, pl. 6, fig. 2, Pahranagat Valley, Nev.
1061. *Apocope falcata* (Eigenmann and Eigenmann).
Columbia River Basin; locally abundant; Boise River at Caldwell, Idaho; Payette River at Payette, Idaho; Columbia River at Pasco and Umatilla; and Mill Creek at Walla Walla.
Agosia falcata Eigenmann and Eigenmann, Amer. Nat., XXVII, Feb. 4, 1893, 153, Boise River, Caldwell, Idaho.
1062. *Apocope umatilla* (Gilbert and Evermann).
Columbia River at Umatilla, Oreg., and Payette and Salmon Rivers, Idaho.
Agosia umatilla Gilbert and Evermann, Bull. U. S. Fish Com., XIV, 1894, 192, pl. 9, fig. 2, Columbia River, Umatilla, Oreg.
- Genus 346. *ERICYMBA* Cope.
Ericymba Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 88 (*E. buccata* Cope).

1063. *Ericymba buccata* Cope. *Silver-mouthed minnow*.
Michigan and western Pennsylvania to Kansas and southward to western Florida
Ericymba buccata Cope, Proc., Acad. Nat. Sci. Phila., XVII, 1865, 88,
Kiskiminitas River, a tributary of the Monongahela, western Pennsylvania.
- Genus 347. **PHENACOBIVS** Cope.
Phenacobius Cope, Proc., Acad. Nat. Sci. Phila., XIX, 1867, 96 (*P. teretulus* Cope).
Sarcidium Cope, Hayden's Geol. Surv. Wyo., 1870, 363 (*S. scopiferum* Cope).
1064. *Phenacobius teretulus* Cope.
Kanawha River, W. Va.
Phenacobius teretulus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 96, Kanawha River, Eggleston Springs, W. Va.
1065. *Phenacobius mirabilis* (Girard). *Sucker-mouthed minnow*.
Illinois to Arkansas.
Exoglossum mirabile Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 191, Arkansas River, Fort Smith, Ark.
1066. *Phenacobius scopifer* (Cope).
Illinois to the Rio Grande.
Sarcidium scopiferum Cope, Hayden's Geol. Surv. Wyo., 1870 (1871), 440, Missouri River near St. Joseph, Mo.
Phenacobius teretulus liosternus Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 46, brooks of McLean County, Ill.
1067. *Phenacobius uranops* Cope.
Upper Tennessee Basin.
Phenacobius uranops Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 96, Holston River, Saltville, Va.
1068. *Phenacobius catostomus* Jordan.
Alabama River Basin.
Phenacobius catostomus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 332, Silver Creek, Etowah River, Rome, Ga.
- Genus 348. **FALCULARIUS** Jordan and Snyder.
Falcula Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 124 (*Falcula chapalæ* Jordan and Snyder); name preoccupied.
Falcularius Jordan, Amer. Nat., XXXVII, 1903, 360 (*Falcula chapalæ* Jordan and Snyder).
1069. *Falcularius chapalæ* (Jordan and Snyder).
Lake Chapala, Mexico
Falcula chapalæ Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1901), 124, Lake Chapala at Ocotlan, Mexico.
- Genus 349. **XYSTROSUS** Jordan and Snyder.
Xystrosus Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1901), 123 (*X. popoche* Jordan and Snyder).
1070. *Xystrosus popoche* Jordan and Snyder.
Basin of the Rio Lerma, Mexico.
Xystrosus popoche Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1901), 123, Lake Chapala at Ocotlan, Jalisco, Mexico.
- Genus 350. **EVARRA** Woolman.
Evarra Woolman, Bull. U. S. Fish Com., XIV, 1894, 64 (*E. eigenmanni* Woolman).
1071. *Evarra eigenmanni* Woolman.
Central Mexico.
Evarra eigenmanni Woolman, Bull. U. S. Fish Com., XIV, 1894, 64, City of Mexico.
1072. *Evarra tlahuacensis* Meek.
Chalco Lake, Tlahuac, Mexico.
Evarra tlahuacensis Meek, Field Col. Mus., Zool., III, 1902, 85, Pl. XVIII, lower figure, Chalco Lake, Tlahuac, Mexico.

Genus 351. TIAROGA Girard.

Tiaroga Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 204 (*T. cobitis* Girard).

1073. *Tiaroga cobitis* Girard.

Gila Basin, Ariz.

Tiaroga cobitis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 204, Rio San Pedro, Ariz.

Genus 352. AZTECULA Jordan and Evermann.

Azteca Jordan and Evermann, Fishes North and Mid. Amer., 1896, 254 (*Notropis aztecus* Woolman); name preoccupied.

Aztecula Jordan and Evermann, loc. cit., 1898, 2799, (*Notropis aztecus* Woolman=*Codoma vittata* Girard).

1074. *Aztecula vittata* (Girard).

Valley of Mexico and headwaters of the Rio Balsas at Pueblo, Mexico.

Codoma vittata Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 195, Valley of Mexico.

Notropis aztecus Woolman, Bull. U. S. Fish Com., XIV, 1894, 63, pl. 2, canals about City of Mexico

1075. *Aztecula mexicana* Meek.

Basin of Rio San Juan on the Mexican plateau.

Aztecula mexicana Meek, Field Col. Mus., Zool., III, 1902, 81, figured; loc. cit., V, 1904, 62, fig. 15; San Juan del Rio, Mexico.

1076. *Aztecula lermæ* (Evermann and Goldsborough).

Basin of Rio Lerma.

Notropis lermæ Evermann and Goldsborough, Bull. U. S. Fish. Com., XXI, 1901 (1902), 147, fig. 3, Lake Lerma, at Lerma, Mexico.

Genus 353. CODOMA Girard.

Codoma Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 194 (*C. ornata* Girard).

1077. *Codoma ornata* Girard.

Headwaters of the Rio Nezquital and Rio Nazas in Durango to upper tributaries of the Rio Yaqui and Rio Conchos in Chihuahua, Mexico.

Codoma ornata Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 195, Chihuahua River, Mexico.

1078. *Codoma topeka* (Gilbert).

Western Iowa and eastern South Dakota to Kansas; locally abundant in the prairie streams.

Notropis topeka Gilbert, Bull. Washburn Lab. Nat. Hist., I, 1884, 13, tributary of Kansas River, Topeka, Kans.

Notropis æneolus Hay, Proc. U. S. Nat. Mus., X, 1887, 245, Solomon River, Wakeeney, Kans.

Genus 354. HYBOGNATHUS Agassiz.

Hybognathus Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 223 (*H. nuchalis* Agassiz).

1079. *Hybognathus nuchalis* Agassiz. Gudgeon; Smelt; Silvery minnow; Smelt minnow.

Clear streams from the upper Missouri southward to Georgia and Texas, west of the Alleghenies.

Hybognathus nuchalis Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 224, Quincy, Ill.

Hybognathus evansi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 182, Fort Pierre, Nebr.

Hybognathus placitus Girard, loc. cit., 1856, 182, sluices of the Arkansas River.

1080. *Hybognathus regius* Girard. Gudgeon; Silvery minnow.

Streams New Jersey to Virginia, east of the Alleghenies.

Hybognathus regius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 209, Potomac River.

Hybognathus osmerinus Cope, Proc. Amer. Phil. Soc., XI, 1870, 466, Raritan River, N. J.

1081. *Hybognathus argyritis* Girard.
Upper Missouri region and Red River of the North
Hybognathus argyritis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 182, Milk River, Mont
1082. *Hybognathus hayi* Jordan.
Lower Mississippi Valley and neighboring rivers in Mississippi.
Hybognathus hayi Jordan, Proc. U. S. Nat. Mus., VII, 1884, 548, Pearl River, Jackson, Miss.; this and three preceding are all very close to *Hybognathus nuchalis*.
- Genus 355. *DIONDA* Girard.
Dionda Girard, Proc. Acad. Nat. Sci. Phila., VIII, 176 (*D. episcopa* Girard)
Algoma Girard, loc. cit., 180 (*A. amara* Girard).
1083. *Dionda episcopa* Girard.
Southern Texas and northern Mexico; south to the Rio Mezquital in Durango and the Rio San Juan in Nuevo Leon.
Dionda episcopa Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 177, headwaters of Pecos River and Comanche Spring, tributary to the Rio Grande, Tex.
Dionda tetensis Girard, loc. cit., 177, Nueces River, Tex.
Dionda papalis Girard, loc. cit., 178, Delaware Creek, a tributary of Pecos River, Tex.
Dionda argentosa Girard, loc. cit., 178, San Felipe Creek and Devils River, tributaries of Rio Grande.
Dionda chrysitis Girard, loc. cit., 178, Live Oak Creek, a tributary of the Pecos.
Dionda couchi Girard, loc. cit., 178, Guajuco, Monterey, and Cadereita, Nuevo Leon
Hybognathus flavipinnis Cope, Bull. U. S. Nat. Mus., XVII, 1880, 36, Johnsons Fork of Llano River, Tex.
Hybognathus nigrotæniata Cope, loc. cit., 37, Wallace Creek, a tributary of Medina River, Tex.
Hybognathus (Dionda) punctifer Garman, Bull. Mus. Comp. Zool., VIII, 1881, 89, Parras and spring near Saltillo, Mexico
Hybognathus civilis Cope, Trans. Amer. Phil. Soc., 1884, 167, Monterey, Nuevo Leon
1084. *Dionda serena* Girard.
Sabinal River, Tex.
Dionda serena Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 177, Sabinal River, Tex.
1085. *Dionda amara* (Girard)
Near Fort Brown, Tex.
Algoma amara Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 180, Lagoon near Fort Brown, Tex.
1086. *Dionda melanops* Girard.
Coahuila, Mexico.
Dionda melanops Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 178, Buena Vista, Coahuila.
1087. *Dionda nubila* (Forbes).
Western Illinois, west to Wyoming and southwest to the Ozark region.
Alburnops nubilus Forbes, Bull. Ill. Lab. Nat. Hist., I, 1878, 56, Rock River, Ogle County, Ill.
1088. *Dionda plumbea* Girard.
Tributaries of Arkansas River, in Oklahoma.
Dionda plumbea Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 178, Canadian River, Okla.
Dionda spadicea Girard, loc. cit., 178, Fort Smith, Ark.
Dionda grisea Girard, Pac. R. R. Surv., X, 1858, 230, Pl. LII, figs. 6-10, near Choctaw Agency.
- Genus 356. *CERATICHTHYS* Baird and Girard.
Ceraticthys Baird and Girard, Proc., Acad. Nat. Sci. Phila., VI, 1853, 391 (*C. vigilax*; misprinted *Ceraticthys*, not of Girard, 1856).
Ciola Girard, loc. cit., VIII, 1856, 192 (*Ceraticthys vigilax* Baird and Girard).
Hypargyrus Forbes, in Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 200 (*Hybopsis tuditanus* Cope).

1089. *Ceratichthys vigilax* Baird and Girard. *Bullhead minnow*; *Fathead minnow*.
Ohio to Georgia, Iowa, and Texas.
Ceratichthys vigilax Baird and Girard, Proc., Acad. Nat. Sci. Phila., VI, 1853, 391, Otter Creek, North fork of Red River, Ark.
Chola velox Girard, loc. cit., VIII, 1856, 192, San Pedro Creek and *Chola vivax*, Rio Leon, tributaries of Rio San Antonio, Tex.
Hybopsis tudilanus Cope, Trans., Amer. Phil. Soc., XIII, 1866, 381, Detroit, Mich.
Alburnops taurocephalus Hay, Proc. U. S. Nat. Mus., 1880, 503, Chickasawha River, Miss.
1090. *Ceratichthys smithi* (Evermann and Cox).
Southern South Dakota; known only from Prairie Creek near Scotland.
Chola smithi Evermann and Cox, Rept. U. S. Fish Com., XIX, 1894 (1896), 400, Prairie Creek, Scotland, S. Dak.
- Genus 357. **HYBORHYNCHUS** Agassiz.
Hyborhynchus Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 223 (*Minnulus notatus* Rafinesque)
1091. *Hyborhynchus notatus* (Rafinesque). *Blunt-nosed minnow*; *Spotted minnow*.
Quebec to Delaware; Kentucky, Alabama, Arkansas, and northward to the Dakotas; generally very abundant in small streams west of the Alleghenies.
Minnulus notatus Rafinesque, Ichth. Ohiensis, 1820, 47, Ohio River.
Hyborhynchus perspicuus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 179, Arkansas River, Fort Smith.
Hyborhynchus tenellus Girard, loc. cit., 179, Choctaw Agency
Hyborhynchus puniceus Girard, loc. cit., 179, Antelope Creek, Canadian River.
Hybopsis hæmaturus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 382, St. Joseph River, Michigan City, Ind.
Hyborhynchus superciliosus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 234, Kanawha River.
- Genus 358. **PIMEPHALES** Rafinesque. *Fatheads*.
Pimephales Rafinesque, Ichth. Ohiensis, 1820, 52 (*P. promelas* Rafinesque).
Coliscus Cope, Hayden's Geol. Surv. Wyo., 1872, 437 (*Coliscus parietalis* Cope; young of *Pimephales promelas*).
Spinicephalus LeSueur, Bull. Sci. Philom., VIII, 1896, 14, pl. 26 (*S. fibulatus* LeSueur = *Pimephales promelas* Rafinesque).
1092. *Pimephales promelas* Rafinesque. *Fathead minnow*; *Blackhead minnow*.
Lake Champlain to Kentucky, Dakota, and the Rio Grande; abundant in sluggish brooks, especially in the Missouri Basin.
Pimephales promelas Rafinesque, Ichth. Ohiensis, 1820, 53, pond near Lexington, Ky.
Pimephales maculosus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 180, sluices of the Arkansas River, Fort Makee, Ark.
Pimephales fasciatus Girard, loc. cit., 180, Yellowstone River.
Plargyrus melanocephalus Abbott, loc. cit., XII, 1860, 325, Lake Whittlesey, Minn.
Pimephales mulesi Cope, loc. cit., XVI, 1864, 282, Grosse Isle, Detroit River, Mich.
Pimephales agassizii Cope, Trans. Amer. Phil. Soc., XIII, 1866, 394, tributary of Ohio River, near Richmond, Ind.
Coliscus parietalis Cope, Hayden's Geol. Surv. Wyo., 1871, 437, St. Joseph, Mo.; young.
Hyborhynchus nigrellus Cope, Wheeler's Surv., V, Zool., 1875, 671, Arkansas River.
Spinicephalus fibulatus LeSueur, Bull. Soc. Philom., VIII, 1896, pl. 26; no locality.
1093. *Pimephales confertus* (Girard).
Tributaries of the Rio Grande.
Hyborhynchus confertus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 179, Hurrah Creek, a tributary of the Pecos River, Tex.

1094. *Pimephales anuli* Kendall.

Thoroughfare between Mud and Grass Lakes, Me.

Pimephales anuli Kendall, Bull. U. S. Fish Com., XXII, 1902, 257, between Mud Lake and Grass Lake, Me.

Genus 359. *COCHLOGNATHUS* Baird and Girard.

Cochlognathus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 158 (*C. ornatus* Baird and Girard).

1095. *Cochlognathus ornatus* Baird and Girard.

Rio Grande Basin.

Cochlognathus ornatus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 158, Brownsville, Tex.

1096. *Cochlognathus biguttatus* Cope.

Trinity River, Tex.

Cochlognathus biguttatus Cope, Bull. U. S. Nat. Mus., XVII, 1880, 37, Trinity River, Fort Worth, Tex.

Genus 360. *EXOGLOSSUM* Rafinesque.

Exoglossum Rafinesque, Jour. Acad. Nat. Sci. Phila., I, 1818, 419

(*E. lesueurianum* Rafinesque = *Cyprinus maxillingua* LeSueur).

Maxillingua Rafinesque, loc. cit., 421 (*Cyprinus maxillingua* LeSueur).

1097. *Exoglossum maxillingua* (LeSueur). *Cut-lips; Nigger chub; Nigger dick; Butter chub; Niggerfish.*

Lake Ontario, St. Lawrence River, Lake Champlain, Hudson River, and Cayuga Lake, and southward to Virginia; abundant in the basins of the Susquehanna, Hudson, Potomac, James, Roanoke, and Kanawha, but not widely distributed.

Cyprinus maxillingua LeSueur. Jour. Acad. Nat. Sci. Phila., I, 1817, 85, Pipe Creek, Md.

Exoglossum annulatum Rafinesque, loc. cit., 1818, 421, Hudson River.

Exoglossum nigrescens Rafinesque, loc. cit., 422, Hudson River.

Exoglossum lesueurianum Rafinesque, loc. cit., 420; after LeSueur.

Genus 361. *CAMPOSTOMA* Agassiz. *Stonerollers.*

Campostoma Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 218 (*Rutilus anomalus* Rafinesque).

1098. *Campostoma anomalum* (Rafinesque). *Stoneroller; Greased chub; Blue sucker; Rotgut minnow; Mammy; Doughbelly; Steel-backed chub.*

Central New York to Tennessee, Wyoming, and Texas.

Rutilus anomalus Rafinesque, Ichth. Ohienensis, 1820, 52, Licking River, Ky.

Exoglossum spinicephalum Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 363 (489), Wabash River.

Leuciscus prolixus Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 48, Alabama.

Exoglossum dubium Kirtland, Bost. Jour. Nat. Hist., V, 1845, 272, Yellow Creek, Ohio.

Chondrostoma pullum Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 357, Burlington, Iowa.

Campostoma nasutum Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 176, Cadereita, and near Monterey, Nuevo Leon, Mexico.

Campostoma callipteryx Cope, loc. cit., XVI, 1864, 284, Flint River, Mich.

Campostoma mormyrus Cope, loc. cit., 284, Bruce, Macomb County, Mich.

Campostoma gobionium Cope, loc. cit., 284, Bruce, Macomb County, and Grosse Isle, Mich.

Campostoma hippops Cope, loc. cit., 284, Platte River, Nebr.

Campostoma brevis Haseman, Proc. Ind. Acad. Sci., 1906 (1906), 161, Wabash River near Wabash, Ind.

1099. *Campostoma ornatum* Girard.

Chihuahua River, and tributaries of the Rio Grande in Mexico.

Campostoma ornatum Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 176, Chihuahua, Mexico.

1100. *Campostoma pricei* Jordan and Thoburn.
Rio Yaqui, Sonora and neighboring streams, tributary to the Gulf of California.
Campostoma pricei Jordan and Thoburn, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 203, Rucker Canyon, Chiricahua Mountains, Ariz., very near *C. ornatum*.
1101. *Campostoma formosulum* Girard.
Texas; San Antonio River to the Rio Grande.
Campostoma formosulum Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 176, Rio Sabinal, near San Antonio, Tex.

Family 91. MEDIDÆ

Genus 362. LEPIDOMEDA Cope.

- Lepidomeda* Cope, Proc. Amer. Phil. Soc. Phila., XIV, 1874, 131 (*L. vittata* Cope).
1102. *Lepidomeda vittata* Cope.
Known only from the Colorado Chiquito River, Ariz., and from Pahrn-agat Valley, Nev.
Lepidomeda vittata Cope, Proc. Amer. Phil. Soc. Phila., XIV, 1874, 131, Rio Colorado Chiquito, Ariz.
1103. *Lepidomeda jarrovii* Cope.
Colorado Chiquito River, Ariz.; also in southern Nevada, in springs in the desert.
Lepidomeda jarrovii Cope, Proc. Amer. Phil. Soc. Phila., XIV, 1874, 132, Rio Colorado Chiquito, Ariz.

Genus 363. MEDA Girard.

- Meda* Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 191 (*M. fulgida* Girard).
1104. *Meda fulgida* Girard.
Rio Gila Basin, Ariz.
Meda fulgida Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 192, Rio San Pedro, Ariz.
- Genus 364. FLAGOPTERUS Cope.
Flagopterus Cope, Proc. Amer. Phil. Soc., XIV, 1874, 130 (*P. argentsimus* Cope).
1105. *Flagopterus argentsimus* Cope.
Colorado Basin in western Colorado; Fort Yuma.
Flagopterus argentsimus Cope, Proc. Amer. Phil. Soc., XIV, 1874, 130, San Luis Valley, Colo.

Order Y. NEMATOGNATHI.²⁸

Family 92. ARIIDÆ. Sea catfishes

Genus 365. BAGRE Oken. *Gafftopsail catfishes*.

- Les Bagres Cuvier, Règne Anim., II, 1817 (*Silurus bagre* L.).
Bagre Oken, Isis, 1817, 1183 (after les Bagres of Cuvier; type by tautonymy *Silurus bagre* Linnæus).
Breviceps Swainson, Nat. Hist. Anim., I, 1838, 328 (*Breviceps filamentosus* Swainson); name preoccupied.
Felichthys Swainson, loc. cit., II, 1839, 305 (*Breviceps filamentosus* Swainson); substitute for *Breviceps*.
Ailurichthys Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 26 (*Silurus marinus* Mitchell).
Ælurichthys, corrected spelling.

²⁸ Until recently all members of this order have been usually included in the single family *Siluridæ*. As properly restricted the *Siluridæ* is a small group of large river fishes of Europe and Asia, resembling our larger catfishes, but without adipose fin, the anal joined to the caudal.

1106. *Bagre marina* (Mitchill). *Sea catfish; Gafftopsail*.
Cape Cod to Texas
Silurus marinus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 433,
New York.
In the investigation of Linnæan types at Upsala Dr. Einar Lönnberg
has found an example of this species labeled *Silurus felis*. On the
strength of this discovery this species has been called *Felichthys felis*.
But Linnæus's own description shows clearly that this was not his
type, which must have been our common sea cat here called *Galeichthys*
felis.
1107. *Bagre isthmensis* (Regan).
Atlantic coast of Panama; Colon.
Elurichthys isthmensis Regan, Biol. Centrali-Amer., 1907, 117, Tab.
XV, fig. 1, and Tab. XIX, fig. 1, Atlantic coast of Panama, Colon.
1108. *Bagre panamensis* (Gill).
Mazatlan to Panama.
Elurichthys panamensis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863,
172, Panama.
1109. *Bagre pinnimaculatus* (Steindachner).
Mazatlan to Panama.
Elurichthys pinnimaculatus Steindachner, Ichth. Beitr., IV, 1875, 15,
Panama, Altata, Costa Rica.
1110. *Bagre eydouxi* (Cuvier and Valenciennes).
Guayaquil.
Galeichthys eydouxi Cuvier and Valenciennes, Hist. Nat. Poiss., XV,
1840, 43, Guayaquil.
1111. *Bagre scutatus* (Regan).
Coast of Panama and Ecuador.
Elurichthys scutatus Regan, Biol. Centrali-Amer., 1907, 116, Tab.
XV, fig. 2, and Tab. XIX, fig. 2, Pacific coast of Panama.
1112. *Bagre bahiensis* (Castelnau).
Mexico to Bahia.
Galeichthys bahiensis Castelnau, Anim. Amér. Sud, 1855, 37, Bahia.
Elurichthys longispinus Gunther, Cat., V, 1864, 178; Mexico; South
America.
- Genus 366. *SCIADÉICHTHYS* Bleeker.
Sciadeichthys Bleeker, Act. Soc. Sci. Indo-Neerl., V, 1858, 99 (*Sciades*
pictus Müller and Troschel); not *Sciadeichthys* Bleeker, 1863.
1113. *Sciadeichthys troscheli* (Gill).
Gulf of California to Panama.
Sciades troscheli Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 171,
Panama.
Arius brandti Steindachner, Ichth. Beitr., IV, 1876, 21, Altata, Panama.
1114. *Sciadeichthys emphysetus* (Müller and Troschel).
Surinam.
Bagrus (Sciades) emphysetus Müller and Troschel, Horæ Ichth., III,
1849, 8, Surinam.
1115. *Sciadeichthys proops* (Cuvier and Valenciennes).
West Indies from Porto Rico to Surinam and south to Pernambuco.
Bagrus proops Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839,
457, Guiana; Porto Rico.
- Genus 367. *SELENASPIS* Bleeker.
Selenaspis Bleeker, Vissch. Ind. Archipel., I, 1858, 62 (*Silurus hertzbergi*
Bloch).
Leptarius Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 170 (*L. dowi*
Gill).
1116. *Selenaspis dowii* (Gill).
Panama to Guayaquil.
Leptarius dowi Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 170, Panama.
Arius alatus Steindachner, Ichth. Beitr., IV, 1876, 19, Panama.

Genus 368. NETUMA Bleeker.

Netuma Bleeker, Vissch. Ind. Archipel., I, 1858, 62 (*N. thalassina*, Bleeker=*N. nasuta* Bleeker).

Gurtinga Bleeker, loc. cit., 62 (*Arius commersoni* Bleeker).

Notarius Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 171 (*Arius grandicassis* Cuvier and Valenciennes).

1117. Netuma planiceps (Steindachner).

Gulf of California to Panama; not rare.

Arius planiceps Steindachner, Ichth. Beitr., IV, 1876, 26, Altata, Panama.

1118. Netuma kessleri (Steindachner).

Gulf of California to Panama; not rare.

Arius kessleri Steindachner, Ichth. Beitr., IV, 1876, 24, Altata, Panama.

1119. Netuma insculpta (Jordan and Gilbert).

Panama.

Arius insculptus Jordan and Gilbert, Bull. U. S. Fish Com., II, 1882 (1883), 41, Panama.

1120. Netuma platypogon (Günther).

Gulf of California to Peru; generally common.

Arius platypogon Günther, Cat., V, 1864, 147, San Jose de Guatemala.

1121. Netuma oscula (Jordan and Gilbert).

Panama, a single example known

Arius osculus Jordan and Gilbert, Bull. U. S. Fish Com., II, 1882 (1883), 46, Panama.

1122. Netuma elattura (Jordan and Gilbert).

Panama; scarce.

Arius elatturus Jordan and Gilbert, Bull. U. S. Fish Com., II, 1882 (1883), 45, Panama.

Genus 369. GALEICHTHYS Cuvier and Valenciennes. *Sea catfish.*

Galeichthys Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 28 (*G. feliceps* Cuvier and Valenciennes).

Hexanematichthys Bleeker, Act. Soc. Sci. Indo-Neerl., V, 1858, 2 (*Bagrus sundacus* Bleeker).

Cephalocassis Bleeker, Vissch. Ind. Archipel., I, 1858, 62 (*Arius truncatus* Cuvier and Valenciennes).

Arropsis Gill, Suppl. Proc. Acad. Nat. Sci. Phila., XIII, 1861, 56 (*Arius milberti*=*Silurus belis* L.).

Hemiaris Bleeker, Nederl. Tijdschr. Dierk., I, 1863, 7 (*Cephalocassis stormi* Bleeker).

Subgenus GALEICHTHYS²⁹ Cuvier and Valenciennes.**1123. Galeichthys lentiginosus** (Eigenmann and Eigenmann).

Panama.

Tachisurus lentiginosus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 139, Panama.

1124. Galeichthys peruvianus Lütken.

Pacific coast of Mexico to Peru.

Galeichthys peruvianus Lütken, Vid. Med. Naturh. Foren. Kjøbenhavn, 1874, 204, Callao, Peru.

Subgenus HEXANEMATICTHYS³⁰ Bleeker.**1125. Galeichthys felis** (Linnæus). *Sea cat.*

Cape Cod to Texas.

Silurus felis Linnæus, Syst. Nat., ed. XII, 1766, 503, Charleston, S. C.

Arius milberti Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 74, New York; Charleston, S. C.

?*Arius equestris* Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII,

1855, 26, Indianola, Tex.

1126. Galeichthys guentheri Regan.

Gulf of Mexico.

Galeichthys guentheri Regan, Biol. Centrali-Amer., 1907, 124, Tab. XVI, fig. 1, and Tab. XIX, fig. 3, Gulf of Mexico.

²⁹ Top of head covered with soft skin, hiding the rugose plates.

³⁰ Top of head with thin skin which does not hide the rugose plates.

1127. *Galeichthys surinamensis* (Bleeker).
Surinam.
Hexanematichthys surinamensis Bleeker, Versl. Akad. Amsterdam, 1862, 380, Surinam.
1128. *Galeichthys seemani* (Gunther).
Panama.
Arius seemani Gunther, Cat., V, 1864, 147, "Central America."
1129. *Galeichthys gilberti* Jordan and Williams.
Coast of Sinaloa.
Galeichthys gilberti Jordan and Williams, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 395, pl. 26, Mazatlan, Mexico.
1130. *Galeichthys jordani* (Eigenmann and Eigenmann).
Panama.
Tachisurus jordani Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 142, Panama.
1131. *Galeichthys caerulescens* (Gunther).
West coast of Guatemala.
Arius caerulescens Gunther, Cat., V, 1864, 149, Rio Huamuchal, Guatemala.
1132. *Galeichthys azureus* Jordan and Williams.
Mazatlan, Mexico.
Galeichthys azureus Jordan and Williams, Proc. Calif. Acad. Sci., 2d ser., IV, 1895, 398, pl. 27, Mazatlan, Mexico.
1133. *Galeichthys xenauchen* Gilbert.
Panama.
Galeichthys xenauchen Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2777, Panama.
1134. *Galeichthys guatemalensis* (Gunther).
Mazatlan to Central America.
Arius guatemalensis Gunther, Cat., V, 1864, 145, Guatemala; Chiapas.
1135. *Galeichthys assimilis* (Günther).
Atlantic coast of Central America, ascending streams.
Arius assimilis Gunther, Cat., V, 1864, 146, Lake Yzabel, Guatemala.
1136. *Galeichthys dasycephalus* (Günther).
Panama (wrongly ascribed to the Hawaiian Islands).
Arius dasycephalus Gunther, Cat., V, 1864, 157, Oahu; apparently an error.
1137. *Galeichthys longicephalus* (Eigenmann and Eigenmann).
Panama.
Tachisurus longicephalus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 143, Panama.
- Genus 370. *ARIUS* Cuvier and Valenciennes.
? *Tachisurus* Lacépède, Hist. Nat. Poiss., V, 1803, 151 (*T. chinensis* Lacépède; from a Chinese drawing, not identifiable as to species; may be a fresh-water fish or a *Netuma*).
Arius Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 52 (*A. arius* Cuvier and Valenciennes, type by autonymy).
Ariodes Müller and Troschel, Horae Ichth., III, 1849, 9 (*Arius arenarius* Müller and Troschel).
Pseudarius Bleeker, Nederl. Tijdschr. Dierk., I, 1862, 91 (*Arius arvus* Cuvier and Valenciennes).
1138. *Arius spixii* (Agassiz).
Trinidad to Brazil.
Pimelodus albidus Spix, in Agassiz, Gen. Spec. Pisc. Brasil., 1829, 19, equatorial Brazil; name preoccupied.
Pimelodus spixii Agassiz, loc. cit., 19; after Spix.
Arius arenatus Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 106, Cayenne.
Arius laticeps Gunther, Cat., V, 1864, 171, British Guiana, Trinidad.
1139. *Arius melanopus* Gunther.
Pacific coast of Central America.
Arius melanopus Günther, Cat., V, 1864, 172, Rio Motagua.

1140. *Arius furthi* Steindachner.
Panama.
Arius furthi Steindachner, Ichth. Beitr., IV, 1876, 29, Panama.
1141. *Arius steindachneri* (Gilbert and Starks).
Pacific coast of Panama.
Tachysurus steindachneri Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 29, Pl. V, fig. 9, Panama Bay.
1142. *Arius evermanni* (Gilbert and Starks).
Panama.
Tachysurus evermanni Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 32, Pl. V, fig. 10, Panama.
1143. *Arius liropus* (Bristol).
Coast of Sonora, Mexico.
Tachysurus liropus Susan Brown Bristol, in Gilbert, Proc. U. S. Nat. Mus., XIX, 1896 (Feb. 5, 1897), 438, San Juan Lagoon, near mouth of Rio Ahome, Sonora, Mexico.
1144. *Arius emmelane* (Gilbert).
Panama.
Tachysurus emmelane Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2785, Panama.
1145. *Arius taylori* Hildebrand. *Bagre*.
Coast of El Salvador
Arius taylori Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925, 250, fig. 10, Rio Lempa, San Marcos.
1146. *Arius multiradiatus* Gunther.
Streams of Isthmus of Panama, Rio Bayano, Panama.
Arius multiradiatus Gunther, Cat., V, 1864, 173, Rio Bayano, Panama.
- Genus 371. CATHOROPS Jordan and Gilbert.
Cathorops Jordan and Gilbert, Bull. U. S. Fish Com., II, 1882, 54
(*Arius hypophthalmus* Steindachner).
1147. *Cathorops hypophthalma* (Steindachner).
Panama; rather scarce.
Arius hypophthalmus Steindachner, Ichth. Beitr., IV, 1875, 31, Pl. X, Panama.
1148. *Cathorops gulosa* (Eigenmann and Eigenmann).
Panama.
Tachysurus gulosus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d. ser., I, 1888, 146, Panama.

Family 93. AMEIURIDÆ. Horned pouts; Channel cats

Genus 372. ISTLARIUS Jordan and Snyder.
Istlarius Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 118
(*I. balsanus* Jordan and Snyder).

1149. *Istlarius balsanus* Jordan and Snyder.
Balsas River Basin, Pacific slope of Mexico.
Istlarius balsanus Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 118; tributary of Rio de las Balsas at Puente de Ixtla, Guerrero, Mexico (Pacific drainage).

Genus 373. ICTALURUS Rafinesque. Channel cats.
Ictalurus Rafinesque, Ichth. Ohiensis, 1820, 61 (*Silurus punctatus* Rafinesque).
Ellips Rafinesque, loc. cit., 62 (*Pimelodus maculatus* Rafinesque).
Synechoglanis Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 39 (*S. beadlei* Gill = *Ictalurus furcatus* Rafinesque).
Ichthyælus, corrected spelling.

1150. *Ictalurus furcatus* (Cuvier and Valenciennes). *Blue cat; Great fork-tailed cat; Mississippi cat, Poisson bleu.*
Pimelodus furcatus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1840, 136, New Orleans.
Pimelodus affinis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 26, Rio Grande.
Ameiurus ponderosus Bean, Proc. U. S. Nat. Mus., II, 1879, 286, St. Louis.
Ictalurus robustus Jordan, Bull. U. S. Nat. Mus., X, 1877, 76, Cairo, Ill.
1151. *Ictalurus anguilla* Evermann and Kendall. *Eel cat, Willow cat; Fulton cat*
 Mississippi River, occasionally as far north as Louisville, Ky., Muscatine and Fairport, Iowa, where it is known as the "Fulton cat."
Ictalurus anguilla Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897 (Feb. 9, 1898), 125, pl. 6, fig. 1, Atchafalaya River, La.
1152. *Ictalurus punctatus* (Rafinesque). *Channel cat; White cat; Fiddler.*
 Rivers of the Great Lakes region, the Mississippi Valley, and streams tributary to the Gulf of Mexico.
Silurus punctatus Rafinesque, Amer. Month. Mag., 1818, 359, Ohio River.
Pimelodus caudafurcatus LeSueur, Mem. Mus., V, 1819, 152, Wabash River; Mississippi River.
Pimelodus maculatus, pallidus, argentinus, argyrus, with varieties *erythroptera, marginatus, lateralis, leucoptera* Rafinesque, Quarterly Jour. Sci., London, 1820, and Ichth. Ohiensis, 1820, 62, etc., Ohio River.
Pimelodus furcifer Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 139, Surinam, by error.
Pimelodus gracilis Hough, Ann. Rept. Regent's, Albany, 1852, 26, Somerville, N. Y.
Pimelodus vulpes Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 170, rivers of Texas.
Synechoglanis beadler Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1859, 40, St. Catherine's, Ont.
Pimelodus houghi Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 159, Somerville; Ogdensburg; Foxburg, Pa.
Pimelodus megalops Girard, loc. cit., 161, Black Warrior River, Bigsby River, Ala.
Pimelodus graciosus Girard, loc. cit., 161, Prairie Mer Rouge, La.
Pimelodus hammondi and *notatus* Abbott, loc. cit., XII, 1860, 568, 569, Fort Riley, Kans.
Ictalurus simpsoni Gill, Proc. Bost. Soc. Nat. Hist., VIII, 1862, 43, Platte River.
1153. *Ictalurus meridionalis* (Gunther).
 Rio Usumacinta, Guatemala.
Amiurus meridionalis Gunther, Cat., V, 1864, 102, and in Trans Zool. Soc. Lond., 1868, 473, Rio Usumacinta, Guatemala.
- Genus 374. **HAUSTOR** Jordan and Evermann.
Haustor Jordan and Evermann, Fishes North and Mid. Amer., 1896, 137 (*Gadus lacustris* Walbaum = *Pimelodus nigricans* LeSueur).
Vallarius Rutter, Proc. Calif. Acad. Sci., VI, Ser. 2, June 23, 1896, 258 (*V. pricei* Rutter).
1154. *Haustor lacustris* (Walbaum).
 Saskatchewan River and the Great Lakes.
Gadus lacustris Walbaum, Artdi Pisc., 1792, 144, Arctic America; based on the mathemeg or land cod of Pennant.
Pimelodus nigricans LeSueur, Mem. Mus., V, 1819, 153, Lake Erie, Lake Ontario.
Pimelodus borealis Richardson, Fauna Bor.-Amer., III, 1836, 135, Pine Island Lake.
1155. *Haustor okeechobeensis* (Heilprin).
 Florida.
Ictalurus okeechobeensis Heilprin, Trans. Wagner Inst. Sci. Phila., I, 1887, pl. 18, Kissimee River, Fla.
Ameiurus lacustris okeechobeensis Fowler, Proc. Acad. Nat. Sci. Phila., LI, 1899, 480; same type.

1156. **Haustor catus** (Linnaeus). *White cat; Channel cat of the Potomac.*
Silurus catus Linnaeus, Syst. Nat., ed. X, 1758, 305, based on the catfish of Catesby (*Bagre secundæ speciei maragraver affinis*), which is a rude figure of the adult or "lophius" form of this species; northern part of America.
Pimelodus albidus LeSueur, Mem. Mus., V, 1819, 148, Delaware River.
Pimelodus lynx Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 160, Potomac River.
Amiurus nweiventris Cope, Proc. Amer. Phil. Soc., XI, 1870, 486, Neuse River.
Ictalurus macaskeyi and *I. keviniskii* Stauffer, Hist. Lancaster County, Pa., 1869, 578, Lancaster County, Pa.
1157. **Haustor lophius** (Cope). *Big-mouthed cat.*
 Potomac River.
Amiurus lophius Cope, Proc. Amer. Phil. Soc., XI, 1870, 486, Potomac River; not observed elsewhere. Known by its very wide flat head; apparently not the adult of *Ameiurus catus*, as has been supposed.
1158. **Haustor lupus** (Girard).
 Rio Neuces and Rio Pecos, Tex.
Pimelodus lupus Girard, Pac. R. R. Surv., X, 1858, Pt. 4, Fishes, 211, Rio Pecos, Tex.
1159. **Haustor pricei** (Rutter).
 Southern Arizona.
Villarius pricei Rutter, Proc., Calif. Acad. Sci., 2d ser., VI, 1896, 257, San Bernardino Creek, a tributary of the Yaqui River, southern Arizona.
1160. **Haustor dugesi** (Bean).
 Rio Turbio, Guanajuato, Mexico, west of the Sierra Madre
Ameiurus dugesi Bean, Proc. U. S. Nat. Mus., II, 1879, 304, Rio Turbio, Guanajuato, Mexico.
- Genus 375. **AMEIURUS** Rafinesque. *Horned pout.*
Ameiurus Rafinesque, Ichth. Ohiensis, 1820, 65 (*Silurus cupreus* Rafinesque = *Pimelodus natalis* LeSueur).
Amiurus, corrected spelling
1161. **Ameiurus erebennus** Jordan.
 Coast swamps and black water streams from New Jersey to Florida.
Ameiurus erebennus Jordan, Bull. U. S. Nat. Mus., X, 1877, 85, St. John's River, Fla.
1162. **Ameiurus natalis** (LeSueur). *Yellow cat.*
 Great Lakes region to Virginia and southward
Pimelodus natalis LeSueur, Mem. Mus., 1819, 154, North America
Silurus lvidus and var. *fuscus* Rafinesque, Quarterly Jour. Sci. Lond., 1820, 48, Ohio River.
Silurus cupreus Rafinesque, loc. cit., 1820, 51, Ohio River.
Pimelodus cupreus Rafinesque, Ichth. Ohiensis, 1820, 65, Ohio River.
Pimelodus cænosus Richardson, Fauna Bor.-Amer., III, 1836, 132, Lake Huron.
Pimelodus felinus et *antoniensis* Girard, Pac. R. R. Surv., X, 1858, 209, Canadian and San Antonio Rivers, Tex.
Pimelodus cupresoides Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 159, Aux Plaines, Ill.
Ameiurus natalis analis Jordan, Bull. U. S. Nat. Mus., X, 1877, 81, Arkansas River.
Ameiurus bolli Cope, Bull. U. S. Nat. Mus., XX, 1880, 35, Little Wichita River, Tex.
1163. **Ameiurus vulgaris** (Thompson).
 Vermont to Minnesota and Illinois, chiefly northward. A doubtful species.
Pimelodus vulgaris Thompson, Hist. Vermont, 1842, 138, Lake Champlain.
P[imelodus] dekeyi Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 160, Oswego, N. Y.
Pimelodus ailurus Girard, Pac. R. R. Surv., X, 1858, 210, Lake Amelia, Minn.

1164. *Ameiurus nebulosus* (LeSueur). *Horned pout; Common bullhead; Small catfish; Schuylkill cat; Sacramento cat.*
Great Lakes, Ohio Valley, eastward to Maine, southwestward to Texas and southwestward to Florida; introduced about 1877 into the Humboldt, Sacramento, San Joaquin, and Gila Rivers, where it is now excessively abundant
Pimelodus nebulosus LeSueur, Mem. Mus., V, 1819, 149, Lake Ontario.
Pimelodus atrarius DeKay, N. Y. Fauna: Fishes, 1842, 185, Wappingers Creek, Dutchess County, N. Y.
Pimelodus felis Agassiz, Lake Superior, 1850, 281, Lake Superior.
Pimelodus vulpeculus Girard, Proc., Acad. Nat. Sci. Phila., XI, 1859, 160, Charleston, S. C.
Pimelodus hoyi Girard, loc. cit., 159, Racine, Wis.
Ameiurus missillensis Cope, Proc., Amer. Phil. Soc., 1870, 486, Missillon Creek, Del.
1165. *Ameiurus nebulosus catulus* (Girard).
Streams of Texas; abundant.
Pimelodus catulus Girard, Pac. R. R. Surv., X, 1858, Pt. 4, Fishes, 208, Fort Smith, Ark.
1166. *Ameiurus prosthistius* Cope.
New Jersey.
Ameiurus prosthistius Cope, Proc. Acad. Nat. Sci. Phila., XXXV, 1883, 133, Batsto River, N. J.
1167. *Ameiurus platycephalus* (Girard). *Mud cat; Brown cat.*
Streams of the Carolinas and eastern Georgia, Cape Fear to the Chatahoochee.
Pimelodus platycephalus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 161, Anderson, S. C.
Ameiurus brunneus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1877, 366, Ocmulgee River, Flat Shoals, Ga.; the young of *platycephalus*.
1168. *Ameiurus brachyacanthus* Cope.
Streams of Texas.
Ameiurus brachyacanthus Cope, Bull. U. S. Nat. Mus., XVII, 1880, 35, Wallace Creek, Bandera County, Tex.
1169. *Ameiurus melas* (Rafinesque). *Black bullhead; Small catfish.*
Northern New York (Genesee River) to Kansas and Texas; common especially west of the Mississippi.
Silurus melas Rafinesque, Quarterly Jour. Sci. Lond., 1820, 51, Ohio River.
Silurus xanthocephalus Rafinesque, loc. cit., 1820, 51, Ohio River.
Pimelodus pullus DeKay, N. Y. Fauna: Fishes, 1842, 184, Lake Pleasant, Janet, N. Y.
Pimelodus confinis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 159, Racine, Wis.
Ameiurus obesus Gill, Proc. Bost. Soc. Nat. Hist., 1862, 45, Nebraska.
Ameiurus cragini Gilbert, Bull., Washburn Lab. Nat. Hist., I, 1884, 10, Garden City, Kans.
1170. *Ameiurus marmoratus* (Holbrook).
Lowland streams and swamps from southwestern Indiana (Switz City) southward and eastward to Florida.
Pimelodus marmoratus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 54, South Carolina.
- Genus 376. *GRONIAS* Cope. *Blind catfishes.*
Gronias Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 231 (*G. nigrilabris* Cope).
1171. *Gronias nigrilabris* Cope.
Cave streams tributary to Conestoga River in eastern Pennsylvania
Gronias nigrilabris Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 231, Conestoga River, Pa.

Genus 377. OPLADELUS Rafinesque.

?*Plodictis* Rafinesque, Jour. Phys., 1819, 422 (*P. limosa* Rafinesque, after Audubon; drawing of an imaginary catfish).

Leptops Rafinesque, Ichth. Ohiensis, 1820, 64 (*Pimelodus viscosus* Rafinesque = *Silurus olivaris* Rafinesque).

Opladelus Rafinesque, loc. cit., 64 (*Pimelodus nebulosus* Rafinesque = *Silurus olivaris* Rafinesque); name chosen by first reviser.

Iuctis Rafinesque, loc. cit., 66 (*Pimelodus nebulosus* Rafinesque = *Silurus olivaris* Rafinesque).

Hopladelus and *Pelodichthys*, corrected spelling.

1172. *Opladelus olivaris* (Rafinesque). *Mud cat*; *Yellow cat*; *Bashaw*; *Russian cat*; "Hoosier" at New Boston, Ill.; *Goujon*.

Silurus olivaris Rafinesque, Amer. Month. Mag., 1818, 355. Ohio River. *Silurus nebulosus*, *viscosus* and *limosus* Rafinesque, Quarterly Jour. Sci. Lond., 1820, 50, Ohio River.

Pyloodictis limosus Rafinesque, Ichth. Ohiensis, 1820, 67, Ohio and Mississippi Rivers; based on a drawing by Audubon

Pimelodus punctulatus Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 134, New Harmony, Ind.

Pimelodus zeneus Cuvier and Valenciennes, loc. cit., 135, New Orleans.

Genus 378. NOTURUS Rafinesque. Stone cats.

Noturus Rafinesque, Amer. Month. Mag., November, 1818, 41 (*N. flavus* Rafinesque).

1173. *Noturus flavus* Rafinesque. *Stone cat*.

Great Lakes region, westward to Montana and Wyoming and south to Texas.

Noturus flavus Rafinesque, Amer. Month. Mag., 1818, 41, falls of Ohio River.

Noturus luteus Rafinesque, Jour. Phys., 1819, 421, Ohio River.

Noturus occidentalis Gill, Proc. Bost. Soc. Nat. Hist., VIII, 1862, 45, Platte River.

Noturus platycephalus Gunther, Cat., V, 1864, 104, North America.

Genus 379. SCHILBEODES Bleeker. Tadpole cats.

Schilbeodes Bleeker, Ichth. Archip. Indici. Prodr. Siluri, 1858, 258 (*Silurus gyrinus* Mitchill).

1174. *Schilbeodes gyrinus* (Mitchill).

Hudson River and westward through almost the entire Mississippi Valley, and Great Lakes region.

Silurus gyrinus Mitchill, Amer. Month. Mag., 1818, 322, Walkill River, N. Y.

Noturus sialis Jordan, Bull. U. S. Nat. Mus., X, 1877, 102, White River, Ind.

Genus 380. TROGLOGLANIS Eigenmann.

Trogloglanis Eigenmann, Proc. Amer. Phil. Soc., LVIII, 1919, 397 (*T. pattersoni* Eigenmann).

1175. *Trogloglanis pattersoni* Eigenmann.

Artesian well in San Antonio, Tex.

Trogloglanis pattersoni Eigenmann, Proc. Amer. Phil. Soc., LVIII, 1919, 397, figs. 1 and 2.

Genus 381. RABIDA Jordan and Evermann. Mad toms.

Rabida Jordan and Evermann, Fishes North and Mid Amer., 1896, 146 (*Noturus furiosus* Jordan and Meek).

Pimelodon (LeSueur) Vaillant, Bull. Soc. Philom., 1896, 28 (*Pimelodon insignarius* LeSueur, unpublished plate of 1819; probably *Pimelodus insigne* Richardson, the *lemniscatus* = *Pimelode insigne* of LeSueur).

Subgenus PIMELODON.

1176. *Rabida leptacantha* (Jordan).

Gulf States, in sandy streams; basins of Ogeechee, Alabama, and Pearl Rivers.

Noturus leptacanthus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 352, Silver Creek, branch of Etowah River, Rome, Ga.

1177. *Rabida nocturna* (Jordan and Gilbert).
Sandy streams, among weeds, in lower Mississippi Valley; known from the lower Wabash in Indiana and from the Poteau, Washita, and Saline Rivers, Ark.
Noturus nocturnus Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1886, 6, Saline River at Benton, Ark.
1178. *Rabida funebris* (Jordan and Swain).
Tributary of Black Warrior River, Tuscaloosa, Ala.
Noturus funebris Jordan and Swain, Bull. U. S. Fish Com., IX, 1889, 153, North River, Tuscaloosa, Ala.
1179. *Rabida exilis* (Nelson).
Wisconsin to Missouri and Kansas; common in prairie streams.
Noturus exilis Nelson, Bull. Ill. Lab. Nat. Hist., 1876, 51, Illinois River.
Noturus classochar Swain and Kalb, Proc. U. S. Nat. Mus., V, 1882, 639, Naperville, Ill.
1180. *Rabida insignis* (Richardson). *Mad tom*.
Pennsylvania to South Carolina; very common in streams east of the Alleghenies.
Pimelodus insignis Richardson, Fauna Bor.-Amer., III, 1836, 32, based on *Pimelode livree* LeSueur, Mem. Mus., V, 1819, 155, locality not known.
Noturus marginatus Baird, in Cope, Jour. Acad. Nat. Sci. Phila., VI, 1869, 237, Pa.
Pimelodon insignarius LeSueur, in Vaillant, Bull. Soc. Philom., 1896, 28.
1181. *Rabida gilberti* (Jordan and Evermann).
Roanoke River in southern Virginia, locally abundant.
Noturus gilberti Jordan and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 352, Roanoke River, Roanoke, Va.
1182. *Rabida miura* (Jordan).
Mississippi Valley and tributaries of Lake Michigan, south to Louisiana; common in the Ohio Valley.
Noturus miurus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1877, 371, White River, Ind.
1183. *Rabida eleuthera* (Jordan).
Mississippi Valley, rare, but widely distributed; tributaries of French Broad River (Big Pigeon River, Clifton, Tenn., French Broad at Hot Springs, N. C.); White River, Gosport, Ind.; Green River, Greensburg, Ky.; Schriner Lake, Columbia City, Ind.; White River, Fayetteville, Ark.; Sallisaw River, Indian Territory [Okla.]. Only a few specimens known.
Noturus eleutherus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1877, 371, Big Pigeon River, Clifton, Tenn.
1184. *Rabida furiosa* (Jordan and Meek).
Eastern North Carolina; known from the Neuse, Tar, and Little Rivers.
Noturus furiosus Jordan and Meek, Proc. U. S. Nat. Mus., XI, 1888, 351, Neuse River, N. C.

Family 94. PIMELODIDÆ. South American catfishes; Bagres de Rio

Genus 382. CONORHYNCHOS Bleeker.

Conorhynchus Bleeker, Vissch. Ind. Archipel. Siluri, 1858, 191 (*Pimelodus conirostris* Cuvier and Valenciennes), not *Conorhynchus* Nozeman, a pre-Linnæan name (1757) revived by Gill (1861).
Conorhynchichthys Regan, *Conorhynchus* being regarded as preoccupied.

1185. *Conorhynchus nelsoni* Evermann and Goldsborough.
Rio Usumacinta, Yucatan.

Conorhynchus nelsoni Evermann and Goldsborough, Bull. U. S. Fish Com., XXI, 1901 (1902), 140, fig. 1, Rio Usumacinta at Monte Cristo, Yucatan.

Genus 383. RHAMDIA Bleeker. *Bagrés de Rio*.

Pteronotus Swainson, Nat. Hist. Fishes, Amphib., etc., II, 1839, 309 (*Heterobranchius septentaculatus* Agassiz, preoccupied in mollusks).

Rhamdia Bleeker, Vissch. Ind. Archipel.: Siluri, 1858, 197 (*Pimelodus quelon* Quoy and Gaimard).

Rhamdella Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 172 (*R. eriarcha* Eigenmann and Eigenmann).

1186. *Rhamdia wagneri* (Gunther). *Barbudo*.
Rivers of Central America, on both sides of the mountains; very abundant.
Pimelodus wagneri Günther, Fish. Cent. Amer., 1869, 474, Atlantic and Pacific rivers of Panama.
Rhamdia bransfordi Gill, Proc. Acad. Nat. Sci. Phila., XXVIII, 1876, 337, Panama.
1187. *Rhamdia oaxacæ* Meek. *Bagre*.
Streams of Mexico on the Atlantic side south of Vera Cruz.
Rhamdia oaxacæ Meek, Field Col. Mus., Zool., III, 74 (figured in plate V, loc. cit., V, 1905) Rio Quiotepec, Cuicatlan, Mexico.
1188. *Rhamdia baronis-mülleri* (Troschel).
"Pacific Ocean" in Mexico.
Pimelodus baronis-mülleri Troschel, in Müller's Beitr. Gesch. Statist. und Zool. Mexico, III, 1865, 636, no definite locality.
1189. *Rhamdia motaguensis* (Gunther).
Rio Motagua, Guatemala.
Pimelodus motaguensis Günther, Cat., V, 1864, 127, Rio Motagua, Guatemala.
1190. *Rhamdia guatemalensis* (Günther).
Huamuchal, Guatemala.
Pimelodus guatemalensis Günther, Cat., V, 1864, 122, Rio Huamuchal, Guatemala.
Pimelodus brachypterus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 404, Orizaba, Mexico.
1191. *Rhamdia salvini* (Gunther).
Rio San Geronimo, Guatemala.
Pimelodus salvini Günther, Cat., V, 1864, 130, Rio San Geronimo, Guatemala.
1192. *Rhamdia hypselura* (Günther).
Mexico.
Pimelodus hypselurus Günther, Cat., V, 1864, 126, Mexico.
1193. *Rhamdia brachycephala* (Regan).
Mexico.
Pimelodus brachycephalus Regan, Ann. Mag. Nat. Hist., XIX, 1907, 258, Guatemala, Rio Nacasil.
1194. *Rhamdia laticauda* (Heckel).
Mexico.
Pimelodus laticaudus Heckel, in Kner, Sitzungsber. Kais. Akad. Wiss. Wien, XXVI, 1857, 420, Mexico.
1195. *Rhamdia godmani* (Gunther).
Valley of Rio Usumacinta, Guatemala.
Pimelodus godmani Günther, Cat., V, 1864, 124, Rio Usumacinta, Guatemala.
1196. *Rhamdia nicaraguensis* (Günther).
Lake Nicaragua.
Pimelodus nicaraguensis Günther, Cat., V, 1864, 125, Lake Nicaragua, Nicaragua.
1197. *Rhamdia microptera* (Gunther).
Rio San Geronimo, Guatemala.
Pimelodus micropterus Günther, Cat., V, 1864, 124, Rio San Geronimo, Guatemala.
1198. *Rhamdia managuensis* (Gunther).
Lake Managua, Nicaragua.
Pimelodus managuensis Günther, Fish. Cent. Amer., 1869, 474, Lake Managua, Nicaragua.
1199. *Rhamdia cabreræ* Meek.
Lake Amatitlan, Guatemala.
Rhamdia cabreræ Meek, Field Mus. Nat. Hist., Zool., VII, 1906, 93, Lake Amatitlan, Nicaragua.

1200. *Rhamdia barbata* Meek. *Chuchin*.
Lakes of Nicaragua.
Rhamdia barbata Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 106,
San Francisco, Nicaragua.
1201. *Rhamdia sacrificii* Barbour and Cole
Chichen-Itza, Yucatan.
Rhamdia sacrificii Barbour and Cole, Bull. Mus. Comp. Zool., L, 1906,
No. 5, 156, Sacrificial Cenote, near Chichen-Itza, Yucatan.
1202. *Rhamdia depressa* Barbour and Cole.
Near Chichen-Itza, Yucatan.
Rhamdia depressa Barbour and Cole, Bull. Mus. Comp. Zool., L, 1906,
No. 5, 155, pl. 1, Ikil Cenote, near Chichen-Itza, Yucatan.
1203. *Rhamdia heteracantha* Regan.
Costa Rica.
Rhamdia heteracantha Regan, Biol. Centrali-Amcr., Pisces, 1907, 134,
Juan Viñas, Costa Rica.
1204. *Rhamdia polycaula* (Gunther).
Rio San Geronimo.
Pimelodus polycaulus Gunther, Cat., V, 1864, 131, Rio San Geronimo.
1205. *Rhamdia petenensis* (Gunther).
Lake Peten, Yucatan.
Pimelodus petenensis Gunther, Cat., V, 1864, 126, Lake Peten, Yucatan.
1206. *Rhamdia parryi* Eigenmann.
Rio Zanalenco, Chiapas.
Rhamdia parryi Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888,
130, Rio Zanalenco, Chiapas.
1207. *Rhamdia underwoodi* Regan.
Costa Rica.
Rhamdia underwoodi Regan, Biol. Centrali-Amcr., Pisces, 1907, 135,
pl. 23, fig. 4, Juan Viñas, Costa Rica.
1208. *Rhamdia rogersi* (Regan).
Pimelodus rogersi Regan, Ann. Mag. Nat. Hist., XIX, 1907, 259,
Irazu, Costa Rica.
Rhamdia regani Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 144,
Rio Tiribi, San José.
- Genus 384. *CÆCORHAMIDIA* Norman.
Cæcorhamdia J. R. Norman, Ann. Mag. Nat. Hist., ser. 9, XVIII, 1926,
324 (*C. urichi* Norman).
1209. *Cæcorhamdia urichi* Norman.
Guacharo Cave, Trinidad.
Cæcorhamdia urichi J. R. Norman, Ann. Mag. Nat. Hist., ser. 9,
XVIII, 1926, 324, Guacharo Cave, Trinidad.
- Genus 385. *PIMELODELLA* Eigenmann and Eigenmann.
Pimelodella Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d
ser., I, 1888, 131 (*Pimelodus cristatus* Muller and Troschel).
1210. *Pimelodella chagresi* (Steindachner).
Rio Chagres.
Pimelodus chagresi Steindachner, Ichth. Beitr., IV, 1876, 34, Rio
Chagres at Obispo, Canal Zone.
- Genus 386. *PIMELODUS* Lacépède.
Pimelodus Lacépède, Hist. Nat. Poiss., V, 1803, 94 (*P. maculatus*
Lacépède, as restricted by authors=*Silurus clarias* Bloch).
Pseudariodes Bleeker, Nederl. Tijdschr. Dierk., 1863, 99 (*Silurus clarias*
Bloch).
Pseudorhamdia Bleeker, Versl. Akad. Amsterdam, XIV, 1862, 384
(*Pimelodus ascita* Bleeker).

1211. *Pimelodus clarias* (Bloch).

Rio Mamoni (Panama) and southeastward to Rio de la Plata.

Silurus clarias Linnaeus, Syst. Nat., ed. X, 1758, 306, in part, not the type.

Pimelodus maculatus Lacépède, Hist. Nat. Poiss., V, 1803, 94, 107, Rio de la Plata.

Piramutana blochii Gunther, Cat., V, 1864, III, Cayenne; Surinam.

Pimelodus rigidus Spix, in Agassiz, Pisc. Brasil, 1829, 19, Brazil.

Pimelodus macronema Bleeker, Silures de Suriname, 1864, 79, pl. 14, Surinam.

Pseudaroides pantherinus Lutken, Meddel. Nat. Hist. Foren., 1874, Nos. 1, 2, p. 194, Venezuela.

Pseudorhamdia piscatrix Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 569, Pebas.

Piramutana macrospila Gunther, Ann. and Mag. Nat. Hist., 5th ser., VI, July, 1880, 10, Pl. II, Pirana, Rio de la Plata.

Megalonema punctatum Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 77, Rio Tuyra, Marrigante, Panama.

Megalonema robustum Meek and Hildebrand, loc. cit., 78, Rio Tuyra, Marrigante, Panama.

Genus 387. TRACHCORYSTES Bleeker.

Trachcorystes Bleeker, Vissch. Ind. Archipel. Siluri, 1858, 20 (*T. typus* Bleeker).

Parauchenipterus Bleeker, Nederl. Tijdschr. Dierk., I, 1863, 7 (*Silurus galeatus* L.).

1212. *Trachcorystes amblops* (Meek and Hildebrand).

Rio Tuyra Basin, Panama.

Felchithys amblops Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 77, Rio Tuyra, Marrigante, Panama.

Genus 388. AGENEIOSUS Lacépède

Ageneiosus Lacépède, Hist. Nat. Poiss., V, 1803, 132 (*A. armatus* Lacépède = *Silurus militaris* Linnaeus).

Pseudogeneiosus Bleeker, Nederl. Tijdschr. Dierk., I, 1863, 14 (*Ageneiosus brevifilis* Cuvier and Valenciennes).

1213. *Ageneiosus caucanus* Steindachner.

Lower Rio Tuyra, Marrigante, Panama.

Ageneiosus caucanus Steindachner, Denkschr. Kais. Akad. Wiss. Wien, XLI, 1880, 61, Pl. VI, figs. 1, 2, Rio Cauca, Panama.

Family 95. PYGIDIIDÆ**Genus 389. PYGIDIUM Meyen.**

Pygidium Meyen, Reise in Peru, I, 1835, 475 (*P. fuscum* Meyen).

1214. *Pygidium striatum* Meek and Hildebrand.

Upper Tuyra Basin, Panama.

Pygidium striatum Meek and Hildebrand, Field. Mus. Nat. Hist., Zool., X, 1913, 78, Rio Cana, Cana, Panama.

Family 96. ASTROBLEPIDÆ**Genus 390. ASTROBLEPUS³¹ Humboldt. Cave catfishes.**

Astroblepus Humboldt, Obs. Zool. Phil. Mag., XXIV, 1806, 19, 331 (*A. graxalva* Humboldt).

Cyclopium Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 305 (*Pimelodus cyclopium* Humboldt = *Cyclopium humboldti* Swainson).

Arges Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 333 (*A. sabalo* Cuvier and Valenciennes).

Brontes Cuvier and Valenciennes, loc. cit., 341 (*B. prenadilla* Cuvier and Valenciennes, preoccupied).

Strephon Gistel, Thierreich., VIII, 1848 (*Brontes prenadilla* Cuvier and Valenciennes, preoccupied).

Stygogenes Gunther, Cat., V, 1864, 223 (*S. humboldti* Gunther).

³¹ According to Eigenmann, all these names belong to the same genus

1215. *Astroblepus pirrensis* (Meek and Hildebrand).

Tuyra River Basin.

Cyclopterus pirrensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 83, Rio Cana, Cana, Panama (Pl. XVI, Field Mus. Nat. Hist., Zool., X).

Family 97. CALLICHTHYIDÆ

Genus 391. HOPILOSTERNUM Gill.

Hoplosternum Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 395 (*Callichthys lævigatus* Cuvier and Valenciennes = *Callichthys littoratus* Hancock).1216. *Hoplosternum punctatus* Meek and Hildebrand.

Rio Marte Arnade, Panama.

Callichthys (*Hoplosternum*) *thoracatus* (not Cuvier and Valenciennes), Steindachner, Sitzungsber., Kais. Akad. Wiss. Wien, XLI, 1880, 66, Cauca, near Caceres.*Hoplosternum punctatum* Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 264, Rio Marte Arnade, 6 miles east of Panama City.

Family 98. LORICARIIDÆ. Mailed catfishes

Genus 392. HYPOSTOMUS (Lacépède).

Plecostomus Gronow, Zoophyl., 1763, 127 (*Plecostomus dorso dipterygion* Gronow); not binomial.*Hypostomus* Lacépède, Hist. Nat. Poiss., V, 1803, 144 (*Hypostomus guacari* Lacépède).1217. *Hypostomus plecostomus* (Linnæus).

Both slopes of Panama, south to Uruguay.

Loricaria plecostomus Linnæus, Syst. Nat., ed. XII, I, 1766, 508, America.*Hypostomus guacari* Lacépède, Hist. Nat. Poiss., V, 1803, 145, America.*Loricaria flava* Shaw, Gen. Zool., V, 1805, 38, pl. 101.*Hypostomus robeni* Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 46, Trinidad.*Hypostomus brasiliensis* Bleeker, Silures de Suriname, 1864, 7, Surinam.*Plecostomus seminudus* Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2nd ser., I, 1888, 169, Brazil.*Plecostomus boulengeri* Eigenmann and Kennedy, Proc. Acad. Nat. Sci. Phila., LV, 1903, 502, Paraguay.

Genus 393. THYSANOCARA Regan.

Thysanocara Regan, Ann. Mag. Nat. Hist., ser. 7, XVII, 1906, 95 (*Hypostomus cirrhosus* Cuvier and Valenciennes).1218. *Thysanocara spinosa* (Meek and Hildebrand).

Rare, male taken at mouth of Rio Yape, the female (type) in Rio Calobre, Panama.

Ancistrus spinosus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 252, Rio Calobre, tributary of the Rio Bayano, Panama.

Genus 394. XENOCARA Regan.

Xenocara Regan, Trans. Zool. Soc. Lond., XVII, 1903, 251 (*Chætostomus latifrons* Gunther).1219. *Xenocara chagresi* (Eigenmann and Eigenmann).

Rio Chagres, Panama.

Ancistrus chagresi Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2nd ser., II, 1889, 47, Rio Chagres, Panama.

Genus 395. CHÆTOSTOMA Heckel.

Chætostoma Heckel, in Tschudi, Fauna Peruana, 1846, 25 (*C. lobarhynchus* Heckel).*Chætostomus* Kner and authors generally; needlessly altered spelling.

1220. *Chaetostoma fischeri* (Steindachner). *Couchu*.
Rio Mamoni, near Panama.
Chaetostomus fischeri Steindachner, Flussfische Sudamerikas, I, 1879, 14,
Rio Mamoni, Panama
- Genus 396. **LASIANCISTRUS** Regan.
Lasiancistrus Regan, Trans. Zool. Soc. Lond., XVII, Pt. III, 1904, 224
and 237 (*Chaetostomus heteracanthus* Gunther).
1221. *Lasiancistrus planiceps* (Meek and Hildebrand).
Rio Tuyra Basin, Panama.
Ancistrus planiceps Meek and Hildebrand, Field Mus. Nat. Hist., Zool.,
X, 1913, 79, Pl. X; loc. cit., 1916, Rio Tuyra, Boca de Cupe, Panama.
- Genus 397. **LEPTOANCISTRUS** Meek and Hildebrand.
Leptoancistrus Meek and Hildebrand, Field. Mus. Nat. Hist., Zool.,
X, 1916, 254 (*Ancanthus canescens* Meek and Hildebrand).
1222. *Leptoancistrus canensis* (Meek and Hildebrand).
Mountain streams near Cana, Panama
Acanthicus canensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool.,
X, 1913, 80, Rio Cana, Cana, Panama.
- Genus 398. **STURISOMA** Swainson.
Sturisoma Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 304
(*Loricaria rostrata* Spix).
Hemiodon Kner, Panzerwelse, Denkschr., Kais. Akad. Wiss. Wien, 1854,
65 (*H. depressus* Kner).
1223. *Sturisoma rostrata* (Spix).
Rivers of eastern South America, from Panama to the Paraguay;
widely distributed and abundant.
Loricaria rostrata Spix, Pisc. Brasil, 1829, 5, Pl. III, Rio Blanco, Brazil
Loricaria acuta Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840,
349 (472), pl. 452, Barra di Rio Negro.
? *Loricaria barbata* Kner, Panzerwelse, Denkschr. Kais. Akad. Wiss.
Wien, 1853, 87, Rio Cujuba.
- Genus 399. **ANCISTRUS** Bleeker.
Ancistrus Kner, Denkschr., Kais. Akad. Wiss. Wien, VII, 1854, 272
(abstract issued in 1853; *Ancistrus medians* Kner, as first restricted
by Gill).
Hemiancistrus Bleeker, Nederl. Tijdschr. Dierk., I, 1862 (1863), 78
(*Ancistrus mediana* Kner).
1224. *Ancistrus guacharote* (Cuvier and Valenciennes).
Porto Rico.
Hypostomus guacharote Cuvier and Valenciennes, Hist. Nat. Poiss., XV,
1840, 375 (508), Porto Rico.
1225. *Ancistrus aspidolepis* (Gunther).
Veragua, Central America.
Chaetostomus aspidolepis Gunther, Trans. Zool. Soc. Lond., VI, 1868
(Nov. 27), 477, Veragua.
- Genus 400. **LORICARIA** Linnæus.
Loricaria Linnæus, Syst. Nat., ed. X, 1753, 307 (*L. cataphracta* Linnæus).
Hemiodon Kner, Denkschr. Akad. Wiss. Wien, 1854, about 65 (*H. de-*
pressus Kner).
Loricariichthys Bleeker, Nederl. Tijdschr. Dierk., I, 1863, 3 (*Loricaria*
maculata Bloch)
Pseudoloricaria Bleeker, loc. cit., 3 (*Loricaria læviuscula* Cuvier and
Valenciennes).
Parahemiodon Bleeker, loc. cit., 3 (*P. typus* Bleeker).
Hemioloricaria Bleeker, loc. cit., 3 (*H. caracasenses* Bleeker).
Pseudohemiodon Bleeker, loc. cit., 3 (*Hemiodon platycephalus* Kner).
1226. *Loricaria variegata* Steindachner.
Rio Mamoni, Panama.
Loricaria variegata Steindachner, Flussfische Sudamerikas, I, 1879, 15,
Rio Mamoni, Panama.

1227. *Loricaria uracantha* Kner and Steindachner.
Rio Chagres, Panama.
Loricaria uracantha Kner and Steindachner, Abhandl., Bayer. Acad. Wiss., 1865, 56, Rio Chagres, Panama.
Loricaria bransfordi Gill, Proc. Acad. Nat. Sci. Phila., XXVIII, 1876, 338, Empire Station, Isthmus Railway, Panama.
1228. *Loricaria latiura* Eigenmann and Vance.
Rio Tuyra and Rio Atrato Basins, Panama; probably not in Pacific streams.
Loricaria filamentosa latiura Eigenmann and Vance, Ind. Univ. Stud., No. 16, 1912, 13, Boca de Certegai, Panama.
Loricaria tuyrensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 81, Rio Capeti, Panama.
1229. *Loricaria capetensis* Meek and Hildebrand.
Rio Tuyra Basin, Panama
Loricaria capetensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 80, Pl. XII; loc. cit., 1916; Rio Capeti, tributary Rio Tuyra, Panama.
1230. *Loricaria fimbriata* Eigenmann and Vance.
Rio Tuyra Basin, Panama.
Loricaria fimbriata Eigenmann and Vance, Ind. Univ. Stud., No. 16, 1912, 12; Boca de Certegai and Bernal Creek, Colombia.
1231. *Loricaria altipinnis* Breder.
Tuyra Basin, Pacific slope of Panama.
Loricaria altipinnis C. M. Breder, jr., Amer. Mus. Novitates, CLXXX, 1925, 1, fig 1, 2, Rio Chucunaque, Panama.
- Genus 401. **RINELORICARIA** Bleeker.
Rineloricaria Bleeker, Nederl. Tijdschr. Dierk., I, 1862 (1863), 80 (*Loricaria lima* Kner).
1232. *Rineloricaria lima* (Kner).
Coastwise streams of Brazil, from the Parahyba northward to Panama; widely distributed and abundant in rocky brooks
Loricaria lima Kner, Panzerwelse, Denkschr. Kais. Akad. Wiss. Wien, 1853, 89, Brazil.
? *Loricaria strigilata* Hensel, Wiegmann's Archiv, 1868, 368, Santa Cruz.
- Genus 402. **OXYLORICARIA** Bleeker.
Oxyloricaria Bleeker, Nederl. Tijdschr. Dierk., I, 1863, 5 (*Loricaria barbata* Kner).
1233. *Oxyloricaria panamensis* (Eigenmann and Eigenmann).
Pacific slope of Panama and Colombia, south to Ecuador.
Loricaria rostrata (non Spix) Steindachner, Denkschr. Kais. Akad. Wiss. Wien, XLI, 1879, 165, Rio Mamoni, Panama.
Loricaria panamensis Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., ser. 2, II, 1889, 34, Panama.
Loricaria aurea Steindachner, Anz. Kais. Akad. Wiss. Wien, 1900, 206, Rio Magdalena.
Loricaria frenata Boulenger, Ann. Mag. Nat. Hist., ser. 7, IX, 1902, 69, northwestern Ecuador. *L. aurea* and *L. frenata* are extralimital and may not belong here.
Oxyloricaria dariensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 81, Rio Tuyra and Rio Bayano Basins, Panama.
1234. *Oxyloricaria citurensis* Meek and Hildebrand.
Rio Bayano and Rio Tuyra Basins, Panama.
Oxyloricaria citurensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 82, Pl. XII, loc. cit., 1916, Rio Cituro, Panama.

Order Z. INIOMI

Family 99. SYNODONTIDÆ. Lizardfishes

Genus 403. TRACHINOCEPHALUS Gill.

Trachinocephalus Gill, Cat. Fish. East Coast N. Amer., Suppl. Proc. Acad. Nat. Sci. Phila., XIII, 1861, 53 (*Saurus myops* Cuvier and Valenciennes).

1235. *Trachinocephalus myops* (Forster). *Ground spearing; Lagarto; Snakefish.*

Tropical parts of the western Atlantic; West Indies and Brazil to coast of South Carolina; occasionally northward to Woods Hole.

Salmo myops Forster, in Bloch and Schneider, Syst. Ichth., 1801, 421, St. Helena.

Osmerus lemniscatus Lacépède, Hist. Nat. Poiss., V, 1803, 236, Martinique; after Plumier.

Saurus truncatus Agassiz, Pisc. Brasil., 1828, 82, Brazil.

Saurus brevirostris Poey, Memorias, II, 1861, 305, Cuba; erroneously stated to have 10 rays.

Genus 404. SYNODUS Gronow. *Lizardfishes.*

Synodus Gronow, in Scopoli, Int. Hist. Nat., 1777, 449 (*S. synodus* L.).

Tirus Rafinesque, Caratteri, 1810, 56 (*T. marmoratus* Rafinesque = *Esox synodus* Linnaeus).

Saurus Cuvier, Règne Anim., ed. 1, 1817, 169 (*Salmo saurus* Linnaeus).

Alpsmaris Risso, Europ. Mérid., III, 1826, 458 (*A. russo* Risso, very young).

1236. *Synodus synodus* (Linnaeus).

Coast of Brazil and Lesser Antilles.

?*Esox synodus* Linnaeus, Syst. Nat., ed. XII, 1766, 516, America; based on *Synodus* of Gronow.

1237. *Synodus saurus* (Linnaeus). *Tarentola; Lacerto; Tiru.*

Coasts of southern Europe and neighboring islands, recorded by Doctor Goode from the Bermudas.

Salmo saurus Linnaeus, Syst. Nat., Ed. XII, 1766, 511, Mediterranean; after Artedi.

Saurus griseus Lowe, Trans., Zool. Soc. Lond., II, 1841, 188, Madena.

Saurida mediterranea Swainson, Nat. Hist. Anim., II, 1839, 287, Mediterranean; name only.

Tirus marmoratus Rafinesque, Caratteri, 1810, 56, Palermo.

Alpsmaris russo Risso, Europ. Mérid., III, 1826, 458; very young.

Synodus lacerta Goode, Bull., U. S. Nat. Mus., V, 1876, 68, Bermuda specimen; not of Risso.

1238. *Synodus foetens* (Linnaeus). *Lizardfish; Gallwasp; Lagarto; Soapfish.*

Cape Cod to Brazil, very common from South Carolina southward on sandy coasts.

Salmo foetens Linnaeus, Syst. Nat., ed. XII, 1766, 513, South Carolina.

Osmerus albidus Lacépède, Hist. Nat. Poiss., V, 1803, 229, Carolinas; after Linnaeus.

Coregonus ruber Lacépède, loc. cit., 263, Martinique; after Plumier.

Esox salmoneus Mitchell, Trans., Lit. Phil. Soc. N. Y., I, 1815, 442, New York.

Saurus longirostris Agassiz, Spix. Pisc. Brasil., 1828, pl. 43, Brazil.

Saurus mexicanus Cuvier, Règne Anim., ed. 2, 1829, 314, Mexico.

Saurus spixianus Poey, Memorias, II, 1861, 304, Cuba.

1239. *Synodus lucioceps* (Ayres).

San Francisco to Santa Barbara, rather common in summer.

Saurus lucioceps Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 69, San Francisco.

1240. *Synodus jenkinsi* Jordan and Bollman.

Gulf of California to Galapagos Islands in 16 to 33 fathoms; common.

Synodus jenkinsi Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 153, off coast of Colombia, Galapagos Islands.

1241. *Synodus scituliceps* Jordan and Gilbert.
Cape San Lucas to Panama.
Synodus scituliceps Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
344, Mazatlan, Mexico.
1242. *Synodus lacertinus* Gilbert.
Acapulco, Mexico; one specimen known.
Synodus lacertinus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 55,
Acapulco, Mexico.
1243. *Synodus poeyi* Jordan.
Cuba; abundant.
Synodus poeyi Jordan, Proc. U. S. Nat. Mus., IX, 1886, 526, Habana.
1244. *Synodus intermedius* (Agassiz). *Sand-dwer*.
Coast of southern Florida to Brazil; not rare.
Saurus intermedius Agassiz, Spix, Pisc. Brasil., 1828, 81, Brazil.
Saurus anolis Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849,
483, Bahia, Martinique.
1245. *Synodus evermanni* Jordan and Bollman.
Mazatlan to coast of Colombia; abundant.
Synodus evermanni Jordan and Bollman, Proc. U. S. Nat. Mus., XII,
1889, 152, Pacific Ocean off coast of Colombia in 33 fathoms.
1246. *Synodus simulans* Garman.
Off Cocos Island.
Synodus simulans Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 251,
Pl. I, fig 3; towings at surface off Cocos Island.
1247. *Synodus acutus* Garman.
Off Panama
Synodus acutus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 252,
off coast of Panama.
- Genus 405. BATHYSAURUS Günther.
Bathysaurus Gunther, Ann. Mag. Nat. Hist., ser. 5, II, 1878, 181 (*B. ferox* Gunther).
1248. *Bathysaurus agassizii* Goode and Bean.
Gulf Stream, in deep water.
Bathysaurus agassizii Goode and Bean, Bull. Mus. Comp. Zool., X,
1882, No. 5, 215, Gulf Stream, 33° N., in 647 fathoms.
1249. *Bathysaurus mollis* Günther.
Southern Pacific to coast of Lower California
Found off Lower California, 23° N., 1,760 fathoms (only N. Amer.
record; Townsend and Nichols, Bull. Amer. Mus. Nat. Hist., LII,
1925, 10).
Bathysaurus mollis Gunther, Ann. Mag. Nat. Hist., 5th ser., II, 1878,
182, middle of South Pacific off Yeddo, in 1,875 and 2,385 fathoms.
- Genus 406. BATHYLACO Goode and Bean
Bathylaco Goode and Bean, Oceanic Ichth., 1895, 57 (*B. nigricans* Goode
and Bean).
1250. *Bathylaco nigricans* Goode and Bean.
Gulf of Mexico; off Santa Cruz, at a depth of 2,393 fathoms; a single
specimen known.
Bathylaco nigricans Goode and Bean, Oceanic Ichth., 1895, 57, fig. 69,
off Santa Cruz, Mexico.
- Genus 407. SAURIDA Cuvier and Valenciennes.
Saurida Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849, 499
(*Salmo tumbil* Bloch).
1251. *Saurida nebulosa* Cuvier and Valenciennes.
West Indies north to St. Eustatius.
Saurida nebulosa Cuvier and Valenciennes, Hist. Nat. Poiss., XXII,
1849, 504, pl. 649.

Family 100. CHLOROPHTHALMIDÆ

Genus 408. CHLOROPHTHALMUS Bonaparte.

Chloropthalmus Bonaparte, Icon. Fauna Italica, III, 1834, 144 (*C. agassizii* Bonaparte).

Hyphalonedrus Goode, Proc. U. S. Nat. Mus., III, 1880, 483 (*H. chalybeius* Goode).

1252. *Chloropthalmus agassizii* Bonaparte.

Atlantic and Mediterranean; western Atlantic.

Chloropthalmus agassizii Bonaparte, Fauna Italica, 1840, pl. 121, Italy.

1253. *Chloropthalmus chalybeius* (Goode).

Gulf Stream, in from 85 to 167 fathoms.

Hyphalonedrus chalybeius Goode, Proc. U. S. Nat. Mus., III, 1880 (Feb. 16, 1881), 484, Gulf Stream off Rhode Island.

1254. *Chloropthalmus truculentus* Goode and Bean.

One specimen, Blake station LII, off Barbados.

Chloropthalmus truculentus Goode and Bean, Oceanic Ichth., 1895, 61, fig. 72, off Barbados in 158 fathoms.

1255. *Chloropthalmus mento* Garman.

Off Panama.

Chloropthalmus mento Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 254, 3, Pl. LIV, figs. 1-1c, Albatross stations 3385, 3386, and 3389 in 210 to 286 fathoms.

Family 101. SUDIDÆ

Genus 409. SUDIS Rafinesque.

Sudis Rafinesque, Caratteri, 1810, 60 (*S. hyalina* Rafinesque).

1256. *Sudis ringens* Jordan and Gilbert.

Santa Barbara Channel, Calif., the only specimen known taken from the stomach of a hake, itself found in the stomach of a long-finned albicore (*Germo*).

Sudis ringens Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 273, Santa Barbara Channel.

1257. *Sudis intermedius* (Poey).

Matanzas, Cuba; one specimen known.

Paralepis intermedius Poey, Repertorio, II, 1867, 416, Mantanzas, Cuba.

Genus 410. ARCTOZENUS Gill.

Arctozenus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 188 (*Paralepis borealis* Reinhardt).

1258. *Arctozenus borealis* (Reinhardt).

Deep waters of Arctic America.

Paralepis borealis Reinhardt, Vidensk. Selsk. Naturv. Math. Afhandl., VII, 1832, 115, Greenland.

1259. *Arctozenus coruscans* (Jordan and Gilbert).

Puget Sound; one specimen known.

Paralepis coruscans Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 411, Port Townsend, Wash.

Genus 411. PARALEPIS Risso.

Paralepis Risso, in Cuvier, Règne Anim., ed. 1, 1817 (*Coregonus paralepis* Risso = *Paralepis coregonoides* Risso).

1260. *Paralepis coregonoides* Risso.

Mediterranean, rare; Gulf stream at Albatross station 2393.

Paralepis coregonoides Risso, Europ. Mérid., III, 1826, 472, Pl. VII, fig. 15, Nice.

1261. *Paralepis barracudina* Fowler and Phillips.

Coast of New Jersey.

Paralepis barracudina Fowler and Phillips, Proc. Acad. Nat. Sci. Phila., LXII, 1910, 403, with figure, Corsons Inlet, Cape May County, N. J.

Genus 412. LESTIDIOPS Hubbs.

Lestidiops Hubbs, Univ. Calif. Publ. Zool., XVI, 1916, 154 (*L. sphyrænops* Hubbs).

1262. Lestidiops sphyrænops Hubbs.

Lestidiops sphyrænops Hubbs, Univ. Calif. Pub. Zool., XVI, 1916, 155, pl. 18, Avalon Bay, Santa Catalina Island.

Family 102. BATHYPTEROIDÆ**Genus 413. BATHYPTEROIS** Günther.

Bathypterois Günther, Ann. Mag. Nat. Hist., ser. 5, II, 1878, 83 (*B. longifilis* Günther).

Synapteretmus Goode and Bean, Oceanic Ichth., 1895, 64 (*Bathypterois quadrifilis* Günther).

Subgenus SYNAPTERETMUS Goode and Bean.**1263. Bathypterois quadrifilis** Günther.

Brazil; St. Vincent.

Bathypterois quadrifilis Günther, Ann. Mag. Nat. Hist., II, 1878, 184, off coast of Brazil in 500 to 770 fathoms.

1264. Bathypterois pectoralis Garman.

Off Central America in deep water.

Bathypterois pectoralis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 257, Pl. K, fig. 1, Albatross stations 3376, 3393, 3407, and 3431 in 885 to 1,132 fathoms.

1265. Bathypterois ventralis Garman.

Off west coast of Mexico.

Bathypterois ventralis Garman, Mem., Mus. Comp. Zool., XXIV, 1899, Pl. LV, fig. 1, 256, Albatross stations 3418 and 3425 in 660 to 680 fathoms.

Family 103. BENTHOSAURIDÆ**Genus 414. BENTHOSAURUS** Goode and Bean.

Benthosaurus Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, No. 5, 168 (*B. grallator* Goode and Bean).

1266. Benthosaurus grallator Goode and Bean.

Gulf of Mexico and Gulf Stream, in lat. 24° 33' N., long. 84° 23' W., and lat. 39° 3' 15" N., long. 70° 50' 45" W.

Benthosaurus grallator Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, No. 5, 168, Gulf Stream.

Family 104. MYCTOPHIDÆ.³² Lanternfishes**Genus 415. MACROSTOMA** Risso.

Macrostoma Risso, Europ. Mérid., III, 1826, 447 (*Macrostoma angustidens* Risso); not *Macrostomus* Wied., a genus of insects.

Notoscopelus Günther, Cat., V, 1864, 405 (*Lampanyctus resplendens* Richardson), to replace *Macrostoma*, regarded as preoccupied.

Catablemella Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., III, 1890, 3, 24, (*C. brachyichir* Eigenmann and Eigenmann).

1267. Macrostoma angustidens Risso. *Prickfish; Maure d'Amplora.*

Atlantic and Indian Oceans, Norway and Greenland.

Macrostoma angustidens Risso, Europ. Mérid., III, 1826, 448, off Nice.

1268. Macrostoma quercinum (Goode and Bean).

Western Atlantic; Mediterranean.

Notoscopelus quercinus Goode and Bean, Oceanic Ichth., 1895, 83, fig. 97, Grand Banks, Newfoundland.

1269. Macrostoma margaritifera (Goode and Bean).

Grand Banks of Newfoundland.

Notoscopelus margaritifera Goode and Bean, Oceanic Ichth., 1895, 84, fig. 98, off Banquereau.

³² Several species recently described as *Myctophum* may prove to belong to other genera.

1270. *Macrostoma castaneum* (Goode and Bean).
Grand Banks.
Notoscopelus castaneus Goode and Bean, *Oceanic Ichth.*, 1895, 84, fig. 95, Grand Banks, Newfoundland.
- Genus 416. **CERATOSCOPELUS** Günther.
Ceratioscopelus Gunther, Cat., V, 1864, 405 and 412 (*Scopelus madeirensis* Lowe).
1271. *Ceratioscopelus madeirensis* (Lowe).
Atlantic and Mediterranean; west to the Grand Banks.
Scopelus madeirensis Lowe, Proc. Zool. Soc. Lond., 1839, 87, Madeira.
- Genus 417 **LAMPANYCTUS** Bonaparte.
Lampanyctus Bonaparte, *Fauna Italica*, Fasc. XXVII, 1840 (*L. bonapartei* Bonaparte).
1272. *Lampanyctus guntheri* Goode and Bean.
Grand Banks
Lampanyctus guntheri Goode and Bean, *Oceanic Ichth.*, 1895, 79, fig. 90, Grand Banks.
1273. *Lampanyctus gemmifer* Goode and Bean.
Gulf Stream.
Lampanyctus gemmifer Goode and Bean, *Oceanic Ichth.*, 1895, 80, fig. 88, Grand Banks, in lat. 39° 40' N., long. 71° 35' W., in 538 fathoms.
1274. *Lampanyctus lacerta* Goode and Bean.
Gulf of Mexico.
Lampanyctus lacerta Goode and Bean, *Oceanic Ichth.*, 1895, 81, fig. 89, Gulf of Mexico, in lat. 28° 38' 30'' N., long. 85° 52' 30'' W.
1275. *Lampanyctus ritteri* Gilbert.
Coast of California; Monterey Bay; San Clemente.
Lampanyctus ritteri Gilbert, Proc. U. S. Nat. Mus., XLVII, 1915, 318, pl. 15, fig. 3, Monterey Bay; San Clemente Island, Calif.
- Genus 418. **NYCTIMASTER** Jordan.
Nyctimaster Jordan, Proc. U. S. Nat. Mus., LIX, 1921, 645 (*Lampanyctus jordani* Gilbert).
Most of the species hitherto referred to *Lampanyctus*, having the scales of the lateral line not larger than the others, should be referred to *Nyctimaster*.
1276. *Nyctimaster townsendi* (Eigenmann and Eigenmann).
Cortez Banks near San Diego, in 45 fathoms
Myctophum townsendi Eigenmann and Eigenmann, *West Amer. Scientist*, 1889, 125, Cortez Banks, off San Diego, Calif.
1277. *Nyctimaster alatus* (Goode and Bean).
Gulf of Mexico.
Lampanyctus alatus Goode and Bean, *Oceanic Ichth.*, 1895, 79, Gulf Stream.
- Genus 419. **LAMPADENA** Goode and Bean.
Lampadena Goode and Bean, *Oceanic Ichth.*, 1895, 85 (*L. speculigera* Goode and Bean).
1278. *Lampadena speculigera* Goode and Bean.
Gulf Stream.
Lampadena speculigera Goode and Bean, *Oceanic Ichth.*, 1895, 85, fig. 99, Gulf Stream, lat. 39° 48' N., long. 70° 36' W., in 551 fathoms.
- Genus 420. **NANNOBRACHIUM** Günther.
Nannobrachium Gunther, *Deep Sea Fishes*, *Challenger*, 1887, 199, (*N. nigrum* Günther).
1279. *Nannobrachium nannochir* (Gilbert).
Alaska to Santa Barbara Islands, in deep water.
Myctophum nannochir Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 51, off Washington, Albatross stations 2925, 3072, and others, in 266 to 685 fathoms.

1280. *Nannobranchium leucopsarum* (Eigenmann and Eigenmann).
Pacific coast of America, from Alaska to San Diego
Myctophum (*Stenobranchius*) *leucopsarum* Eigenmann and Eigenmann,
Proc. Calif. Acad. Sci., 2d ser., III, 1890, 5, off Point Loma, Calif.,
in stomachs of *Sebastodes*.
1281. *Nannobranchium macdonaldi* Goode and Bean.
Gulf Stream.
Nannobranchium macdonaldi Goode and Bean, Oceanic Ichth., 1895, 94,
fig. 110, Gulf Stream.
1282. *Nannobranchium regale* (Gilbert).
Santa Barbara Channel, in deep water.
Myctophum regale Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 544,
Santa Barbara Channel, Albatross station 2923, in 822 fathoms.
1283. *Nannobranchium mexicanum* (Gilbert).
Coast of Lower California, in deep water.
Myctophum mexicanum Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 51,
off Lower California, at Albatross stations 3008 and 3009 in 306
and 857 fathoms.
- Genus 421. *ÆTHOPRORA* Goode and Bean.
Æthoprora Goode and Bean, Oceanic Ichth., 1895, 86 (*Scopelus meto-*
poclampus Cocco).
1284. *Æthoprora effulgens* Goode and Bean.
Western Atlantic.
Æthoprora effulgens Goode and Bean, Oceanic Ichth., 1895, 87, fig.
103, Browns Bank and Albatross station 2127, 19° 45' N., 75° W.,
in 689 fathoms.
1285. *Æthoprora lucida* Goode and Bean.
Gulf Stream.
Æthoprora lucida Goode and Bean, Oceanic Ichth., 1895, 87, fig. 102,
Gulf Stream, lat. 19° 45' N., long. 75° 4' W.
- Genus 422. *COLLETTIA* Goode and Bean.
Collettia Goode and Bean, Oceanic Ichth., 1895, 88 (*Myctophum rafinesquei*
Cocco).
1286. *Collettia rafinesquei* (Cocco).
Mediterranean and Atlantic; Gulf Stream off the New England coast.
Nyctophus rafinesquei Cocco, Alcuni Salmon, etc., 1820, 20, Messina.
1287. *Collettia nocturna* (Poey).
Coast of Cuba.
Myctophum nocturnum Poey, Memorias, II, 1861, 426, Cuba.
- Genus 423. *DIAPHUS* Eigenmann and Eigenmann.
Diaphus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser.,
III, 1890, 3 (*Scopelus engraulis* Gunther).
1288. *Diaphus theta* Eigenmann and Eigenmann.
Point Loma, near San Diego, Calif., to coast of Oregon.
Diaphus theta Eigenmann and Eigenmann, Proc. Calif. Acad. Sci.,
2d ser., III, March, 1890, 4, Point Loma, near San Diego.
Myctophum protoculus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 52,
July 2, coast of Washington, in 584 fathoms, Albatross station 3072,
in 584 fathoms.
- Genus 424. *RHINOSCOPELUS* Lütken.
Rhinoscopelus Lütken, Vidensk. Selsk. Naturv. Copenhagen, VII,
1892, 237 (*Scopelus coccoi* Cocco).
Alysia Lowe, Proc. Zool. Soc. Lond., 1839, 87 (*A. lorica* Lowe=
Scopelus coccoi Cocco; name preoccupied).
1289. *Rhinoscopelus andreae* (Lütken).
Open Atlantic and Indian Ocean; Gulf Stream.
Scopelus andreae Lütken, Spolia Atlantica, 1892, 25, North Atlantic.
1290. *Rhinoscopelus rarus* (Lütken).
Open Atlantic, west to 50° W., 33° N.
Scopelus rarus Lütken, Spolia Atlantica, II, 1892, 26, North Atlantic.

- 1291. *Rhinoscopelus coccoi* (Cocco).**
 West Atlantic, Gulf Stream, Mediterranean, ranging from Newfoundland to Africa.
Scopelus coccoi Cocco, Giorn. Sci. Litt. Art. Sicilia (No. 77), Palermo, 1829, 143 ("Scopelo de Cocco"), Palermo.
Alysia loricata Lowe, Proc. Zool. Soc. Lond., 1839, 87, Madeira.
- Genus 425. MYCTOPHUM Rafinesque. Lanternfishes.**
Myctophum Rafinesque, Indice, 1810, 56 (*M. punctatum* Rafinesque)
Scopelus Cuvier, Règne Anim., ed. 1, 1817, 56 (*Gasteropelecus humboldti* Risso).
Myctophus Cocco, Giorn. Sci. Litt. Art. Sicil., 1829, 44 (*N. rafinesquei*, substitute for *Myctophum*).
- 1292. *Myctophum punctatum* Rafinesque.**
 Warmer parts of the Atlantic, Grand Banks, and from the Gulf Stream to the Mediterranean.
Myctophum punctatum Rafinesque, Indice, 1810, 56, Pl. II, fig. 5, Palermo.
Scopelinus caninianus Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849, 445, coast of Italy.
- 1293. *Myctophum affine* (Lütken).**
 Open Atlantic west to 63° W., 38° N., and in the open Pacific. Perhaps more than one species included.
Scopelus affine Lutken, Spolia Atlantica, II, 1892, 32, open Atlantic.
Myctophum nitidulum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 266, Pl. LVI, fig. 3, lat. 27° 50' N., long. 145° 45' 30" W.
Rhinoscopelus oceanicus Jordan and Evermann, Bull. U. S. Fish Com., XX, 1902, 168, Hawaiian Islands.
Myctophum margaritatum Gilbert, loc. cit., XXIII, pt. 2, 1905, p. 596, pl. 68, fig. 2, off south coast of Molokai, Hawaiian Islands.
- 1294. *Myctophum humboldti* (Risso).**
 Open seas, Mediterranean, and Atlantic on both shores.
Gasteropelecus humboldti Risso, Ichth. Nice, 1810, 358, Nice.
- 1295. *Myctophum californiense* Eigenmann and Eigenmann.**
 Cortez Banks, San Diego, Calif.
 ?*Myctophum boops* Richardson, Voy. *Erebus* and *Terror*, 1845, 39, open sea between Australia and New Zealand.
Myctophum californiense Eigenmann and Eigenmann, West Amer. Scientist, Nov. 9, 1889, 124, San Diego, Calif.
- 1296. *Myctophum gracile* (Lütken).**
 Open Atlantic, west to 48° W., 23° N.
Scopelus gracilis Lütken, Spolia Atlantica, II, 1892, 35, open Atlantic.
 ?*Myctophum hians* Richardson, Voy. *Erebus* and *Terror*, 1845, 41, pl. 27, open sea, locality unknown.
- 1297. *Myctophum benoiti* (Cocco).**
 Mediterranean to Norway and Greenland
Scopelus benoiti Cocco, Lett. su Salmon., 1838, 12, pl. 2, fig. 4, Messina.
- 1298. *Myctophum hygomi* (Lutken).**
 North Atlantic, occasional in our waters.
Scopelus hygomi Lutken, Spolia Atlantica, 1892, 257, North Atlantic.
Myctophum remiger Goode and Bean, Oceanic Ichth., 1895, 75, western Atlantic.
- 1299. *Myctophum oculateum* Garman.**
 Off Panama in deep water.
Myctophum oculateum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 260, Pl. LVI, fig. 2, numerous stations of the *Albatross*.
- 1300. *Myctophum atratum* Garman.**
 Off western Mexico in the open sea.
Myctophum atratum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 268, *Albatross* station 3433, lat. 25° 26' 15" N., long. 109° 48' W., in 1,218 fathoms.

1301. *Myctophum lateratum* Garman.
Off Pacific coast of Mexico in open sea
Myctophum lateratum Garman, Mem Mus Comp. Zool., XXIV, 1899,
267, Pl. LVI, fig. 7, several stations of the *Albatross*.
1302. *Myctophum aurolaternatum* Garman.
Off Panama.
Myctophum aurolaternatum Garman, Mem Mus. Comp. Zool., XXIV,
1899, 264, Pl. LV, fig. 3, *Albatross* station 3382, lat. 6° 21' N., long.
80° 41' W. at surface.
1303. *Myctophum luminosum* Garman.
Off Panama.
Myctophum luminosum Garman, Mem. Mus. Comp. Zool., XXIV,
1899, 263, Pl. LV, fig. 2, *Albatross* station 3042, lat. 0° 57' 30'' S.,
long. 89° 3' 30'' W., in 421 fathoms.
1304. *Myctophum tenuiculum* Garman.
Off Panama in deep water
Myctophum tenuiculum Garman, Mem Mus. Comp. Zool., XXIV, 1899,
262, pl. 5, fig. 5, *Albatross* station 3382, lat. 6° 21' N., 80° 41' W.,
at surface.
- Genus 426. **BENTHOSEMA** Goode and Bean.
Benthoosema Goode and Bean, Oceanic Ichth., 1895, 76 (*Scopelus muelleri*
Gmelin).
1305. *Benthoosema arcticum* (Lutken).
Davis Straits, Greenland.
Scopelus arcticus Lutken, Spolia Atlantica, 1892, 29, Davis Straits.
1306. *Benthoosema mulleri* (Gmelin).
North Atlantic, occasional from Norway to Greenland, more abundant
southward to South Carolina; probably always in very deep water.
Salmo mulleri Gmelin, Syst. Nat., 1788, 1378, Norway; after Strom.
Scopelus glacialis Reinhardt, Overs., 1835, 36, Greenland.
- Genus 427. **SCOPELENGYS** Alcock.
Scopelengys Alcock, Ann. Mag. Nat. Hist., ser. 6, VI, 1890, 302 (*S.*
tristis Alcock).
1307. *Scopelengys dispar* Garman.
Off Panama in deep water.
Scopelengys dispar Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
254, Pl. LIV, figs. 2-2d, *Albatross* stations 3353, 3360, 3382, and
3383, in from 695 to 1,832 fathoms.
- Genus 428. **TARLETONBEANIA** Eigenmann and Eigenmann.
Tarletonbeania Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., IV,
1891, 6 (*T. tenua* Eigenmann and Eigenmann).
1308. *Tarletonbeania crenularis* (Jordan and Gilbert).
San Diego and vicinity
Myctophum crenulare Jordan and Gilbert, Proc. U. S. Nat. Mus., III,
1880, 274, Santa Barbara, Calif.
Myctophum procellarium Bean, in Jordan and Gilbert, loc. cit., 467, off
Straits of Fuca.
Tarletonbeania tenua Eigenmann and Eigenmann, Proc. Calif. Acad.
Sci., IV, 1890, 7, Coronado Islands near San Diego, Calif.

Family 105. IPNOPIDÆ

Genus 429. IPNOPS Günther.

Ipnops Günther, Ann. Mag. Nat. Hist., II, 1878, 187 (*I. murrayi*
Günther).

1309. *Ipnops murrayi* Günther.

Bottom fish, at about 2,000 fathoms; coast of Brazil, Tristan da Cunha,
Celebes, in tropical America (24° 36' N., 84° W.) and off Bequia,
Windward Islands.

Ipnops murrayi Günther, Ann. Mag. Nat. Hist., II, 1878, 187, coast
of Brazil, etc.

1310. *Ipnots agassizi* Garman.

Pacific coast of Central America in deep water.

Ipnots agassizi Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 259, Pl. H, figs. 2, 2a, Albatross station 3413, lat. 2° 34' N., long. 92° 6' W., in 1,360 fathoms.**Family 106. OMOSUDIDÆ****Genus 430. OMOSUDIS Günther.***Omosudis* Günther, *Challenger* Rept., XXII, 1887, 201 (*O. lowii* Günther).**1311. *Omosudis lowii* Günther.**

Open sea; Gulf Stream at Albatross station 2392.

Omosudis lowii Günther, *Challenger* Rept., XXII, 1887, 201, pl. 52, figs. C, C', Philippine Islands, Magdalena.**Family 107. ALEPISAURIDÆ. Handsawfishes****Genus 431. ALEPISAURUS Lowe.***Plagyodus* Steller, in Pallas, Zoogr. Rosso-Asiat., III, 1811, 383 (no type named, doubtless *P. æsculapius* Bean); given in oblique form only, as *Plagyodontem*.*Alepisaurus* Lowe, Proc. Zool. Soc. Lond., I, 1833 (*A. ferox* Lowe).**1312. *Alepisaurus ferox* Lowe. Lancelfish.**

Deep waters of the Atlantic; occasional off the coasts of Nova Scotia, Massachusetts, and on the Grand Banks.

Alepisaurus ferox Lowe, Trans. Zool. Soc. Lond., I, 1833, 395, Madeira.*Alepisaurus azureus* Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849, 530, Canaries.**1313. *Alepisaurus æsculapius* (Bean). Wolffish; Subotka.**

Coast of Alaska southward to California.

Plagyodus Steller in Pallas, Zoogr. Rosso-Asiat., III, 1811, 383, Kurile Islands.*Alepidosaurus æsculapius* Bean, Proc. U. S. Nat. Mus., V, 1882, 661, Iliuliuk, Unalaska.**Genus 432. CAULOPUS Gill. Handsawfishes.***Caulopus* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 128 (*Alepidosaurus borealis* Gill).**1314. *Caulopus altivelis* (Poey).**

Cuba, in deep water.

Alepisaurus altivelis Poey, Memorias, II, 1861, 302, Cuba.*Alepidosaurus (Caulopus) poeyi* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 131, Cuba.**1315. *Caulopus borealis* (Gill). Handsawfish; Serra.**

Pacific coast of America from Alaska to Puget Sound.

Alepidosaurus borealis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 128, Puget Sound.*Alepidosaurus serra* Gill, loc. cit., 129. "Monterey, Lower California."**Order AA. CETUNCULI****Family 108. RONDELETIIDÆ****Genus 433. RONDELETIA Goode and Bean.***Rondeletia* Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894, 454 (*R. bicolor* Goode and Bean).**1316. *Rondeletia bicolor* Goode and Bean.**

Gulf Stream.

Rondeletia bicolor Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894 (1895), 454, pl. 17, fig. 7, and Oceanic Ichth., 1895, 68, fig. 77, at Albatross station 2724 in 1,641 fathoms.

Family 109. CETOMIMIDÆ

Genus 434. CETOMIMUS Goode and Bean.

Cetomimus Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894, 453
(*C. gillii* Goode and Bean).

1317. *Cetomimus gillii* Goode and Bean.

Gulf Stream.

Cetomimus gillii Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894
(1895), 452, pl. 17, fig. 2, and in Oceanic Ichth., 1895, 69, fig. 78,
Gulf Stream, in lat. 39° 35' N., long. 71° 24' 30'' W.

1318. *Cetomimus storeri* Goode and Bean.

Gulf Stream.

Cetomimus storeri Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894
(1895), 453, pl. 17, fig. 3, and in Oceanic Ichth., 1895, 69, fig. 79,
Gulf Stream at Albatross station 2222 in 1,535 fathoms.

Order BB. XENOMI

Family 110. DALLIIDÆ. Alaska blackfish

Genus 435. DALLIA Bean.

Dallia Bean, Proc. U. S. Nat. Mus., II, 1879, 358 (*D. pectoralis* Bean).

1319. *Dallia pectoralis* Bean. Alaska blackfish; *Charma ryba*.

Streams and ponds of northern Alaska and Siberia.

Dallia pectoralis Bean, Proc. U. S. Nat. Mus., II, 1879, 358, St.
Michaels, Alaska.

Umbra delicatissima Smitt, Offer. Kongl. Vetensk. Akad. Forh. Arg.,
Stockholm, 1881, 38, pl. 5, fig. 1, Pittlekag, Siberia.

Order CC. HAPLOMI

Family 111. ESOCIDÆ. Pikes

Genus 436. ESOX Linnæus.

Esox Linnæus, Syst. Nat., ed. X, 1758, 313 (*E. lucius* Linnæus).

Lucrus Geoffroy, Descr. de 719 Plantes, etc., 1767, 407 (*Esox lucius*
Linnæus).

Lucrus Rafinesque, Caratteri, 1810, 59 (*Esox lucius* Linnæus = *L. vorax*
Rafinesque).

Picorellus Rafinesque, Ichth. Ohiensis, 1820, 70 (*Esox vittatus* Rafinesque,
a mythical species).

Mascalongus Jordan, Klippart's rept., Ohio Fish Com., 1878, 92 (*Esox*
masquinongy Mitchill = *Esox nobilior* Thompson).

Kenoza Jordan and Evermann, Fishes North and Mid. Amer., 1896,
626 (*Esox americanus* Gmelin).

Subgenus MASCALONGUS Jordan.

1320. *Esox masquinongy* Mitchill. *Muskallunge*,³³ *Muskellunge*; *Maskinongy*;
Muscalonge; *Great pike*.

Great Lakes region; Upper Mississippi Valley and northward.

Esox masquinongy Mitchill, "Mirror," 1824, 297, Lake Erie.

Esox masquinongy Mitchill, in Kirtland, Fishes of Ohio, 1838, 194,
Lake Erie.

Esox nobilior Thompson, Proc. Bost. Soc. Nat. Hist., III, 1850, 163,
Lake Champlain.

Esox nobilis Kirtland, Proc. Cleveland Acad. Nat. Sci., 1854, 84, Lake
Erie; name a slip of the pen for *Esox nobilior*.

1321. *Esox masquinongy ohiensis* Kirtland.

Ohio River and its tributaries; Chautauqua Lake, N. Y.

Esox salmoneus Rafinesque, Amer. Month. Mag., III, 1818, 447, Ohio
River; not *Esox salmoneus* Mitchill 1815 = *Synodus fatens*.

Esox ohiensis Kirtland, Proc. Cleveland Acad. Nat. Sci., Feb. 7, 1854,
Mahoning River, Ohio.

³³ The common names of the fishes of the pike family, and of the pike perches, with which they are confused, are too numerous to be given here. As Weed, who gives many of them (Field Mus. Nat. Hist., Zool. leaflet No. 9, 1827) remarks, "There are as many as 80 names for 1 fish and 7 fish for one name."

1322. *Esox masquinongy immaculatus* Garrard. *Great northern pike*.
Lakes of Wisconsin and Minnesota.
Esox immaculatus Garrard, MS; noticed in various fishing journals;
Eagle Lake, northern Wisconsin.

Subgenus **ESOX** Linnæus.

1323. *Esox estor* ³⁴ LeSueur. *American pike*.
Northern North America; eastern United States south to New York
and the Ohio River.
Esox estor LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 413, Lake
Erie at Buffalo.
??*Esox vittatus* Rafinesque, Amer. Month. Mag., III, 1818, 447, Ohio
River; mythical; described from hearsay.
??*Esox australis* Cuvier and Valenciennes, Hist. Nat. Poiss., XVII,
1846, 323, locality unknown; thought to be Van Diemen's Land
(Tasmania).
Esox depraudus LeSueur in Cuvier and Valenciennes, Hist. Nat. Poiss.,
XVIII, 1846, 336, Wabash River, New Harmony, Ind.
Esox lugubris LeSueur, loc. cit., 338, Crab Orchard, Ky.
Esox boreus Agassiz, Lake Superior, 1850, 317, Lake Superior.
Esox lucoides Agassiz and Girard, in Herbert Forrester's Fish and
Fishing, 1850, 154, Lake Superior.

Subgenus **KENOZA** Jordan and Evermann.

1324. *Esox niger* LeSueur. *Eastern pickerel; Green pike; Jack*.
Massachusetts to Florida, in lowland streams and swamps.
Esox reticulatus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 414,
Connecticut River, Adams, Mass.; Philadelphia; name preoccupied
in Europe.
?*Esox phaleratus* Say in LeSueur, loc. cit., 416, St. Augustine, Fla.;
unidentifiable.
Esox niger LeSueur, loc. cit., 415, South Carolina.
Esox tridecemlineatus Mitchell, "Mirror," 1825, 361, Oneida Lake,
N. Y.
1325. *Esox vermiculatus* LeSueur. *Little pickerel; Grass pike*.
Mississippi Valley and tributaries of Lake Erie and Lake Michigan,
south to Mississippi and Arkansas.
Esox vermiculatus LeSueur in Cuvier and Valenciennes, Hist. Nat.
Poiss., XVIII, 1846, 333, Wabash River, New Harmony, Ind.
Esox lineatus LeSueur, loc. cit., 335, Wabash River, New Harmony, Ind.
Esox crassus Agassiz, Amer. Jour. Sci. Arts, 1854, 308, Tennessee
River, Huntsville, Ala.
Esox umbrosus Kirtland, Proc. Cleveland Acad. Nat. Sci., 1854, 79,
small Bay of Rocky River, Rockport, near Cleveland, Ohio.
Esox cypho Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 78, Water-
ford, Oakland County, Mich.
Esox porosus Cope, Trans. Amer. Phil. Soc. Phila., XIII, 1866, 408,
Waterford, Oakland County, Mich.; substitute for *cypho* regarded
as an inept name.
1326. *Esox americanus* Gmelin. *Barred pickerel*.
Esox lucius americanus Gmelin, Syst. Nat., 1788, 1390, Long Island,
N. Y.; after Schöpfung.
Esox scomberius Mitchell, Amer. Month. Mag., 1818, 322, Murderers
Creek, N. Y.
Esox fasciatus DeKay, New York Fauna: Fishes, 1842, 224, Murderers
Creek and other streams near New York.
Esox ornatus Girard, "Proc. Bost. Soc. Nat. Hist., V, 1854, 41," Proc.
Acad. Nat. Sci., Phila., XXXVII, 1885, 369, 41; Charles River,
Mass.
Esox raveneli Holbrook, Ichth. South Carolina, 1860, 201, Charleston,
S. C.

³⁴ Not clearly separated from *Esox lucius* L. of Europe.

Family 112. UMBRIDÆ. Mud minnows

Genus 437. UMBRA Müller.

Umbra Kramer, MS., in Muller, Anim. Austr. Infer., 1756; Müller, Abhandl. Akad. Wiss. Wien, 1842, 188 (*crameri*).

Melanura Agassiz, Amer. Jour. Sci. Arts, XVI, 1853, 135 (*M. annulata* Agassiz, not *Ezoglossum annulatum* Rafinesque, etc. = *Leuciscus pygmæus* DeKay).

1327. *Umbra limi* (Kirtland). Mud minnow; Dogfish.

Great Lakes region and upper Mississippi Valley in weedy streams and ponds; Lake Champlain to Minnesota. A hardy and attractive aquarium fish.

Hydrargira lima Kirtland, Bost. Jour. Nat. Hist., III, 1840, 277, streams in northern Ohio.

Hydrargira fusca Thompson, Hist. Vermont, 1842, 137, Lake Champlain.

Hydrargira atricauda DeKay, N. Y. Fauna: Fishes, 1842, 220, Lake Champlain.

1328. *Umbra pygmæa* (DeKay). Striped mud minnow; Eastern mud minnow. Lowland streams and swamps coastwise, from Long Island to the Neuse River, N. C.; locally abundant.

Leuciscus pygmæus DeKay, New York Fauna: Fishes, 1842, 214, Tappan, Rockland County, N. Y.

Fundulus fuscus Ayres, Bost. Jour. Nat. Hist., IV, 1843, 296, Brookhaven, Long Island.

Umbra pygmæa bilineata Shufeldt, Aquatic Life, July, 1917, tributaries of Chesapeake Bay, slight color variation.

Order DD. CYPRINODONTES. (Microcyprini)

Family 113. CYPRINODONTIDÆ. Killfishes

Genus 438. PROFUNDULUS Hubbs.

Profundulus Hubbs, Misc. Pub. 13, Mus. Zool. Univ. Mich., Jan. 18, 1924, 12 (*Fundulus punctatus* Günther).

1329. *Profundulus punctatus* (Günther).

Southern Mexico and Central America in streams and lakes, south to Ecuador.

Fundulus punctatus Günther, Cat., VI, 1866, 320, Chiapas.

Fundulus guatemalensis Günther, loc. cit., 321, Lake Duenas

Fundulus pachycephalus Günther, loc. cit., 321, Lake Atitlan, Guatemala.

1330. *Profundulus labialis* (Günther).

Rivers of Guatemala in Atlantic drainage, Lake Yzabal, Rio Chisoy and Rio San Geronimo.

Fundulus labialis Günther, Cat., VI, 1866, 319, Rio San Geronimo, Guatemala, Yzabal.

1331. *Profundulus oaxacæ* (Meek).

Streams of Oaxaca, Mexico.

Fundulus oaxacæ Meek, Pub., Field Col. Mus., Zool., III, 1902, 90, Oaxaca, Mexico.

1332. *Profundulus candalarius* Hubbs.

Profundulus candalarius Hubbs, Misc. Pub. 13, Mus. Zool., Univ. Mich., Jan. 18, 1924, 15, Candalaria, Guatemala, Rio Chiapas Basin, Atlantic drainage.

Genus 439. CUBANICHTHYS Hubbs.

Cubanichthys Hubbs, Misc. Pub. Univ. Mich. Mus. Zool., No. 16, July 9, 1926, 4 (*Fundulus cubensis* Eigenmann).

1333. *Cubanichthys cubensis* (Eigenmann).

Streams of Pinar del Rio, Cuba.

Fundulus cubensis Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903), 222, Pinar del Rio, Cuba.

Genus 440. LEPTOLUCANIA Myers.

Leptolucania Myers, Amer. Mus. Novitates, June 6, 1924, 8 (*Heterandria ommata* Jordan = *Zygoneustes manni* Hay).

1334. *Leptolucania ommata* (Jordan).
Florida; Indian River and Santa Fe River, and their tributaries.
Heterandria ommata Jordan, Proc. U. S. Nat. Mus., VII, 1884, 323,
Indian River, Fla.
Zygonectes manni Hay, loc. cit., VIII, 1885, 555, Yellow Water River,
Fla.
- Genus 441. **CHRIOPEOPS** Fowler.
Chriopeops Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 425
(*Lucania goodei* Jordan).
1335. *Chriopeops goodei* (Jordan).
Waters of the Florida Everglades; locally abundant.
Lucania goodei Jordan, Proc. U. S. Nat. Mus., II, 1879, 240, Arlington
River, Fla., a tributary of the St. Johns.
- Genus 442. **LUCANIA** Girard.
Lucania Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 118 (*Lamia
venusta* Girard); not *Lucanus*, a genus of beetles.
1336. *Lucania parva* (Baird and Girard). *Rainwaterfish*; *Little killifish*.
Atlantic coast from Connecticut to Key West, Texas, and Mexico.
Cyprinodon parvus Baird and Girard, Ninth Smithsonian. Rept., 1854
(1855), 345, Long Island.
Lamia venusta Girard, U. S. Mex. Bound. Surv., Ichth., 1859, 71, pl.
39, figs. 20-23, Indianola, Tex.
Lucania affinis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 118,
Matamoras, Mexico.
- Genus 443. **FUNDULUS** Lacépède.
Fundulus Lacépède, Hist. Nat. Poiss., V, 1803, 37 (*Fundulus mudfish
Lacépède*=*Cobitis heteroclitus* Linnæus).
Hydrargyra Lacépède, loc. cit., 378 (*H. swampina* Lacépède=*Fundulus
majalis* Walbaum).
Boiborys (Broussonet) Goode and Bean, Proc. U. S. Nat. Mus., VIII,
1885, 205 (*Cobitis heteroclitus* Linnæus).
Gambusia Jordan and Evermann, Fish. North and Mid Amer.,
1896, 633 (*Fundulus rathbuni* Jordan and Meek).
Galasaccus Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 417
(*Hydrargyra similis* Baird and Girard).
1337. *Fundulus heteroclitus* (Linnæus). *Common killifish*; *Minnow*; *Pike
minnow*; *Mummichog*; *Mud-minnow*; *Muddabblers*; *Cobbler*.
Coast of Maine to the Rio Grande; Gulf coast; common in brackish
waters.
Cobitis heteroclitus Linnæus, Syst. Nat., ed. XII, 1766, 500, Charleston,
S. C.
Pacilia canicola Bloch and Schneider, Syst. Ichth., 1801, 452, Carolinas.
Fundulus mudfish Lacépède, Hist. Nat. Poiss., V, 1803, 37, Carolinas.
Hydrargyra swampina Lacépède, loc. cit., 378, South Carolina.
Fundulus nisorius Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 456,
reported from Gaboon, West Africa, appears to be *heteroclitus*, wrong
locality.
1338. *Fundulus heteroclitus fonticola* Cuvier and Valenciennes.
West Indies.
Fundulus fonticola Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII,
1846, 148 (198), mountain springs of Porto Rico.
Fundulus antillarum Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII,
1916, 418, fig. 1, St. Martins, West Indies.
1339. *Fundulus heteroclitus bermudæ* Gunther. *Mangrove minnow*.
Bermudas.
Fundulus bermudæ Günther, Ann. Mag. Nat. Hist., ser. 4, XIV, 1874,
370, Bermudas.
Fundulus rhizophoræ Goode, Amer. Jour. Sci. Arts, XIII, 1877, 298,
Basden Pond, Bermudas.

1340. *Fundulus heteroclitus macrolepidotus* (Walbaum). *Common cobbler; Killifish, Catfish; Dogfish; Kully; Killie; River minnow; Mummichog; Blunthead*
Maine to Virginia; common in brackish waters. Hardly distinct from *F. heteroclitus*, of which it is the northern representative.
Cobitis macrolepidota Walbaum, *Artedi Pisc.*, III, 1792, 11, Long Island; after Schopf.
Cobitis killifish Walbaum, loc. cit., 12, Long Island; after Schopf.
Esox pisciculus Mitchell, *Trans. Lit. Phil. Soc. N. Y.*, I, 1815, 440, New York.
Esox pisculentus Mitchell, loc. cit., 441, New York.
Hydrargyra nigrofasciata LeSueur, *Jour. Acad. Nat. Sci. Phila.*, I, 1817, 133, Newport, R. I.
Fundulus viridescens DeKay, *New York Fauna: Fishes*, 1842, 217, New York.
Fundulus zebra DeKay, loc. cit., 218, New York.
Hydrargyra ornata LeSueur, loc. cit., 131, Delaware River, near Philadelphia.
1341. *Fundulus heteroclitus grandis* Baird and Girard.
Gulf coast of the United States.
Fundulus grandis Baird and Girard, *Proc. Acad. Nat. Sci. Phila.*, VI, 1853, 389, Indianola, Tex.
Fundulus floridensis Girard, loc. cit., XI, 1859, 157, Charlotte Bay, Fla.
Zygonectes funduloides Evermann, *Bull. U. S. Fish Com.*, XI, 1891, 85, pl. 35, fig. 3, Dickinson Bayou, Dickinson, Tex.
1342. *Fundulus heteroclitus badius* Garman.
Gulf of St. Lawrence.
Fundulus heteroclitus badius Garman, *Mem. Mus. Comp. Zool.*, XIX, 1895, Grand Manan.
1343. *Fundulus majalis* (Walbaum). *Killifish; Mayfish; Rockfish; Catfish; Bass killy.*
Cape Cod to Florida, chiefly marine in shallow bays.
Cobitis majalis Walbaum, *Artedi Pisc.*, II, 1792, 12, Long Island; after Mayfish Schopf.
Esox flavulus Mitchell, *Trans. Lit. Phil. Soc. N. Y.*, I, 1815, 439, New York.
Esox zonatus Mitchell, loc. cit., 443, New York.
Hydrargyra trifasciata Storer, *Jour. Bost. Soc. Nat. Hist.*, I, 1837, 417, Massachusetts.
Hydrargyra vernalis Cuvier and Valenciennes, *Hist. Nat. Poiss.*, XVIII, 1846, 154 (206), pl. 531; probably South Carolina.
1344. *Fundulus ocellaris* Jordan and Gilbert.
Pensacola to Fort Myers, Fla.
Fundulus ocellaris Jordan and Gilbert, *Proc. U. S. Nat. Mus.*, V, 1882, 254, Pensacola, Fla.
1345. *Fundulus parvipinnis* Girard.
Coast of California from Point Conception southward to Lower California.
Fundulus parvipinnis Girard, *Proc. Acad. Nat. Sci. Phila.*, VII, 1854, 154, San Diego, Calif.
1346. *Fundulus parvipinnis brevis* Osburn and Nichols.
Magdalena Bay, Lower California.
Fundulus parvipinnis brevis Osburn and Nichols, *Bull. Amer. Mus. Nat. Hist.*, XXV, 1916, 150, fig. 5, Magdalena Bay, Lower California.
1347. *Fundulus pallidus* Evermann.
Galveston Bay near Swan Lake, Tex.
Fundulus pallidus Evermann, *Bull. U. S. Fish Com.*, XI, 1891 (1892), 84, pl. 35, Galveston Bay near Swan Lake, Tex.
1348. *Fundulus vinctus* Jordan and Gilbert.
Coast of Lower California.
Fundulus vinctus Jordan and Gilbert, *Proc. U. S. Nat. Mus.*, V, 1882, 355, Cape San Lucas, Lower California.

1349. *Fundulus confluentus* Goode and Bean.
Streams of Florida.
Fundulus confluentus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 118, Lake Monroe, Fla.
1350. *Fundulus albolineatus* Gilbert.
Tennessee Basin in Alabama.
Fundulus albolineatus Gilbert, Bull. U. S. Fish Com., IX, 1889 (1891), 149, Spring Creek, Huntsville, Ala.
Subgenus *GALASACEUS* Fowler.
1351. *Fundulus similis* (Baird and Girard). *Sac-à-lait*.
Coast of the Gulf States.
Hydrargyra similis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 389, Indianola, Tex.
- Genus 444. *ZYGONECTES* Agassiz.
Zygonectes Agassiz, Amer. Jour. Sci. Arts, 1854, 135 (*Semotilus olivaceus* Raf. = *Semotilus notatus* Raf.).
1352. *Zygonectes notatus* (Rafinesque). *Top minnow*.
Michigan to Alabama, Mississippi, and Texas; generally abundant in ponds and canals.
Semotilus notatus Rafinesque, Ichth. Ohiensis, 1820, 86, tributaries of Ohio River in Kentucky; vaguely described but recognizable.
Pacilia olivacea Storer, Proc. Bost. Soc. Nat. Hist., II, 1845 (1848), 51, Florence, Ala.
Fundulus tenellus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 389, Prairie Mer Rouge, La., and Russellville, Ky.
Zygonectes lateralis Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 353, Mobile, Ala.
Zygonectes zonatus Agassiz, loc. cit., 353, St. Louis, Mo.
Zygonectes pulchellus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 113, Sugar Loaf Creek, Ark.
Fundulus aureus Cope, loc. cit., XVII, 1865, 78, Sugar Loaf Creek, Ark.
1353. *Zygonectes sciadicus* (Cope).
Middle Missouri River Basin; abundant in ponds and sluggish creeks in eastern Nebraska and southeastern South Dakota.
Fundulus sciadicus Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 78, Platte River, Nebr.
Haplochilus floripinnus Cope, Proc. Amer. Phil. Soc., XIV, 1874, 138, Platte River, at Denver, Colo.
Zygonectes lineatus Garman, Bull. Mus. Comp. Zool., VIII, 1881, 88, northeastern Wyoming.
Zygonectes macdonaldi Meek, Bull. U. S. Fish Com., IX, 1889, 122, pl. 42, fig. 1, Jones Creek, Dixon, Mo., Osage fork of Gasconade River at Mansfield, Mo.
Fundulus scartes Meek, loc. cit., XV, 1895, 347, St. Francis River, Big Bay, Ark.
1354. *Zygonectes chrysotus* (Holbrook).
Coastwise swamps, South Carolina to Florida; rather common.
Fundulus chrysotus Holbrook, in Gunther, Cat., VI, 1866, 317, Charleston, S. C.
Zygonectes henshalli Jordan, Proc. U. S. Nat. Mus., II, 1879, 237, San Sebastian River, Fla.
Gambusia arlingtonia Goode and Bean, loc. cit., 118, Arlington River, Fla.
1355. *Zygonectes cingulatus* (Cuvier and Valenciennes).
New Jersey to Florida, in coastwise swamps, not rare; Escambia River at Flomaton, Ala.
Fundulus cingulatus Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 147 (197), United States.
Zygonectes rubrifrons Jordan, Proc. U. S. Nat. Mus., II, 1879, 237, San Sebastian River, Fla.
Zygonectes auroguttatus Hay, loc. cit., VIII, 1885, 556, Westville, Fla.

1356. *Zygonectes dispar* Agassiz. *Top minnow*.
Northern Ohio to Illinois, south to Mississippi; Maumee, Wabash, Big Black and Pearl Rivers.
Zygonectes dispar Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 353, creeks opposite St. Louis; Beardstown, Ill.
1357. *Zygonectes nottii* Agassiz. *Star-headed minnow*; *Top minnow*.
Swamps and streams of Florida and neighboring States; not rare.
Zygonectes nottii Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 353, Mobile, Ala.
Zygonectes lineolatus Agassiz, loc. cit., 353, Augusta, Ga.
Zygonectes hieroglyphicus Agassiz, loc. cit., 353, Mobile, Ala.
Fundulus zonatus Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 146 (196), interior of South Carolina; not *Esox zonatus* Mitchell.
Zygonectes craticula Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 433, Elbow Creek, a tributary of Indian River, eastern Florida.
Zygonectes zonifer Jordan and Meek, loc. cit., VII, 1884, 482, Allapaha River, Nashville, Ga.
1358. *Zygonectes guttatus* Agassiz.
Alabama.
Zygonectes guttatus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 353, Mobile, Ala.
Zygonectes escambiae Bollman, Proc. U. S. Nat. Mus., IX, 1886, 463, Escambia River, Flomaton, Ala.
1359. *Zygonectes luciae* (Baird).
Atlantic coast from Long Island to Virginia; rare.
Hydrargyra luciae Baird, Ninth Smithson. Rept., 1854, 344, Beesleys Point, N. J.
1360. *Zygonectes pulvereus* Evermann.
Coast of Texas.
Zygonectes pulvereus Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 85, Dickinson Bayou; Buffalo Bayou at Houston; Oso Creek at Corpus Christi, Tex.
1361. *Zygonectes jenkinsi* Evermann.
Coast of Texas.
Zygonectes jenkinsi Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 86, Dickinson Bayou, arm of Galveston Bay, Tex.
1362. *Zygonectes limi* (Vaillant).
Fundulus limi Vaillant, Bull. Soc. Philom. (8), VI, 1894, 71, Lower California.
Fundulus meeki Evermann, Proc. Biol. Soc. Wash., XXI, 1908 (Jan. 23), 25, fig. 1, San Ignacio, central Lower California.
1363. *Zygonectes melapleurus* (Gosse). *Ticky-ticky*.
Streams of Jamaica.
Pæcilia melapleura Gosse, Naturalists' Sojourn in Jamaica, 1851, 84, Jamaica.
- Subgenus FONTICOLA Jordan and Evermann.
1364. *Zygonectes seminolis* (Girard).
Florida.
Fundulus seminolis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 60, Palatka, eastern Florida.
- Subgenus MENONA Jordan and Evermann.
1365. *Zygonectes diaphanus* (LeSueur).
Maine to North Carolina in river mouths; New York to northern Illinois in lakes.
Hydrargyra diaphana LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 130, Saratoga Lake, N. Y.
Hydrargyra multifaciata LeSueur, loc. cit., 131, Saratoga Lake, N. Y.
1366. *Zygonectes diaphanus menona* (Jordan and Copeland).
Streams and lakes of Ohio to Iowa, Minnesota, and South Dakota.
Fundulus menona Jordan and Copeland, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 68, Catfish River, Dane County, Wis., outlet of Lake Menona, Madison, Wis.

Subgenus GAMBUSINUS Jordan and Evermann.

1367. *Zygonectes rathbuni* (Jordan and Meek).
Eastern North Carolina, abundant in spring brooks, not descending to salt water.

Fundulus rathbuni Jordan and Meek, Proc. U. S. Nat. Mus., XI, 1888, 356, Reedy Fork, Allemanee Creek, Buffalo Creek, and other tributaries of the Cape Fear River, about Greensboro, N. C.

1368. *Zygonectes extensus* (Jordan and Gilbert).

Coast of Lower California.

Fundulus extensus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 355, Cape San Lucas, Lower California.

Genus 445. XENISMA Jordan.

Xenisma Jordan, Bull. Buffalo Soc. Nat. Hist., 1876, 142 (*X. stellifera* Jordan).

1369. *Xenisma catenatum* (Storer). *Studfish*.

Tennessee and Cumberland Rivers and clear streams of the Ozark Mountains.

Pacilia catenata Storer, Synopsis, 1846, 430, Tennessee River, Florence, Ala.

1370. *Xenisma stellifer* Jordan. *Studfish*.

Alabama River and tributaries.

Xenisma stellifera Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 322, Etowah and Oostanula Rivers, Rome, Ga.

Genus 446. PLANCTERUS Garman.

Plancterus Garman, Mem. Mus. Comp. Zool., XIX, 1895, 96 (*Fundulus kansæ* Garman).

1371. *Plancterus zebra* (Girard).

Kansas, western Iowa and South Dakota to Texas and New Mexico, in upper tributaries of Kansas and Red Rivers.

Hydrargyra zebra Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 61, tributaries of Rio Grande "between Fort Defiance and Fort Union, N. Mex.," name preoccupied in *Fundulus*.

Fundulus adina Jordan and Gilbert, Synopsis, 1883, 335, Rio Grande at Brownsville, Tex.

1372. *Plancterus kansæ* (Garman).

Kansas.

Fundulus kansæ Garman, Mem. Mus. Comp. Zool., XIX, 1895, 103, pl. 2, fig. 10, Kansas.

Fundulus zebrinus Jordan and Gilbert, Synopsis, 1882, 891; not of Girard.

Genus 447. OXYZYGONECTES Fowler.

Oxyzygonectes Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 425 (*Haplochilus dovii* Günther).

1373. *Oxyzygonectes balboæ* (Fowler).

Panama; only one example known.

Fundulus balboæ Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 423, fig. 2, Panama.

1374. *Oxyzygonectes dovii* (Günther).

Coast of Costa Rica.

Haplochilus dovii Günther, Cat., VI, 1866, 316, Punta Arenas, Costa Rica.

Genus 448. RIVULUS Poey.

Rivulus Poey, Memorias, II, 1861, 307, (*R. cylindraceus* Poey).

Cynodonichthys Meek, Field Col. Mus. Zool., V, 1904, 101 (*C. tenuis* Meek).

1375. *Rivulus cylindraceus* Poey.

Cuba.

Rivulus cylindraceus Poey, Memorias, II, 1861, 308, stream at Mardazo, near Habana, Cuba.

Rivulus marmoratus Poey, Anal. Soc. Española Hist. Nat., IX, 1880, 248, Cuba.

1376. *Rivulus heynei* Nichols.

Haiti.

Rivulus heynei Nichols, Bull. Amer. Mus. Nat. Hist., XXXII, 1914, 143, Haiti.1377. *Rivulus isthmensis* Garman.

Streams of Panama and Costa Rica.

Rivulus isthmensis Garman, Mem. Mus. Comp. Zool., XIX, 1895, 140, Río San José, Costa Rica.1378. *Rivulus brunneus* Meek and Hildebrand.

Panama.

Rivulus brunneus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 86, Toro Point, Canal Zone, Panama.1379. *Rivulus flabellicauda* Regan

Streams of Costa Rica.

Rivulus flabellicauda Regan, Ann. Mag. Nat. Hist., XIX, 1907, 64, Juan Vinas, Costa Rica.1380. *Rivulus godmanni* Regan.*Rivulus godmanni* Regan, Ann. Mag. Nat. Hist., 7th ser., XIX, 1907, 65, Guatemala.1381. *Rivulus tenuis* (Meek).*Cynodontichthys tenuis* Meek, Field Col. Mus., Zool., V, 1904, 101, El Hule, Oaxaca, Mexico.1382. *Rivulus harti* (Boulenger). *Rainfish; Small guabín.*

Mountain streams of Trinidad and Granada in rain-water pools.

Haplochilus harti Boulenger, Proc. Zool. Soc. Lond., I, 1906, 389, Pl. XXI, fig. 2, Trinidad.1383. *Rivulus chucunaque* Breder.

Tuyra Basin, Pacific slope of Panama.

Rivulus chucunaque Breder (C. M.), Amer. Mus. Novitates, No. CLXXX, July 6, 1925, fig. 6, Río Chucunaque, near Yavisa, Darien, Panama.*Rivulus chucunaque sucubti* Breder, loc. cit., 8, fig. 7, Río Sucubti, Darien, Panama.Genus 449. *ADINIA* Girard.*Adinia* Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 117 (*A. multifasciata* Girard).1384. *Adinia multifasciata* Girard.

Gulf Coast, west Florida to Texas; locally abundant in shallow lagoons.

Adinia multifasciata Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 118, Galveston; St. Joseph Island, Indianola, Tex.*Fundulus xenicus* Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 255, Laguna Grande, Pensacola, Fla.; substitute for *multifasciata*, preoccupied in *Fundulus* but not in *Adinia*.Genus 450. *FLORIDICHTHYS* Hubbs.*Floridichthys* Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 16 (*Cyprinodon carpio* Günther).1385. *Floridichthys carpio* (Günther).

Florida to Texas.

Cyprinodon carpio Günther, Cat., VI, 1866, 306, America.*Cyprinodon mydrus* Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 433, Pensacola, Fla.Genus 451. *JORDANELLA* Goode and Bean.*Jordanella* Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 117 (*Jordanella floridæ* Goode and Bean).1386. *Jordanella floridæ* Goode and Bean.

Streams from Florida to Yucatan.

Jordanella floridæ Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 117, Lake Monroe, Fla.Genus 452. *CYPRINODON* Lacépède.*Cyprinodon* Lacépède, Hist. Nat. Poiss., V, 1803, 486 (*C. variegatus* Lacépède).*Prinodon* Rafinesque, Anal. Nat., 1815, 88 (*Cyprinodon variegatus* Lacépède).*Encrotes* Gistel, Thierreich, IX, 1848 (*Cyprinodon variegatus* Lacépède).*Trifarcus* Poey, Memorias, II, 1861, 306 (*T. riverendi* Poey).

1387. *Cyprinodon variegatus* Lacépède. Killifish; Porsy minnow; Sheeps-head minnow; Variegated minnow; Mummichog.
Cape Cod to Rio Grande, in bays, rarely entering streams
Cyprinodon variegatus Lacépède, Hist. Nat. Poiss., V, 1803, 486, South Carolina.
Esoc ovinus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 441, New York.
Lebias ellipsoidea LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 6, Florida.
Lebias rhomboidalis Valenciennes, in Humboldt, Obs. Zool. Phil. Mag., II, 1810 (1811), 100, Lake Pontchartrain, La.
Cyprinodon gibbosus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 390, Indianola, Tex.
1388. *Cyprinodon riverendi* (Poey).
Cuba and Florida Keys; common at Key West in shallow shore waters.
Trifarcus riverendi Poey, Memorias, II, 1861, 306, Habana; scarcely distinct from *C. variegatus*.
1389. *Cyprinodon dearborni* Meek.
Curaçao, Dutch West Indies.
Cyprinodon dearborni Meek, Field Mus. Nat. Hist., Zool., VII, 1909, 208, Willemstad, Curaçao, Dutch West Indies.
1390. *Cyprinodon felicianus* (Poey).
Cuba.
Trifarcus felicianus Poey, Synopsis, 1867, 412, Habana, Cuba.
1391. *Cyprinodon latifasciatus* Garman.
Parras, Coahuila, Mexico.
Cyprinodon latifasciatus Garman, Bull. Mus. Comp. Zool., VIII, 1881, No. 3, 92, Parras, Coahuila, Mexico.
1392. *Cyprinodon bovinus* Baird and Girard.
Leon Spring in southwestern Texas.
Cyprinodon bovinus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 389, Leon Spring, Tex.
1393. *Cyprinodon martæ* Steindachner.
Santa Marta, coast of Colombia, South America.
Cyprinodon martæ Steindachner, Ichth. Beitr., IV, 1875, 61, Santa Marta, Rio Cauca.
1394. *Cyprinodon rubrofluvialis* Fowler.
Streams of Texas.
Cyprinodon bovinus rubrofluvialis Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 430, fig. 4, Rio Brazos, Seymour, Tex., and Red River, Tulip and Clarendon, Tex.
1395. *Cyprinodon elegans* Baird and Girard.
Tributaries of the Rio Grande.
Cyprinodon elegans Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 389, Comanche Spring, Tex.
Cyprinodon eximius Girard, loc. cit., XI, 1859, 158, Chihuahua River, Mexico.
1396. *Cyprinodon macularius* Baird and Girard.
Springs and streams of the desert, from southern Nevada to Sonora, locally abundant, often found in Artesian water and desert water holes in southwestern California.
Cyprinodon macularius Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 389, Rio San Pedro, Ariz.
Cyprinodon californiensis Girard, loc. cit., XI, 1859, 158, San Diego County, Calif., probably from Salt Springs in the desert (now Imperial County).
Lucania browni Jordan and Richardson, Proc. U. S. Nat. Mus., XXXIII, 1908, 319, hot spring in southeastern California.
1397. *Cyprinodon nevadensis* Eigenmann.
Death Valley.
Cyprinodon nevadensis Eigenmann, Proc. Calif. Acad. Sci., I, 1889, 270, Saratoga Springs, Death Valley, Inyo County, Calif.

1398. *Cyprinodon baileyi* Gilbert.

Pahranaagat Valley, Nev.

Cyprinodon macularius baileyi Gilbert, Death Valley Exped., 1893, 233, Pahranaagat Valley, Nev.

Family 114. EMPETRICHTHYIDÆ

Genus 453. EMPETRICHTHYS⁴⁴ Gilbert.*Empetrichthys* Gilbert, Death Valley Exped., Fishes, 1893, 233 (*E. merriami* Gilbert).1399. *Empetrichthys merriami* Gilbert.

Death Valley in southeastern California.

Empetrichthys merriami Gilbert, Death Valley Exped., in North American Fauna, No. 7, May 31, 1893, 234, Ash Meadows, Amargosa Desert, on boundary between California and Nevada.

Family 115. GOODEIDÆ

Genus 454. ZOOGONETICUS Meek.

Zoogoneticus Meek, Field Col. Mus., Zool., III, 1902, 91 (*Z. diazi* Meek).1400. *Zoogoneticus dugesii* (Bean).

Guanajuato, Mexico.

Fundulus dugesi Bean, Proc. U. S. Nat. Mus., X, 1887, 373, pl. 20, Guanajuato, Mexico.1401. *Zoogoneticus diazi* Meek.

Lakes of Michoacan, Mexico.

Zoogoneticus diazi Meek, Field Col. Mus., Zool., III, 1902, 93, figured in Field Col. Mus., Zool., V, 1904, 114, fig. 32, Lake Patzcuaro, Mexico.*Zoogoneticus miniatus* Meek, Field Mus. Nat. Hist., Zool., V, 1914, 115, fig. 33, Lake Zirahuen, Mexico.1402. *Zoogoneticus robustus* (Bean).

Rio Lerma, Guanajuato.

Fundulus robustus Bean, Proc. U. S. Nat. Mus., XV, 1892, 285, pl. 44, fig. 2, Guanajuato, Mexico.*Zoogoneticus maculatus* Regan, Ann. Mag. Nat. Hist. (7), XIII, 1904, 256, Rio Santiago (Lerma), Mexico.1403. *Zoogoneticus quitzeensis* (Bean)

Rio Lerma, Mexico.

Platypacilus quitzeensis Bean, Proc. U. S. Nat. Mus., XXI, 1899 (Nov. 21, 1898), 540, fig. 1, Lake Quitzeo, Lerma Basin.

Genus 455. GIRARDINICHTHYS Bleeker.

Girardinichthys Bleeker, Act. Soc. Sci. Indo-Nederl., VII, 1860, 481 (*G. innominatus* Bleeker, after a "*Lucania* sp." of Girard).*Limnurgus* Günther, Cat., VI, 1866, 309 (*L. variegatus* Günther = *Girardinichthys* Bleeker).1404. *Girardinichthys innominatus* Bleeker.

Valle of Mexico.

Lucania sp., Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 118, City of Mexico.*Girardinichthys innominatus* Bleeker, Act. Soc. Sci. Indo-Nederl., VII, 1860, 481, City of Mexico.*Limnurgus variegatus* Günther, Cat., VI, 1866, 309, City of Mexico.*Characodon geddesi* Regan, Ann. Mag. Nat. Hist., (7), XIII, 1904, 257, Xochomilco, Morelos, Mexico.

Genus 456. CHARACODON Günther.

Characodon Günther, Cat., VI, 1866, 308 (*C. lateralis* Günther).⁴⁴ This genus apparently a desert derivative of the extinct genus, *Parafundulus*, has probably no affinity with *Orestias* of Peru.

1405. *Characodon variatus* Bean.
Tributaries of Rio Lerma about Guanajuato, Mexico; locally common.
Characodon variatus Bean, Proc. U. S. Nat. Mus., X, 1887, 370, pl. 20, fig. 1, Guanajuato, Mexico.
Characodon ferrugineus Bean, loc. cit., 372, pl. 20, figs. 3 and 4, Guanajuato, Mexico.
Characodon eiseni Rutter, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 266, branch of the Rio Grande de Santiago at Tepic, Mexico.
1406. *Characodon lateralis* Günther.
Streams of Central America.
Characodon lateralis Günther, Cat., VI, 1866, 308, Central America.
1407. *Characodon garmani* Jordan and Evermann.
Tributaries of Rio Grande in Coahuila, Mexico.
Characodon garmani Jordan and Evermann, Fish. North and Mid. Amer., III, 1898, 2831, Parras, Coahuila, Mexico.
1408. *Characodon fuscoides* Jordan and Gilbert.
Cape San Lucas southward to Colima, Mexico.
Characodon fuscoides Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 354, Cape San Lucas, Lower California.
- Genus 457. *LERMICHTHYS* Hubbs.
Lermichthys Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 18 (*Characodon multiradiatus* Meek).
1409. *Lermichthys multiradiatus* (Meek).
Rio Lerma Basin, Mexico.
Characodon multiradiatus Meek, Field Col. Mus., Zool., V, 1904, 119, Rio Lerma, Mexico.
Girardinichthys limnurgus Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 502, Lake Lerma, Mexico.
- Genus 458. *BALSADICHTHYS* Hubbs.
Balsadichthys Hubbs, Univ. Mich. Mus. Zool., Misc. Pub. No. 16, July 9, 1926, 19 (*Goodea whitei* Meek).
1410. *Balsadichthys whitei* (Meek).
Rio de las Balsas Basin, Mexico.
Goodea whitei Meek, Field Col. Mus., Zool., V, 1904, 137, fig. 40, Rio Balsas, Yantepec, Cuautla, Mexico.
- Genus 459. *SKIFFIA* Meek.
Skiffia Meek, Field Col. Mus., Zool., III, 1902, 102 (*S. lermæ* Meek).
1411. *Skiffia variegata* Meek.
Lake Zirahuen, Mexico.
Skiffia variegata Meek, Field Col. Mus., Zool., V, 1904, 143, Lake Zirahuen, Mexico.
1412. *Skiffia lermæ* Meek.
Skiffia lermæ Meek, Field Col. Mus., Zool., III, 1902, 102, Lake Patzcuaro (Rio Lerma), Mexico.
1413. *Skiffia bilineata* (Bean).
Rio Lerma, Guanajuato, Mexico.
Characodon bilineatus Bean, Proc. U. S. Nat. Mus., X, 1887, 371, Guanajuato, Rio Lerma, Mexico.
1414. *Skiffia multipunctata* (Pellegrin).
Rio Lerma system; Lake Chapala, Mexico.
Xenendum multipunctatum Pellegrin, Bull. Mus. Hist. Nat., Paris, 1901, 205, Jalisco, Mexico.
- Genus 460. *GOODEA* Jordan.
Goodea Jordan, Proc. U. S. Nat. Mus., II, 1879, 299 (*G. atripinnis* Jordan).
Xenendum Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 128, fig. 9 (1906) (*X. zaleicone* = *Characodon luitpoldi* Steindachner).
1415. *Goodea atripinnis* Jordan.
Tributaries of Rio Lerma, Pacific slope, Mexico.
Goodea atripinnis Jordan, Proc. U. S. Nat. Mus., II, 1879, 299, Leon, Guanajuato, Mexico.

1416. *Goodea calientis* (Jordan and Snyder).
Streams of central Mexico.
Xenododon caliente Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 127, fig. 8, Aguas Calientes, Mexico.
1417. *Goodea luitpoldi* (Steindachner).
Basin of Rio Lerma, Mexico.
Characodon luitpoldi Steindachner, Einige Fischarten Mexicos, 1895, 12, pl. 3, figs. 3-36, Lago de Patzeuaro, Michoacan, Mexico.
Xenododon zaliscone Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1901), 128, fig. 9, Laguna de Chapala, near Ocotlan, Jalisco, Mexico.
1418. *Goodea toweri* Meek.
Rio Panuco, San Luis Potosi, Mexico.
Goodea toweri Meek, Field Col. Mus., Zool., V, 1904, 138, fig. 41, Rio Verde, San Luis Potosi, Mexico.
1419. *Goodea captiva* Hubbs.
Panuco Basin, Mexico.
Goodea captiva Hubbs, Occ. Papers, Mus. Zool., Univ. Mich., No. 148, 1924, 4; upper tributary of the Rio Panuco at Jesus Maria, Mexico.

Family 116. PÆCILIIDÆ. Top minnows

- Genus 461. *BELONESOX* Kner.
Belonesox Kner, Sitzungsber., Kais. Akad. Wiss. Wien, 1860, XL, 419 (*B. belizanus* Kner).
1420. *Belonesox belizanus* Kner.
Southern Mexico, Honduras, and Guatemala.
Belonesox belizanus Kner, Sitzungsber., Kais. Akad. Wiss. Wien, XL, 1860, 419, with figure, Belize; largest of the Pæciliidæ.
- Genus 462. *CHAPALICHTHYS* Meek.
Chapalichthys Meek, Field Col. Mus., Zool., III, 1902, 97 (*Characodon encaustus* Jordan and Snyder).
1421. *Chapalichthys encaustus* (Jordan and Snyder).
Lake Chapala, Mexico.
Characodon encaustus Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1901), 126, Laguna de Chapala, near Ocotlan, Jalisco, Mexico.
- Genus 463. *HETEROPHALLUS* Regan.
Heterophallus Regan, Ann. Mag. Nat. Hist., (8), XV, 1914, 66 (*H. rachovii* Regan).
1422. *Heterophallus rachovii* Regan.
Streams of eastern Mexico.
Heterophallus rachovii Regan, Ann. Mag. Nat. Hist., (8), XV, 1914, 66, Fig. A, Vera Cruz, Mexico.
- Genus 464. *GAMBUSIA* Poey. *Guagacones*.
Gambusia Poey, Memorias, I, 1854, 382 (*G. punctata* Poey).
Hemixiphophorus Bleeker, Nat. Tijdschr. Neder.-Indie, XX, 1854, 440; type not named.
Paragambusia Meek, Field Col. Mus., Zool., V, 1904, 133 (*Gambusia nicaraguensis* Gunther).
Heterophallina Hubbs (subgenus), Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926 (*Gambusia regani* Hubbs).
Arthrophallus Hubbs (subgenus), loc. cit., 38 (*Heterandria patruelsi* Baird and Girard).
Schizophallus Hubbs (subgenus), loc. cit., 40 (*Gambusia holbrooki* Girard).

- 1423. *Gambusia patruelis* (Baird and Girard).** *Top minnow; Mosquitofish.*
Ditches, marshes, and lagoons, South Atlantic and Gulf States, lowland streams of the southwest; lower Mississippi from St. Louis to Louisiana and the Rio Grande. Introduced from Galveston into Hawaii, California, Formosa, Luzon, etc., a very effective mosquito killer.
Heterandria patruelis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 390, Rio Sabinal, Rio Nueces, and Elm Creek, Tex.
Heterandria affinis Baird and Girard, loc. cit., 390, Rio Medina and Rio Salado, Tex.
? *Gambusia speciosa* Girard, loc. cit., 1859, 121, Rio San Diego, Nuevo Leon, Mexico.
? *Gambusia gracilis* Girard, loc. cit., 1859, 121, Matamoros, Mexico; name preoccupied.
Gambusia humilis Gunther, Cat., VI, 1866, 334; substitute for *gracilis*.
Haplochilus melanops Cope, Proc. Amer. Phil. Soc., 1870, 457, Neuse River, N. C.
Zygonectes (Micristius) brachypterus Cope, Bull., U. S. Nat. Mus., XX, 1880, 34, Trinity River, Fort Worth, Tex.
Zygonectes inurus Jordan, Proc., U. S. Nat. Mus., 1882, 143, Cache Creek, southern Illinois.
Gambusia modesta Ahl, Blatter für Aquarium- und Terrariakunde, XXXVI, 1923, 220.
Gambusia myersi Ahl, loc. cit., 1925, 36, with figure (substitute for *modesta* preoccupied).
- 1424. *Gambusia holbrooki* (Girard).** *Holbrook's top minnow.*
New Jersey to Florida and Alabama.
Heterandria holbrooki Agassiz, in Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 61, Charleston, S. C.
Haplochilus melanops Cope, Proc. Amer. Phil. Soc., XI, 1870, 457, Neuse River, N. C.
Zygonectes atrilatus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 84, Neuse River, Goldsboro, N. C.
- 1425. *Gambusia nobilis* (Baird and Girard).**
Heterandria nobilis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 390, Leona and Comanche Spring, Rio Grande, Tex.
- 1426. *Gambusia senilis* Girard.**
Rio Grande Basin.
Gambusia senilis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 122, Chihuahua, Mexico.
- Subgenus **HEMIXIPHOPHORUS** Bleeker.
- 1427. *Gambusia gracilis* (Heckel).**
Lowland streams of eastern Mexico.
Xiphophorus gracilis Heckel, Sitzungsber. Kais. Akad. Wiss. Wien, I, 1848, 300, taf. 9, figs. 3 and 4, Orizaba, Mexico.
Gambusia heckeli Bleeker, Fam. Cypr., 1860, 485.
- Subgenus **GAMBUSIA** Poey.
- 1428. *Gambusia punctata* Poey.** *Guajacon.*
Cuba.
Gambusia punctata Poey, Memorias, I, 1854, 384, Rio Almendares, Cuba.
Gambusia picturata Poey, Repertorio, 1868, 410, San Diego de los Baños, Cuba.
Gambusia punctata punctulata Garman, Mem. Mus. Comp. Zool., XIX, 1895, 87, Remedios, Cuba.
- 1429. *Gambusia puncticulata* Poey.**
Streams of Cuba.
Gambusia puncticulata Poey, Memorias, I, 1854, 386, Cuba.
- 1430. *Gambusia melapleura* (Gosse).** *Ticky-ticky.*
Streams of Jamaica; locally common.
Pacilia melapleura Gosse, Naturalist's Sojourn in Jamaica, 1851, 84, Jamaica.

1431. *Gambusia nicaraguensis* Günther.
Southern Mexico and Central America.
Gambusia nicaraguensis Gunther, Cat., VI, 1866, 366, Lake of Nicaragua.
Gambusia dovri Regan, Proc. Zool. Soc. Lond., 1913, 986, Lake Nicaragua.
Gambusia yucalana Regan, Ann. Mag. Nat. Hist., 8th ser., XIV, 1914, 67, Progreso, Yucatan.
1432. *Gambusia mcnieli* Fowler.
Panama
Gambusia mcnieli Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 433, Panama
1433. *Gambusia wrayi* Regan.
Jamaica.
Gambusia wrayi Regan, Proc. Zool. Soc. Lond., 1913, 988, Pl. XCIX, figs. 3, 4 and text fig. 168B, Jamaica.
1434. *Gambusia caymanensis* Regan.
West Indies.
Gambusia caymanensis Regan, Proc. Zool. Soc. Lond., 1913, 990, Grand Cayman, West Indies.
1435. *Gambusia gracilior* Regan.
Jamaica.
Gambusia gracilior Regan, Proc. Zool. Soc. Lond., 1913, 989, Pl. XCIX, figs. 5, 6, and text fig. 168C, Jamaica.
1436. *Gambusia dominicensis* Regan.
Haiti.
Gambusia dominicensis Regan, Proc. Zool. Soc. Lond., 1913, 989, Pl. XCIX, fig. 7 and text fig. 169C, San Domingo.
1437. *Gambusia oligosticta* Regan.
Jamaica.
Gambusia oligosticta Regan, Proc. Zool. Soc. Lond., 1913, 988, Pl. XCIX, figs. 1, 2, and text fig. 169B, Jamaica.
1438. *Gambusia manni* Hubbs.
Bahamas.
Gambusia manni Hubbs, Copeia, 1927, 62, New Providence, one of the Bahamas.
- Subgenus **HETEROPHALLINA** Hubbs.
1439. *Gambusia regani* Hubbs.
Rio Panuco, northwestern Mexico.
Gambusia regani Hubbs, Univ. Mich. Mus. Zool., Misc. Pub. No. 16, July 9, 1926, 28, Forlon, Tamaulipas, Mexico, in the River Forlon.
1440. *Gambusia vittata* Hubbs.
Santa Engracia, Tamaulipas, Mexico.
Gambusia vittata Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 26, Rio Forlon, at Forlon, Tamaulipas, Mexico.
1441. *Gambusia panuco* Hubbs.
Rio Panuco, Mexico.
Gambusia panuco Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 30, Rio Valles, at Valles, San Luis Potosi, Mexico.
- Genus 465. **BRACHYRHAPHIS** Regan.
Brachyrhaphis Regan, Proc. Zool. Soc. Lond., 1913, 997 (*Gambusia rhabdophora* Regan).
1442. *Brachyrhaphis terrabensis* (Regan).
Pacific streams of Costa Rica.
Gambusia terrabensis Regan, Ann. Mag. Nat. Hist., (7), XIX, 1907, 260, Rio Grande de Terraba, Costa Rica.
1443. *Brachyrhaphis rhabdophora* (Regan).
Costa Rica.
Gambusia rhabdophora Regan, Ann. Mag. Nat. Hist., (8), II, 1908, 457 Volcano of Tenorio and Rio Grande de Terraba, Costa Rica.

1444. *Brachyrhaphis olomina* (Meek).
Valley of the Rio Grande de Tarcoles, Costa Rica.
Priapichthys olomina Meek, Field Mus. Nat. Hist., Zool., X, 1912, 114,
Las Cañas, Alajuela, Costa Rica.
1445. *Brachyrhaphis episcopi* (Steindachner).
Isthmus of Panama at Obispo station, and streams of the Pacific slope
of Panama.
Gambusia episcopi Steindachner, Ichth. Beitr., VI, 1878, 9, Obispo near
Panama.
Gambusia latipunctata Meek and Hildebrand, Field Mus. Nat. Hist.,
Zool., X, 1913, 87, Arrijan, Panama.
1446. *Brachyrhaphis cascajalensis* (Meek and Hildebrand).
Streams of Atlantic slope of Panama.
Gambusia cascajalensis Meek and Hildebrand, Field Mus. Nat. Hist.,
Zool., X, 1913, 86, Rio Cascajal, Panama.
1447. *Brachyrhaphis parismina* (Meek).
Costa Rica.
Gambusia parismina Meek, Field Mus. Nat. Hist., Zool., X, 1912, 71,
Costa Rica.
- Genus 466. **TRIGONOPHALLUS** Hubbs.
Trigonophallus Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16,
July 9, 1926, 48 (*T. punctifer* Hubbs).
1448. *Trigonophallus punctifer* Hubbs.
Streams of Pacific coast of Panama.
Trigonophallus punctifer Hubbs, Univ. Mich. Mus. Zool. Misc. Pub.
No. 16, July 9, 1926, 49, Quibari Creek, Rio Secamola at Kakintun.
- Genus 467. **PRIAPICHTHYS** Regan.
Priapichthys Regan, Proc. Zool. Soc. Lond., 1913, 991 (*Gambusia*
annectans Regan).
1449. *Priapichthys annectens* (Regan).
Streams of Atlantic slope of Costa Rica.
Gambusia annectens Regan, Ann. Mag. Nat. Hist., (7), XIX, 1907,
250, Costa Rica.
1450. *Priapichthys annectens hesperis* Hubbs.
Streams of the Pacific slope of Costa Rica.
Priapichthys annectens hesperis Hubbs, Univ. Mich. Mus. Zool. Misc.
Pub. No. 13, Jan. 18, 1924, 22, Rio Maria Aguilar, Rio Grande de
Tancoles, and their mountain tributaries.
1451. *Priapichthys huberi* Fowler.
Nicaragua.
Priapichthys huberi Fowler, Proc. Acad. Nat. Sci. Phila., LXXV, 1923,
27, Marceligo Creek, tributary of the Tunky River, at Miranda,
Nicaragua.
- Genus 468. **PANAMICHTHYS** Hubbs.
Panamichthys Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 13,
Jan. 18, 1924, 8 (*Priapichthys panamensis* Meek and Hildebrand).
1452. *Panamichthys panamensis* (Meek and Hildebrand).
Priapichthys panamensis Meek and Hildebrand, Field Mus. Nat. Hist.,
Zool., X, 1916, 322, Panama.
- Genus 469. **PSEUDOXIPHOPHORUS** Bleeker.
Pseudoxiphophorus Bleeker, Ichth. Indice Prodr. Cypr., in Acata Societ.
Scient. Indo-Neer., VII, 1859, 483 (*Xiphophorus bimaculatus* Heckel).
Poecilodes Steindachner, Sitzungsber. Kais. Akad. Wiss. Wien, XLVIII,
1863, 176 (*P. bimaculatus* Steindachner).
1453. *Pseudoxiphophorus bimaculatus* (Heckel).
Central Mexico; Rio Blanco, Orizaba.
Xiphophorus bimaculatus Heckel, Sitzungsber. Kais. Akad. Wiss. Wien,
I, 1848, pt. 3, 169, taf. 6, figs 1 and 2, Mexico.
Pseudoxiphophorus reticulatus Troschel, in Miller's Reisen Vereinigten
Staaten, Canada, Mexico, III, 1865, 638, no locality.

1454. *Pseudoxiphophorus bimaculatus tæniatus* Regan.
Streams of the Atlantic slope of southeastern Mexico.
Pseudoxiphophorus bimaculatus tæniatus Regan, Ann. Mag. Nat. Hist., (7), XVI, 1905, 303, Rio Touto and San Domingo de Guzman.
1455. *Pseudoxiphophorus bimaculatus jonesi* (Günther).
Streams of southeastern Mexico.
Mollenesia jonesi Günther, Ann. Mag. Nat. Hist., (4), XIV, 1874, 371,
Lake Alehuaca, Huamanita, Mexico.
Pseudoxiphophorus pauciradatus Regan, Ann. Mag. Nat. Hist., (7),
X, 1904, 14, 296, Orizaba, Mexico.
- Genus 470. *HETERANDRIA* Agassiz.
Heterandria Agassiz, Amer. Jour. Sci. Arts, XVI, 1853, 135 (*H. formosus*
Agassiz MS.; *H. holbrooki* Agassiz MS.).
1456. *Heterandria formosa* Agassiz.
Streams of South Carolina to Florida; one of the smallest of fishes.
Heterandria formosa Agassiz, in Girard, Proc. Acad. Nat. Sci. Phila.,
XI, 1859, 62, Charleston, S. C.; Palatka, Fla
- Genus 471. *PRIAPELLA* Regan.
Priapella Regan, Proc. Zool. Soc. Lond., 1913, 992 (*Gambusia bonita*
Meek).
1457. *Priapella bonita* (Meek).
Streams of western Mexico; Rio Papaloapam, Motzorongo.
Gambusia bonita Meek, Field Col. Mus., Zool., V, 1904, 132, fig. 39,
104, Refugio, Vera Cruz, Mexico.
- Genus 472. *ALLOGAMBUSIA* Hubbs.
Allogambusia Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 13, Jan.
18, 1924, 8 (*Gambusia tridentiger* Garman).
1458. *Allogambusia cana* (Meek and Hildebrand).
Streams of the Pacific slope of Panama.
Gambusia cana Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X,
1913, 87, Rio Satiganti, Cana, Panama.
1459. *Allogambusia tridentiger* (Garman).
Rivers of Panama.
Gambusia tridentiger Garman, Mem. Mus. Comp. Zool., XIX, 1895, 89
(Pl. IV, fig. 10, teeth), rivers of Panama.
- Genus 473. *TOXUS* Eigenmann.
Toxus Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903), 226 (*T.*
riddlei Eigenmann).
1460. *Toxus riddlei* Eigenmann.
Streams of western Cuba.
Toxus riddlei Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903),
226, San Cristobal, Cuba.
1461. *Toxus cubensis* (Eigenmann).
Streams of western Cuba.
Heterandria cubensis Eigenmann, Bull. U. S. Fish Com., XXII, 1902
(1903), 227, Los Palacios, Cuba.
- Genus 474. *GLARIDICHTHYS* Garman.
Glaridodon Garman, Mem. Mus. Comp. Zool., XIX, 1895, 40 (*latidens*);
name preoccupied.
Glaridichthys Garman, Amer. Nat., XXX, 1896, 232 (same type).
1462. *Glaridichthys uninotatus* (Poey).
Cuba.
Girardinus uninotatus Poey, Memorias, II, 1861, 309, Cuba.
1463. *Glaridichthys torralbasi* Eigenmann.
Streams of Pinar del Rio, Cuba.
Glaridichthys torralbasi Eigenmann, Bull. U. S. Fish Com., XXII, 1902
(1903), 225, fig. 4, Pinar del Rio, Cuba.
1464. *Glaridichthys falcatus* Eigenmann.
Streams of western Cuba.
Glaridichthys falcatus Eigenmann, Bull. U. S. Fish Com., XXII, 1902
(1903), 224, San Cristobal, Palacios, Pinar del Rio, Cuba.

Genus 475. GIRARDINUS Poey.*Girardinus* Poey, Memorias, I, 1854, 383 (*G. metallicus* Poey).**1465. *Girardinus metallicus* Poey.**

Streams of Cuba.

Girardinus metallicus Poey, Memorias, I, 1854, 387, Cuba.*Girardinus denticulatus* Garman, Mem. Mus. Comp. Zool., XIX, 1895, 47, Remedios, Cuba.*Girardinus creolus* Garman, loc. cit., 47, Cuba.*Girardinus garmani* Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903), 296, Pinar del Rio, Palacios, Cuba.**Genus 476. DARIENICHTHYS Hubbs.***Darienichthys* Hubbs, Univ. Mich. Mus. Zool., Misc. Pub. No. 13, Jan. 18, 1924, 8 (*Gambusia darienensis* Meek and Hildebrand).**1466. *Darienichthys darienensis* (Meek and Hildebrand).**

Streams of Pacific slope, Panama.

Gambusia darienensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 116, 88, Rio Capeti (Tuyra River Basin), Panama.**Genus 477. PÆCILIOPSIS Regan.***Pæciliopsis* Regan, Proc. Zool. Soc. Lond., 1913, 996 (*Pæcilia presidionis* Jordan and Culver).*Leptorhaphis* Regan, loc. cit., 980, 998 (*Gambusia infans* Woolman).**1467. *Pæciliopsis occidentalis* (Baird and Girard).**

Gila Basin, Ariz., and basins of Yaqui and Sonora Rivers, Sonora.

Heterandria occidentalis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 390, Rio Santa Cruz, near Tucson, Ariz.*Girardinus sonoriensis* Girard, loc. cit., XI, 1859, 120, San Bernardino Creek, a tributary of the Rio Yaqui, Sonora.**1468. *Pæciliopsis infans* (Woolman).**

Rio Lerma, Salamanca, Mexico.

Gambusia infans Woolman, Bull. U. S. Fish Com., XIV, 1894, 62, Rio Lerma, Salamanca, Mexico.**1469. *Pæciliopsis balsas* Hubbs.**

Balsas Basin, Mexico.

Pæciliopsis balsas Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 66, Rio Balsas, Guerrero, Mexico.**1470. *Pæciliopsis latidens* (Garman).**

Chihuahua.

Glaridodon latidens Garman, Mem. Mus. Comp. Zool., XIX, 1895, 40, Chihuahua, Mexico.**1471. *Pæciliopsis fasciata* (Meek).**

Atlantic streams of southern Mexico.

Gambusia fasciata Meek, Field Col. Mus., Zool., V, 1904, 129, fig. 37, San Geronimo, Oaxaca, Mexico.**1472. *Pæciliopsis turrubarensis* (Meek).**

Costa Rica.

Gambusia turrubarensis Meek, Field Mus. Nat. Hist., Zool., X, 1912, 71, Rio Turrubara, Costa Rica.*Heterandria colombianus* Eigenmann and Henn, Indiana Univ. Stud., No. 8, 1912, 27, brackish water, mouth of Rio Dagua, Colombia.*Priapichthys fosteri* Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925, 260, figs. 14 and 15, Rio Lempa, San Marcos, El Salvador, Central America.**1473. *Pæciliopsis presidionis* (Jordan and Culver).**

Rio Presidio, Sinaloa, Mexico.

Pæcilia presidionis Jordan and Culver, in Jordan, Fishes of Sinaloa, Proc., Calif. Acad. Sci., V, 1895, 413, pl. 29, Rio Presidio, Sinaloa, Mexico.**Genus 478. PÆCILISTES Hubbs.***Pæcilistes* Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 68 (*Heterandria lutzi* Meek).

1474. *Pæcilistes pleurospilus* (Günther).

Streams of Guatemala.

Gerrardinus pleurospilus Günther, Cat., VI, 1866, 353, Lake Dueñas, Guatemala.*Heterandria lutzi* Meek, Field Col. Mus., Zool., V, 1904, 148, fig. 47, Orizaba, Mexico.*Præpichthys letona* Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925, 258, Rio San Miguel, San Salvador.Genus 479. *AULOPHALLUS* Hubbs.*Aulophallus* Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 69 (*Pæcilia elongata* Gunther).1475. *Aulophallus elongatus* (Gunther).Panama, a large marine species quite unlike *Pæcilia* in habits.*Pæcilia elongata* Gunther, Cat., VI, 1866, 342, Panama.Genus 480. *PHALLICHTHYS* Hubbs.*Phallichthys* Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. 13, Jan. 18, 1924, 10, (*Pæciliopsis isthmensis* Regan).1476. *Phallichthys amates* (Miller).

Motagua Basin, Guatemala.

Pæcilia amates Miller, Bull. Amer. Mus. Nat. Hist., XXIII, 1907, 108, fig. 1, pond and its outlet, Los Amates, an affluent of the Motagua River, Guatemala.1477. *Phallichthys pittieri* (Meek).

Costa Rica.

Pæcilia pittieri Meek, Field Mus. Nat. Hist., Zool., X, 1912, 71, La Junta, Costa Rica.*Pæciliopsis isthmensis* Regan, Proc., Zool. Soc. Lond., 1913, 997, Pl. C, figs. 3, 4, and text fig. 171B, Colon, Panama.Genus 481. *XENOPHALLUS* Hubbs.*Xenophallus* Hubbs, Mus. Zool. Univ. Mich. Misc. Pub. 13, Jan. 18, 1924, 10 (*Gambusia umbratilis* Meek.)1478. *Xenophallus umbratilis* (Meek).

Streams of Costa Rica.

Gambusia umbratilis Meek, Field Mus. Nat. Hist., Zool., X, 1912, 70, Costa Rica.Genus 482. *PÆCILIA* Schneider.*Pæcilia* Schneider, ed. Bloch, Syst. Ichth. 1801, 452 (*P. vivipara* Bloch and Schneider); not *Pæcilus* of genus of beetles.*Alazon* Gistel, Thierreich, X, 1848 (*Pæcilia vivipara* Schneider); substitute for *Pæcilia*, regarded as preoccupied.*Acropæcilia* Hilgendorf, Sitzungsab. Naturf. Freunde, 1889, 52 (*Pæcilia tridens* Hilgendorf).*Neopæcilia* Hubbs, Univ. Mich. Mus. Zool., Misc. Pub. 13, 1924 (*N. holacanthus* Hubbs).1479. *Pæcilia vivipara* Bloch and Schneider.

Rivers of Brazil; Guiana; Martinique; Porto Rico.

Pæcilia vivipara Bloch and Schneider, Syst. Ichth., 1801, 452, pl. 86, fig. 2, Surinam.*Pæcilia surinamensis* Valenciennes, Humboldt, Observ. Zool., II, 1817, 158, pl. 51, fig. 1, Surinam.*Pæcilia schneideri* Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 102 (135), Surinam.*Neopæcilia holacanthus* Hubbs, Univ. Mich. Mus. Zool., Misc. Pub. No. 13, Jan. 18, 1924, 11, based on material recorded as *Pæcilia vivipara* Evermann and Marsh, Bull., U. S. Fish Com., XX, 1900, first part (1902), 97, from Porto Rico (Ponce, Fijaro, Arroyo, and Hucare).1480. *Pæcilia tridens* Hilgendorf.

Streams of Haiti.

Pæcilia (*Acropæcilia*) *tridens* Hilgendorf, Sitzungsab. Naturf. Freunde, 1889, 52, Port au Prince.

Genus 483. LIMIA Poey.

Limia Poey, Memorias, I, 1854, 388 (*L. cubensis* Poey=*Pæcilia vittata* Guichenot).

1481. *Limia heterandria* Regan.

Santo Domingo.

Limia heterandria Regan, Proc. Zool. Soc. Lond., 1913, 1017, Pl. CI, "Venezuela"; locality probably wrong.

Heterandria zonata Nichols, Bull. Amer. Mus. Nat. Hist., XXXIV, 1915, 603, fig. 3, Santo Domingo.

1482. *Limia vittata* (Guichenot). *Faguto; Guajica.*

Streams of Cuba.

Pæcilia vittata Guichenot, in Ramon de la Sagra, Hist. Nat. Cuba, Poiss., 1853, 146, pl. 5, fig. 1, Cuba.

Limia cubensis Poey, Memorias, I, 1854, 388, Habana.

Limia pavonina Poey, Enumeratio, 1876, 142, Habana.

1483. *Limia perugiæ* (Evermann and Clark).

Santo Domingo.

Platyæcilus perugiæ Evermann and Clark, Proc. U. S. Nat. Mus., XXX, 1906, 851, Santo Domingo.

1484. *Limia caudofasciata* Regan.

Jamaica.

Limia caudofasciata Regan, Proc., Zool. Soc. Lond., 1913, 1017, Jamaica.

1485. *Limia nigrofasciata* Regan.

Limia nigrofasciata Regan, Proc., Zool. Soc. Lond., 1913, 1015, Pl. CI, figs. 1, 2, 1913, Haiti.

Limia arnoldi Regan, loc. cit., 1016, Miragoâne, Haiti; young of *nigrofasciata*.

1486. *Limia ornata* Regan.

Limia ornata Regan, Proc. Zool. Soc. Lond., 1913, 1016, Haiti.

1487. *Limia dominicensis* (Cuvier and Valenciennes).

Santo Domingo and Barbados.

Pæcilia dominicensis Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 99 (131), pl. 525, Santo Domingo.

Pæcilia melanogaster Gunther, Cat., VI, 1866, 345, locality unknown.

Platyæcilus dominicensis Evermann and Clark, Proc. U. S. Nat. Mus., XXX, 1906, 852, fig. 2, streams in San Francisco Mountains, Santo Domingo.

1488. *Limia melanonotata* Nichols and Myers.

Santo Domingo.

Limia melanonotata Nichols and Myers, Amer. Mus. Novitates, No. 79, 1923, 1, Hispaniola.

1489. *Limia versicolor* (Günther).

Rivers of Haiti.

Garardinus versicolor Günther, Cat., VI, 1866, 352, Santo Domingo.

Genus 484. LEBISTES De Filippi

Lebistes de Filippi, Arch. Zool. Anat. Fisiol., I, 1862, 69 (*L. pæciloides* de Filippi).

Acanthophaelus Eigenmann, Proc. U. S. Nat. Mus., XXXII, 1907, 426 (*Pæcilia reticulatus* Peters).

1490. *Lebistes reticulatus* (Peters). *Milmons; Bellyfish; Guppy and rainbow-fish of aquarists.*

Pools and streams of Barbados, Trinidad, and Venezuela.

Pæcilia reticulata Peters, Monatsber. Akad. Wiss. Berlin, 1859, 412, Caracas, Venezuela.

Lebistes pæciloides de Filippi, Arch. Zool. Anat. Fisiol., I, 1861, 69, Pl. IV, fig. 6, Barbados.

Garardinus guppia Günther, Cat., VI, 1866, 353, Trinidad; Venezuela.

Acanthophaelus melanzonus Eigenmann, Ann. Carnegie Mus., VI, 1909, 51, Georgetown Trenches, British Guiana.

Genus 485. MOLLIESIA LeSueur.

Molliesia LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 2 (*Molliesia latipinna* LeSueur).

1491. *Mollienesia latipinna* LeSueur. *Mudfish*.
 South Carolina to northern Mexico, in lowland swamps and streams.
Mollienesia latipinna LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 3,
 New Orleans.
Pæcilia multilineata LeSueur, loc. cit., 4, New Orleans.
Pæcilia lineolata Girard, U. S. Mex. Bound. Surv., Ichth., 1859, 70,
 pl. 35, figs. 9 to 11, Rio Grande near Brownsville, Tex.
Lima pæciloides Girard, loc. cit., 70, pl. 38, figs. 8 to 14, Indianola, Tex.
Lima matamorenensis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859,
 117, Matamoras.
1492. *Mollienesia gracilis* Regan.
 Lake Peten, Guatemala.
Mollienesia petenensis Gunther, Cat., VI, 1866, 348, Lake Peten; not
Pæcilia petenensis Günther, also a *Mollienesia*.
Mollienesia gracilis Regan, Proc. Zool. Soc. Lond., 1913, 1012; substitute
 name.
1493. *Mollienesia petenensis* (Gunther).
 Lake Peten, Guatemala.
Pæcilia petenensis Günther, Cat., VI, 1866, 342, Lake Peten, Guatemala;
 not *Mollienesia petenensis* Gunther, loc. cit., 348.
1494. *Mollienesia formosa* (Girard).
 Mexico; Lake Alcohuaca, Huamantla, Mexico; Palo Alto.
Lima formosa Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 115, Palo
 Alto, Mexico.
1495. *Mollienesia spilura* (Günther).
 Streams of Central America.
Pæcilia spilurus Günther, Cat., VI, 1866, 345, Central America.
1496. *Mollienesia velifera* Regan.
 Yucatan.
Mollienesia velifera Regan, Ann. Mag. Nat. Hist., 8th ser., XIII, 1914,
 338, Progreso, Yucatan.
1497. *Mollienesia sphenops* (Cuvier and Valenciennes).
 Lowland streams of Mexico and Central America.
Pæcilia sphenops Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII,
 1836, 98 (130), 525, Vera Cruz.
Mollienesia fasciata Muller and Troschel, Monatsber. Akad. Wiss. Berlin,
 1844, 36, Mexico.
Pæcilia mexicana Steindachner, Sitzungsb., Kais. Akad. Wiss. Wien,
 XLVIII, 1863, 178, taf. 4, fig. 2, southern Mexico.
Pæcilia thermalis Steindachner, loc. cit., 181, warm springs in Central
 America.
Xiphophorus gillii Kner and Steindachner, Abhandl. Bayer. Akad.
 Wiss., X, 1864, 28, taf. 4, fig. 1 (1866), Rio Chagres, Panama.
Gambusia modesta Troschel, in Müller's Reisen Vereinigten Staaten,
 Canada and Mexico, III, 1865, 639, Mexico.
Pæcilia chisoyensis Gunther, Cat., VI, 1866, 342, Rio Chisoy, Vera
 Paz.
Gambusia plumbea Troschel, loc. cit., III, 1865, 640, Mexico.
Pæcilia dovii Gunther, loc. cit., VI, 1866, 344, Lake Nicaragua; Lake
 Amatitlan.
Platyæcilius mentalis Gill, Proc. Acad. Nat. Sci. Phila., XXVIII, 1876,
 335, Isthmus of Panama.
Pæcilia boucardi Steindachner, Ichth. Beitr., VI, 1878, 8, Colon (or
 Aspinwall).
Pæcilia limantouri Jordan and Snyder, Bull. U. S. Fish Com., XIX,
 1899, 129, fig. 10, Rio Tamesin, Tampico, Mexico.
Pæcilia butleri Jordan, Proc. U. S. Nat. Mus., XI, 1888, 330, Rio
 Presidio, Mazatlan.
Platyæcilius tropicus Meek, Field Col. Mus., Zool., VII, 1907, 146, Tur-
 rialba, Costa Rica.
Pæcilia tenuis Meek, loc. cit., 147, San Jose, Costa Rica.
Pæcilia salvatoris Regan, Ann. Mag. Nat. Hist., (7), XIX, 1907, 65,
 Salvador.
Pæcilia spilota Regan, loc. cit., 8th ser., II, 1908, 460, San Jose.

1498. *Mollienesia sphenops cuneata* (Garman).
Turbo, Gulf of Darien.
Pacilia cuneata Garman, Mem., Mus. Comp. Zool., XIX, 1895, 62, Pl. V, fig. 3, teeth, Turbo, Gulf of Darien.
1499. *Mollienesia sphenops vandepolli* (Van Lindth de Jeude).
Tropical America, abundant in fresh-water pools.
Pacilia vandepolli Van Lindth de Jeude, Notes, Leyden Mus., IX, 1887, 137, Freshwater, Curaçao.
Pacilia vandepolli arubensis Van Lindth de Jeude, loc. cit., 138, Aruba, one of the Leeward Islands.
- Genus 486. **ALFARO** Meek.
Petalosoma Regan, Ann. Mag. Nat. Hist., (8), II, 1908, 462 (*P. cultratum* Regan, name preoccupied).
Alfaro Meek, Field Mus. Nat. Hist., Zool., X, September, 1912, 72 (*A. acutiventris* Meek).
Petalurichthys Regan, Ann. Mag. Nat. Hist., X, November 1912, 494 (*Petalosoma cultratum* Regan; substitute for *Petalosoma* preoccupied).
1500. *Alfaro cultratus* (Regan).
Streams of Costa Rica.
Petalosoma cultratum Regan, Ann. Mag. Nat. Hist., XI, 1908, 458, Rio Iroquois, Costa Rica.
Alfaro acutiventris Meek, Field Mus. Nat. Hist., Zool., X, 1912, 72, Costa Rica.
- Genus 487. **PLATYPECILUS** Günther.
Platypacilus Günther, Cat., VI, 1866, 350 (*P. maculatus* Günther).
1501. *Platypacilus maculatus* Günther. *Moonfish of the aquarists*.
Mexico.
Platypacilus maculatus Günther, Cat., VI, 1866, 350, Mexico.
Platypacilus variatus Meek, Field Col. Mus., Zool., V, 1904, 145, 146, San Luis Potosi, Mexico.
Platypacilus maculatus immaculatus Myers, Copeia, No. 113, Dec. 20, 1922, 89, a form developed in an aquarium, therefore without standing in scientific nomenclature.
1502. *Platypacilus couchianus* (Girard).
Rio Grande Basin.
Limia couchiana Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 116, Rio San Juan, at Cadereita and Monterey, in Nuevo Leon, Mexico.
- Genus 488. **XIPHOPHORUS** Heckel.
Xiphophorus Heckel, Sitzungs. Kais. Akad. Wiss. Wien, I, 1848, pt. 3, 163 (*X. helleri* Heckel).
1503. *Xiphophorus helleri* Heckel. *Swordtail*.
Lowland streams of southern Mexico and Guatemala, of Atlantic drainage.
Xiphophorus helleri Heckel, Sitzungs. Kais. Akad. Wiss. Wien, I, 1848, pt. 3, 163, pl. 5, Rio Chisoy, Guatemala.
Xiphophorus jalapæ Meek, Field Col. Mus., Zool., III, 1902, 107, Jalapa.
1504. *Xiphophorus helleri guntheri* (Jordan and Evermann)
Rivers of eastern slope of Guatemala.
Xiphophorus guntheri Jordan and Evermann, Fish. North and Mid. Amer., 1896, 702, Rio Chisoy (Usumacinta), Guatemala.
1505. *Xiphophorus helleri strigatus* (Regan).
Atlantic slope of southern Mexico.
Xiphophorus strigatus Regan, Ann. Mag. Nat. Hist., (7), XIX, 1907, 65; streams of Atlantic slope of southern Mexico; Cordoba, Oaxaca, etc.
1506. *Xiphophorus helleri rachovii* (Regan).
Atlantic slope of Guatemala.
Xiphophorus rachovii Regan, Ann. Mag. Nat. Hist., (8), VIII, 1911, 374, Puerto Barrios between Lake Yzabal and Rio Motagua.

1507. *Xiphophorus helleri brevis* (Regan).
British Honduras.
Xiphophorus brevis Regan, Ann. Mag. Nat. Hist., (7), XIX, 1907, 65,
British Honduras.
1508. *Xiphophorus montezumæ* Jordan and Snyder.
Rio Panuco, eastern Mexico.
Xiphophorus montezumæ Jordan and Snyder, Bull. U. S. Fish Com.,
XIX, 1899 (1900), 131, fig. 11, Rio Panuco, Mexico.

Family 117. ANABLEPIDÆ. Four-eyed fishes

- Genus 489. ANABLEPS (Gronow) Scopeli.
Anableps Gronow, in Scopeli, Introd. Hist. Nat., 1777, 450 (no type
mentioned; *A. anableps* L. understood).
1509. *Anableps anableps* Linnæus. *Cuatro ojos*; *Four-eyed fish*.
Trinidad to Brazil; in streams.
Anableps anableps Linnæus, Syst. Nat., ed. 9, 1756, 55.
Anableps tetraphthalmus Bloch, Naturgesch. Ausl. Fische, VIII, 1794,
7, pl. 361.
1510. *Anableps dowei* Gill. *Cuatro ojos*.
Streams of Central America, locally common (Salvador-Chiapas,
Guatemala)
Anableps dowei Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 4, La
Union, Salvador.

Family 118. AMBLYOPSIDÆ. Cave blindfishes

- Genus 490. CHOLOGASTER Agassiz.
Chologaster Agassiz, Amer. Jour. Sci. Arts, XVI, 1853, 135 (*C. cornutus*
Agassiz)
1511. *Chologaster cornutus* Agassiz. *Fish of the rice ditches of South Atlantic*
States.
Lowland swamps of the Southern States, from the Dismal Swamp to
the Okefinokee Swamp; locally very abundant.
Chologaster cornutus Agassiz, Amer. Jour. Sci. Arts, XVI, 1853, 135,
ditch in a rice field in South Carolina.
1512. *Chologaster avitus* Jordan and Jenkins. *Fish of Dismal Swamp*.
Streams and swamps from southeast Virginia southward, probably
varying into *C. cornutus*.
Chologaster avitus Jordan and Jenkins, Proc. U. S. Nat. Mus., XI, 1888,
356, pl. 44, fig. 8, outlet of Lake Drummond, Dismal Swamp, near
Suffolk, Va.
- Genus 491. FORBESELLA Jordan and Evermann.
Forbesella Jordan and Evermann, Proc., Calif. Acad. Sci., 4th ser., XVI,
1927 (Apr. 27), 503 (*Chologaster papilliferus* Forbes).
This genus stands between the ancient type of *Chologaster* well colored
and with developed eyes, and the colorless blind fishes of the caves
of the middle west. From *Chologaster*, *Forbesella* differs in the pres-
ence of tactile papillæ as in *Amblyopsis*.
1513. *Forbesella papillifera* (Forbes). *Cavefish*.
Cave springs, Union and Pope Counties, Ill.
Chologaster papilliferus Forbes, Amer. Nat., XVI, January, 1882, 1,
cave spring in Union County, southern Illinois.
1514. *Forbesella agassizii* (Putnam).
Underground streams of Tennessee and Kentucky.
Chologaster agassizii Putnam, Amer. Nat., VI, 1872, 30, Pl. 2, figs.
4, 4a, 4b, from a well at Lebanon, Tenn.
- Genus 492. TYPHLICHTHYS Girard.
Typhlichthys Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 63 (*T.*
subterraneus Girard).

1515. *Typhlichthys subterraneus* Girard. *Small blindfish*.
Subterranean streams in limestone regions of Indiana, Kentucky, Tennessee, Missouri, and Alabama.
Typhlichthys subterraneus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 63, well at Bowling Green, Ky.
1516. *Typhlichthys osborni* Eigenmann.
Horse Cave, near Mammoth Cave, Ky.
Typhlichthys osborni Eigenmann, Biol. Bull., VIII, 1905, 65, figs. 3 and 4, Horse Cave, Ky.
1517. *Typhlichthys wyandotte* Eigenmann.
Well near Corydon, Ind.
Typhlichthys wyandotte Eigenmann, Biol. Bull., VIII, 1905, 63, Corydon, Ind.
- Genus 493. **TROGLICHTHYS** Eigenmann.
Troglichthys Eigenmann, A Case of Convergence, Science, new series, IX, 1899, 282 (*Typhlichthys rosæ* Eigenmann).
1518. *Troglichthys rosæ* (Eigenmann). *Missouri cavefish*.
Typhlichthys rosæ Eigenmann, Science, new series, IX, 1899, No. 217, 282, caves in Missouri.
- Genus 494. **AMBLYOPSIS** DeKay.
Amblyopsis DeKay, New York Fauna: Fishes, 1842, 187 (*A. spelæus* DeKay).
1519. *Amblyopsis spelæus* DeKay. *Mammoth Cave blindfish*.
Subterranean streams of the limestone regions of Kentucky and Indiana; River Styx, Mammoth Cave; the largest species of the group.
Amblyopsis spelæus DeKay, New York Fauna: Fishes, 1842, 187, Mammoth Cave, Ky.

Order EE. SYNENTOGNATHI

Family 119. BELONIDÆ. Houndfishes; Needlefishes

- Genus 495. **STRONGYLURA** Van Hasselt.
Strongylura Van Hasselt, Bull. Sci. Nat. Ferussac., II, 1824, 374 (*S. caudimaculata* Van Hasselt=*Belone strongylura* Van Hasselt, 1823).
1520. *Strongylura notata* (Poey). *Needlefish; Longjaws; Gar*.
West Indies, north to Pensacola; common about Key West.
Belone notata Poey, Memorias, II, 1861, 293, Cuba.
1521. *Strongylura scapularis* (Jordan and Gilbert).
Panama.
Tylosurus scapularis Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 307, Panama.
1522. *Strongylura timucu* (Walbaum). *Timucu; Peixe agulha*.
Florida Keys to Brazil; not rare.
Timucu or *Peixe agulha* Maregrave, Hist. Pisc. Brasil., 1648, 168, Brazil.
Esox timucu Walbaum, Artedi Pisc., III, 1792, 88, Brazil; after *Timucu* of Maregrave.
Belone subtruncata Poey, Memorias, II, 1860, 295, Habana.
Belone depressa Poey, loc. cit., 296, Cuba.
Tylosurus sagitta Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 25, Key West.
Belone ardeola Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 315 (425), Porto Rico.
Belone cigonella Cuvier and Valenciennes, loc. cit., 323 (436), Porto Rico.
Belone almeida Quoy and Gaimard, Voy. Uranie, Zool., 1824, 226, Bay of Rio Janeiro.
Belone timucu Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 316 (426), Rio de Janeiro.

1523. *Strongylura argalus* (LeSueur).
West Indies.
Belone argalus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 326 (439), Guadeloupe, West Indies.
1524. *Strongylura euryops* (Bean and Dresel). *Longgaw*.
Cuba and Jamaica
Tylosurus euryops Bean and Dresel, Proc. U. S. Nat. Mus., VII, 1884, 168, Jamaica.
1525. *Strongylura diplotænia* (Cope).
St. Martins Island, West Indies.
Belone diplotænia Cope, Trans. Amer. Phil. Soc., XIV, 1871, 481, St. Martins
1526. *Strongylura microps* (Gunther).
Coast of Guiana.
Belone microps Gunther, Cat., VI, 1866, 237, Surinam, British Guiana.
1527. *Strongylura angusticeps* (Gunther).
Coast of Ecuador.
Belone angusticeps Gunther, Cat., VI, 1866, 238, Ecuador.
1528. *Strongylura stolzmanni* (Steindachner). *Sierrita*.
Pacific coast of America from Guaymas and Mazatlan to Peru.
Belone stolzmanni Steindachner, Ichth. Beitr., VII, 1878, 21, Tumbes, Peru
Tylosurus sierrita Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 458, Mazatlan.
1529. *Strongylura exilis* (Girard). *Needlefish; California garfish*.
Coast of southern California, from Point Conception southward to Cedros Island, Mexico.
Belone exilis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 149, San Diego.
1530. *Strongylura pterura* (Osburn and Nichols).
Gulf of California.
Tylosurus pterurus Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 152, fig. 6, Carmen Island, Gulf of California.
1531. *Strongylura marina* (Walbaum). *Garfish; Bullfish; Needlefish; Aguñón; Salt-water gar; Silver gar; Swordfish; Green gar; Skippick; Sea pike; Harvest pike; Silvery gar*.
Coast of Maine to Texas; Atlantic and Gulf coasts, entering rivers.
Esox marinus Walbaum, Arted. Pisc., III, 1792, 88; after Schopf.
Esox longirostris Mitchill, Amer. Month. Mag., II, 1818, 322, Hudson River.
Belone truncata LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 126, New York Bay.
Belone sculator Girard, U. S. Mex. Bound. Surv., Ichth., 1859, 30, pl. 13, Brazos Santiago and St. Joseph Island, Tex.
1532. *Strongylura guianensis* (Müller and Troschel).
Tylosurus guianensis Muller and Troschel in Schomburgk's Reise, III, 1848, 626, coast of Guiana.
- Genus 496. *TYLOSURUS* Cocco.
Tylosurus Cocco, Giorn. Sci. Litt. Art. Sicilia, XVII, 1829, 18 (*T. cantrani* Cocco = ? *Sphyræna acus* Lacépède).
1533. *Tylosurus fodiator* Jordan and Gilbert. *Aguñón*.
Pacific coast of Mexico.
Tylosurus fodiator Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 459, Mazatlan.
1534. *Tylosurus raphidoma* (Ranzani). *Houndfish; Aguja de Casta; Guardfish*.
West Indies, Florida Keys to Brazil; north to Ocean City, N. J., and North Carolina.
Belone raphidoma Ranzani, Nov. Com. Acad. Sci. Bonon., V, 1842, 359, pl. 37, fig. 1, Brazil.
Belone gerania Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 325 (437), Martinique.
Belone crassa Poey, Memorias, II, 1861, 291, Cuba.
Belone melanochira Poey, loc. cit., 294, Cuba.
Tylosurus gladius Bean, Proc. U. S. Nat. Mus., V, 1882, 239, 430, Pensacola, Fla.

1535. *Tylosurus galeatus* (Cuvier and Valenciennes).
Cayenne, French Guiana.
Belone galeata Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 319 (429), Cayenne, French Guiana.
1536. *Tylosurus pacificus* (Steindachner).
Acapulco to Panama.
Belone pacifica Steindachner, Ichth. Beitr., III, 1875, 65, Panama.
1537. *Tylosurus acus* (Lacépède). *Houndfish; Agujón*.
West Indies and occasionally northward.
Sphyræna acus Lacépède, Hist. Nat. Poiss., V, 1803, 6, pl., 1 fig. 3, Martinique.
? *Esox imperialis* Rafinesque, Caratteri, 1810, 59, Palermo.
? *Tylosurus cantrainii* Cocco, Giorn. Sci. Litt. Art. Sicilia, XVII, 1829, 18, pl. 1, fig. 4, Messina.
Belone latimana Poey, Memorias, II, 1861, 290, Habana.
Belone jonesi Goode, Amer. Jour. Sci. Arts, XIII, 1877, 295, Bermuda.
1538. *Tylosurus caribbæus* (LeSueur).
West Indies.
Belone caribbæa LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 127, Caribbean Sea.
Belone altipinna Poey, Memorias, II, 1861, 293, Cuba.
- Genus 497. ABLENNES Jordan and Fordice.
Athlennes Jordan and Fordice, Proc. U. S. Nat. Mus., IX, 1886, 342 (*Belone hians* Cuvier and Valenciennes); error of transcription for *Ablennes*.
1539. *Ablennes hians* (Cuvier and Valenciennes).
West Indies, ranging from Florida to Brazil; generally common; also recorded from Acapulco and Hawaii.
Belone hians Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 321 (432), pl. 543, Bahia; common at Habana.
Belone maculata Poey, Memorias, II, 1861, 290, Cuba.

Family 120. SCOMBERESOCIDÆ

- Genus 498. SCOMBERESOX Lacépède. *Sauries*.
Scomberesox Lacépède, Hist. Nat. Poiss., V, 1803, 344 (*S. camperi* Lacépède).
Sayris Rafinesque, Caratteri, 1810, 60 (*S. recurvirostra* Rafinesque = *Esox saurus* Walbaum).
- *Grammiconotus* Costa, Ann. Mus. Zool. Univ. Napoli, I, 1862, 55 (*G. bicolor* Costa; said to lack the air bladder).
1540. *Scomberesox saurus* (Walbaum). *Saury; Skipper; Billfish; Skipjack*.
Temperate parts of Atlantic Ocean; common in schools especially north of Cape Cod and France.
Esox saurus Walbaum, Artdi Pisc., III, 1792, 93, Cornwall; after the saury pike of Pennant.
Scomberesox camperi Lacépède, Hist. Nat. Poiss., V, 1803, 345, locality unknown.
Sayris recurvirostra Rafinesque, Caratteri, 1810, 61; substitute for *camperi*.
Sayris hians Rafinesque, loc. cit., 61, Palermo.
Sayris bimaculatus Rafinesque, loc. cit., 62, Palermo.
Scomberesox scutellatum LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821 (1823), 132, Newfoundland.
Scomberesox storeri DeKay, N. Y. Fauna: Fishes, 1842, 229, pl. 35, fig. 3, Banks of Newfoundland; coast of Massachusetts; New York.
Scomberesox equirostrum LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821 (1823), 132, locality unknown.
Scomberesox rondeletii Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 350 (472), Mediterranean.
Grammiconotus bicolor Costa, Ann. Mus. Zool. Univ. Napoli, I, 1862, 55, Naples.
? *Scomberesox forsteri* Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 357 (481), New Zealand.
Sayris serrata Rafinesque, Caratteri, 1810, 61, after Rondelet.

Genus 499. COLOLABIS Gill.

Cololabis Gill, Proc. U. S. Nat. Mus., XVIII, 1895, 176 (*Scomberesox brevirostris* Peters).

1541. *Cololabis brevirostris* (Peters).

Coast of California, south to San Diego in the open sea; often in schools. *Scomberesox brevirostris* Peters, Monatsberichte Akad. Wiss. Berlin, July, 1866, 521, Tomales Bay, Calif.; perhaps not separable from the Japanese *Cololabis sawa* Brevoort.

Family 121. HEMIRAMPHIDÆ. Halfbeaks

Genus 500. EULEPTORHAMPHUS Gill.

Euleptorhamphus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 156 (*E. brevoortii* Gill = probably *E. velox* Poey).

1542. *Euleptorhamphus longirostris* (Cuvier).

East Indies; Galapagos Islands.

Hemirhamphus longirostris Cuvier, Règne Anim., ed. 2, II, 1829, 286, warm seas of both hemispheres.

1543. *Euleptorhamphus velox* Poey. *Slender halfbeak*.

West Indies, occasionally northward in the Gulf Stream to Massachusetts.

? *Euleptorhamphus brevoortii* Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 156, no locality.

Euleptorhamphus velox Poey, Synopsis, 1867, 383, Cuba; perhaps identical with the preceding.

Genus 501. HYPORHAMPHUS Gill. *Halfbeaks*

Hyporhamphus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 131 (*H. tricuspidatus* Gill = *Hemirhamphus unifasciatus* Ranzani).

1544. *Hyporhamphus unifasciatus* (Ranzani). *Common halfbeak*; *Escribano*; *Pajarito*.

Tropical seas on both coasts of America from Brazil north to Rhode Island, New Jersey, Guaymas, and Mazatlan; generally common from Key West to Rio de Janeiro; also from Guaymas to the Galapagos; variable; but the distinction into two or more species not trustworthy.

Hemirhamphus unifasciatus Ranzani, Nov. Com. Acad. Sci. Bonon., V, 1842, 326, Brazil.

? *Hemirhamphus picarti* Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 18 (25), Bono, Algiers.

Hemirhamphus richardi Cuvier and Valenciennes, loc. cit., 19 (26), St. Croix, Antilles.

Hyporhamphus tricuspidatus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 131, Barbados.

Hemirhamphus fasciatus Poey, Memorias, II, 1861, 299, Cuba; not of Bleeker.

1545. *Hyporhamphus roberti* ³⁵ (Cuvier and Valenciennes).

Both coasts of tropical America; West Indies and Brazil north to Rhode Island, and on the Pacific coast from Gulf of California to the Galapagos.

Hemirhamphus roberti Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 18 (24), Cayenne.

1546. *Hyporhamphus gilli* Meek and Hildebrand.

Acapulco to the Galapagos.

Hyporhamphus gilli Meek and Hildebrand, Field Mus. Nat. Hist. Pub. 215, Zool. Ser., XV, Dec. 20, 1923, 240, Pl. XVII, fig. 1, Panama Bay, Balboa.

1547. *Hyporhamphus rosæ* (Jordan and Gilbert).

Coast of southern California and probably southward.

Hemirhamphus rosæ Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 335, San Diego, Calif.

³⁵ This species is slenderer than *H. unifasciatus*, with longer beak and more numerous gill rakers, 28 to 31 on lower limb (21 in *H. unifasciatus*).

1548. *Hyporhamphus snyderi* Meek and Hildebrand.

Lower California to Panama.

Hyporhamphus snyderi Meek and Hildebrand, Field Mus. Nat. Hist. Pub. 215, Zool., Ser. XV, Dec. 20, 1923, 240, Pl. XVII, fig. 2, Balboa, Panama Bay.**Genus 502. *HEMIRAMPHUS* Cuvier.***Hemirampus* Cuvier, Règne Anim., ed. 1, II, 1817, 186 (*Esox brasiliensis* Linnæus).**1549. *Hemirampus brasiliensis* (Linnæus). *Balao*; *Escribano*; *Red-tailed balao*.**West Indies, generally abundant from Key West southward to Bahia
Esox brasiliensis Linnæus, Syst. Nat., ed. X, 1758, 314, Jamaica; after Browne; the *Timucu* of Macgrave wrongly included in the synonymy; Bloch, Ichth., 1801, 391, corrected synonymy and description.*Hemirampus browni* Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 10 (13), Guadeloupe, Martinique.*Hemirampus pleii* Cuvier and Valenciennes, loc. cit., 15 (19), Martinique, San Domingo.*Macrogathus brevirostris* Gronow, Cat. Fish., 1854, 148, America; after Browne.*Hemirampus filamentosus* Poey, Memorias, II, 1861, 297, Cuba.**1550. *Hemirhamphus balao* LeSueur. *Balao*; *Piper*; *Blue-tailed balao***

Cuba; also known from Panama; Guadeloupe; Martinique.

Hemirhamphus balao LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1823, 136, Lesser Antilles, Guadeloupe, Martinique.*Hemirhamphus macrochirus* Poey, Memorias, II, 1861, 299, Cuba.**1551. *Hemirhamphus saltator* Gilbert and Starks.**

Panama; Galapagos Islands.

Hemirhamphus saltator Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 53, Pl. IX, fig. 16, Panama.**Genus 503. *CHRIODORUS* Goode and Bean. *Hardheads*.***Chriodorus* Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 431 (*C. atherinoides* Goode and Bean).**1552. *Chriodorus atherinoides* Goode and Bean. *Hardhead*.**

Florida Keys and vicinity.

Chriodorus atherinoides Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 432, Key West, Fla.**Genus 504. *HEMIEXOCETUS* Fowler.***Hemioxocetus* Fowler, Proc. Acad. Nat. Sci. Phila., LIII, 1901, 293 (*H. caudimaculatus* Fowler).**1553. *Hemioxocetus caudimaculatus* Fowler.**

Mazatlan, Mexico.

Hemioxocetus caudimaculatus Fowler, Proc. Acad. Nat. Sci. Phila., LIII, 1901, 294, Mazatlan, Mexico.**Family 122. *EXOCETIDÆ*. Flyingfishes****Genus 505. *FODIATOR* Jordan and Meek.***Fodiator* Jordan and Meek, Proc. U. S. Nat. Mus., VIII, 1885, 45 (*Exocetus acutus* Cuvier and Valenciennes).**1554. *Fodiator acutus* (Cuvier and Valenciennes). *Sharp-nosed flyingfish*.**

Both shores of tropical America; Gulf of California; San Luis Gonzales; Panama; Nice.

Exocetus acutus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 91 (125), Surinam; Nice.**Genus 506. *EVOLANTIA* Heller and Snodgrass.***Evolantia* Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 189 (*Exocetus micropterus* Cuvier and Valenciennes).**1555. *Evolantia microptera* (Cuvier and Valenciennes). *Malolo*.**

Galapagos Islands.

Exocetus micropterus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 92 (127), Port King George, New Holland, etc.

Genus 507. *PAREXOCETUS* Bleeker.

Parexocetus Bleeker, Nederl. Tijdschr. Dierk., III, 1866, 126 (*Exocetus mento* Cuvier and Valenciennes).

1556. *Parexocetus mesogaster* (Bloch).

Tropical seas; East Indies and West Indies; Hawaiian Islands; Newport, Rhode Island; Carolina coast.

Exocetus mesogaster Bloch Naturgesch. Ausl. Fisch., IX, 1795, 17, Martinique; on a drawing by Plumier, in which the pectorals and ventrals are much too long.

Exocetus orbignianus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 95 (131), Montevideo

Exocetus hillianus Gosse, Naturalist's Sojourn in Jamaica, II, 1851, Pl. I, fig. 1, Jamaica.

Exocetus gryllus Klunzinger, Verhandl. Zool. Botan. Gesell. Wien, 1871, 586, Red Sea.

Genus 508. *EXOCETUS* Linnæus. *Flyingfishes.*

Exocetus Linnæus, Syst. Nat., ed. X, 1758, 316 (*E. volitans*).

1557. *Exocetus volitans* Linnæus. *Sharp-finned flyingfish.*

Warm seas; Atlantic coast north to Newfoundland; England; Hawaiian Islands and Japan, largely pelagic.

Exocetus volitans Linnæus, Syst. Nat., ed. X, 1758, 316, Atlantic Ocean; not of Cuvier and Valenciennes.

Exocetus evolans Linnæus, loc. cit., ed. XII, 1766, 521; after Gronow.

Exocetus splendens Abel, Narr. Voy. China, 1818, 4, China

Exocetus georgianus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 139, equatorial America.

Exocetus chiliensis Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 472, Chile.

1558. *Exocetus obtusirostris* Gunther.

Tropical Atlantic; recorded from Cape Verde Islands, New Orleans, and Miami, Fla.

Exocetus obtusirostris Gunther, Cat., VI, 1866, 283, Cape Verde Islands, etc.; according to Nichols (Copeia, No. 108, 1922, 49), this is different from *E. volitans*.

Genus 509. *EXONAUTES* Jordan and Evermann.

Exonautes Jordan and Evermann, Rept. U. S. Fish Com., 1895, 322 (*Exocetus exsiliens* P. L. S. Muller).

1559. *Exonautes exsiliens* (P. L. S. Müller).

Open seas; rare on our Atlantic coast.

Exocetus exsiliens Philip Ludwig Statius Muller, Nürnberg, ed. Linnæus, Syst. Nat., 1776, 209, Carolinas

Exocetus fasciatus LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 8, pl. 4, fig. 2, Atlantic.

Exocetus lamellifer Kner and Steindachner, Neue Fische Godeffroy Mus., 1866, 29, Atlantic 12° S., 39° W.

1560. *Exonautes rondeletii* (Cuvier and Valenciennes).

Tropical seas; Florida; France; Acapulco; West Indies; southern Europe; perhaps the adult of *E. exsiliens*.

Exocetus rondeletii Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 84 (115), pl. 562, Naples; Sicily; Canaries.

Exocetus brachycephalus Günther, Cat., VI, 1866, 297, China.

Exocetus volador Jordan, Proc. U. S. Nat. Mus., VII, 1884, 34 Pensacola, Fla.

1561. *Exonautes vinciguerræ* (Jordan and Meek).

Open Atlantic; southern Europe; Newfoundland; St. Martins; Gulf of Mexico.

Exocetus vinciguerræ Jordan and Meek, Proc. U. S. Nat. Mus., VIII, 1885, 56, open sea off Newfoundland, 46° N., 61° W.

- 1562. *Exonautes rubescens* (Rafinesque).**
 Atlantic Ocean, north to Newfoundland; also in the Pacific if *E. speculiger* is the same.
Exocetus rubescens Rafinesque, Amer. Month. Mag., 1818, 205, Banks of Newfoundland.
 ? *Exocetus speculiger* Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 69 (63), Friendly Islands, Pacific coast of South America, etc.
Exocetus roberti Muller and Troschel, Schomburgk's Hist. Barbados, 1848, 675, Barbados.
 ? *Exocetus quadrremus* Gronow, Cat. Fish., 1854, 145, Spain and India.
Exocetus volitans of various authors; not of Linnæus.
- 1563. *Exonautes affinis* (Günther).**
 West Indies to Africa; locally common.
Exocetus affinis Günther, Cat., VI, 1866, 288, West Africa; Cuba.
- 1564. *Exonautes rufipinnis* (Cuvier and Valenciennes).**
 Tropical America; Payta; Panama; Tabasco; Barbados.
Exocetus rufipinnis Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 72 (99), Payta, Peru.
Exocetus downi Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 167, near Panama.
Exocetus scylla Cope, Trans. Amer. Phil. Soc., XIV, 1871, 481, Tabasco, Mexico.
- Genus 510. CYPSELURUS Swainson.**
Cypselurus Swainson, Class. Fishes, etc., II, 1839, 296 (*Exocetus nuttalli* LeSueur); misprint for *Cypselurus*, according to decision Int. Com. Zool. Nom.
- 1565. *Cypselurus heterurus* (Rafinesque). *Flyingfish; Single-bearded flyingfish.***
 Atlantic Ocean, generally common southward to Newfoundland and to England.
Exocetus heterurus Rafinesque, Caratteri, 1810, 58, Palermo.
Exocetus comatus Mitchill, Trans. Lit. Phil. Soc. N. Y., 1815, 448, pl. 5, fig. 1, New York.
Exocetus noveboracensis Mitchill, Amer. Month. Mag., II, 1818, 333, New York.
Exocetus appendiculatus Wood, Jour. Acad. Nat. Sci. Phila., IV, 1824, 283, pl. 17, fig. 2, locality unknown.
Exocetus melanurus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 74 (101), New York.
- 1566. *Cypselurus lutkeni* (Jordan and Evermann).**
 Coast of Cuba.
Exocetus lutkeni Jordan and Evermann, Fishes North and Mid. Amer., 1896, 736, Cape San Antonio, Cuba.
- 1567. *Cypselurus furcatus* (Mitchill). *Double-bearded flyingfish.***
 Warm seas, north to Cape Cod and the Mediterranean, generally common.
Exocetus furcatus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 449, New York.
Exocetus nuttalli LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 10, Pl. IV, fig. 1, Gulf of Mexico.
Exocetus (*Cypselurus*) *procne* De Filippie Verany, Mem. Acad. Sci. Torino, ser. 2, XVIII, 1857, 10, fig. 5, Nice.
Exocetus maculipinnis Vinciguerra, Ann. Mus. Genov., XVIII, 1883, pl. 1, fig. 6, 577, Tunis.
- 1568. *Cypselurus nigricans* (Bennett).**
 Tropical seas, north to Cuba, Central America, and France.
Exocetus nigricans Bennett, Whaling Voyage, II, 1840, 287, taken in both the Atlantic and Pacific Oceans, in lat. 5° N.
Exocetus bicolor Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 81 (111), Atlantic.
Exocetus spilopus Cuvier and Valenciennes, loc. cit., 86 (118), La Rochelle; St. Helena; West Indies; India; Arabia; De Witt Land.

1569. *Cypselurus xenopterus* (Gilbert).
Clarion Island, one of the Revillagigedo Islands.
Exocetus xenopterus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 58,
Clarion Island.
1570. *Cypselurus lineatus* (Cuvier and Valenciennes).
Tropical Atlantic; Gorea; the Canaries; Madeira and Bermuda.
Exocetus lineatus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX,
1846, 67 (92), Gorea; Canaries.
1571. *Cypselurus cyanopterus* (Cuvier and Valenciennes).
Coasts of Brazil and Caribbean Sea; not common.
Exocetus cyanopterus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX,
1846, 71 (98), American seas.
Exocetus albidactylus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 167,
Caribbean Sea.
1572. *Cypselurus bahiensis* (Ranzani).
Tropical Atlantic; north to Cuba.
Exocetus bahiensis Ranzani, Nov. Com. Acad. Sci. Bonon., V, 1842,
362, pl. 38, Bahia.
Exocetus vermiculatus Poey, Memorias, II, 1861, 300, Cuba.
? *Exocetus parræ* Poey, Synopsis, 1868, 385, Cuba.
1573. *Cypselurus spilonotopterus* (Bleeker).
Tropical Pacific, north to Hawaii and Galapagos Islands.
Exocetus spilonotopterus Bleeker, Nederl. Tijdschr. Dierk., III, 1863,
113, Sumatra.
1574. *Cypselurus californicus* (Cooper). *Great flyingfish; Volador*.
Southern California; Point Conception to Cape San Lucas; Santa
Barbara Islands; locally very abundant
Exocetus californicus Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 93,
fig. 20, Santa Catalina Island.
1575. *Cypselurus callopterus* (Gunther).
Panama
Exocetus callopterus Gunther, Cat., VI, 1866, 292, Panama.
1576. *Cypselurus fucorum* (Zugmayer).
Mid-Atlantic.
Exocetus fucorum Zugmayer, Bull. 193, Inst. Oceanogr., Jan. 20, 1911,
in 31° 38' N., 42° 38' W., at surface.
1577. *Cypselurus holubi* (Steindachner).
Exocetus holubi Steindachner, in Collett, Voy. *Hirondelle*, 1896, 123,
Pl. IV, fig. 18, Newfoundland.
1578. *Cypselurus gibbifrons* (Cuvier and Valenciennes).
Atlantic Ocean; Newport, R. I.
Exocetus gibbifrons Cuvier and Valenciennes, Hist. Nat. Poiss., XIX,
1846, 86 (118), Atlantic.

Order FF. ANACANTHINI

Family 123. MACROURIDÆ (Coryphænoididæ). Grenadiers; Rat-tails

Genus 511. GADOMUS Regan.

Gadomus Regan, Ann. Mag. Nat. Hist., XII, 1903, 459, 466 (*Bathygadus longifilis* Goode and Bean).

1579. *Gadomus arcuatus* (Goode and Bean).
Gulf of Mexico, in deep water.
Bathygadus arcuatus Goode and Bean, Bull. Mus. Comp. Zool., XII,
1886, No. 5, 158, off Martinique in 334 fathoms.
1580. *Gadomus longifilis* (Goode and Bean).
Deep waters of the Gulf of Mexico.
Bathygadus longifilis Goode and Bean, Proc. U. S. Nat. Mus., VIII,
1885, 599, Gulf of Mexico at Albatross station 2392, lat. 28° 47' 30"
N., long. 87° 27' W., in 724 fathoms.

Genus 512. BATHYGADUS Günther.

Bathygadus Günther, Ann. Mag. Nat. Hist., II, 1878, 23 (*B. cottoides* Günther).

Melanobranchus Regan, loc. cit., (7), XI, 1903, 459 (*Bathygadus melanobranchus* Vaillant).

1581. Bathygadus favosus Goode and Bean.

Off Martinique, in deep water.

Bathygadus favosus Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, No. 5, 160, off Martinique at Blake station 80 in 472 fathoms.

1582. Bathygadus macrops Goode and Bean.

Deep waters of the Gulf of Mexico.

Bathygadus macrops Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 598, at Albatross station 2396 in lat. 28° 34' N., long. 86° 48' W., in 335 fathoms.

Genus 513. STEINDACHNERIA Goode and Bean.

Steindachneria Goode and Bean, in Agassiz, Three Cruises of the *Blake*, II, 1888, 26 (diagnosis but no type named); not *Steindachneria* Eigenmann 1888=*Steindachneridion* Eigenmann and Eigenmann, 1919.

Steindachnerella Eigenmann and Eigenmann, Amer. Nat., XXXI, 1897, 159 (*Steindachneria argentea* Goode and Bean).

1583. Steindachneria argentea Goode and Bean.

Gulf of Mexico.

Steindachneria argentea Goode and Bean, Oceanic Ichth., 1896, 419, fig. 351, off delta of Mississippi River in 68 fathoms.

Genus 514. TRACHYRINCUS Giorna.

Trachyrincus Giorna, Mem. Acad. Imp. Turin, XVI, 1805, 4, 179, (*Lepidolepsus trachyrincus* Rasso, not mentioned).

1584. Trachyrincus helolepis (Gilbert).

Off the Pacific coast of Central America.

Trachyrhynchus helolepis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 562, Pacific coast of Central America, at Albatross station 2818, in deep water.

Genus 515. DOLLOA Jordan.

Moseleya Goode and Bean, Oceanic Ichth., 1895, 417 (*Macrurus longifilis* Günther); name preoccupied.

Dolloa Jordan, Amer. Nat., XXXIV, 1900, 877 (*Macrurus longifilis* Günther).

1585. Dolloa cyclolepis (Gilbert).

Coast of British Columbia.

Nematonurus cyclolepis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 458, off Queen Charlotte Island in 1,588 fathoms.

Genus 516. NEMATONURUS Günther.

Nematonurus Günther, *Challenger* Rept., Deep Sea Fishes, XXII, 1887, 124, 150 (*N. armatus* Hector).

1586. Nematonurus goodii (Günther).

Gulf Stream from Cape Cod to Habana; generally abundant.

Macrurus asper Goode and Bean, Bull. Mus. Comp. Zool., X, 1883, No. 5, 196, Gulf Stream, south of New England in 1,242 fathoms.

Macrurus goodii Günther, *Challenger* Report, XXII, 1887, 36; substitute for *Macrurus asper*, preoccupied.

1587. Nematonurus suborbitalis (Gill and Townsend).

Bering Sea.

Macrurus suborbitalis Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 234, Bering Sea, southwest of Pribilof Islands, in 1,771 fathoms.

1588. Nematonurus acrolepis (Bean).

Straits of Juan de Fuca; off Bogoslof Island, north of Aleutian Islands; once taken at Monterey (W. F. Thompson).

Macrurus acrolepis Bean, Proc. U. S. Nat. Mus., VI, 1883, 362, Port Townsend, Wash.

1589. *Nematonurus abyssorum* Gilbert.
Off southern California in deep water.
Nematonurus abyssorum Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 374, pl. 21, off Santa Catalina in 1,350 to 2,182 fathoms.
- Genus 517. ALBATROSSIA Jordan and Evermann.
Albatrossia Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2573 (*Macrourus pectoralis* Gilbert).
1590. *Albatrossia pectoralis* (Gilbert).
Off coast of Oregon.
Macrourus (*Malacocephalus*) *pectoralis* Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 563, off coast of Oregon, at *Albatross* stations 3071, 3074, and 3075, in 685 to 877 fathoms
Macrurus (*Nematonurus*) *magnus* Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 234, Bering Sea, southwest of Pribilof Islands.
- Genus 518. BOGOSLOVIUS Jordan and Evermann.
Bogoslovius Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2574 (*B. clarki* Jordan and Gilbert).
1591. *Bogoslovius clarki* Jordan and Gilbert.
Bering Sea.
Bogoslovius clarki Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2575, Bering Sea off Bogoslof Island in 664 fathoms.
1592. *Bogoslovius firmisquamis* (Gill and Townsend).
Bering Sea.
Macrurus firmisquamis Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 234, Bering Sea southwest of Pribilof Islands.
- Genus 519. CHALINURA Goode and Bean.
Chalinura Goode and Bean, Bull. Mus. Comp. Zool., X, 1882, No. 5, 198 (*C. simula* Goode and Bean).
Chalinurus Gunther, altered spelling.
1593. *Chalinura occidentalis* (Goode and Bean).
Atlantic Ocean off Cape Hatteras and Caribbean Sea.
Malacocephalus occidentalis Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 597, off Cape Hatteras in 132 fathoms.
1594. *Chalinura serrula* Bean.
East of Prince of Wales Island in deep water.
Chalinura serrula Bean, Proc. U. S. Nat. Mus., XIII, 1890, 37, east of Prince of Wales Island, at *Albatross* station 2859 in 1,569 fathoms.
1595. *Chalinura filifera* Gilbert.
Off Queen Charlotte Island, in deep water.
Chalinura filifera Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1896), 458, off Queen Charlotte Island, at *Albatross* station 3342.
1596. *Chalinura simula* Goode and Bean.
Gulf Stream.
Chalinura simula Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 199, Gulf Stream, about latitude 40°.
1597. *Chalinura brevibarbis* Goode and Bean.
Atlantic Basin.
Chalinura brevibarbis Goode and Bean, Oceanic Ichth., 1895, 413, Atlantic Ocean in 1,255 fathoms.
1598. *Chalinura spinulosa* Gilbert and Burke.
East coast of Kamchatka.
Chalinura spinulosa Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 92, fig. 36, Avatcha Bay.
- Genus 520. CORYPHÆNOIDES³⁶ Gunner.
Coryphænoides Gunner, Trondhj. Selsk. Skrift., III, 1765, 50 (*C. rupestris* Gunner).

³⁶ The name *Coryphænoides* is older than *Macrourus*, the term usually applied to the species of this genus. The distinction between *Coryphænoides* and *Macrourus* is rather slight, the mouth in *Macrourus* being smaller and more inferior. Gilbert and Hubbs unite the two groups as *Coryphænoides*. Many species have been described from the deep waters of America. These we leave in *Coryphænoides* or in *Macrourus* as their describers have placed them. Some of the genera accepted here are of questionable value.

1599. *Coryphænoides rupestris* Gunner.
Banks of Newfoundland to Norway.
Coryphænoides rupestris Gunner, Trondhj. Selsk. Skrift., III, 1765, 50.
Lepidolepus norvegicus Nilsson, Prodr. Ichth. Scand., 1832, 51, Norway.
1600. *Coryphænoides carapinus* Goode and Bean.
Gulf Stream.
Coryphænoides carapinus Goode and Bean, Bull., Mus. Comp. Zool., X, 1883, No. 5, 197, Gulf Stream, about lat. 40°, in deep water.
- Genus 521. *MACROURUS* ³⁷ Bloch.
Macrourus Bloch, Naturgesch. Ausl. Fische, II, 1786, 150 (*Coryphæna rupestris* Fabricius = *Macrourus berglax* Lacépède); not *Coryphænoides rupestris* Gunner.
Macrurus Gunther, and of most authors; corrected spelling.
1601. *Macrourus berglax* Lacépède.
Massachusetts to Greenland and Norway.
? *Coryphæna rupestris* Fabricius, Fauna Groenlandica, 1780, 154, Greenland.
Macrourus berglax Lacépède, Hist. Nat. Poiss., III, 1802, 170, Greenland.
1602. *Macrourus bairdi* Goode and Bean. *Common rat-tail*.
West Indies to Massachusetts Bay; excessively common north in deep water.
Macrourus bairdi Goode and Bean, Amer. Jour. Sci. Arts., XIII, 1877, 471, Massachusetts Bay.
1603. *Macrourus stelgidolepis* Gilbert.
Pacific Ocean off Point Conception, Calif.
Macrurus (*Macrourus*) *stelgidolepis* Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 116, Albatross station 2960, lat. 34° 10' 45" N., long. 120° 16' 45" W., in 267 fathoms.
1604. *Macrourus cinereus* Gilbert. *Popeye*.
North Pacific and Bering Sea; very abundant.
Macrurus cinereus Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1896), 457, at Albatross stations 3307, 3329, and 3340, lat. 54° to 55° N. and long. 155° to 171° W., in 399 to 1,033 fathoms.
1605. *Macrourus bulbiceps* Garman.
Off Cocos Islands in deep water.
Macrurus bulbiceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 194, Pl. XLIV, off Cocos Islands.
1606. *Macrourus bucephalus* Garman.
Off Galapagos Islands to Gulf of California, in deep water.
Macrurus bucephalus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 195, Pl. XLIV, Gulf of California.
1607. *Macrourus barbiger* Garman.
Gulf of California in deep water.
Macrurus barbiger Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 197, Pl. XLV, Gulf of California.
1608. *Macrourus capito* Garman.
Off Panama in deep water.
Macrurus capito Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 200, off Bay of Panama.
1609. *Macrourus leucophæus* Garman.
Off Panama in deep water.
Macrurus leucophæus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 201, off Bay of Panama.
1610. *Macrourus boöps* Garman.
Off Panama in deep water.
Macrurus boöps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 202, off Bay of Panama.

³⁷ The many deep-sea forms described by Garman are nearly all referred to *Macrourus* without reference to the closely related minor genera proposed for forms allied to *Macrourus* and *Coryphænoides*, the two principal genera of the family, and themselves barely separable.

1611. *Macrourus fragilis* Garman.
Off Panama in deep water.
Macrurus fragilis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 203,
Pl. XLV, off coast of Panama.
1612. *Macrourus carminifer* Garman.
Off Panama in deep water.
Macrurus carminifer Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
204, Pl. XLVI, off Bay of Panama.
1613. *Macrourus gracilicauda* Garman.
Off Panama in deep water.
Macrurus gracilicauda Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
206, Pl. H, off Bay of Panama.
1614. *Macrourus orbitalis* Garman.
Off Panama in deep water.
Macrurus orbitalis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 207,
Pl. XLVII, off Bay of Panama.
1615. *Macrourus cuspidatus* Garman.
Gulf of California in deep water.
Macrurus cuspidatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
209, Gulf of California.
1616. *Macrourus convergens* Garman.
Off Panama in deep water.
Macrurus convergens Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
210, Pl. XLVIII, off Bay of Panama.
1617. *Macrourus latirostratus* Garman.
Off Panama in deep water.
Macrurus latirostratus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
211, Pl. XLVIII, off Bay of Panama.
1618. *Macrourus latinus* Garman.
Gulf of California in deep water.
Macrurus latinus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
214, Gulf of California.
1619. *Macrourus anguliceps* Garman.
From Galapagos Islands to Gulf of California in deep water.
Macrurus anguliceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
212, Pl. G; off Bay of Panama.
1620. *Macrourus trichurus* Garman.
Off Panama in deep water.
Macrurus trichurus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
215, off Bay of Panama.
1621. *Macrourus tenuicauda* Garman.
Off Panama in deep water.
Macrurus tenuicauda Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
216, Pl. XLIX, off Bay of Panama.
1622. *Macrourus canus* Garman.
Off Panama in deep water.
Macrurus canus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 217,
Pl. XLIX, off Bay of Panama.
1623. *Macrourus albatrossus* Townsend and Nichols.
Off Lower California in deep water.
Macrurus albatrossus Townsend and Nichols, Bull. Amer. Mus. Nat.
Hist., LII, 1925, 17, lat. 31° W., southwest of San Diego in 1,076
fathoms.
1624. *Macrourus ingolfi* Lütken.
Iceland and Greenland in deep water.
Macrurus ingolfi Lütken, Ichth. Ingolf Exped., 1898, 27, south of Iceland.

1625. *Macrourus lepturus* Gill and Townsend.
Bering Sea, St. Paul Island, Yunaska Island.
Macrourus lepturus Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 233, Bering Sea, southwest of the Pribilof Islands, in 1,401 fathoms.
Macrurus dorsalis Gill and Townsend, loc. cit., 233, Bering Sea, southwest of Pribilof Islands, in 1,401 fathoms.
- Genus 522. *COELORHYNCHUS* Giorna.
Cœlorhynchus Giorna, Mem. Acad. Nat. Sci. Turin, XVI, 1805, 179, pl. 1, figs. 3 and 4 (*C. laville* Giorna).
1626. *Cœlorhynchus occa* (Goode and Bean).
Gulf of Mexico in deep water.
Macrurus occa Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 595, Gulf of Mexico in 335 fathoms.
1627. *Cœlorhynchus carminatus* (Goode). *Grenadier*.
Caribbean Sea, north in the Gulf Stream to Long Island.
Macrurus carminatus Goode, Proc. U. S. Nat. Mus., III, 1880, 346, Gulf Stream, 40° 02' 54" N., 70° 23' 40" W., in 115 fathoms.
1628. *Cœlorhynchus caribbæus* (Goode and Bean).
West Indies to northern part of Gulf of Mexico.
Macrurus caribbæus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 594, northern part of Gulf of Mexico in 210 fathoms.
1629. *Cœlorhynchus scaphopsis* (Gilbert).
Gulf of California in deep water.
Macrurus (Cœlorhynchus) scaphopsis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 115, Gulf of California at Albatross station 3015, lat. 29° 19' N., long. 112° 50' W., in 145 fathoms.
- Genus 523. *TRACHONURUS* Günther.
Trachonurus Günther, *Challenger* Rept., Deep-sea Fishes, XXII, 1887, 124 (*Macrurus villosus* Günther).
1630. *Trachonurus sulcatus* (Goode and Bean).
West Indies and Gulf of Mexico in deep water.
Coryphænoides sulcatus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 596, Gulf of Mexico in 420 fathoms.
- Genus 524. *LIONURUS* Günther.
Lionurus Günther, *Challenger* Rept., Deep-sea Fishes, XXII, 1887, 124 (*Coryphænoides filicauda* Günther).
1631. *Lionurus liolepis* (Gilbert).
Off southern California in deep water.
Macrurus (Lionurus) liolepis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 117, coast of California at Albatross station 2980, lat. 33° 49' 45" N., long. 119° 24' 30" W. in 600 fathoms.
- Genus 525. *HYMENOCEPHALUS* Giglioli and Issel.
Hymenocephalus Giglioli and Issel, Pelagos, Genoa, 1884, 228 (*H. italicus* Giglioli and Issel).
1632. *Hymenocephalus cavernosus* (Goode and Bean).
Gulf of Mexico in deep water.
Bathygadus cavernosus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 598, Albatross station 2398, lat. 28° 45' N, long. 86° 26' W. in 227 fathoms.
- Genus 526. *ATELEOBRACHIUM* Gilbert and Burke.
Ateleobrachium Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 94 (*A. pterotum* Gilbert and Burke).
1633. *Ateleobrachium pterotum* Gilbert and Burke.
East coast of Kamchatka
Ateleobrachium pterotum Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 94, Avatcha Bay.

Family 124. BREGMACEROTIDÆ

Genus 527. BREGMACEROS (Cantor) Thompson.

Bregmaceros Thompson, in Charlesworth's Mag. Nat. Hist., IV, 1840, I, 184 (*B. maclellandii* Thompson).

Calloptilum Richardson, Voy. Sulphur, 94, 1843, 16 (*C. mirum* Richardson).

Asihenurus Tickell, Jour. Asiat. Soc. Bengal, 1865, 32 (*A. atripinnus* Tickell).

1634. *Bregmaceros bathymaster* Jordan and Bollman.

South Pacific and Indian Oceans; also off coast of Panama.

?*Bregmaceros maclellandii* Cantor, in Charlesworth's Mag. Nat. Hist., IV, 1840, 184, fig. 185, India.

?*Asihenurus atripinnus* Tickell, Jour., Asiat. Soc. Bengal, 1865, 32, with plate, Bay of Bengal.

Bregmaceros bathymaster Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 173, southwest of Panama.

1635. *Bregmaceros atlanticus* Goode and Bean.

Blake stations 99, off Granada, 113, off Nevis; and 185.

Bregmaceros atlanticus Goode and Bean, Bull., Mus. Comp. Zool., XII, 1886, No. 5, 165, off Nevis and Granada, West Indies.

1636. *Bregmaceros longipes* Garman.

Off coast of Acapulco, Mexico, in shallow water.

Bregmaceros longipes Garman, Mem., Mus. Comp. Zool., XXIV, 1899, 191, Pl. XLIII, off Pacific coast of Mexico.

Family 125. GAIDROPSARIDÆ. Rocklings

Genus 528. GAIDROPSARUS Rafinesque. Three-bearded rocklings.

Gaidropsarus Rafinesque, Indice, II, 1810, 51 (*G. mustellurus* = *Enchelyopus mediterraneus* Schneider).

Dropsarus Rafinesque, Anal. Nat., 1815, 88, shortened form; same type.

Les Mustèles Cuvier, Règne Anim., ed. 1, II, 1817, 215 (*tricurthatus* = *mediterraneus*).

Mustela Oken, Isis, 1817 (for *les Mustèles*, not *Mustela* L., a genus of weasels).

Onos Risso, Hist. Europ. Mérid., III, 1826, 214 (*Gadus mustela* Bloch).

Mustela Stark, Elem. Nat. Hist., I, 1828, 425; after *les Mustèles*.

Motella Cuvier, Règne Anim., ed. 2, II, 1829, 334 (*Gadus mustela* L.); substitute for *Mustela* Cuvier.

Onus Günther, corrected spelling; not of Rafinesque.

1637. *Gaidropsarus ensis* (Reinhardt). Rockling; Three-bearded rockling.

New York to Greenland.

Motella ensis Reinhardt, Dansk. Vidensk. Selsk. Afhandl., VII, 1838, 116, 128, Greenland.

Onos rufus Gill, Proc. U. S. Nat. Mus., VI, 1883, 259, Gulf Stream.

1638. *Gaidropsarus septentrionalis* (Collett).

Coasts of Norway and Greenland.

Motella septentrionalis Collett, Ann. Mag. Nat. Hist., 4th ser., XV, 1875, 82, Lofoten, Florø, Bergen, coast of Norway and Bodo, lat. 67° 15' N., in 30 fathoms.

Genus 529. CILIATA Couch. Five-bearded rocklings.

Ciliata Couch, Mag. Nat. Hist., V, 1822, 15 (*C. glauca* Couch, the "mackerel midge," larva of *Gadus mustela* L.)

Couchia Thompson, Nat. Hist. Ireland, IV, 1856, 288 (*C. minor* Thompson = larva of *Gadus mustela* L.).

Molvella Kaup, Archiv Natur., XXIV, 1858, I, 90 (*Molvella borealis* Kaup = *Gadus mustela* L.).

1639. *Ciliata argentata* (Reinhardt). Silvery rockling; Mackerel midge; Five-bearded rockling.

Coasts of Greenland, south to Farøe and Bear Islands.

Motella argentata Reinhardt, Dansk. Vidensk. Selsk. Afhandl., VII, 1838, 116, 128, Greenland.

Motella reinhardii Krøyer, in Collett, Forhandl. Vidensk. Selsk. Chr., No. 14, 1878, 83, Greenland.

Genus 530. ENCHELYOPUS³⁸ Bloch and Schneider. *Four-bearded rocklings*.
Enchelyopus Bloch and Schneider, Syst. Ichth., 1801, 50 (*Gadus cimbrius* L.); not of Gronow, which is nonbinomial.
Rhinnonemus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 241 (*Gadus cimbrius* L.).

1640. *Enchelyopus cimbrius* (Linnæus). *Four-bearded rockling*.
 North Atlantic on both coasts, south in deep water to the Gulf Stream; common in Massachusetts Bay.
Gadus cimbrius Linnæus, Syst. Nat., ed. 12, I, 1766, 440, Atlantic Ocean; Scania.
Motella caudacuta Storer, Proc. Bost. Soc. Nat. Hist., III, 1848, 5, Long Point, Provincetown, Mass.

Family 126. GADIDÆ. Codfishes

Genus 531. BOREOGADUS Günther.
Boreogadus Günther, Cat., IV, 1862, 336 (*Gadus fabricii* Richardson = *Gadus saida* Lepechin).

1641. *Boreogadus saida* (Lepechin).
 Arctic seas, Greenland to Alaska and northern Russia; common in the far north.
Gadus saida Lepechin, Nov. Comm. Acad. Sci. Petropol., XVIII, 1774, 512, Russia.
Merlangus polaris Sabine, Suppl. Parry's Voy., CCXI, 1824, Baffin's Bay.
Gadus fabricii Richardson, Fauna Bor.-Amer., III, 1836, 245, northern bays of Greenland.
Gadus agilis Reinhardt, Danske Vidensk. Selsk. Afhandl., VII, 1838, 126, Greenland.

Genus 532. POLLACHIUS Nilsson. *Pollacks*.
Pollachius Nilsson, in Bonaparte, Cat. Metod. Pesci. Europ., 1846, 45 (*Gadus pollachius* Linnæus).

1642. *Pollachius virens* (Linnæus). *Pollack; Coalfish; Green cod*.
 North Atlantic; common northward on both coasts, south to Cape Cod.
Gadus virens Linnæus, Syst. Nat., ed. X, 1758, 253, oceans of Europe.
Gadus carbonarius Linnæus, loc. cit., 254, seas of Europe.
Gadus colinus Lacépède, Hist. Nat. Poiss., II, 1800, 416, England, etc.
Gadus purpureus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 370, New York.
Merlangus leptocephalus DeKay, New York Fauna: Fishes, 1842, 288, pl. 45, fig. 146, Long Island.

Genus 533. THERAGRA Lucas.
Theragra Lucas, in Jordan and Gilbert, Rept. Fur-seal Invest., III, 1896, 486 (*Gadus chalcogrammus* Pallas).

1643. *Theragra chalcogramma* (Pallas). *Alaska pollack*.
 North Pacific, south to Sitka.
Gadus chalcogrammus Pallas, Zoogr. Rosso.-Asiat., III, 1811, 198, Kamchatka.
Gadus periscopus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 30, Unalaska.
 1644. *Theragra fucensis* (Jordan and Gilbert). *Wall-eyed pollack; Puget Sound pollack*.
 Puget Sound.
Pollachius chalcogrammus fucensis Jordan and Gilbert, Proc. U. S. Nat. Mus., XVI, 1893, 315, Puget Sound at Tacoma, Wash.

Genus 534. ELEGINUS Fischer.
Eleginus Fischer, Mem. Soc. Nat. Moscow, 2d ed., V, 1813, 4, 252-257 (*Gadus nawaga* Kolreuter).

³⁸ The name *Enchelyopus* ("eels without ventral fins") has been used variously since first proposed by Gronow; Gronow applied the name to *Zoarces*, Klein to *Trichurus*; as both these writers were nonbinomial we recognize its first binomial use, that of Schneider, for *Gadus cimbrius* L.

1645. *Eleginus nawaga* (Kölreuter). *Wachna cod*.

Puget Sound to Bering Sea.

Gadus nawaga Kölreuter, Nov. Com. Acad. Sci. Petropol., XIV, 1770, 484, pl. 12, coast of northern Russia.

Gadus gracilis Tilesius, Mém. Acad. Sci. Imp. St. Petersb., II, 1810, 354, Kamchatka.

Gadus wachna Pallas, Zoogr. Rosso.-Asiat., III, 1811, 182, Kamchatka.

Genus 535. *MICROGADUS* Gill. *Tomcods*.

Microgadus Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 69 (*Gadus proximus* Girard).

1646. *Microgadus tomcod* (Walbaum). *Tomcod*; *Frostfish*.

East coast of North America, from Virginia to Labrador.

Gadus tomcod Walbaum, Artedi Pisc., III, 1792, 133, Long Island; after Schöpfung.

Gadus frost Walbaum, loc. cit., 134, North America.

Gadus tomcodus pruinosus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 269, New York.

Gadus tomcodus fuscus Mitchill, loc. cit., 369, New York.

Gadus tomcodus luteus Mitchill, loc. cit., 369, New York.

Gadus tomcodus mixtus Mitchill, loc. cit., 369, New York.

Gadus polymorphus Mitchill, loc. cit., 369, New York.

Morrhua americana Storer, Rept. Fish. Mass., 1839, 120, coast of Massachusetts.

1647. *Microgadus proximus* (Girard). *California tomcod*.

West coast of North America, from Monterey to Alaska.

Gadus proximus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 141, San Francisco.

Morrhua californica Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 8, San Francisco.

Genus 536. *GADUS* Linnæus. *Codfishes*.

Gadus (Artedi) Linnæus, Syst. Nat., ed. X, 1758, 251 (*Gadus morrhua* L.).

Morrhua Oken, Isis, 1817, 1182, after "Les Morrhues" Cuvier (*Gadus morrhua* L.).

Cephus Swainson, Nat. Hist. Class. Fish., II, 1839, 300 (*Gadus macrocephalus* Tilesius).

Cerdo Gistel, Thierreich, X, 1848 (*Gadus morrhua* L., by implication).

1648. *Gadus morrhua* Linnæus. *Common codfish*.

North Atlantic, south to Virginia.

Gadus callarias Linnæus, Syst. Nat., ed. X, 1758, 252, Baltic Sea and ocean off Europe; young.

Gadus morrhua Linnæus, loc. cit., seas of Europe; name preferred by first reviser.

Gadus vertagus Walbaum, Artedi Pisc., III, 1792, 143; after Jagershen.

?*Gadus heteroglossus* Walbaum, loc. cit., 144; after Hornbogen of Klein.

Gadus arenosus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 368, New York.

Gadus rupestris Mitchill, loc. cit., 368, New York.

?*Gadus nanus* Faber, Natur. Fische Islands, 1829, 112, 182, Iceland.

1649. *Gadus macrocephalus* Tilesius. *Alaska codfish*; *Pacific cod*.

Bering Sea, south to coast of Oregon.

Gadus macrocephalus Tilesius, Mém. Acad. Sci. St. Petersb., II, 1810, 360, Bering Sea.

Gadus pygmaeus Pallas, Zoogr. Ross.-Asiat., III, 1811, 199, Kamchatka.

Gadus auratus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 30, Unalaska.

1650. *Gadus ogac* Richardson. *Greenland codfish*; *Ogac*.

Greenland; Godhaven.

Gadus ogac Richardson, Fauna Bor.-Amer., III, 1836, 246, Greenland.

Genus 537. *MELANOGRAMMUS* Gill.

Melanogrammus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 280 (*Gadus æglefinus* Linnæus).

Æglefinus Malm, Göteborg och Bohusläns Fauna, 1877, 481 (*Gadus æglefinus* L. = *Æglefinus linnæi* Malm).

1651. *Melanogrammus æglefinus* (Linnæus). *Haddock*.
North Atlantic, on both coasts.
Gadus æglefinus Linnæus, Syst. Nat., ed. X, 1758, 251, ocean off Europe.
Morrhua punctatus Fleming, Brit. Anim., 1828, 191.
Æglefinus linnæi Malm, Göteborg Bohusläns Fauna, 1877, 481.
- Genus 538. **LEPIDION** Swainson.
Lepidion Swainson, Nat. Hist. Class'n Anim., I, 1838, 318, and II, 1839, 300 (*L. rubescens* Swainson=*Gadus lepidion* Risso); not *Lepidia* Savigny, 1817,
Haloporphyrus Günther, Cat., IV, 1862, 358 (*Gadus lepidion* Risso).
1652. *Lepidion verecundum* Jordan and Cramer.
Off Pacific coast of Mexico.
Lepidion verecundum Jordan and Cramer, Proc. U. S. Nat. Mus., XIX, 1896, 456, Clarion Island, Mexico.
- Genus 539. **MICROLEPIDIDIUM** Garman.
Microlepidium Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 180, (*M. grandiceps* Garman).
1653. *Microlepidium grandiceps* Garman.
Gulf of California in deep water.
Microlepidium grandiceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 181, Pl. XLIII, Gulf of California.
- Genus 540. **ANTIMORA** Günther.
Antimora Günther, Ann. Mag. Nat. Hist., 5th ser., II, 1878, 18 (*A. rostrata* Gunther).
1654. *Antimora viola* (Goode and Bean).
Banks of Newfoundland and southward.
Haloporphyrus viola Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 257, LaHave Bank in 400 fathoms.
1655. *Antimora microlepis* Bean.
Pacific Ocean, off Queen Charlotte Islands.
Antimora microlepis Bean, Proc. U. S. Nat. Mus., XIII, 1890, 38, lat. 51° 00' 23" N., long. 130° 34' W., off Cape St. James, Queen Charlotte Islands, in 875 fathoms.
1656. *Antimora rhina* Garman.
Off Panama in deep water.
Antimora rhina Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 185, off Bay of Panama.
- Genus 541. **GADELLA** Lowe.
Gadella Lowe, Proc. Zool. Soc. Lond., 1843, 91 (*G. gracilis* Lowe).
Uraleptus Costa, Wiegmann's Archiv Naturgesch., 1858, 87 (*Gadus maraldi* Risso).
1657. *Gadella maraldi* (Risso).
Nice; Madeira; Naples and Catania; also off island of Nevis in the West Indies.
Gadus maraldi Risso, Ichth. Nice, 1810, 123, pl. 6, fig. 13, Nice.
Gadella gracilis Lowe, Proc. Zool. Soc. Lond., 1843, 91, Madeira.
- Genus 542. **LOTELLA** Kaup.
Lotella Kaup, Wiegmann's Archiv, 1858, 88 (*Lota phycis* Temminck and Schlegel).
1658. *Lotella maxillaris* Bean.
Gulf Stream.
Lotella maxillaris Bean, Proc. U. S. Nat. Mus., VII, 1884, 241, lat. 39° 55' N., long. 70° 28' W., in 396 fathoms.
- Genus 543. **PHYSICULUS** Kaup.
Physiculus Kaup, Wiegmann's Archiv, 1858, 88 (*P. dalwigkii* Kaup).
1659. *Physiculus kaupi* Poey.
Cuba.
Physiculus kaupi Poey, Repertorio, I, 1865, 186, Matanzas, Cuba.

1660. *Physiculus fulvus* Bean.
Caribbean Sea and north in the Gulf Stream to latitude 35°.
Physiculus fulvus Bean, Proc. U. S. Nat. Mus., VII, 1884, 240, lat. 40° 1' N., long. 69° 56' W., 79 fathoms.
1661. *Physiculus nematopus* Gilbert.
On both coasts of Lower California, in deep water.
Physiculus nematopus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 114, on both coasts of Lower California, at Albatross stations 2997, 3011, 3015, and 3016, in 71 to 221 fathoms.
1662. *Physiculus rastrelliger* Gilbert.
Off Panama to southern California in deep water.
Physiculus rastrelliger Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 113, coast of southern California.
1663. *Physiculus longipes* Garman.
Off Panama in deep water.
Physiculus longipes Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 188, Pl. XLII, off Bay of Panama.
- Genus 544. *LOTA* Cuvier. *Burbot*s.
Les Lottes Cuvier, Règne Anim., ed. 1, II, 1817, 215 (*Gadus lota* L.).
Lota Oken, Isis, 1817, 1182 (*Gadus lota* L.).
1664. *Lota maculosa* (LeSueur). *Burbot*; *Lake lawyer*; *Ling*; *Alekey trout*; *El pout* and *Long-tailed cat* (trade names at Sandusky, Ohio); *Mud blower*.
Rivers and lakes of New England and the Great Lakes region, north to Arctic seas, and west to the upper Missouri and Columbia River Basins.
Gadus maculosus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 83, Lake Erie.
Gadus compressus LeSueur, loc. cit., 84, Connecticut River.
Gadus lacustris Mitchell, Amer. Month. Mag., II, 1818, 244, Sebago Lake, Maine; not of Walbaum.
Molva hunta LeSueur, Mém. Mus. Hist. Nat. Paris, V, 1819, 161, Connecticut River.
Lota inornata DeKay, New York Fauna: Fishes, 1842, 283, pl. 45, fig. 145, Hudson River, Lansingburgh, N. Y.
Lota brosmiana Storer, Bost. Jour. Nat. Hist., IV, 1841, 58, pl. 5, fig. 1, Lake Winnipissiogee, N. H.
- Genus 545. *MOLVA* Fleming. *Lings*.
Molva Fleming, Hist. Brit. Anim., 1828, 192 (*Gadus molva* L. = *Molva vulgaris* Fleming).
Molva Nilsson, Scandinavian Fauna, IV, 1832, 573 (*Gadus molva* L.).
1665. *Molva molva* (Linnæus). *Ling*; *Iverksoak*.
Spitzbergen to Gulf of Gascony, Arcachon; San Juan de Luz; Iceland; Greenland; Faroe Islands.
Gadus molva Linnæus, Syst. Nat., ed. X, 1758, 254, seas of Europe.
- Genus 546. *PHYCIS* Röse. *Codlings*.
Phycis Röse, Synonymia, etc., ed. II, 1793 (*Φυκίς* Aristotle = *Phycis tinca* Schroeder); not *Phycis* Fabricius, 1798, a genus of insects.
Phycis Rafinesque, Amer. Month. Mag., 1818, 242 (*Phycis marginatus* Rafinesque = *Blennius chisso* Walbaum).
1666. *Phycis earlii* Bean.
Charleston, S. C.
Phycis earlii Bean, Proc. U. S. Nat. Mus., III, 1880, 69, Charleston, S. C.
1667. *Phycis cirratus* Goode and Bean.
Deep waters of the Gulf of Mexico.
Phycis cirratus Goode and Bean, Oceanic Ichth., 1896, 358, Gulf of Mexico, at lat. 29° 03' 15'' N., long. 88° 16' W.

Genus 547. UROPHYCIS Gill.

Phycis Bloch and Schneider, Syst. Ichth., 1801, 56 (*tinca*=*blennioides*), not *Phycis* Fabricius, 1798, a genus of Lepidoptera.

Urophycis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 240 (*Blennius regius* Walbaum).

Emphycus Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2552 (*Gadus tenuis* Mitchell).

1668. Urophycis floridanus (Bean and Dresel).

Pensacola, Fla.

Phycis floridanus Bean and Dresel, Proc. Biol. Soc. Wash., II, 1884, 100, Pensacola, Fla.

1669. Urophycis regius (Walbaum). Spotted codling, Spotted hake.

North Atlantic, south to Cape Fear.

Blennius regius Walbaum, Artedi Pisc., III, 1792, 186; after Schöpf.

Enchelyopus regalis Bloch and Schneider, Syst. Ichth., 1801, 53; after Schöpf.

Gadus punctatus Mitchell, Trans., Lit. Phil. Soc. N. Y., I, 1815, 372, New York.

1670. Urophycis tenuis (Mitchill). Codling; Squirrel hake; Hake; Lang, Thimble-eyed lang; White hake; Old English hake.

North Atlantic, south to Virginia; abundant northward.

Gadus tenuis Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 371, New York.

Phycis dekayi Kaup, Wiegmann's Archiv, 1858, Pt. I, 89, North America.

?*Phycis rostratus* Gunther, Cat., IV, 1862, 353, no locality.

1671. Urophycis chuss (Walbaum). Codling, Squirrel hake; Thimble-eyed lang; Old English hake; Chuss; Fork-beard; Red hake.

Atlantic coast, chiefly northward.

Blennius chuss Walbaum, Artedi Pisc., III, 1792, 186, Long Island; after Schöpf.

Enchelyopus americanus Bloch and Schneider, Syst. Ichth., 1801, 53; after Schöpf.

Gadus longipes Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 372, pl. 1, fig. 4, New York.

Phycis marginatus Rafinesque, Amer. Month. Mag., II, 1818, 205, Point Judith, R. I.

1672. Urophycis chesteri (Goode and Bean).

Off Massachusetts, in deep water; abundant.

Phycis chesteri Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 256, off Cape Ann, in 140 fathoms.

Genus 548. LÆMONEMA Günther.

Læmonema Günther, Cat., IV, 1862, 356 (*Phycis yarrellii* Lowe).

1673. Læmonema barbatulum Goode and Bean.

Gulf Stream.

Læmonema barbatula Goode and Bean, Bull. Mus. Comp. Zool., X, 1883, No. 5, 204, Gulf Stream, in lat. 32° 43' 25" N., long. 77° 20' 30" W., in 230 fathoms, and 38° 35' N., 73° 13' W.

1674. Læmonema melanurum Goode and Bean.

Caribbean Sea, north to New York.

Læmonema melanurum Goode and Bean, Oceanic Ichth., 1895, 363, lat. 30° 44' N., long. 79° 26' W., in 440 fathoms.

1675. Læmonema gracillipes Garman.

Galapagos Islands to Bay of Panama in deep water.

Læmonema gracillipes Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 187, Pl. XLII; off Galapagos Islands.

Genus 549. BROSME Oken. Cusk.

Brosme Oken, Isis, 1817, 1182 (*Gadus brosme* Müller); after Les Brosmes Cuvier.

Brosmerus LeSueur, Mém. Mus. Hist. Nat. Paris, 1819, 158, Pl. XVI (*Brosmerus flavesny* LeSueur=*Gadus brosme* Müller).

Brosmius Cuvier, Regne Anim., ed. 2, II, 1829, 354 (*Gadus brosme* Müller).

1676. *Brosme brosme* (Muller). *Cusk; European cusk; Nejoerpallujak; Torsk; Tusk.*
 Polar regions, south to Cape Cod.
Gadus brosme Müller, Prodr. Zool. Dan., 1776, 41, Denmark.
Gadus torsk Bonnaterre, Encycl. Meth. Ichth., 1788, 51, Sundmer, Norway; after Ström.
Brosmerus flavescens LeSueur, Mém., Mus. Hist. Nat. Paris, V, 1819, 158, Banks of Newfoundland; Marblehead, Mass.

Family 127. MERLUCCIIDÆ. Hakes

- Genus 550. MERLUCCIUS Rafinesque. *Hakes.*
Merluccius Rafinesque, Caratteri, 1810, 25 (*M. smiridus* Rafinesque = *Gadus merluccius* L.).
Onus Rafinesque, Indice, 1810, 12; substitute for *Merluccius*.
Merlangus Rafinesque, loc. cit., 167, another substitute for *Merluccius*.
Merlus Gay, in Guichenot, Pesces de Chile, II, 1848, 238 (*Merlus gayi* Guichenot).
Epicopus Gunther, Cat., II, 1860, 248 (*Merlus gayi* Guichenot).
1677. *Merluccius merluccius* (Linnæus). *European hake.*
 Coasts of Europe, straying to Greenland.
Gadus merluccius Linnæus, Syst. Nat., ed. X, 1758, 254, no locality; Europe.
Gadus ruber Lacépède, Hist. Nat. Poiss., V, 1803, 673, Scotland; Dieppe.
Gadus merlus Risso, Ichth. Nice, 1810, 122, Nice; after Linnæus.
Onus riali Rafinesque, Indice, 1810, 12, 67; substitute for *Gadus merluccius*.
Merlucius esculentus Risso, Europ. Mérid., III, 1826, 220, Nice.
Merluccius ambiguus Lowe, Proc. Zool. Soc. Lond., 1840, 37, Madeira.
Merluccius lanatus Gronow, Cat. Fishes, ed. Gray, 1854, 130, Mediterranean.
1678. *Merluccius bilinearis* (Mitchill). *Silver hake; New England hake; Whiting; Old England hake; Silverfish; American hake; Stockfish.*
 Coast of New York and northward.
Stomodon bilinearis Mitchill, Rept. Fish. N. Y., 1814, 7, New York.
Gadus albidus Mitchill, Jour. Acad. Nat. Sci. Phila., I, 1817, 409, New York.
1679. *Merluccius productus* (Ayres).
 Pacific coast, from San Diego northward to Washington.
Merlangus productus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 64 (65), no locality given; types from San Francisco market, probably from San Francisco Bay.
Homalopomus troubridgii Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 132, Astoria, Oreg.
1680. *Merluccius angustimanus* Garman.
 Off Panama, in deep water.
Merluccius angustimanus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 183, Pl. XLI, fig. 1, off Bay of Panama.

Order GG. SALMOPERCÆ

Family 128. PERCOPSIDÆ. Sand rollers; Trout perch

- Genus 551. PERCOPSIS Agassiz.
Percopsis Agassiz, Lake Superior, 1850, 284 (*P. guttatus* Agassiz = *Salmo omiscomaycus* Walbaum).
Salmoperca Thompson, App. Hist. Vermont, 1853, 33 (*S. pellucida* Thompson = *Salmo omiscomaycus* Walbaum).

1681. *Percopsis omiscomaycus* (Walbaum). *Sand roller; Trout perch.*
 Lake Champlain and Delaware River to Ohio River, Kansas, and
 northward; Great Lakes; Hudson Bay; Red River of the North;
 Saskatchewan River.
Salmo omiscomaycus Walbaum, Artedi Pisc., 1792, 65, Hudson Bay.
Percopsis guttatus Agassiz, Lake Superior, 1850, 286, Lake Superior
Salmoperca pellucida Thompson, App. Hist. Vermont, 1853, 33, Lake
 Champlain.
Percopsis hammondii Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864,
 151, Kansas.
- Genus 552. COLUMBIA Eigenmann and Eigenmann.
Columbia Eigenmann and Eigenmann, Science, XX, Oct. 21, 1892, 233
 (*C. transmontana* Eigenmann and Eigenmann).
1682. *Columbia transmontana* Eigenmann and Eigenmann.
 Columbia River Basin; Umatilla and Wallula Rivers.
Columbia transmontana Eigenmann and Eigenmann, Science, XX,
 1892, 233, mouth of Umatilla River, Oreg.

Order HH. XENARCHI

Family 129. APHREDODERIDÆ. Pirate perches

- Genus 553. APHREDODERUS LeSueur. *Pirate perches.*
Aphredoderus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss.,
 IX, 1833, 329 (445), (*A. gibbosus* LeSueur=*Scolopsis sayanus*
 Gilliams).
Sternotremia Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 39 (*S. isolepis*
 Nelson); young with the vent posterior.
Asternotremia Nelson, in Jordan, Bull. U. S. Nat. Mus., X, 1877, 51.
Aphododerus, attempted correction.
1683. *Aphredoderus sayanus* (Gilliams). *Pirate perch; Pirate; Mud perch.*
 From New York in coastwise streams to Texas; also in the Mississippi
 Basin, north to Michigan; abundant in sluggish waters.
Scolopsis sayanus Gilliams, Jour. Acad. Nat. Sci. Phila., IV, 1824, 81,
 near Philadelphia.
Aphredoderus gibbosus LeSueur, in Cuvier and Valenciennes, Hist. Nat.
 Poiss., IX, 1833, 331 (448), Lake Pontchartrain, La.
Sternotremia isolepis Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 39.
 Calumet River, Ill.; young with vent posterior
Aphododerus cookranus Jordan, Proc. Acad. Nat. Sci. Phila., XXIX,
 1877, 60, Sawyers Creek, Kendallville, Ind.
Asternotremia mesotrema Jordan, Bull. U. S. Nat. Mus., X, 1877, 52,
 Flint River, Ga.

Order II. ALLOTRIognathi

Suborder HISTICHTHYES

Family 130. LOPHOTIDÆ

- Genus 554. LOPHOTES Giorna.
Lophotes Giorna, Mém. Acad. Sci. Torino, XVI, 1805, 19 (*L. cepedianus*
 Giorna).
Leptopus Rafinesque, Specchio Scienze Palermo, I, 1814, 16 (*L. peregrinus*
 Rafinesque=*Lophotes cepedianus* Giorna).
Podoleptus Rafinesque, Anal. Nat., 1815, 93; a substitute for *Leptopus*;
 same type.
1684. *Lophotes cepedianus* Giorna.
 Open seas north to Long Beach, Calif. (W. F. Thompson), and to
 Japan, if *Lophotes capellei* is the same.
Lophotes cepedianus Giorna, Mém. Acad. Sci. Torino, XVI, 1803, 19,
 Mediterranean.
Leptopus peregrinus Rafinesque, Specchio Scienze Palermo, I, 1814,
 16, Palermo.
 ?*Lophotes capellei* Schlegel, Fauna Japonica, 1845, 132, pl. 71, Japan.

Suborder TÆNIOSOMI

Family 131. TRACHIPTERIDÆ. King of the herrings

Genus 555. TRACHIPTERUS Gouan.

Trachipterus Gouan, Hist. Nat. Poiss., 1770, 103 (*T. gouani* Gouan = *Cephola trachiptera* Gmelin).

Trachipterus of authors; corrected spelling.

Bogmarus Schneider, Syst. Ichth., 1801, 518 (*B. islandicus* Schneider = *Trachipterus arcticus*).

Vogmarus Reid, Ann. Mag. Nat. Hist., III, 1849, 456 (*B. islandicus* Schneider); substitute name as the Danish name is *Vaagmær*.

1685. *Trachipterus arcticus* (Brünnich). *Dealfish; Vaagmar.*

Arctic Sea, Iceland to Norway.

Gymnogaster arcticus Brünnich, Nya. Handl., Dansk. Vidensk. Selsk. Skrifter, III, 1788, 408, Tab. B, figs. 1-3.

Bogmarus islandicus Schneider, Svst. Ichth., 1801, 518, Tab. 101.

Trachipterus bogmarus Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1827?, 346.

Trachipterus vogmarus Reinhardt, Dansk. Vidensk. Selsk. Skrifter, VII, 1838, 71, with fig.

1686. *Trachipterus rex-salmonorum* Jordan and Gilbert. *King of the salmon.*

Off coast of California.

Trachipterus rex-salmonorum Jordan and Gilbert, Proc. Calif. Acad. Sci., 2d ser., IV, 1894, 144, pl. 9, open sea, outside Bay of San Francisco.

Trachipterus selenius Snyder, Proc. Acad. Nat. Sci. Phila., LX, 1908, 319, Monterey; adult form.

1687. *Trachipterus trachyurus* Poey.

Cuba.

Trachipterus trachyurus Poey, Memorias, II, 1861, 420, Cuba.

Suborder ATELAXIA

Family 132. STYLEPHORIDÆ

Genus 556. STYLEPHORUS Shaw.

Stylephorus Shaw, Trans. Linn. Soc. Lond., I, 1791, 90 (*S. chordatus* Shaw).

1688. *Stylephorus chordatus* Shaw.

Open sea, Atlantic; very rare.

Stylephorus chordatus Shaw, Trans., Linn. Soc. Lond., I, 1791, 90, pl. 6, Atlantic, between Cuba and Martinique.

Order JJ. SELENICHTHYES

Family 133. LAMPRIDÆ. Opah; Moonfish

Genus 557. LAMPRIUS Retzius.

Lampris Retzius, Vetensk. Akad. Nya Handl., XX, 1799, 91 (*Zeus guttatus* Brünnich = *Zeus regius* Bonnaterre).

Chrysotosus Lacépède, Hist. Nat. Poiss., III, 1803 (*Zeus luna* Gmelin).

Echemythes Gistel, Thierreich, X, 1848 (*Zeus luna* Gmelin).

1689. *Lampris regius* (Bonnaterre). *Mariposa; Opah; Moonfish; San Pedro fish; Cravo; Jerusalem haddock; Glancefish; Gudlax; Poisson lune.*

Open seas, Atlantic and Pacific; Madeira; Newfoundland; Maine; Cuba; Monterey; Hawaii; Japan; recently taken off San Pedro.

*Zeus regius*³⁹ Bonnaterre, Encycl. Meth. Ichth., 1788, 72, pl. 39, Torbay, England.

Zeus guttatus Brünnich, Danske Skrifter, III, 1788, 403, Elsinore, Denmark.

Zeus luna Gmelin, Syst. Nat., ed. XIII, 1789, 1225, Normandy.

³⁹ The name *regius* certainly has priority over *luna* and probably over *guttatus* also.

Zeus stroemii Walbaum, Artedi Pisc., ed. 2, 1792, 398, Norway; after Ström.

Scomber gunneri Bloch and Schneider, Syst. Ichth., 1801, 38, Thronhjem.

Zeus imperialis Shaw, Nat. Misc., IV, 1803, pl. 140, Torbay; after Pennant.

Zeus opah Shaw, Gen. Zool., IV, 1803, 287, Torbay; after Pennant.

Lampris lauta Lowe, Fishes Madeira, 1843, 27, Madeira.

Order KK. HETEROSOMATA. Flounders; Flatfishes

Family 134. BOTHIDÆ

Genus 558. LOPHOPSETTA Gill. *Windowpanes*.

Lophopsetta Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 220 (*Pleuronectes maculatus* Mitchill).

1690. *Lophopsetta maculata* (Mitchill). *Windowpane*.

Atlantic coast of United States, from Casco Bay to South Carolina; common.

Pleuronectes maculatus Mitchill, Rept. in part, Fish. N. Y., 1814, 9, New York.

Pleuronectes aquosus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 389, pl. 2, fig. 3, New York.

Genus 559. PLATOPHRYS Swainson.

Platophrys Swainson, Nat. Hist. Fish., Reptiles and Amphib., II, 1839, 302 (*Pleuronectes ocellatus* Spix and Agassiz).

Peloria Cocco, Intern. Alcuni Pesci, etc., 1844, 21 (*P. heckels* Cocco).

Coccolus Bonaparte loc. cit., 21 (*C. annectens* Cocco, a larva of *Platophrys*).

Bothus Bonaparte, Cat. Metod., 1846, 49 (*podas*); not of Rafinesque. *Rhomboidichthys* Bleeker, Act. Soc. Sci. Indo-Neder., I, 1856, 67 (*R. myriaster* Bleeker).

1691. *Platophrys spinosus* (Poey).

West Indies.

Rhomboidichthys spinosus Poey, Synopsis, 1868, 409, Cuba.

1692. *Platophrys constellatus* Jordan.

Galapagos Islands.

Platophrys constellatus Jordan, in Jordan and Goss, Rept. U. S. Fish Com., XIV, 1886 (1889), 266, James Island, Galapagos Islands.

1693. *Platophrys ocellatus* (Agassiz).

Tropical America; sandy shores from Long Island to Rio Janeiro.

Rhombus ocellatus Agassiz, in Spix, Pisc. Brasil., 1829, 85, pl. 46, Brazil.

Rhombus bahianus Castelnau, Anim. Amer. Sud, 1855, Bahia.

Platophrys nebularius Jordan and Gilbert, Proc. U. S. Nat. Mus., VII,

1884, 31, Key West, Fla.

1694. *Platophrys maculifer* (Poey).

West Indies.

?*Pleuronectes maculiferus* Poey, Memorias, II, 1861, 316, Cienfuegos, Cuba.

1695. *Platophrys ellipticus* (Poey).

West Indies.

Pleuronectes ellipticus Poey, Memorias, II, 1861, 315, Cuba.

1696. *Platophrys lunatus* (Linnæus). *Peacock flounder*.

West Indies.

Solea lunata et punctata (the sole) Catesby, Nat. Hist. Carolinas, 1725, Tab. 27, Bahamas.

Pleuronectes lunatus Linnæus, Syst. Nat., ed. X, 1758, 269, Bahamas; based on Catesby and the various copyists.

Pleuronectes argus Bloch, Ichthyologia, 1783, Tab. 48, Martinique; after Plumier.

?*Pleuronectes surinamensis* Bloch and Schneider, Syst. Ichth., 1801, 156, Surinam; *satis parva et glabra*.

1697. *Platophrys leopardinus* (Günther).
 Gulf of California.
Rhomboidichthys leopardinus Gunther, Cat., IV, 1862, 434, locality unknown.
Platophrys leopardinus Jordan, Proc. U. S. Nat. Mus., VII, 1884, 260, specimen from Guaymas, Sonora, Mexico.
- Genus 560. *SYACIUM* Ranzani.
Syacium Ranzani, Nov. Spec. Pisc., 1840, 20 (*S. micrurum* Ranzani).
Hemirhombus Bleeker, Versl. Akad. Amsterdam, XIII, 1862, 425 (*H. gunneensis* Bleeker).
Aramaca Jordan and Goss, in Jordan, Cat. Fish. N. Amer., 1885, 133 (*Citharichthys pætilus* Jordan and Gilbert=*Pleuronectes aramaca* Dönndorf=*Pleuronectes papillosus* Linnæus).
1698. *Syacium papillosum* (Linnæus).
 West Indies.
Aramaca Marcgrave, Hist. Pisc. Brasil., 1648, 181, Brazil.
Pleuronectes papillosus Linnæus, Syst. Nat., ed. X, 1758, 271, Brazil; based on Marcgrave.
Rhombus solexformis Agassiz, in Spix, Pisc. Brasil., 1829, 86, Tab. 47, Atlantic Ocean.
1699. *Syacium micrurum* Ranzani.
 West Indies; Key West to Rio Janeiro.
Syacium micrurum Ranzani, Nov. Spec. Pisc. Diss. Sec., 1840, 20, pl. 5, Brazil.
Hypoglossus ocellatus Poey, Memorias, II, 1861, 314, Cuba.
Hemirhombus aramaca Gunther, Cat., IV, 1862, 42, Cuba; Jamaica; not *Rhombus aramaca* Cuvier.
Citharichthys æthalon Jordan, Proc. U. S. Nat. Mus., IX, 1886, 52, Habana.
1700. *Syacium latifrons* (Jordan and Gilbert).
 Pacific coast of tropical America; Panama.
Citharichthys latifrons Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 334, Panama.
1701. *Syacium ovale* (Günther).
 Pacific coast of tropical America; Mazatlan to Panama.
Hemirhombus ovalis Günther, Proc. Zool Soc. Lond., 1864, 154, Panama.
- Genus 561. *CITHARICHTHYS* Bleeker. *Whiffs*.
Citharichthys Bleeker, Versl. Akad. Amsterd., XIII, 1862, 427 (*C. cayennensis* Bleeker=*C. spilopterus* Günther).
Orthopsetta Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Psettichthys sordidus* Girard).
Metonops Gill, loc. cit., XVI, 1864, 198 (*M. cooperi* Gill=*Psettichthys sordidus* Girard).
- Subgenus *ORTHOPSETTA* Gill.
1702. *Citharichthys sordidus* (Girard).
 Pacific coast of North America, in water of moderate depth from British Columbia to Lower California.
Psettichthys sordidus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 142, San Francisco; Tomales Bay.
Metaponops cooperi Gill, loc. cit., XVI, 1864, 198, Santa Barbara; shriveled specimen.
1703. *Citharichthys fragilis* Gilbert.
 Gulf of California.
Citharichthys fragilis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 120, Gulf of California, east coast of Lower California.
1704. *Citharichthys xanthostigma* Gilbert.
 Magdalena Bay, west coast of Lower California to Hermosa Beach, southern California, and in Gulf of California.
Citharichthys xanthostigma Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 120, Gulf of California, west coast of Lower California, and Magdalena Bay.

1705. *Citharichthys stigmæus* Jordan and Gilbert.
Coast of southern California.
Citharichthys stigmæus Jordan and Gilbert, Proc. U. S. Nat. Mus., V,
1882, 410, 411, Santa Barbara, Calif.
- Subgenus CITHARICHTHYS.
1706. *Citharichthys dinoceros* Goode and Bean.
Deep waters of Gulf of Mexico.
Citharichthys dinoceros Goode and Bean, Bull. Mus. Comp. Zool., XII,
No. 5, 157, off Martinique, St. Lucia, and Barbados.
1707. *Citharichthys platophrys* Gilbert.
Southwest of Panama; one specimen known.
Citharichthys platophrys Gilbert, Proc. U. S. Nat. Mus., XIII, 1890,
454, Albatross station 2799, southwest of Panama.
1708. *Citharichthys arctifrons* Goode.
Deep waters of Gulf Stream.
Citharichthys arctifrons Goode, Proc. U. S. Nat. Mus., III, 1880, 341,
472, Gulf Stream off southern coast of New England.
1709. *Citharichthys unicornis* Goode.
Deep waters of Gulf Stream.
Citharichthys unicornis Goode, Proc. U. S. Nat. Mus., III, 1880, 342,
Gulf Stream off southeast of New England.
1710. *Citharichthys uhleri* Jordan.
West Indies.
Citharichthys uhleri Jordan, Rept. U. S. Fish Com., XIV, 1886, 273,
Haiti.
1711. *Citharichthys macrops* Dresel.
South Atlantic and Gulf coasts of the United States.
Citharichthys macrops Dresel, Proc. U. S. Nat. Mus., VII, 1884, 539,
Pensacola, Fla.
1712. *Citharichthys spilopterus* Günther.
Both coasts of tropical America, north to New Jersey and Mazatlan.
Citharichthys spilopterus Günther, Cat., IV, 1862, 421, New Orleans,
San Domingo, Jamaica.
Citharichthys cayennensis Bleeker, Compt. Rend. Acad. Sci. Amsterdam,
XIII, 1862, 6, Cayenne, name only.
Citharichthys guatemalensis Bleeker, Nederl. Tijdschr. Dierk., 1864, 73,
Guatemala.
Hemirhombus fuscus Poey, Synopsis, 1868, 406, Habana.
1713. *Citharichthys arenaceus* Evermann and Marsh.
Porto Rico and Santo Domingo.
Citharichthys arenaceus Evermann and Marsh, Bull. U. S. Fish Com.,
XX, first part, 1900 (1902), 326, fig. 106, Mayaguez, Porto Rico.
1714. *Citharichthys gilberti* Jenkins and Evermann.
Pacific coast of tropical America, from Gulf of California to Panama.
Citharichthys gilberti Jenkins and Evermann, Proc. U. S. Nat. Mus.,
XI, 1888, 157, Guaymas, Mexico.
Citharichthys sumichrasti Jordan and Goss, Rept. U. S. Fish Com.,
XIV, 1886 (1889) 276, Rio Zanatenco, Chiapas, Mexico.
- Genus 562. *ETROPUS* Jordan and Gilbert.
Etropus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 364 (*E.*
crossotus Jordan and Gilbert).
1715. *Etropus microstomus* (Gill).
Coast of New Jersey.
Citharichthys microstomus Gill, Proc. Acad. Nat. Sci. Phila., XVI,
1864, 223, Beesleys Point, N. J.
1716. *Etropus micros* (Fowler).
Citharichthys micros Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1911,
200, with figure, Corsons Inlet, Cape May, N. J.

1717. *Etropus rimosus* Goode and Bean.
Gulf of Mexico.
Etropus rimosus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 593, coast of Florida between Pensacola and Cedar Keys, in 21 fathoms.
1718. *Etropus crossotus* Jordan and Gilbert.
Tropical America on both coasts, north to Guaymas, Mexico, and Cape Hatteras, N. C.
Etropus crossotus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 364, Mazatlan, Mexico.
- Genus 563. **PERISSIAS** Jordan and Evermann.
Perissias Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2667 (*Platophrys tæniopterus* Gilbert).
1719. *Perissias tæniopterus* (Gilbert).
Gulf of California and western coast of Lower California.
Platophrys tæniopterus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 118, Gulf of California north of La Paz, at *Albatross* station 2298, lat. 24° 51' N., long. 110° 39' W., in 40 fathoms.
- Genus 564. **ENGYPHRYS** Jordan and Bollman.
Engyphrys Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889 (1890), 176 (*E. sancti-laurentii* Jordan and Bollman).
1720. *Engyphrys sancti-laurentii* Jordan and Bollman.
Pacific Ocean off coast of Colombia.
Engyphrys sancti-laurentii Jordan and Bollman, Proc., U. S. Nat. Mus., XII, 1889, 176, Pacific Ocean off coast of Colombia, at *Albatross* station 2805, 7° 56' N., 79° 41' 30" W., and station 2795, lat. 7° 57' N., long. 78° 55' W., in 33 and 51 fathoms.
- Genus 565. **TRICHOPSETTA** Gill.
Trichopsetta Gill, Proc. U. S. Nat. Mus., XI, 1888, 601 (*Arnoglossus ventralis* Goode and Bean).
1721. *Trichopsetta ventralis* (Goode and Bean).
Gulf of Mexico, in deep water.
Citharichthys ventralis Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 592, Gulf of Mexico, *Albatross* station 2386, lat. 29° 15' N., long. 88° 6' W., in 60 fathoms.
- Genus 566. **MONOLENE** Goode.
Monolene Goode, Proc., U. S. Nat. Mus., III, 1880, 338 (*M. sessilicauda* Goode).
Thyris Goode, loc. cit., 344 (*T. pellucidus* Goode); larva of *Monolene*.
Delothyris Goode, loc. cit., VI, 1883, 110; substitute for *Thyris*, preoccupied.
1722. *Monolene sessilicauda* Goode.
Deep waters of the Gulf Stream.
Monolene sessilicauda Goode, Proc. U. S. Nat. Mus., III, 1880, 337, 338, deep sea south of New England.
Thyris pellucidus Goode, loc. cit., 344, Gulf Stream off the coast of Rhode Island.
1723. *Monolene atrimana* Goode and Bean.
Deep waters of the Caribbean Sea.
Monolene atrimana Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, 155, deep waters off Barbados.
- Genus 567. **CYCLOPSETTA** Gill.
Cyclopsetta Gill, Proc. U. S. Nat. Mus., XI, 1888, 601 (*Hemirhombus fimbriatus* Goode and Bean).
1724. *Cyclopsetta fimbriata* (Goode and Bean).
Deep waters of the Gulf of Mexico.
Hemirhombus fimbriatus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 591, deep waters of the Gulf of Mexico, between the Mississippi Delta and Cedar Keys, Fla.

1725. *Cyclopsetta querna* (Jordan and Bollman).
Pacific coast of America, off coast of Colombia.
Azevia querna Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889,
174, Pacific Ocean off coast of Colombia, at Albatross station 2802,
lat. 8° 38' N., long. 79° 31' 30'' W.
1726. *Cyclopsetta chittendeni* B. A. Bean.
Trinidad Island
Cyclopsetta chittendeni B. A. Bean, Proc. U. S. Nat. Mus., XVII, 1894,
635, Trinidad Island.
- Genus 568. *AZEVIA* Jordan and Goss.
Azevia Jordan and Goss, Rept. U. S. Fish Com., XIV, 1886 (1888),
271 (*Citharichthys panamensis* Steindachner).
1727. *Azevia panamensis* (Steindachner).
Pacific coast of Central America.
Citharichthys panamensis Steindachner, Ichth. Beitr., III, 1875, 62,
Panama.
- Genus 569. *PLATOTICHTHYS* Nichols.
Platotichthys Nichols, Bull. Amer. Mus. Nat. Hist., XLIV, 1921, 21
(*P. charles* Nichols).
1728. *Platotichthys charles* Nichols.
Bahamas.
Platotichthys charles Nichols, Bull. Amer. Mus. Nat. Hist., XLIV,
1921, 21, pl. III, Turk Island, Bahamas.
- Family 135. HIPPOGLOSSIDÆ. Halibuts
- Genus 570. *ATHERESTHES* Jordan and Gilbert.
Atheresthes Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 51
(*Platysomatichthys stomias* Jordan and Gilbert).
1729. *Atheresthes stomias* (Jordan and Gilbert). Arrow-toothed halibut.
San Francisco to Alaska.
Platysomatichthys stomias Jordan and Gilbert, Proc. U. S. Nat. Mus.,
III, 1880, 301, off San Francisco, Calif.
- Genus 571. *REINHARDTIUS* Gill.
Reinhardtius Gill, Cat. Fishes East Coast N. Amer., Suppl. Proc. Acad.
Nat. Sci. Phila., XIII, 1861, 50 (*Pleuronectes cynoglossus* Fabricius; not
of Linnæus = *Pleuronectes hippoglossoides* Walbaum).
Platysomatichthys Bleeker, Versl. Akad. Amsterdam, XIII, 1862, 426
(*Pleuronectes pinguis* Fabricius).
1730. *Reinhardtius hippoglossoides* (Walbaum). Greenland halibut.
Arctic parts of the Atlantic, south to Finland and Grand Banks.
Pleuronectes cynoglossus Fabricius, Fauna Grœnlandica, 1780, 163,
Greenland; not of Linnæus.
Pleuronectes hippoglossoides Walbaum, Artedi Pisc., 1792, 115; based
on Fabricius.
Pleuronectes pinguis Fabricius, Zoologiske Bidrag., 1824, 43, Greenland.
Hippoglossus grœnlandicus Günther, Cat., IV, 1862, 404, Greenland.
- Genus 572. *HIPPOGLOSSUS* Cuvier. Halibut.
Hippoglossus Cuvier, Règne Anim., ed. 1, II, 1817, 221 (*Pleuronectes
hippoglossus* Linnæus).
1731. *Hippoglossus hippoglossus* (Linnæus). Common halibut.
All northern seas, southward in deep waters to France, Sandy Hook,
and northern California.
Pleuronectes hippoglossus Linnæus, Syst. Nat., ed. X, 1758, 269,
European oceans.
Hippoglossus maximus Gottsche, Archiv für Naturgesch., 1835, 164,
no locality.
Hippoglossus gigas Swainson, Nat. Hist. Class'n Anim., II, 1839, 302,
no locality.
Hippoglossus ponticus Bonaparte, Cat. Metod., 1846, 47, Black Sea;
after Pallas.

Genus 573. LYOPSETTA Jordan and Goss.

Lyopsetta Jordan and Goss, in Jordan, Cat. Fish. N. Amer., 1885, 135
(*Hippoglossoides exilis* Jordan and Gilbert).

1732. *Lyopsetta exilis* (Jordan and Gilbert).

North Pacific, in rather deep water; San Francisco to Alaska.

Hippoglossoides exilis Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, off San Francisco, between the Golden Gate and Point Reyes, Calif.

Genus 574. HIPPOGLOSSOIDES Gottsche.

Hippoglossoides Gottsche, Archiv für Naturgesch., 1835, 164 (*Pleuronectes limanda* Linnaeus).

Citharus Reinhardt, Kong. Dansk. Vid Selsk., 1838, 116 (*Pleuronectes platessoides* Fabricius; not *Citharus* Bleeker, 1862).

Drepanopsetta Gill, Cat. Fish. East Coast N. Amer., Suppl. Proc. Acad. Nat. Sci. Phila., XIII, 1861, 50 (*Pleuronectes platessoides* Fabricius).

Pomatopsetta Gill, Proc., Acad. Nat. Sci. Phila., XVI, 1864, 217 (*Platessa dentata* = *Pleuronectes platessoides* Fabricius).

1733. *Hippoglossoides platessoides* (Fabricius). *Sand dab*; *Dab*.

North Atlantic, south to Cape Cod, and coasts of England and Scandinavia.

Pleuronectes linguatula Müller, Prodr. Zool. Dan., 1776, 45; not of Linnaeus.

Pleuronectes platessoides Fabricius, Fauna Groenlandica, 1780, 164, Greenland.

Pleuronectes limandoides Bloch, Naturgesch. Ausl. Fische, III, 1787, 24, Tab. 186, Europe.

1734. *Hippoglossoides elassodon* Jordan and Gilbert

Puget Sound and northward.

Hippoglossoides elassodon Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 278, Puget Sound at Seattle and Tacoma, Wash.

1735. *Hippoglossoides robustus* Gill and Townsend.

Bering Sea.

Hippoglossoides robustus Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17), Bering Sea, lat. 56° 14' N., long. 164° 08' W., Albatross station 3541 in 49 fathoms.

1736. *Hippoglossoides hamiltoni* Jordan and Gilbert.

Off Dalnoi Point, Kamchatka.

Hippoglossoides hamiltoni Jordan and Gilbert, Rept. Fur-seal Invest., 1898, 489, Dalnoi Point, Avatcha Bay, Kamchatka, in 16 fathoms.

Genus 575. EOPSETTA Jordan and Goss.

Eopsetta Jordan and Goss, in Jordan, Cat. Fish. N. Amer., 1885, 135
(*Hippoglossoides jordani* Lockington).

1737. *Eopsetta jordani* (Lockington). *California sole*.

West coast of United States, from Puget Sound to Monterey

Hippoglossoides jordani Lockington, Proc. U. S. Nat. Mus., II, 1879, 73, San Francisco.

Genus 576. PSETTICHTHYS Girard.

Psettichthys Girard, Proc. Acad. Nat. Sci. Phila., VI, 1854, 140 (*P. melanostictus* Girard).

1738. *Psettichthys melanostictus* Girard.

Pacific coast of North America from Alaska south to Monterey.

Psettichthys melanostictus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 140, San Francisco; Astoria, Oreg.

Family 136. PARALICHTHYIDÆ. Bastard halibuts**Genus 577. HIPPOGLOSSINA** Steindachner.

Hippoglossina Steindachner, Ichth. Beitr., V, 1876, 13 (*H. macrops* Steindachner).

1739. *Hippoglossina stomata* Eigenmann and Eigenmann.
Off San Diego, Calif.
Hippoglossina stomata Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., III, 1890, 22, San Diego, Calif.
1740. *Hippoglossina macrops* Steindachner.
Pacific coast of Mexico at Mazatlan.
Hippoglossina macrops Steindachner, Ichth. Beitr., V, 1876, 13, pl. 3, Mazatlan, Mexico.
1741. *Hippoglossina bollmani* Gilbert.
Pacific coast off Colombia.
Hippoglossina macrops Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 175; not of Steindachner.
Hippoglossina bollmani Gilbert, loc. cit., XIII, 1890, 122, off coast of Colombia at Albatross station 2805, lat. 7° 56' N., long. 79° 41' 30" W. in 51½ fathoms.
1742. *Hippoglossina sabanensis* Boulenger.
Isthmus of Panama.
Hippoglossina sabanensis Boulenger, Bolletino Mus. Zool. Univ. Torino, XIV, 1899, 4, Rio Sabana, Darien.
- Genus 578. *LIQGLOSSINA* Gilbert.
Lioglossina Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 122 (*L. tetrophthalmia* Gilbert).
1743. *Lioglossina tetrophthalmia* Gilbert.
Gulf of California.
Lioglossina tetrophthalmus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 122, Gulf of California, Albatross stations 3014 and 3016, in 29 and 76 fathoms.
- Genus 579. *VERASPER* Jordan and Gilbert.
Verasper Jordan and Gilbert, Rept. Fur-seal Invest., 1898, 490 (*V. moseri* Jordan and Gilbert).
1744. *Verasper moseri* Jordan and Gilbert.
Kuril Islands.
Verasper moseri Jordan and Gilbert, Rept. Fur-seal Invest., 1898, 490, Pl. LXXXV, Shana Bay, Iturup Island, Kuril Group.
- Genus 580. *XYSTREURYS* Jordan and Gilbert.
Xystreureys Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 34 (*X. liolepis* Jordan and Gilbert).
1745. *Xystreureys liolepis* Jordan and Gilbert.
Coast of California, from Point Conception southward.
Xystreureys liolepis Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 34, Santa Barbara, Calif.
- Genus 581. *PARALICHTHYS* Girard. *Bastard halibuts*.
Paralichthys Girard, Pac. R. R. Surv., X, 1859, 146 (*P. maculosus* Girard).
Chænopsetta Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 218 (*Platessa oblonga* Storer).
Pseudorhombus Bleeker, Compt. Rend. Acad. Sci. Amsterdam, XIII, 1862, 5 (*Rhombus polyspilus* Bleeker).
Uropsetta Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Hippoglossus californicus* Ayres).
1746. *Paralichthys californicus* (Ayres). *Bastard halibut*; *Monterey halibut*.
Coast of California, Tomales Bay to San Diego
Hippoglossus californicus Ayres, Proc. Calif. Acad. Nat. Sci., ser. 1, II, 1859, 29, and 1860, fig. 10, San Francisco.
1747. *Paralichthys magdalenæ* Abbott.
Coast of Lower California.
Paralichthys magdalenæ Abbott (James Francis), in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2871, Magdalen Bay, Lower California.

1748. *Paralichthys æstuarius* Gilbert and Scofield.
Mouth of Colorado River, Gulf of California.
Paralichthys æstuarius Gilbert and Scofield, Proc. U. S. Nat. Mus., XX, 1898, 499, Pl. XXIX, Shoal Point, mouth of the Colorado River.
1749. *Paralichthys brasiliensis* (Ranzani).
Atlantic coast of South America; said to range northward to Guatemala.
Hippoglossus brasiliensis Ranzani, Nov. Spec. Pisc., 1840, 10, Tab. 3, Brazil.
Pseudorhombus vorax Gunther, Cat., IV, 1862, 429, South America.
1750. *Paralichthys woolmani* Jordan and Williams.
Guaymas to Panama.
Paralichthys woolmani Jordan and Williams, in Gilbert, Proc. U. S. Nat. Mus., XIX, 1896, 457, Galapagos Islands.
1751. *Paralichthys sinaloæ* Jordan and Abbott.
Pacific coast of Mexico.
Paralichthys sinaloæ Jordan and Abbott (J. F.), in Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2872, Mazatlan and La Paz.
1752. *Paralichthys dentatus* (Linnæus). *Summer flounder*; *Fluke*.
Atlantic coast of United States from Cape Cod to Florida.
Pleuronectes dentatus Linnæus, Syst. Nat., ed. XII, 1766, and of numerous copyists.
Pleuronectes melanogaster Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 390, New York; doubled example.
Platessa ocellaris DeKay, N. Y. Fauna: Fishes, 1842, 300, p. 47, fig. 152, New York.
Paralichthys ophryas Jordan and Gilbert, Synopsis, 1883, 822, Charleston, S. C.
1753. *Paralichthys lethostigmus* Jordan and Gilbert. *Southern flounder*.
South Atlantic and Gulf coasts of the United States, north to New York.
Platessa oblonga DeKay, N. Y. Fauna: Fishes, 1842, 299, pl. 48, fig. 156, New York; not *Pleuronectes oblongus* Mitchill.
Paralichthys lethostigma Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 237, Jacksonville, Fla.
1754. *Paralichthys oblongus* (Mitchill). *Four-spotted flounder*.
Coasts of New England and New York.
Pleuronectes oblongus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 391, New York.
1755. *Paralichthys squamilentus* Jordan and Gilbert.
South Atlantic and Gulf coasts of United States.
Paralichthys squamilentus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 303, Pensacola, Fla.
1756. *Paralichthys albiguttus* Jordan and Gilbert. *Gulf flounder*.
South Atlantic and Gulf coasts of the United States.
Paralichthys albigutta Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 302, Pensacola, Fla., and Beaufort, N. C.
- Genus 582. **ANCYLOPSETTA** Gill.
Ancylosetta Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 224 (*A. quadrocellatus* Gill).
1757. *Ancylosetta quadrocellatus* Gill.
South Atlantic and Gulf coasts of the United States.
Ancylosetta quadrocellatus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 224, Pensacola; not *Platessa quadrocellata* Storer.
Paralichthys ommatus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 616, Charleston, S. C.
- Genus 583. **RAMULARIA** Jordan and Evermann.
Ramularia Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2633 (*Ancylosetta dendritica* Gilbert).
1758. *Ramularia dendritica* (Gilbert).
Gulf of California.
Ancylosetta dendritica Gilbert, Proc., U. S. Nat. Mus., XIII, 1890, 121, Gulf of California, at *Albatross* station 3022, lat. 30° 58' 30" N., long. 113° 17' 15" W., in 11 fathoms.

Genus 584. NOTOSEMA Goode and Bean.

Notosema Goode and Bean, Bull. Mus. Comp. Zool., X, 1882, 193 (*N. dilecta* Goode and Bean).

1759. Notosema dilectum Goode and Bean.

Gulf Stream.

Notosema dilecta Goode and Bean, Bull. Mus. Comp. Zool., X, 1883, 193, Gulf Stream off the coast of South Carolina.

Genus 585. GASTROPSETTA B. A. Bean.

Gastropsetta Bean, Proc. U. S. Nat. Mus., XVII, 1894, 633 (*G. frontalis* Bean).

1760. Gastropsetta frontalis B. A. Bean.

Key West and Apalachicola, Fla.

Gastropsetta frontalis B. A. Bean, Proc. U. S. Nat. Mus., XVII, 1894, 633, Key West and Apalachicola, at Albatross stations 2317 and 2783, in 45 fathoms.

Family 137. PLEURONECTIDÆ**Genus 586. PLEURONICHTHYS** Girard.

Pleuronichthys Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1854, 139 (*P. cænosus* Girard).

Heteroprosopon Bleeker, Versl. Akad. Amsterdam, XIII, 1862 (*Pleuronectes cornutus* Temminck and Schlegel).

Parophrys Gunther, Cat., IV, 1862, 454; not of Girard.

1761. Pleuronichthys decurrens Jordan and Gilbert.

Pacific coast of United States, south to Monterey.

Pleuronichthys decurrens Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 453, San Francisco; Monterey Bay.

1762. Pleuronichthys verticalis Jordan and Gilbert.

Coast of California, in deep water.

Pleuronichthys verticalis Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 49, San Francisco.

1763. Pleuronichthys cænosus Girard.

Alaska, southward to San Diego; most common about Puget Sound.

Pleuronichthys cænosus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 139, San Francisco.

1764. Pleuronichthys nephelus Starks and Thompson.

Puget Sound.

Pleuronichthys nephelus Starks and Thompson, Proc. U. S. Nat. Mus., XXXVIII, 1911, 282, fig. 1, San Juan Islands, Wash.

1765. Pleuronichthys ritteri Starks and Morris.

San Diego Bay.

Pleuronichthys ritteri Starks and Morris in Starks & Thompson, Proc. U. S. Nat. Mus., XXXVIII, 1911, 287, San Diego, Calif.

1766. Pleuronichthys ocellatus Starks and Thompson.

Gulf of California.

Pleuronichthys ocellatus Starks and Thompson, Proc. U. S. Nat. Mus., XXXVIII, 1911, 285, fig. 2, Gulf of California.

Genus 587. HYP SOPSETTA Gill.

Hypsopsetta Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Pleuronichthys guttulatus* Girard).

1767. Hypsopsetta guttulata (Girard). *Diamond flounder*.

Coast of California, Cape Mendocino to Magdalena Bay.

Pleuronichthys guttulatus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 137, Tomales Bay, Calif.

Parophrys ayresi Gunther, Cat., IV, 1862, 457, San Francisco.

Genus 588. PAROPHRYS Girard.

Parophrys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 139 (*P. vetulus* Girard).

1768. *Parophrys vetulus* Girard.

Pacific coast of North America, Alaska to Santa Barbara.

Parophrys vetulus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 140, California.

Pleuronectes diagrammus Günther, Cat., IV, 1862, 445, Victoria, British Columbia.

Parophrys hubbardii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 281, San Francisco.

Genus 589. *PLEURONECTES* Linnæus. *Plaice*.

Pleuronectes Artedi, Genera Pisc., in part, 1738, 16.

Pleuronectes Linnæus, Syst. Nat., ed. X, 1758, 268; after Artedi (*P. platessa* Linnæus).

Platessa Cuvier, Règne Anim., ed. I, II, 1817, 220 (*Pleuronectes platessa* L.).

1769. *Pleuronectes quadrituberculatus* Pallas.

Coasts of Alaska and Kamchatka.

Pleuronectes quadrituberculatus Pallas, Zoogr. Rosso-Asiat., III, 1811, 423, sea between Kamchatka and Alaska.

Pleuronectes pallasi Steindachner, Ichth. Beitr., VIII, 1879, Kamchatka.

Genus 590. *PLATICHTHYS* Girard. *Starry flounders*.

Platichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 139 (*P. rugosus* Girard = *Pleuronectes stellatus* Pallas).

1770. *Platichthys stellatus* (Pallas). *Great flounder; California flounder.*

Pacific coast of America, from Point Conception to the Arctic Ocean, and south to Sakhalin Island.

Pleuronectes stellatus Pallas, Zoogr. Rosso-Asiat., III, 1811, 416, Kamchatka, Aleutian and Kuril Islands.

Platichthys rugosus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 139, 155, San Francisco, Presidio; Petaluma, Calif.

Genus 591. *INOPSETTA* Jordan and Goss.

Inopsetta Jordan and Goss, Rept. U. S. Fish Com., 1885 (1887), 924 (136 of inside paging) (*Parophrys ischyryus* Jordan and Gilbert).

1771. *Inopsetta ischyra* (Jordan and Gilbert).

Puget Sound, probably northward to Alaska.

Parophrys ischyryus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 276, 453, Puget Sound.

Genus 592. *LEPIDOPSETTA* Gill.

Lepidopsetta Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Psetichthys umbrosus* Girard).

1772. *Lepidopsetta bilineata* (Ayres).

Pacific coast of North America, Alaska to Monterey and northern Japan.

Platessa bilineata Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 40 (39 of reprint), San Francisco.

Platichthys umbrosus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 136, Puget Sound.

Pleuronectes perarcuatus Cope, Proc. Amer. Phil. Soc. Phila., XIII, 1873, 32, Unalaska.

Genus 593. *ISOPSETTA* Lockington.

Isopsetta Lockington, in Jordan and Gilbert, Bull. U. S. Nat. Mus., XVI, 1882, 832 (*Lepidopsetta isolepis* Lockington).

1773. *Isopsetta isolepis* (Lockington).

Puget Sound to Point Conception, in rather deep water.

Lepidopsetta isolepis Lockington, Proc. U. S. Nat. Mus., III, 1880, 325, San Francisco.

Genus 594. *LIMANDA* Gottsche. *Mud dabs*.

Limanda Gottsche, Archiv für Naturgesch., 1835, 100 (*Pleuronectes limanda* Linnæus).

Myzopsetta Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 217 (*Platessa ferruginea* Storer).

1774. *Limanda ferruginea* (Storer).
Atlantic coast of North America, Labrador to New York.
Platessa ferruginea D. H. Storer, Rept. Fish. Mass., 1839, 141, pl. 2, Cape Ann.
Platessa rostrata H. R. Storer, Bost. Jour. Nat. Hist., VI, 1850, 268, Labrador.
1775. *Limanda aspera* (Pallas).
Coasts of Alaska and Kamchatka.
Pleuronectes asper Pallas, Zoogr. Rosso.-Asiat., III, 1811, 425, east coast of Siberia.
1776. *Limanda proboscidea* Gilbert.
Bristol and Herendeen Bays, Alaska.
Limanda proboscidea Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1896), 460, pl. 33, Bristol and Herendeen Bays, Alaska.
1777. *Limanda beanii* Goode.
Deep water, off coast of New England.
Limanda beanii Goode, Proc. U. S. Nat. Mus., III, 1880 (Feb. 16, 1881), 473 southern coast of New England, *Fish Hawk* stations 875, 876.
- Genus 595-596. PSEUDOPLEURONECTES Bleeker. *Winter flounders*.
Pseudopleuronectes Bleeker, Versl. Akad. Amsterdam, XIII, 1862, 428 (*Platessa plana* Mitchell=*Pleuronectes americanus* Walbaum).
1778. *Pseudopleuronectes americanus* (Walbaum). *Common flatfish; Winter flounder; Roughback*.
Atlantic coast of North America, from Labrador to Cape Lookout.
Flounder Schopf, Schrift. Gesellsch. Naturf. Freunde, VIII, 1788, 148, New York.
Pleuronectes americanus Walbaum, Artedi Pisc., 1792, 113; based on the *Flounder* of Schöpf.
Pleuronectes planus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 387, New York.
Platessa pusilla DeKay, New York Fauna: Fishes, 1842, 296, pl. 47, fig. 153, New York.
1779. *Pseudopleuronectes dignabilis* Kendall. *Lemon-yellow flounder*.
North Atlantic coast, Newfoundland to Cape Cod.
Pseudopleuronectes dignabilis Kendall, Bull. U. S. Fish Com., XXX, 1910 (1912), 392, Georges Bank.
- Genus 597. LIOPSETTA Gill.
Liopsetta Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 216 (*Platessa glabra* Storer).
Euchalarodus Gill, loc. cit., 221 (*E. putnami* Gill, the male of *Liopsetta glabra* Storer).
1780. *Liopsetta glacialis* (Pallas). *Arctic flounder*.
Arctic Ocean and Bering Sea south to St. Michaels, Alaska.
Pleuronectes glacialis Pallas, Itin., III, App, 1776, 706; mouth of the Obi River.
Pleuronectes cicatricosus Pallas, Zoogr. Rosso.-Asiat., III, 1811, 424, sea between Kamchatka and Alaska; male.
? *Platessa dvinensis* Lilljeborg, Veb. Ab. Handl, 1850, 360, Tab. 20, mouth of River Dwina.
Pleuronectes franklini Günther, Cat., IV, 1862, 442, Arctic seas of America; female.
1781. *Liopsetta putnami* (Gill). *El-back flounder*.
Atlantic coast of North America, from Cape Cod northward to Labrador and beyond.
Platessa glabra Storer, Proc. Bost. Soc. Nat. Hist., 1843, 130, Massachusetts; female; not *Platessa glabra* of Rathke, 1837, a species of *Flesus*.
Euchalarodus putnami Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 216, 221, Salem, Mass.

1782. *Liopsetta obscura* (Herzenstein).

Sea of Okhotsk.

Pleuronectes obscurus Herzenstein, *Mélanges Biologiques*, 1890, 127, Manchuria.Genus 598. MICROSTOMUS Gottsche. *Smear dabs*.*Microstomus* Gottsche, *Archiv für Naturgesch.*, 1835, 150 (*M. latidens* Gottsche); not *Microstoma* Risso, 1826.*Cynoglossus* Bonaparte, *Fauna Italica*, 1837, fasc. XIX (*Pleuronectes cynoglossus* Nilsson); not of Linnæus = *Pleuronectes kitt* Walbaum.*Cynoglossa* Bonaparte, *Cat. Metod*, 1845, 48 (*Pleuronectes cynoglossus* Linnæus).*Brachyprosopon* Bleeker, *Versl. Akad. Amsterdam*, XIII, 1862, 428 (*Pleuronectes microcephalus* Donovan = *P. kitt* Walbaum).*Cynoglossus* Jordan and Gilbert, *Synopsis*, 1883, 460 (*Pleuronectes microcephalus* Donovan), *microstomus* being regarded as preoccupied.1783. *Microstomus kitt* (Walbaum). *Smear dab*.

Arctic Europe, west to the sea between Greenland and Iceland.

Rhombus lævis cornubiensis Jago in Ray, "Syn. Pisc.", 1713, 162, Tab. 1, fig. 1."*Smear dab* Pennant, *British Zoology*, III, 1776, 230, pl. 41.*Pleuronectes kitt* Walbaum, *Artedi Pisc.*, III, 1792, 120; after Ray.*Pleuronectes quenseli* Holboll, *Bohusläns*, Sweden.*Pleuronectes quadridens* Fabricius, *Afhandl., Kongl. Dansk. Vid. Selsk.*, I, 1824, 39, Iceland.*Microstomus latidens* Gottsche, *Archiv für Naturgesch.*, 1835, 150, Zealand.*Pleuronectes gilli* Steindachner, *Ichth. Notizen*, VII, 1868, polar sea, north of Iceland.1784. *Microstomus pacificus* (Lockington).

Pacific coast of North America, Monterey to Vancouver Island and probably northward.

Glyptocephalus pacificus Lockington, *Rept. Calif. Com. Fish.*, 1878-79, 43, off Point Reyes, Calif.

Genus 599. EMBASSICHTHYS Jordan and Evermann.

Embassichthys Jordan and Evermann, *Cheek List*, 1895 (1896), 506 (*Cynoglossus bathybius* Gilbert).1785. *Embassichthys bathybius* (Gilbert).

Coast of California, south of Point Conception.

Cynoglossus bathybius Gilbert, *Proc. U. S. Nat. Mus.*, XIII, 1890, 123, Santa Barbara Channel at Albatross station 2980, lat. 33° 49' 45" N., long. 119° 24' 30" W., in 603 fathoms.

Genus 600. GLYPTOCEPHALUS Gottsche.

Glyptocephalus Gottsche, *Archiv für Naturgesch.*, 1835, 156 (*Pleuronectes saxicola* Fabricius = *P. cynoglossus* Linnæus).1786. *Glyptocephalus cynoglossus* (Linnæus). *Craig fluke; Pole flounder; Gray sole; Lemon sole*.North Atlantic, chiefly in deep water, south to Cape Cod and France. *Pleuronectes cynoglossus* Linnæus, *Syst. Nat.*, ed. X, 1758, 269; after Gronow.*Pleuronectes saxicola* Faber, *Tidsskr. für Naturv.*, 5 B., 1828, 244, Denmark.*Glyptocephalus acadianus* Gill, *Proc. Acad. Nat. Sci. Phila.*, XXV, 1873, 360, Nova Scotia.

Genus 601. ERREX Jordan.

Errex Jordan, *Proc. Acad. Nat. Sci. Phila.*, LXXI, 1919, 343 (*Glyptocephalus zachirus* Lockington).1787. *Errex zachirus* (Lockington). *Rex sole; Long-finned sole*.

Waters of moderate depth of the northern Pacific; from San Francisco northward.

Glyptocephalus zachirus Lockington, *Proc. U. S. Nat. Mus.*, II, 1879, 88, San Francisco.

Family 138. ACHIRIDÆ. Broad soles

Subfamily GYMNACHIRINÆ

Genus 602. GYMNACHIRUS Kaup.

Gymnachirus Kaup, Wiegmann's Archiv, I, 1858, 101 (*Gymnachirus nudus* Kaup).

1788. *Gymnachirus nudus* Kaup.

Coast of Brazil; once taken at Tisbury Great Pond, Martha's Vineyard Island, Mass. (Vinal N. Edwards).

Gymnachirus nudus Kaup, Wiegmann's Archiv, I, 1858, 101, Brazil

1789. *Gymnachirus fasciatus* Günther.

West Indies; very rare, north to Long Key, Fla. (W. C. Kendall).

Gymnachirus fasciatus Günther, Cat., IV, 1862, 488, locality unknown.

Subfamily ACHARINÆ

Genus 603. ACHIRUS Lacépède.

Achirus Lacépède, Hist. Nat. Poiss., IV, 1803, 658 (*Pleuronectes achirus* Linnæus).

Trinectes Rafinesque, Atlantic Jour. and Friend of Knowledge, I, 1832 (*T. scabra* Rafinesque=*Pleuronectes mollis* Mitchill)

Grammichthys Kaup, Arch. Naturgesch., 1858, 94 (*Achirus lineatus* Linnæus).

Baistoma Bean, Proc. U. S. Nat. Mus., V, 1882, 413 (*B. brachialis* Bean).

Subgenus ACHIRUS Lacépède.

1790. *Achirus achirus* (Linnæus).

Atlantic coasts of tropical America.

Pleuronectes oculus dextris corpore glabro, pinnis pectoralibus nullus Gronow, Museum, I, No. 42, Surinam.

Pleuronectes achirus Linnæus, Syst. Nat., ed. X, 1758, 268, Surinam; based on Gronow.

Solea gronovi Günther, Cat., IV, 1862, 472, Surinam.

1791. *Achirus inscriptus* Gosse.

West Indies, north to Key West.

Achirus inscriptus Gosse, Nat. Sojourn Jamaica, 1851, 52, pl. 1, fig. 4, Jamaica.

Monochir reticulatus Poey, Memorias, II, 1861, 317, Cuba.

Subgenus GRAMMICHTHYS Kaup.

1792. *Achirus klunzingeri* (Steindachner).

Pacific coast of tropical America, Panama to Guayaquil.

Solea klunzingeri Steindachner, Zur Fische des Cauca und der Flüsse bei Guayaquil, 1879, 44, Guayaquil.

Subgenus TRINECTES Rafinesque.

1793. *Achirus fasciatus* Lacépède. *American sole*; *Hogchoker*.

Atlantic coast of United States, from Cape Cod to Texas; often ascending streams.

Achirus fasciatus Lacépède, Hist. Nat. Poiss., IV, 1803, 659, Charleston, S. C.

Pleuronectes mollis Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 388, pl. 2, fig. 4, New York.

Pleuronectes apoda Mitchill, Amer. Month. Mag. and Crit. Rev., Feb., 1818, 244, Straits of Bahama.

Trinectes scabra Rafinesque, Atlantic Jour. and Friend of Knowledge, I, 1832, Pennsylvania, in fresh water.

Solea browni Günther, Cat., IV, 1862, 477, New Orleans; Texas.

1794. *Achirus scutum* (Günther).

Pacific coast of Central America.

Solea scutum Günther, Cat., IV, 1862, 474, Gulf of Fonseca, Panama.

1795. *Achirus fonsecensis* (Gunther).
Pacific coast of tropical America, Mazatlan to Fonseca.
Solea fonsecensis Günther, Cat., IV, 1862, 475, Gulf of Fonseca.
Solea panamensis Steindachner, Ichth. Beiträge, V, 1876, 10, Taf. II, Panama.
Solea fischeri Steindachner, Beiträge Kenntniss Fluss-Fische Sudamer., I, 1879, 13, Rio Mamone near Panama.
1796. *Achirus mazatlanus* (Steindachner). *Mexican sole; Teipalcate*.
Pacific coast of tropical America.
Solea mazatlanus Steindachner, Ichth. Notizen, IX, 1869, 23, Mazatlan, Mexico.
Solea pilosa Peters, Berliner Monatsber., 1869, 709, Mazatlan, Mexico.
1797. *Achirus comifer* Jordan and Gilbert
Key West, Fla.
Achirus comifer Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 31, Key West, Fla.
1798. *Achirus lineatus* (Linnæus).
Florida to Brazil.
Passer lineis transversis notatus Sloane, Jamaica, 2, 1725, 77, pl. 246, fig. 2, Jamaica.
Pleuronectes fuscus subrotundus glaber Browne, Jamaica, 1756, 445, Jamaica.
Pleuronectes lineatus Linnæus, Syst. Nat., ed. X, 1758, 268, Jamaica; based on Browne and Sloane; not of ed. XII, which is *Achirus fasciatus*.
Monochir maculipinnis Agassiz, Spix, Pisc. Brasil., 1829, 88, pl. 49, Brazil.
Baostoma brachialis Bean, Proc., U. S. Nat. Mus., V, 1882, 413, Appalachicola Bay and south Florida.
1799. *Achirus barnharti* Jordan.
Coast of southern California; Coronado Islands.
Achirus barnharti Jordan, Univ. Calif. Pub. Zool., XXVI, 1923, 10, Pl. I, Coronado Islands in about 400 feet depth of water.
- Genus 604. *APIONICHTHYS* Kaup.
Apionichthys Kaup, Wiegmann's Archiv, 1858, 104 (*A. dumerili* Kaup).
Soleotalpa Günther, Cat., IV, 1862, 489 (*S. unicolor* Günther).
1800. *Apionichthys dumerili* Kaup.
West Indies.
Apionichthys dumerili Kaup, Wiegmann's Archiv, 1858, 104, no locality; description of species included in that of the genus.
Soleotalpa unicolor Günther, Cat., IV, 1862, 489, West Indies.
Apionichthys nebulosus Peters, Berliner Monatsb., 1869, 709, Surinam.
- Family 139. CYNOGLOSSIDÆ. Tonguefishes
- Genus 605. *SYMPHURUS* Rafinesque. *Tonguefishes*.
Symphurus Rafinesque, Indice, 1810, 13, 52 (*S. nigrescens* Rafinesque).
Bibronia Cocco, Alcorni Pesci del Mare di Messina, 1844, 15 (*B. ligulata* Cocco, larval form of *Symphurus*).
Plagusia Cuvier, Règne Animal, ed. 2, II, 1829, 344; based on *Plagusia* of Browne; name preoccupied in crustaceans, Latreille, 1806.
Plagusia Bonaparte, Cat. Metod., 1846, 51 (*P. lactea*); name a substitute for *Plagusia*.
Aphoristia Kaup, Archiv fur Natur., 1858, 106 (*Achirus ornatus* Lacépède).
Glossichthys Gill, Cat. Fish. E. Coast N. Amer., 1861, 51 (*Pleuronectes plagusia* Linnæus).
Ammopleurus Günther, Cat., IV, 1862, 490 (*Plagusia lactea* Bonaparte).
? *Bascanius* Schiötte, Naturhist. Tijdskr., V, 1867, 269 (*B. lædifera* Schiötte; larval).
1801. *Symphurus piger* (Goode and Bean).
West Indies and Gulf of Mexico in deep water.
Aphoristia pigra Goode and Bean, Bull. Mus. Comp. Zool., XII, No. 5, 1886, 154, St. Kitts, in about 250 fathoms.

1802. *Symphurus marginatus* (Goode and Bean).
West Indies.
Aphoristia marginata Goode and Bean, Bull. Mus. Comp. Zool., XII, No. 5, 1886, 153, off St. Vincent, West Indies.
1803. *Symphurus atramentatus* Jordan and Bollman.
Off coast of Colombia.
Symphurus atramentatus Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 177, off coast of Colombia at Albatross station 2795, lat. 7° 57' N., long. 78° 55' W., in 33 fathoms.
1804. *Symphurus fasciolaris* Gilbert.
Gulf of California.
Symphurus fasciolaris Gilbert, Proc. U. S. Nat. Mus., XIII, 1891, 506, Gulf of California.
1805. *Symphurus elongatus* (Günther).
Pacific coast of tropical America.
Aphoristia ornata var. *elongata* Günther, Fishes Cent. Amer., 1869, 473, Panama.
1806. *Symphurus atricaudus* (Jordan and Gilbert). *San Diego sole*.
Lower California, north to San Diego.
Aphoristia atricauda Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 23, San Diego, Calif.
1807. *Symphurus leei* Jordan and Bollman.
Off the coast of Colombia.
Symphurus leei Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 178, off coast of Colombia at Albatross station 2804, lat. 8° 16' 30'' N., long. 79° 37' 45'' W., in 47 fathoms.
1808. *Symphurus plagusia* (Bloch and Schneider). *Acedia*.
West Indies, south to Rio Janeiro.
Plagusia Browne, Jamaica, No. 1, 1756, 445, Jamaica.
Pleuronectes plagusia Bloch and Schneider, Syst. Ichth., 1801, 162, Jamaica; after Browne.
Achirus ornatus Lacépède, Hist. Nat. Poiss., IV, 1803, 659; on a specimen "presented by Holland to France."
Plagusia tessellata Quoy and Gaimard, Voyage Uranie, Zoology, 1824, 240, Rio Janeiro.
Plagusia brasiliensis Agassiz, in Spix, Pisc. Brasil., 1827, 89, tab. 50, Brazil.
1809. *Symphurus plagiusa* (Linnæus). *Tonguefish*.
South Atlantic and Gulf coasts of United States.
Pleuronectes plagiusa Linnæus, Syst. Nat., ed. XII, 1766, 455, on a specimen from Doctor Garden, probably from Charleston.
Plagusia fasciata Holbrook, in DeKay, New York Fauna: Fishes, 1842, 304, Charleston, S. C.
1810. *Symphurus pusillus* (Goode and Bean).
Off Atlantic coast of United States in deep water.
Aphoristia pusilla Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 590, Gulf Stream, lat. 40° N., in deep water.
1811. *Symphurus diomedeanus* (Goode and Bean).
Gulf of Mexico, off Key West, Fla.
Aphoristia diomedea Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 589, Gulf of Mexico, lat. 25° 4' 30'' N., long. 82° 59' 15'' W.
1812. *Symphurus williamsi* Jordan and Culver.
West coast of Mexico.
Symphurus williamsi Jordan and Culver, in Jordan, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 506, pl. 55, Mazatlan, Mexico.
- Genus 606. *ACEDIA* Jordan.
Acedia Jordan, in Jordan and Goss, Rept. U. S. Fish Com., XIV, 1886 (1888) (*Aphoristia nebulosa* Goode and Bean).
1813. *Acedia nebulosa* (Goode and Bean).
Gulf Stream.
Aphoristia nebulosa Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 192, Gulf Stream, off the coast of the Carolinas.

Superorder ACANTHOPTERYGII

Order LL. ZEOIDEI

Family 140. ZEIDÆ. John dories

Genus 607. ZENOPSIS Gill.

Zenopsis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 126 (*Zeus nebulosus* Temminck and Schlegel).

1814. *Zenopsis ocellata* (Storer). *Ocellated dory*.

Off New England coast.

Zeus ocellatus Storer, Proc. Bost. Soc. Nat. Hist., VI, 1858, 386, off Provincetown, Mass.

Genus 608. ZENION Jordan and Evermann.

Zenion Jordan and Evermann, Rept. U. S. Fish Com., XXI, 1895 (1896), 418 (*Cyttus hololepis* Goode and Bean).

1815. *Zenion hololepis* (Goode and Bean).

West Indies, in deep water.

Cyttus hololepis Goode and Bean, Oceanic Ichth., 1896, 225, figs. 233, 233a, 233b, off Yucatan and Little Bahama Bank.

Genus 609. OREOSOMA Cuvier and Valenciennes.

Oreosoma Cuvier, Règne Animal, ed. 2, II, 1829 (type not named, *Oreosoma atlanticum* Cuvier).

Oreosoma Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1830, 377 (515) (*O. atlanticum* Cuvier and Valenciennes).

1816. *Oreosoma atlanticum* Cuvier and Valenciennes.

Open Atlantic; only a few specimens known.

Oreosoma atlanticum Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 377 (515), open Atlantic.

Oreosoma conferrum Cuvier and Valenciennes, loc. cit., pl. 99, same type.

Family 141. GRAMMICOLEPIDÆ

Genus 610. GRAMMICOLEPIS Poey.

Grammicolepis Poey, Anal. Soc. Esp. Hist. Nat., II, 1873 (*G. brachiusculus* Poey).

1817. *Grammicolepis brachiusculus* Poey.

Cuba; in deep water.

Grammicolepis brachiusculus Poey, Anal. Soc. Española Hist. Nat., II, 1873, Cuba.

Order MM. XENOBERYCES

Family 142. STEPHANOBERYCIDÆ

Genus 611. STEPHANOBERYX Gill.

Stephanoberyx Gill, Proc. U. S. Nat. Mus., VI, 1883, 258 (*S. monæ* Gill).

1818. *Stephanoberyx monæ* Gill.

Gulf Stream, in deep water.

Stephanoberyx monæ Gill, Proc. U. S. Nat. Mus., VI, 1883, 258, Gulf Stream at Albatross station 2077 in 1,255 fathoms.

1819. *Stephanoberyx gillii* Goode and Bean.

Gulf Stream, in deep water.

Stephanoberyx gillii Goode and Bean, Oceanic Ichth., 1895, 187, fig. 206, Gulf Stream at Albatross station 2099 in 2,949 fathoms.

Family 143. MELAMPHAIDÆ

Genus 612. CAULOLEPIS Gill.

Caulolepis Gill, Forest and Stream, XXI, Aug. 30, 1883, and Proc. U. S. Nat. Mus., VI, 1883, 258 (*C. longidens* Gill).

1820. *Caulolepis longidens* Gill.
Gulf Stream and Pacific Ocean at Cortez Banks off San Diego, Calif.
Caulolepis longidens Gill, Proc. U. S. Nat. Mus., VI, 1883, 258, Atlantic Ocean, lat. 39° 27' N., long. 69° 56' 20" W., in 1,346 fathoms.
- Genus 613. *ANOPLOGASTER* Gunther.
Anoplogaster Gunther, Cat., I, 1859, 12 (*Hoplostethus cornutus* Cuvier and Valenciennes).
1821. *Anoplogaster cornutus* (Cuvier and Valenciennes).
Atlantic, in deep water, north and south.
Hoplostethus cornutus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 347 (270); from stomach of a shark, taken at 26° lat. S; others taken by the *Albatross* 31° N., 40° W., from the stomach of an albacore.
- Genus 614. *POROMITRA* Goode and Bean.
Poromitra Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882, 214 (*P. capito* Goode and Bean).
1822. *Poromitra capito* Goode and Bean.
Deep water in Gulf Stream in latitude 34°
Poromitra capito Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882, 215, Gulf Stream.
- Genus 615. *PLECTROMUS* Gill.
Plectromus Gill, Proc. U. S. Nat. Mus., VI, 1883, 257 (*P. suborbitalis* Gill).
1823. *Plectromus suborbitalis* Gill.
Gulf Stream, in deep water.
Plectromus suborbitalis Gill, Proc. U. S. Nat. Mus., VI, 1883, 258, lat. 38° 52' N., long. 69° 24' W., at *Albatross* stations 2036, 2190, and 2535 in 1,149 to 1,800 fathoms.
1824. *Plectromus beani* (Günther).
Gulf Stream in deep water.
Plectromus crassiceps Bean, Proc. U. S. Nat. Mus., VIII, 1885, 73, Gulf Stream; name preoccupied.
Melamphaes beani Günther, *Challenger* Rept., XXII, 1887, 29, Gulf Stream, in about latitude 40°.
1825. *Plectromus crassiceps* (Günther).
Mid-Atlantic; Brazil.
Scopelus crassiceps Günther, Ann. Mag. Nat. Hist., II, 1878, 185, deep water in mid-Atlantic and off Pernambuco.
1826. *Plectromus cristiceps* (Gilbert).
Coast of Washington and Oregon and southeastern Alaska.
Melamphaes cristiceps Gilbert, Proc. U. S. Nat. Mus., XIII, 1890 (1891), 60, coasts of Oregon and Washington and *Albatross* station 3075 in 859 fathoms.
1827. *Plectromus lugubris* (Gilbert).
Coast of California; off Bay of Panama; Bering Sea.
Melamphaes lugubris Gilbert, Proc. U. S. Nat. Mus., XIII, 1890 (1891), 59, coast of California south of Point Conception, at *Albatross* station 2923 in 822 fathoms.
1828. *Plectromus mizolepis* (Günther).
Off Bay of Panama in deep water; south of New Guinea and Bay of Bengal.
Scopelus mizolepis Günther, Ann. Mag. Nat. Hist., 11, 1878, 185, south of New Guinea.
1829. *Plectromus nycterinus* (Gilbert).
Coast of southern California, in deep water.
Melamphaes nycterinus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 326, pl. 16, off Santa Catalina, 2,113 to 2,259 fathoms.
1830. *Plectromus bispinosus* (Gilbert).
Coronado Islands, near San Diego, Calif.
Melamphaes bispinosus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 325, pl. 15, fig. 5; off the Coronado Islands, near San Diego, 642 to 666 fathoms.

Order NN: BERYCOIDEI

Family 144. POLYMIXIDÆ. Barbudos

Genus 616. POLYMIXIA Lowe.

Polymixia Lowe, Trans., Cambr. Phil. Soc., VI, 1838, 198 (*P. nobilis* Lowe).

Nemabrama Valenciennes, Ichth. Îles Canaries, 1844, 40 (*N. webbi* Valenciennes = *Polymixia nobilis* Lowe).

Dinemus Poey, Memorias, II, 1861, 160 (*D. venustus* Poey = *Polymixia lowei* Günther).

1831. *Polymixia lowei* Günther. Barbudo.

Caribbean Sea; Cuba.

Polymixia lowei Günther, Cat., I, 1859, 17, Caribbean Sea.

Dinemus venustus Poey, Memorias, II, 1861, 161, Cuba.

Family 145. BERYCIDÆ

Genus 617. BERYX Cuvier. *Alfonsinos*.

Beryx Cuvier, Règne Anim., ed. 2, II, 1829, 151 (*B. decadactylus* Cuvier and Valenciennes).

1832. *Beryx decadactylus* Cuvier and Valenciennes. *Alfonsin a Casta Larga*.
Cosmopolitan, open seas, in rather deep water; Portugal, Madeira, Japan, and Cuba.

Beryx decadactylus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 165 (222), Madeira or Portugal.

1833. *Beryx splendens* Lowe. *Alfonsin a Casta; Cumpri-da*.

Cosmopolitan, in the open seas, at moderate depth; Madeira, Japan, and Gulf Stream at 35° 49' 30'' N., 74° 34' 45'' W. in 424 fathoms.

Beryx splendens Lowe, Proc. Zool. Soc. Lond., 1833, 142, Madeira.

Family 146. ANOMALOPIDÆ

Genus 618. KRYPTOPHANARON Silvester and Fowler.

Kryptophanaron Silvester and Fowler, Proc. Acad. Nat. Sci. Phila., LXXVIII, 1926, 245 (*K. alfredi* Silvester and Fowler).

1834. *Kryptophanaron alfredi* Silvester and Fowler.

Off coast of Jamaica.

Kryptophanaron alfredi Silvester and Fowler, Proc. Acad. Nat. Sci. Phila., LXXVIII, 1926, 246, Pl. XVIII, fig. 1; Pl. XIX, figs. 2 and 4.

Off Kingston Harbor, Jamaica, floating on surface.

Family 147. TRACHICHTHYIDÆ

Genus 619. HOPISTETHUS Cuvier and Valenciennes.

Hoplostethus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 344 (469) (*H. mediterraneus* Cuvier and Valenciennes).

1835. *Hoplostethus mediterraneus* Cuvier and Valenciennes.

Coasts of southern Europe; Gulf Stream.

Hoplostethus mediterraneus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 344 (469), Mediterranean Sea.

Trachichthys pretiosus Lowe, Proc. Zool. Soc. Lond. 1839, 77, Madeira.

Family 148. HOLOCENTRIDÆ. Soldierfishes; Squirrelfishes

Genus 620. OSTICHTHYS Jordan and Evermann.

Ostichthys Langsdorf, in Cuvier and Valenciennes, Hist. Nat. Poiss., III, 130 (174), 1829; name only, passing reference under *Myripristis japonicus* Cuvier.

Ostichthys Jordan and Evermann, Fish. North and Mid. Amer., 1896, 846 (*Myripristis japonicus* Cuvier and Valenciennes).

1836. *Ostichthys trachypomus* (Günther).

West Indies; Cuba.

Myripristis trachypoma Günther, Cat., I, 1859, 25, Cuba.

Myriopristis fulgens Poey, Memorias, II, 1860, 160, Cuba.

Genus 621. MYRIPRISTIS Cuvier.

Myripristis Cuvier, Règne Anim., ed. 2, II, 1829, 150 (*M. jacobus* Cuvier and Valenciennes).

Rhamphoberyx Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 817 (*R. pæcilopus* Gill); young.

Myripristes of authors; altered spelling.

Myriopristis of authors; altered spelling.

1837. *Myripristis jacobus* Cuvier and Valenciennes. *Frère Jacques; Candil*. West Indies to Brazil; common.

Myripristes jacobus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 121 (162), Martinique.

Rhinoberyx chryseus Cope, Proc. Amer. Phil. Soc., XIV, 1871, 464, St. Croix Island.

1838. *Myripristis occidentalis* (Gill).

West coast of America, from Cape San Lucas to Panama; Galapagos; Cocos Island.

Myriopristis occidentalis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 87, Cape San Lucas.

Rhamphoberyx leucopus Gill, loc. cit., 88, Cape San Lucas; immature.

1839. *Myripristis pæcilopus* (Gill).

West coast of America, from Cape San Lucas to Panama.

Rhamphoberyx pæcilopus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 87, Cape San Lucas; immature example.

Genus 622. HOLOTRACHYS Günther.

Holotrachys Günther, Jour. Mus. Godeffroy, Band I, 1873, 93 (*Myripristes lima* Cuvier and Valenciennes).

1840. *Holotrachys lima* (Cuvier and Valenciennes).

South Seas, east to Hawaii and Cocos Island.

Myripristis lima Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 493, Île de France.

Myripristes humilis Kner and Steindachner, Sitz. Akad. Wiss. Wien, LIV, 1866, 357, pl. 1, fig. 1, Samoa.

Genus 623. HOLOCENTRUS Bloch. Squirrelfishes; Soldierfishes.

Holocentrus Gronow, Zoophyl., 1763, 65; nonbinomial.

Holocentrus Bloch, Naturgeschich. Ausland. Fische, IV, 1790, 61 (*H. sogo* Bloch).

Holocentrum Cuvier, and of most authors; amended spelling.

Rhynchichthys Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 503 (*R. pelamidis* Cuvier and Valenciennes); young.

Rhinoberyx Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 237 (*Rhynchichthys brachyrhynchus* Bleeker); larval form.

1841. *Holocentrus ascensionis*³⁹ (Osbeck). *Matajuelo; Squirrelfish; Soldado; Welchman*.

Islands of the Atlantic and northward from St. Helena and Ascension Island to Florida and Cuba.

?*Perca ascensionis* Osbeck, Iter Chinensis, 1757; reprinted as Reise Ost-Ind. China, 1765, 388, Ascension Island; reprint of a pre-Linnæan paper, hence ineligible.

?*Bodianus pentacanthus* Bloch, Nat. Ausl. Fische, IV, 1790, 40, pl. 225, Brazil; after *Saguaraca* of Maregrave.

?*Holocentrus sogo* Bloch, loc. cit., 61, pl. 232, Africa.

Sciæna rubra Schneider, Ed. Bloch, Syst. Ichth., 1801, 82; after Catesby.

Amphiprion matejuelo Bloch and Schneider, Syst. Ichth., 1801, 206, Cuba; after Parra.

?*Bodianus jaguar* Lacépède, Hist. Nat. Poiss., IV, 1802, 286, Brazil.

?*Holocentrum longipinne* Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 138 (185), Brazil; Martinique; Porto Rico; St. Thomas; San Domingo; Habana.

Holocentrus striatus Gronow, Cat. Fishes, 1854, 173, Antilles.

Holocentrus rostratus Gronow, loc. cit., 173, near the Equator.

³⁹ The specific name of the common matajuelo of Cuba is still wholly uncertain if the two forms (the one from Cuba, the other from Brazil) are really distinct. The earliest specific name, *ascensionis*, is apparently ineligible, occurring in reprints of a pre-Linnæan paper. The pertinence of most of the other names is more or less uncertain. Here, as elsewhere where no official decision has been made by the International Commission on Zoological Nomenclature, we follow the rule of not changing current nomenclature unless, as in matters of priority, the case is reasonably clear.

1842. *Holocentrus rufus* (Walbaum).
The northern form probably varying into *H. ascensionis*, which ranges from the Bahamas to Brazil.
Perca rufa Walbaum, Artedi Pisc., 1792, 351, Bahamas; after *Perca marina rubra* of Catesby.
1843. *Holocentrus coruscus* (Poey).
Cuba; Bahama Islands; Bermudas; Tortugas.
Holocentrum coruscum Poey, Memorias, II, 1861, 158, Cuba.
Holocentrum sicciferum Cope, Trans. Amer. Phil. Soc., XIV, 1871, 465, Bahamas, New Providence
? *Holocentrum puncticulatus* Barbour, Bull. Mus. Comp. Zool., XLVI, 1905, 117, Bermudas.
? *Holocentrus tortugæ* Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904 (1905), 236, fig. 1, Garden Key, Tortugas.
1844. *Holocentrus brachypterus* Poey.
Cuba.
Holocentrus brachypterus Poey, Repertorio, I, 1866, 184, Cuba.
1845. *Holocentrus vexillarius* (Poey).
Cuba; Porto Rico
Holocentrum vexillarium Poey, Memorias, II, 1861, 158, Cuba.
? *Holocentrum riparium* Poey, Enumeratio, 1875, 37, Cuba.
1846. *Holocentrus osculus* (Poey).
Cuba.
Holocentrum osculum Poey, Memorias, II, 1861, 156, Cuba.
Holocentrum perlatum Poey, loc. cit., 157, Cuba.
1847. *Holocentrus sancti-pauli* (Günther).
St. Paul Rocks, mid-Atlantic.
Holocentrum sancti-pauli Günther, Shore Fishes, *Challenger* Rept., Zoology, I, part 6, 1880, 4, St. Paul Rocks, mid-Atlantic.
1848. *Holocentrus meeki* Bean.
Bermudas.
Holocentrus meeki Bean, Proc. Biol. Soc. Wash., XIX, 1906, 31, St. Davids Island, Bermudas.
- Genus 624. FLAMMEO Jordan and Evermann.
Flammeo Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2871 (*Holocentrum marianum* Cuvier and Valenciennes).
1849. *Flammeo marianus* (Cuvier and Valenciennes). *Marian*.
West Indies; Cuba; Jamaica; Martinique.
Holocentrum marianum Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 164 (219), Martinique.
Holocentrum rostratum Poey, Memorias, II, 1861, 157, Cuba.
- Genus 625. ADIORYX Starks.
Adioryx Starks, Science, XXVIII, 1908, 614 (*Holocentrum suborbitale* Gill).
1850. *Adioryx suborbitalis* (Gill). *Mojarra cardinal*.
Mazatlan to Panama; Revillagigedo, Cocos, and Galapagos Islands.
Holocentrum suborbitale Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 86, Cape San Lucas, Lower California.
- Genus 626. PLECTRYPOPS Gill.
Plectrypops Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 237 (*Holocentrum retrospinis* Guichenot).
1851. *Plectrypops retrospinis* (Guichenot).
Cuba.
Holocentrum retrospinis Guichenot, in Ramon de la Sagra, Hist. Cuba, 1853, 35, pl. 1, fig. 3, Cuba.

Order 00. THORACOSTEI

Suborder HEMIBRANCHII

Family 149. GASTEROSTEIDÆ. Sticklebacks

Genus 627. GASTEROSTEUS Linnæus. *Sticklebacks*.*Gasterosteus* Linnæus, Syst. Nat., ed. X, 1758, 295 (*G. aculeatus* Linnæus).*Gasteracanthus* Pallas, Zoog. Rosso-Asiat., III, 1811 (*G. cataphractus* Pallas).*Leivurus* Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 242 (*Gasterosteus aculeatus* Linnæus).*Merrimella* Jordan, Univ. Calif. Publ. Geol., V, 1907, 131 (*M. dorysia* Jordan); fossil form1852. *Gasterosteus aculeatus* Linnæus *Common stickleback*.

Arctic and subarctic regions, in shallow bays and estuaries, very common; also ascending streams, where it loses partially or wholly its bony plates, in Europe, Asia, and America alike. The development of these plates and in some degree of dorsal spines and caudal keel depends on the salinity of the water. Reported *G. williamsoni microcephalus* from irrigation ditch near El Rosario, Lower California, 255 miles south of U. S. border, May 21, 1925, Clinton G. Abbott, collector.

(1. Forms completely mailed)

Gasterosteus aculeatus Linnæus, Syst. Nat., ed. X, 1758, 489, Europe.*Gasterosteus cataphractus* Pallas, Mém. Acad. Petersb., III, 1811, 325, Kamchatka.*Gasterosteus trachura* Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 481, France.*Gasterosteus teraculeatus* Lacépède, Hist. Nat. Poiss., III, 1802, 296.*Gasterosteus obolarius* Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 500, Kamchatka.*Gasterosteus noveboracensis* Cuvier and Valenciennes, loc. cit., 502, pl. 98, fig. 3.*Gasterosteus biaculeatus* Shaw, Zool., IV, 1839, 608; after Pennant.*Gasterosteus dimidiatus* Reinhardt, Fauna Grœnlandica, 1837, 32, 37, Greenland.*Gasterosteus loricatus* Reinhardt, loc. cit., 32, 37, Greenland.*Gasterosteus baillon* Blanchard, Poiss. des eaux douces de France, 1866, 231, Abbeville.*Gasterosteus insculptus* Richardson, Last Arctic Voyage, 1854, 10, Pl. XXV, figs. 1, 2, and 3, Northumberland and Puget Sounds.*Gasterosteus serratus* Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 47, San Francisco.

(2. Forms armed anteriorly only)

Gasterosteus gymnurus Cuvier, Règne Animal, ed. II, 1829, 170, waters of England and northward.*Gasterosteus leiurus* Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 481, Seine.*Gasterosteus semiarmatus* Cuvier and Valenciennes, loc. cit., 493, Havre, France.*Gasterosteus semuloricatus* Cuvier and Valenciennes, loc. cit., 494, Baillon.*Gasterosteus spinulosus* Jenyns and Yarrell, Man. Brit. Vert., 1835, 350, Edinburgh.*Gasterosteus ponticus* Nordmann, in Demidoff, Voy. Russ. Merid., III, 1840, 357, Tauria and Black Sea.*Gasterosteus cuvieri* Girard, in Storer's Fishes of Nova Scotia and Labrador, 1849, 254, p. VII, fig. 1, Bras d'Or, Red Bay, Labrador.*Gasterosteus microcephalus* Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 133, Four Creek (Kaweah River), a tributary to Tule Lake (Tulare Lake), San Joaquin Valley, Calif.*Gasterosteus plebeius* Girard, loc. cit., 147, Presidio, Calif.

- Gasterosteus inopinatus* Girard, loc. cit., 147, Presidio, Calif.
Gasterosteus pugetti Girard, loc. cit., 1856, 135, Fort Steilacoom, Wash.
Gasterosteus neustranus Blanchard, Poiss. des eaux douces de France, 1866, 220, Harfleur.
Gasterosteus argentissimus Blanchard, loc. cit., 232, Avignon.
Gasterosteus elegans Blanchard, loc. cit., 234, Cadillac and Langou, 6 pls.
Gasterosteus islandicus Sauvage, Revision des Épinoches, 1874, 20, Iceland.
Gasterosteus atkinsi Bean, Proc. U. S. Nat. Mus., II, 1879, 67, Schoodic Lakes, Maine.

(3. Forms naked or with one or two plates)

- Gasterosteus willamsoni* Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 133, Williamsons Pass, Calif.
Gasterosteus santa-annæ Regan, Ann. Mag. Nat. Hist., 8th ser., IV, 1909, 437, Santa Anna River, Calif.

Genus 628. *GLADIUNCULUS* Jordan and Evermann.

- Gladiunculus* Jordan and Evermann, Proc. Calif. Acad. Sci., ser. 4, XVI, 1927 (Apr. 27), 504 (*Gasterosteus gladiunculus* Kendall=*G. bispinosus* Walbaum).

1853. *Gladiunculus bispinosus* (Walbaum). *Common eastern stickleback*.
 Labrador to New Jersey, near the coast, in salt water; very common in New England.
Gasterosteus bispinosus Walbaum, Arted. Pisc., 1792, 450; after Pennant; said to be "New York," but more likely Hudson's Bay.
Gasterosteus wheatlandi Putnam, Proc., Essex Inst., V, 1867, 4, Nahant, Mass.
Gasterosteus gladiunculus Kendall, Proc. U. S. Nat. Mus., XVIII, 1896, off coast of Maine, near Seguin Island.

Genus 629. *EUCALIA* Jordan.

- Eucalia* Jordan, Man. Vert., ed. 1, 1876, 248 (*Gasterosteus inconstans* Kirtland).

1854. *Eucalia inconstans* (Kirtland). *Brook stickleback*.
 New York to Kansas and northward to the Saskatchewan; Great Lakes region; central Ohio, Indiana, and Illinois.
Gasterosteus inconstans Kirtland, Bost. Jour. Nat. Hist., III, 1841, 273, brooks of Trumbull County, Ohio.
Gasterosteus micropus Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 81, Fort Riley, Kans.
Gasterosteus globiceps Sauvage, Revision des Épinoches, 1874, 35, North America.
Eucalia inconstans cayuga Jordan, Man. Vert., ed. 1, 1876, 249, Cayuga Lake, Ithaca, N. Y.
1855. *Eucalia pygmaea* (Agassiz).
 Lake Superior.
 A valid species if the fin rays are D. IV-I, 6; A. I, 6. In *E. inconstans* they are D. IV-I, 10; A. I, 10.
Gasterosteus pygmaeus Agassiz, Lake Superior, 1850, 314, Lake Superior.

Subfamily GASTEROSTEINÆ

Genus 630. *PUNGITIUS* Costa. *Nine-spined sticklebacks*.

- Pungitius* Costa, Mém. Savants Etrangers, Paris, X, 1846 (*Gasterosteus pungitius* L.).
Pygosteus (Brevoort) Gill, Can. Nat., and Cat. Fish. E. Coast N. A., supplement to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 39 (*Gasterosteus occidentalis* Cuvier and Valenciennes=*G. pungitius* Linnæus).
Gasterosteus Sauvage, Révision des Épinoches, 1874, 7 (*Gasterosteus pungitius* Linnæus).

1856. *Pungitius pungitius* (Linnæus). *Nine-spined stickleback*.
Gasterosteus pungitius Linnæus, Syst. Nat., ed. X, 1758, 296, Europe;
 after *Gasterosteus aculeis* in dorso decem Artedi.
Gasterosteus lævis Cuvier, Règne Animal, ed. 2, II, 1829, 170, streams of
 France.
Gasterosteus occidentalis Cuvier and Valenciennes, Hist. Nat. Poiss.,
 IV, 1829, 509, Newfoundland; not *G. occidentalis* Linnæus, unrec-
 ognized, probably a species of *Carangidæ*.
Gasterosteus concinnus Richardson, Fauna Bor.-Amer., III, 1836, 57,
 Saskatchewan River and Great Bear Lake.
Gasterosteus nebulosus Agassiz, Lake Superior, 1850, 310, pl. 4, fig. 4,
 Lake Superior.
Gasterosteus dekayi Agassiz, Lake Superior, 1850, 311, New York.
Gasterosteus lotharingus Blanchard, Poiss. France, 1866, 244, Lorraine.
Gasterosteus breviceps Blanchard, loc. cit., 1866, 245, Caen, France.
Gasterosteus blanchardi Sauvage, Révision des Epinoches, 1874, 32, New
 York.
1857. *Pungitius mainensis* (Storer).
 Kennebec River, Me.; a form or species with seven dorsal spines.
Gasterosteus mainensis Storer, Bost. Jour. Nat. Hist., I, 1837, 464,
 Kennebec County, Me.
1858. *Pungitius brachypoda* (Bean).
 Baffins Bay, Greenland. A form with 10 dorsal spines and the ventral
 spines very small.
Gasterosteus pungitius brachypoda Bean, Bull. U. S. Nat. Mus., XV,
 1879, 129, Oosoodlin Mountain, Cumberland Gulf, Greenland.

Subfamily APELTINÆ

- Genus 631. *APELTES* DeKay. *Four-spined stickleback*.
Apeltes DeKay, N. Y. Fauna: Fishes, 1842, 87 (*Gasterosteus quadracus*
 Mitchell).
1859. *Apeltes quadracus* (Mitchill).
 Maine to New Jersey, in salt water only.
Gasterosteus quadracus Mitchell, Trans. Lit. and Phil. Soc., I, 1815,
 430, New York.
Gasterosteus apelles Cuvier and Valenciennes, Hist. Nat. Poiss., IV,
 1829, 505, no locality.
Gasterosteus millipunctatus Ayres, Bost. Jour. Nat. Hist., I, 1842, 294,
 Old Mans Harbor, Long Island.

Subfamily SPINACHIIINÆ

- Genus 632. *SPINACHIA* Cuvier.
Spinachia Cuvier, Règne Anim., ed I, 1817, 320 (*Gasterosteus spinachia*
 Linnæus).
Polycanthus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 242
 (*Gasterosteus spinachia* Linnæus).
Gastræa Sauvage, Révision des Epinoches, 1874, 7 (*Gasterosteus*
spinachia Linnæus).
1860. *Spinachia spinachia* (Linnæus).
 Coasts of northern Europe; recorded from Newfoundland.
Gasterosteus spinachia Linnæus, Syst. Nat., ed. X, 1758, 442.
Spinachia vulgaris Fleming, Brit. Anim., 1828, 210.
Gasterosteus marinus Gronow, Cat. Fish., 1854, 168.

Family 150. AULORHYNCHIDÆ

- Genus 633. *AULORHYNCHUS* Gill.
Aulorhynchus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 169 (*A.*
flavidus Gill).
Auliscops Peters, Berliner Monatsberichte, 1866, 570 (*A. spinescens*
 Peters).

1861. *Aulorhynchus flavidus* Gill.
Coast of California, from San Nicolas Island and Monterey northward to Sitka, Alaska.
Aulorhynchus flavidus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 169, coast of Washington.

Suborder LOPHOBRANCHII

Family 151. SYNGNATHIDÆ. Pipefishes and seahorses

Subfamily SYNGNATHINÆ

- Genus 634. *DERMATOSTETHUS* Gill.
Dermatostethus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 283 (*D. punctipinnis* Gill).
1862. *Dermatostethus punctipinnis* Gill.
San Diego, California.
Dermatostethus punctipinnis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 283, San Diego, Calif.; not lately recognized; the genus perhaps of doubtful value.
- Genus 635. *SYNGNATHUS* Linnæus.
Syngnathus Linnæus, Syst. Nat., ed. X, 1858, 336 (*Syngnathus acus* L., the proper type as decided by the Internat. Com. Zool. Nomencl.).
Siphostoma Rafinesque, Caratteri, 1810, 18 (*Syngnathus pelagicus* Linnæus).
1863. *Syngnathus californiensis* Storer. *Great pipefish*.
Santa Barbara to Monterey and northward.
Syngnathus californiensis Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 73, California.
1864. *Syngnathus griseolineatus* Ayres.
Pacific coast of the United States, from Monterey northward to Puget Sound and southeastern Alaska.
Syngnathus griseolineatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 14, San Francisco Bay.
Syngnathus abboti Girard, Pac. R. R. Survey, Fishes, X, 1858, 346, San Francisco.
1865. *Syngnathus leptorhynchus* Girard.
West coast of America from San Francisco to San Diego.
Syngnathus brevirostris Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 156, San Diego; name preoccupied.
Syngnathus leptorhynchus Girard, loc. cit., 156, San Diego, Calif.
Syngnathus dimidiatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 284, San Diego; substitute for *brevirostris*, preoccupied.
1866. *Syngnathus arundinaceus* Girard.
Coast of California.
Syngnathus arundinaceus Girard, Pac. R. R. Surv., X, 1858, 346, coast of California.
1867. *Syngnathus carinatus* (Gilbert).
Gulf of California.
Siphostoma carinatum Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 547, Gulf of California, lat. 31° 31' N., long. 114° 19' W., at Albatross stations 3027 and 3028.
1868. *Syngnathus auliscus* (Swain).
Southern California.
Siphostoma auliscus Swain, Proc. U. S. Nat. Mus., V, 1882, 310, Santa Barbara and San Diego, Calif.
1869. *Syngnathus barbaræ* (Swain).
California coast at Santa Barbara. Rare; not reported recently.
Siphostoma barbaræ Swain, Proc. U. S. Nat. Mus., VII, 1884, 238, Santa Barbara, Calif.

1870. *Syngnathus mackayi* (Swain and Meek).
Key West to Yucatan.
Siphostoma mackayi Swain and Meek, Proc. U. S. Nat. Mus., VII,
1884, 239, Key West, Fla.
1871. *Syngnathus floridæ* (Jordan and Gilbert).
Virginia to Texas and Porto Rico.
Siphostoma floridæ Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
263, Pensacola, Fla.
1872. *Syngnathus poeyi* (Jordan and Evermann).
Habana.
Syngnathus tenuis Poey, Synopsis, 1867, 444, Cuba; name preoccupied.
Siphostoma poeyi Jordan and Evermann, Fish. North and Mid. Amer.,
1896, 766, Habana, Cuba; substitute for *S. tenuis*, preoccupied.
1873. *Syngnathus elucens* ⁴⁰ Poey. *Pipefish*.
Cuba; Florida Keys; Porto Rico
Syngnathus elucens Poey, Synopsis, 1867, 443, Habana.
Syngnathus flavirostris Poey, Enumeratio, 1876, 178, Habana.
Syngnathus picturatus Poey, loc. cit., 178, Habana.
Syngnathus linea Poey, loc. cit., 178, Habana.
Syngnathus marmoreus Poey, loc. cit., 178, Habana.
Syngnathus ascendens Poey, loc. cit., 179, Habana.
1874. *Syngnathus rousseau* Kaup.
West Indies; known from Jamaica, St. Lucia, and Martinique.
Syngnathus rousseau Kaup, Lophobranchii, 1856, 40, Martinique.
1875. *Syngnathus jonesi* Gunther. *West Indian pipefish*.
Bermudas and Porto Rico.
Syngnathus jonesi Gunther, Ann. Mag. Nat. Hist., ser. 4, XIV, 1874,
8, 455, Bermudas.
1876. *Syngnathus fistulatus* Peters.
Puerto Cabello, near Colon.
Syngnathus fistulatus Peters, Monatsber., Ak. Wiss. Berl., 1868, 456,
Puerto Cabello, near Aspinwall.
1877. *Syngnathus pelagicus* Linnæus.
Open sea in the Atlantic and the Mediterranean; doubtfully recorded
from the West Indies; some other species, perhaps *S. fistulata*, being
mistaken for it.
Syngnathus pelagicus Linnæus, Syst. Nat., ed. I, 1758, open sea in
floating seaweed.
Syngnathus ethon Risso, Eur. Mérid., III, 1826, 182, Nice, France.
1878. *Syngnathus robertsi* (Jordan and Rutter).
West Indies and Jamaica; Caribbean Sea.
Siphostoma robertsi Jordan and Rutter, Proc. Acad. Nat. Sci. Phila.,
XLIX, 1897, 97, Jamaica.
1879. *Syngnathus brachycephalus* Poey.
Tortugas; Cuba.
Syngnathus brachycephalus Poey, Synopsis, 1867, 444, Habana, Cuba.
1880. *Syngnathus brevicaudus* (Meek).
Boca del Rio, Vera Cruz.
Siphostoma brevicaudum Meek, Field Col. Mus., Zool., V, 1904, 163,
Boca del Rio, Mexico.
1881. *Syngnathus crinigerum* (Bean and Dresel).
Pensacola and Key West, to Abrolhos Reef, Brazil.
Siphostoma crinigerum Bean and Dresel, Proc. Biol. Soc. Wash., II,
1884, 99, Pensacola, Fla.
1882. *Syngnathus sinaloæ* (Jordan and Starks).
Pacific coast of Mexico.
Siphostoma sinaloæ Jordan and Starks, Proc. Calif. Acad. Sci., 3d ser.,
VI, 1896, 268, Mazatlan, Mexico.

* Some of the nominal species here included may prove to be valid.

1883. *Syngnathus arctus* (Jenkins and Evermann).
Gulf of California, Guaymas.
Siphostoma arctum Jenkins and Evermann, Proc. U. S. Nat. Mus., XI,
1888, 137, Bay of Guaymas, Sonora, Mexico.
1884. *Syngnathus starksi* (Jordan and Culver).
Estuaries about Mazatlan.
Siphostoma starksi Jordan and Culver, Proc. Calif. Acad. Sci., 3d ser.,
V, 1895, 416, Pl. XXX, Rio Presidio, Mazatlan, Mexico.
1885. *Syngnathus dendriticus* (Barbour).
Ireland Island, Bermudas.
Siphostoma dendriticum Barbour, Bull. Mus. Comp. Zool., XLVI,
No. 7, 1904-1906, 115, Ireland Island, Bermudas.
1886. *Syngnathus mindii* Meek and Hildebrand.
Panama.
Syngnathus mindii Meek and Hildebrand, Field Mus. Nat. Hist., Zool.,
XV, 1923, 261, Pl. XVIII, fig. 2; brackish creek near Mindi, Canal
Zone, Panama.
1887. *Syngnathus elcapitensis* Meek and Hildebrand.
Streams, west coast of Panama.
Syngnathus elcapitanense Meek and Hildebrand in Meek, Field Mus.
Nat. Hist., Zool., X, 1914, 119, Rio Mamona at El Capitan, Panama.
1888. *Syngnathus tweedliei* Meek and Hildebrand.
Panama.
Syngnathus tweedliei Meek and Hildebrand, Field Mus. Nat. Hist.,
Zool., XV, 1923, 259, Pl. XVIII, fig. 1, Chame Point, Panama.
1889. *Syngnathus torrei* (Nichols).
Cuba.
Siphostoma torrei Nichols, Bull. Amer. Mus. Nat. Hist., XXXI, 1912,
183, fig. 1, Matanzas, Cuba.
- Genus 636. SYRICTES Jordan and Evermann.
Syricies Jordan and Evermann Proc. Calif. Acad. Sci., XVI, Apr. 27,
1927, 504 (*Syngnathus fuscus* Storer).
1890. *Syricies fuscus*⁴¹ (Storer). *Common pipefish; Green pipefish; Billfish.*
Atlantic coast of United States from Maine to Virginia; very common
northward.
Syngnathus fuscus Storer, Rept. Fish. Mass., 1839, 162, Nahant, Mass.
Syngnathus peckianus Storer, Rept. Fish. Mass., 1839, 163, Holms
Hole, Marthas Vineyard.
Syngnathus peckii Storer, Rept. Fish. Mass., 1839, 3; name only.
Syngnathus fasciatus DeKay, N. Y. Fauna: Fishes, 1842, 319, pl. 54,
fig. 174, New York.
Syngnathus viridescens DeKay, loc. cit., 321, pl. 54, fig. 176, New York.
Syngnathus dekeyi Duméril, Hist. Nat. Poiss., II, 1870, 569, New York.
Syngnathus milbertianus Duméril, loc. cit., 573, New York.
1891. *Syricies affinis* (Günther).
Gulf of Mexico, on coast of Louisiana.
Syngnathus affinis Günther, Cat., VIII, 1870, 163, Louisiana.
1892. *Syricies scovelli* (Evermann and Kendall).
Coast of Texas and both coasts of Florida.
Siphostoma scovelli Evermann and Kendall, Proc. U. S. Nat. Mus.,
XVIII, 1895, 113, Shamrock Point, Corpus Christi, Tex.
1893. *Syricies louisianæ* (Günther).
Atlantic and Gulf coasts of the United States; North Carolina to Texas
and south to Key West.
Syngnathus pelagicus, var., Linnæus, Syst. Nat., ed. XII, 1766, 416,
Charleston, S. C.
Syngnathus louisianæ Günther, Cat., VIII, 1870, 160, New Orleans, La.
1894. *Syricies bairdianus* (Duméril).
West coast of Lower California.
Syngnathus bairdianus Duméril, Hist. Nat. Poiss., II, 1870, 574, "Coast
of Mexico, near California."

⁴¹ The names *fuscus* and *peckianus* are of the same date. The author who first perceived this identity chose the second name in spite of the page priority of *fuscus*.

- 1895. *Syrristes exilis*** (Osborn and Nichols).
Gulf of California.
Siphostoma exile Osborn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 153, fig. 7, west San Benito Island, Port San Bartolome.
- Genus 637. HIPPICHTHYS** Bleeker.
Hippichthys Bleeker, Verh. Batav. Genootsch., XXII, 1849, 15 (*H. heptagonus* Bleeker); not *Hippocampus heptagonus* Rafinesque.
Corythoichthys Kaup, Archiv Naturgesch., 1853, 231 (*C. albirostris* Heckel MS. Kaup).
- 1896. *Hippichthys albirostris*** (Heckel).
West Indies, north to Pensacola and Key West; south to Bahia, about coral reefs.
Corythoichthys albirostris Heckel in Kaup, Archiv Naturgesch., 1853, 231, Mexico; Bahia; type examined.
Siphostoma zatropis Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 264, Snapper Banks, Pensacola.
- 1897. *Hippichthys cayorum*** (Evermann and Kendall).
Key West, Fla.; Porto Rico.
Corythoichthys cayorum Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 128, pl. 7, fig. 7, near Crawfish Bar, Key West, Fla.
- 1898. *Hippichthys spicifer*** (Rüppell).
Tropical Pacific; recorded from Tehuantepec.
Syngnathus spicifer Rüppell, Neue Wirb., Fische, about 1830, 143.
- Genus 638. DORYCHTHYS** Kaup.
Doryichthys Kaup, Archiv Naturgesch., 1853, 233 (*D. bilineatus* Heckel).
- 1899. *Doryichthys lineatus*** Valenciennes.
Tropical parts of Atlantic from Cuba to Brazil.
Doryichthys lineatus Valenciennes, in Kaup, Lophobranchii, 1856, 59, Bahia, Mexico, and Guadeloupe.
- Genus 639. DORYRHAMPHUS** Kaup.
Doryrhampus Kaup, Archiv Naturgesch., 1853, 233 (*D. excisus* Kaup).
Choeroichthys Kaup, loc. cit., 1853, 233 (*C. valenciennesi* Kaup).
- 1900. *Doryrhampus californiensis*** Gill.
Gulf of California.
Doryrhampus californiensis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 284, Cape San Lucas, Lower California.
- 1901. *Doryrhampus sierra*** Nichols.
Porto Rico.
Doryrhampus sierra Nichols, Bull. Amer. Mus. Nat. Hist., XXXIV, 1915, 145, fig. 2, Porto Rico.
- Genus 640. ACUS**⁴² Swainson.
Acus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 333 (*Syngnathus æquoreus* "Yarrell").
Osphyolax Cope, Proc. Acad. Nat. Sci. Phila., XXVII, 1875, 450 (*O. pellucidus* Cope).
- 1902. *Acus æquoreus*** (Linnæus). *Ocean pipefish*.
Open sea along coasts of Europe; recorded from New Orleans, some other species probably mistaken for it.
Syngnathus æquoreus Linnæus, Syst. Nat., ed. X, 1758, 417, open sea.
Syngnathus sabbaldi Walbaum, Artedi Pisc., 1792, 6, Scotland.
Nerophis maculata Rafinesque, Indice, 1810, 57, Palermo, Italy.
Nerophis anguineus Kaup, Lophobr., 1856, 65, Brownsey, Dorsetshire; Berwick.
- 1903. *Acus martinicensis*** (Bibron) Kaup.
West Indies.
Nerophis martinicensis (Bibron) Kaup, Lophobr., 1856, 67, Martinique.

⁴² We may adopt the generic name *Acus* for the species of the old genus *Nerophis* (Rafinesque, Indice, Sicil., 1810, 57, type *Syngnathus ophidion* L.) in which a small caudal fin is present, the tail not ending in a filament, as is the case in the type of *Nerophis* (*Syngnathus ophidion* L.). The name *Acus*, variously used by authors, is not preoccupied in binomial nomenclature.

1904. *Acus heckeli* (Kaup).
Caribbean Sea.
Nerophis heckeli Kaup, Lophobr., 1856, 66, Bogota; evidently not the exact locality.
1905. *Acus pellucidus* (Cope).
Open Atlantic.
Osphrolax pellucidus Cope, Proc. Acad. Nat. Sci. Phila., XXVII, 1875, 450, pl. 25, figs. 1 to 4, open Atlantic Ocean; a larval form of doubtful relationships and differing from *Acus* in the very short dorsal (D. 16; about 40 in *Acus*).
- Subfamily HIPPOCAMPINÆ
- Genus 641. **NANNOCAMPUS** Gunther.
Nannocampus Gunther, Cat., VIII, 1870, 178 (*N. subosseus* Gunther).
1906. *Nannocampus nanus* Rosen.
Bahama Islands.
Nannocampus nanus Rosen, Arsskr. Univ. Lund, VII, 1912, 50, Bahamas.
- Genus 642. **HIPPOCAMPUS** Rafinesque.
Hippocampus Rafinesque, Caratteri, 1810, 18 (*Syngnathus hippocampus* Linnaeus).
1907. *Hippocampus hudsonius* DeKay. *Common American seahorse; Horsefish.*
Atlantic coast of United States, from Cape Cod to the Tortugas.
Hippocampus hudsonius DeKay, N. Y. Fauna: Fishes, 1842, 322, p. 53, fig. 171, New York.
Hippocampus lævicaudatus Heckel, in Kaup, Lophobranchii, 1856, 16, North America.
1908. *Hippocampus ingens* Girard. *Caballito del mar.*
Pacific coast from San Diego to Mazatlan; rare.
Hippocampus ingens Girard, Pac. R. R. Surv., Fishes, X, 1858, 342, San Diego, Calif.
Hippocampus gracilis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 282, Cape San Lucas, Lower California.
1909. *Hippocampus brunneus* Bean.
Bermudas.
Hippocampus brunneus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 32, Long Bird Island, Bermudas.
1910. *Hippocampus punctulatus* Guichenot. *Caballito del mar; Seahorse; Horsefish.*
Tropical parts of the Atlantic; common in the West Indies, Brazil, and western Africa; Gulf Stream; Beaufort, N. C.
Hippocampus punctulatus Guichenot in Ramon de la Sagra, Hist. Île. Cuba, Poiss., 1853, 174, pl. 5, fig. 2, Cuba.
Hippocampus marginalis Heckel, in Kaup, Lophobr., 1856, 15, Mexico.
Hippocampus fasciularis Heckel, in Kaup, loc. cit., 15, Mexico.
1911. *Hippocampus stylifer* Jordan and Gilbert.
Coasts of Florida; Snapper Banks off Pensacola and Tampa; Gulf Stream.
Hippocampus stylifer Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 265, Snapper Banks, Gulf of Mexico.
1912. *Hippocampus zosteræ* Jordan and Gilbert.
Pensacola Bay, Fla.
Hippocampus zosteræ Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 265, Grand Lagoon, Pensacola, Fla.; smallest of the seahorses,

Order PP. AULOSTOMI

Family 152. AULOSTOMIDÆ. Trumpetfishes

- Genus 643. **AULOSTOMUS** Lacépède.
Aulostomus Lacépède, Hist. Nat. Poiss., V, 1803, 357 (*A. chinensis* Lacépède).
Polyptericthys Bleeker, Nat. Tijdschr., Neder-Indie, IV, 1853, 608 (*P. valentini* Bleeker).
Aulostoma Schlegel, change of spelling.

1913. *Aulostomus maculatus* Valenciennes. *Trompetero*; *Trumpetfish*.
Caribbean Sea, north to southern Florida.
? *Aulostomus maculatum* Valenciennes, in Cuvier, Règne Anim., III,
Poiss., 1845, pl. 92, fig. 2.
Aulostoma coloratum Muller and Troschel, in Schomburgk, History
Barbados, 1848, 673, Barbados.
1914. *Aulostomus cinereus* (Poey).
Cuba.
Aulostoma cinereum Poey, Synopsis, 1867, 386, Cuba.

Family 153. FISTULARIIDÆ. Cornetfishes

- Genus 644. FISTULARIA Linnæus.
Fistularia Linnæus, Syst. Nat., ed. X, 1758, 312 (*F. tabacaria* Linnæus).
Cannorhynchus Cantor, Jour. Roy. Asiat. Soc. Bengal, XVIII, 1849
(*Fistularia tabacaria* Linnæus); substitute name, *Fistularia* having
been used in 1750 as a pre-Linnæan name for a group of polyps.
Flagellaria Gronow, Cat. Fishes, 1854 (*F. fistularis* Gronow = *Fistularia*
tabacaria Linnæus).
Solenostomus (Gronow) Gill, Cat. Fish. East Coast. N. A., Supl. to
Proc. Acad. Nat. Sci. Phila., XIII, 1861, 38 (*Fistularia tabacaria*
Linnæus).
1915. *Fistularia tabacaria* Linnæus *Trumpetfish*; *Trompetero*; *Tobacco*
trumpetfish; *Unarmed trumpetfish*.
West Indies and neighboring seas; northward to Carolina and Florida,
and occasionally to Long Island.
Fistularia tabacaria Linnæus, Syst. Nat., ed. X, 1758, 312, tropical
America.
Fistularia neo-boracensis Mitchell, Trans. Lit. Phil. Soc. Phila., I,
1815, 437, New York.
Aulastome marcgravi Castelnau, Anim. Nouv. Amer. Sud, 1855, 30,
Bahia, Rio Janeiro.
Flagellaria fistularis Gronow, Cat. Fish., 1854, 146, American Ocean.
1916. *Fistularia serrata* ⁴³ Cuvier. *Flutemouth*.
Western Pacific and other warm seas; Bermuda; Cuba.
Fistularia serrata Cuvier, Règne Anim., ed. 1, II, 1817, 389, 34; after Bloch.
? *Fistularia immaculata* Cuvier, loc. cit., 349, Sea of the Indies.
Fistularia commersoni Ruppell, Neue Wirbelthiere, 1834, 142, Red Sea.
Fistularia villosa Klunzinger, Abh. Zool. Ges. Wien, XXI, 1871, 516,
Red Sea.
1917. *Fistularia petimba* Lacépède. *Corneta*.
East Indies; Australia; China; Panama; Lower California; Japan;
Hawaii.
Fistularia petimba Lacépède, Hist. Nat. Poiss., V, 1803, 349, New
Britain, Isle of Reunion, equatorial Pacific.
Fistularia depressa Günther, *Challenger* rept. Shore Fishes, Zoology, I,
pt. 6, 1880, 69, pl. 32, fig. D, Sulu Archipelago, station 200, 255
fathoms.
1918. *Fistularia corneta* Gilbert and Starks.
Pacific coast of tropical America; Panama.
Fistularia corneta Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904,
56, Pl. X, Bay of Panama; Mazatlan.

Family 154. MACRORHAMPHOSIDÆ. Snipefishes

- Genus 645. MACRORHAMPHOSUS Lacépède. *Snipefishes*.
Macrorhamphosus Lacépède, Hist. Nat. Poiss., V, 1803, 136 (*Silurus*
cornutus Forskål = *Balistes scolopax* Linnæus).
Centriscus Cuvier, Règne Animal, ed. 1, II, 1817, 350; not *Centriscus*
Linnæus.
Macrognathus Gronow, Cat. Fishes, 1854, 147 (*Centriscus scolopax*
Linnæus).

⁴³ Weber calls this species *villosa* but quotes *serrata*? in synonymy. The synonymy of *serrata* and *petimba* seem hopelessly confused.

1919. *Macrorhamphosus scolopax* (Linnæus). *Snipefish; Trumpetfish; Bellowsfish*.
Mediterranean Sea, northward to southern England, accidental on our North Atlantic Coast.
Balistes scolopax Linnæus, Syst. Nat., ed. X, 1758, 329, Mediterranean.
Silurus cornutus Forskål, Descr. Anim., 1775, 66, Marseilles.
1920. *Macrorhamphosus hawaiiensis* Gilbert.
Deep waters about Hawaii; once taken off San Pedro, Calif. (W. F. Thompson).
Macrorhamphosus hawaiiensis Gilbert, Bull. U. S. Fish Com., XXIII, 1903 (1905), 613, Laysan Island.

Order QQ. PERCOMORPHI

Suborder PERCESOCES

Family 155. ATHERINIDÆ. Silversides; Hardheads; Pescados del Rey; Peixe Rey

Genus 646. HEPSETIA Bonaparte.

Hepsetia Bonaparte, Fauna Italica, Fasc. 91 (no pages), 1837 (*Atherina boyeri* Risso).

Athernomorus Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 730 (*Atherina laticeps* Poey = *Atherina stipes* Muller and Troschel).

1921. *Hepsetia stipes* (Muller and Troschel). *Cabezote, Brisle herring.*

West Indies; common north to western Florida.

?*Atherina tæmata* Spix in Agassiz, Pisc. Bras., 1829, 135, pl. 30, fig. 2, Atlantic Ocean, off Brazil.

Atherina stipes Muller and Troschel, in Schomburgk, Hist. Barbados, 1847, 671, Barbados.

Atherina laticeps, Poey, Memorias, II, 1861, 265, Habana.

Atherina velutana Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 342, Clear Water Harbor, Fla.

1922. *Hepsetia evermanni* (Eigenmann).

Cuba; San Cristobal; Pinar del Rio.

Atherina evermanni Eigenmann, Bull. U. S. Fish Com., XXII, 1902, 228, fig. 9, San Cristobal, Cuba.

Genus 647. ATHERINA (Artedi) Linnæus. *Friars.*

Atherina Artedi in Linnæus, Syst. Nat., ed. X, 1758, 315 (*Atherina hepsetus* Linnæus).

Aphia Risso, Europ. Mérid. 1826 (*Aphia meridionalis* Risso = apparently very young of *Atherina hepsetus* L.), but Risso describes their laying of eggs, and may refer to some other species.

Tæniomembras Ogilby, Proc., Linn. Soc. New South Wales, XXIII, 1898, 41 (*Atherina microstoma* Günther).

1923. *Atherina aræa* Jordan and Gilbert.

Gulf of Mexico at Key West, Cozumel, and south to Caribbean Sea.

Atherina aræa Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 27, Key West, Fla.

1924. *Atherina harringtonensis* Goode.

Bermudas and south to Curaçao.

Atherina harringtonensis Goode, Amer. Jour. Sci. Arts, 3d ser., XIV, No. 82, 1877, 297, Bermuda Islands.

1925. *Atherina microps* Poey.

Habana.

Atherina microps Poey, Memorias, II, 1861, 266, Habana, Cuba.

1926. *Atherina carolina* Cuvier and Valenciennes.

South Carolina (reference doubtful).

Atherina carolina Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1835, 330 (445), "Carolinas."

Genus 648. MENIDIA Bonaparte.

Menidia Bonaparte, Fauna Italica, Tome III, Pesci, in description of *Atherina hepsetus* p. 91, 1836, no type, *Atherina menidia* Linnaeus intended.

Ischnomembras Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 730, pl. 42, "Gabun River, Gabun County, West Africa," probably an error.

Phozargyrea Fowler, loc. cit., 1903, 732, *Phozargyrea dayi* Fowler from "India," doubtless an error.

Subgenus ISCHNOMEMBRAS Fowler.

1927. *Menidia beryllina* (Cope). *Silverside; Fresh-water silverside.*
Coastwise streams, Massachusetts to Florida and lower Mississippi, north to Tennessee.
? *Atherinichthys gracilis* Günther, Cat., III, 1861, 405, no locality given.
Christoma beryllinum Cope, Trans. Amer. Phil. Soc., 1866, 403, Potomac River, near Washington.
Menidia audens Hay, Bull. U. S. Fish Com., II, 1882, 64, Memphis, Tenn., and Vicksburg, Miss.
Ischnomembras gabunensis Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 731, pl. 42, Gabon River, West Africa; probably an error.
1928. *Menidia beryllina cerea* Kendall.
Atlantic coast, Massachusetts to South Carolina.
Menidia beryllina cerea Kendall, Rept. U. S. Fish Com., XXVII, 1901, 261, Waquoit Bay, Mass.
1929. *Menidia peninsulæ* (Goode and Bean).
Florida and Gulf coast, in salt water.
Chirostoma peninsulæ Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 148, Pensacola.
1930. *Menidia peninsulæ atrimentis* Kendall.
Florida; South Lake, Lake Monroe, Salt Lake, etc.
Menidia peninsulæ atrimentis Kendall, Rept. U. S. Fish Com., XXVII, 1901, 258, South Lake, Fla.
1931. *Menidia menidia* (Linnaeus).
Atlantic coast, Cape Hatteras to Florida.
Atherina menidia Linnaeus, Syst. Nat., ed. XII, 1766, 519, Charleston, S. C.
Atherina boscii Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1835, 345 (465), Charleston.
Menidia dentex Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 429, St. Johns River, Fla.
1932. *Menidia conchorum* Hildebrand and Ginsberg.
Key West, Fla.
Menidia conchorum Hildebrand and Ginsberg, Bull. U. S. Bur. Fish., XLII, 1926 (1927), 207, fig. 1, Boca Chica, Key West, Fla.
1933. *Menidia notata* (Mitchill). *Silversides; Friar; Whitebait.*
Atlantic coast of United States, Nova Scotia to Cape May, intergrading south with *M. menidia*.
Atherina notata Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 446, New York.
Atherina viridescens Mitchell, loc. cit., 447, New York.
Phozargyrea dayi Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 732, pl. 41, "India"; no doubt an error in locality.
1934. *Menidia starksi* Meek and Hildebrand.
Panama.
Menidia starksi Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 267, Pl. XX, fig. 2, Taboga Island, Panama.
1935. *Menidia sardina* (Jenkins and Evermann). *Pez del rey.*
Gulf of California.
Atherina sardina Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 137, Guaymas, Sonora, Mexico.

1936. *Menidia clara* Evermann and Jenkins.
Gulf of California.
Menidia clara Evermann and Jenkins, Proc. U. S. Nat. Mus., XIII, 1891, 136, Guaymas, Sonora, Mexico.
- Genus 649. ARCHOMENIDIA Jordan and Hubbs.
Archomenidia Jordan and Hubbs, Rev. Silversides, 1919, 54 (*Atherinichthys saltæ* Regan).
1937. *Archomenidia sallei* (Regan).
Mexico.
Atherinichthys sallei Regan, Proc. Zool. Soc. Lond., 1903, 60, Mexico.
- Genus 650. HUBBESIA Jordan.
Hubbesia Jordan, Proc. U. S. Nat. Mus., LV, 1920 (May 2, 1919), 310 (*Menidia gilberti* Jordan and Bollman).
1938. *Hubbesia gilberti* (Jordan and Bollman).
Panama.
Menidia gilberti Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 155, Panama.
- Genus 651. MEMBRAS Bonaparte.
Membras Bonaparte, Fauna Italica, fasc. 91 (not paged), 1837 (type not named directly, *Atherina martinica* Cuvier and Valenciennes being indicated).
1939. *Membras martinicus* (Cuvier and Valenciennes).
Martinique.
Atherina martinica Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1835, 340 (459), Martinique.
1940. *Membras beani* (Meek and Hildebrand).
Atlantic waters of Panama.
Kirtlandia beani Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 270, Pl. XXI, Fox Bay, Colon.
- Genus 652. KIRTLANDIA Jordan and Evermann.
Kirtlandia Jordan and Evermann, Fishes North and Mid. Amer., 1896, 794 (*Chirostoma vagrans* Goode and Bean).
1941. *Kirtlandia vagrans* (Goode and Bean). *Rough silverside*.
Gulf of Mexico; Florida to Texas; occasionally northward.
Chirostoma vagrans Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 148, Pensacola, Fla.
1942. *Kirtlandia lacinata* (Swain). *Silverfish; Whitebait; Silversides*.
Lower Chesapeake to South Carolina, intergrading with *K. vagrans*.
Menidia vagrans lacinata Swain, in Jordan and Gilbert, Synopsis, 1882, 908, Beaufort, N. C.
- Genus 653. THYRINA Jordan and Culver.
Thyrina Jordan and Culver, in Jordan, Proc. Calif. Acad. Sci., 3d ser., V, 1895, 419 (*Atherinella evermanni* Jordan and Culver).
Melaniris Meek, Field Mus.—Zool., III, 1902, 117 (*Melaniris balsanus* Meek).
1943. *Thyrina crystallina* Jordan and Culver.
Rio Presidio near Mazatlan, Sinaloa.
Thyrina crystallina Jordan and Culver, in Jordan, Proc. Calif. Acad. Sci., 3d ser., V, 1895, 420, Rio Presidio, below Presidio, Sinaloa.
1944. *Thyrina evermanni* Jordan and Culver.
West coast of Mexico; Mazatlan.
Thyrina evermanni Jordan and Culver, Proc. Calif. Acad. Sci., 3d ser., V, 1895, 419, Mazatlan, Mexico.
1945. *Thyrina guija* Hildebrand. *Pepesca; Manjuda; Alfiler; Rovalote*.
Lake Guija, Rio del Desague near Lake Guija and Rio Lempa at Suchitoto and San Marcos, El Salvador.
Thyrina guija Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925, 264, fig. 18, Lake Guija, El Salvador.

1946. *Thyrina balsana* (Meek).
Mexico, Rio Balsas, Guerrero.
Melaniris balsanus Meek, Pub. Field Col. Mus., Zool., III, 1902, 117,
Rio Balsas, at Balsas.
1947. *Thyrina guatemalensis* (Gunther).
Lakes of Huamachal.
Atherinichthys guatemalensis Gunther, Proc. Zool. Soc. London, 1864,
151, Lakes of Huamachal, Pacific coast of Guatemala.
1948. *Thyrina sardina* (Meek). *Sardina*.
Lakes Nicaragua and Managua.
Melaniris sardina Meek, Field Col. Mus., Zool., VII, 1907, 114, Lake
Managua, Nicaragua.
Chirostoma guatemalensis Gill, Proc. Acad. Nat. Sci. Phila., XXIX,
1877, 187, Lake Nicaragua; not of Gunther.
1949. *Thyrina chagresi* (Meek and Hildebrand).
Streams of Atlantic slope of Panama.
Menidia chagresi Meek and Hildebrand, Field Mus. Nat. Hist., Zool.,
X, 1914, 119, Rio Chagres at Gorgona, Zent and Parismina, Costa
Rica.
1950. *Thyrina meeki* Miller.
Rio Motagua, Guatemala.
Thyrina meeki Miller, Bull. Amer. Mus. Nat. Hist., XXIII, 1907,
110, fig. 2, Rio Motagua, east coast of Guatemala.
- Genus 654. **THYRINOPS** Hubbs.
Thyrinops Hubbs, Proc. Acad. Nat. Sci. Phila., LXIX, 1918, 306
(*Atherinichthys pachylepis* Günther).
1951. *Thyrinops pachylepis* (Gunther).
Panama.
Atherinichthys pachylepis Günther, Proc. Zool. Soc. Lond., 1864, 25,
Panama.
- Genus 655. **MELANORHINUS** Metzelaar.
Melanorhinus Metzelaar, Tropisch Atl. Visschen, 1919, 38 (*M. boeki*
Metzelaar).
1952. *Melanorhinus boeki* Metzelaar.
West Indies.
Melanorhinus boeki Metzelaar, Trop. Atl. Visschen, 1919, 38, Oyster
Pond, St. Martin Island.
- Genus 656. **XENATHERINA** Regan.
Xenatherina Regan, Biologia Centrali Amer., Pisces, 1906, 64 (*Menidia*
lisa Meek).
1953. *Xenatherina lisa* (Meek).
Streams of Vera Cruz, Mexico.
Menidia lisa Meek, Field Col. Mus., Pub. Zool., V, 1904, 182, Refugio,
Vera Cruz, Basin of Rio Papaloapam, Mexico.
- Genus 657. **ATHERINELLA** Steindachner.
Atherinella Steindachner, Ichth. Beitr., II, 1875, 35 (*A. panamensis*
Steindachner).
1954. *Atherinella panamensis* Steindachner.
Panama.
Atherinella panamensis Steindachner, Ichth. Beitr., II, 1875, 35,
Panama.
- Genus 658. **EURYSTOLE** Jordan and Evermann.
Eurystole Jordan and Evermann, Fishes North and Mid. Amer., 1896,
802 (*Atherinella eriarcha* Jordan and Gilbert).
1955. *Eurystole eriarcha* (Jordan and Gilbert).
West coast of Mexico; Mazatlan.
Atherinella eriarcha Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
348, Mazatlan, Mexico.

Genus 659. CHIROSTOMA Swainson. *Pescados blancos.*

Chirostoma Swainson, Class. Fishes, etc., II, 1839, 243 (*Atherina humboldtiana* Cuvier and Valenciennes).

Atherinoides Bleeker, Verh. Bat. Gen., Japan, XXV, 1853, 40 (*Chirostoma vomerina* Cuvier and Valenciennes=*Atherina humboldtiana* Cuvier and Valenciennes).

Atherinichthys Bleeker, loc. cit., 40 (*Atherina humboldtiana* Cuvier and Valenciennes).

Heterognathus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 198 (*Atherina humboldtiana* Cuvier and Valenciennes).

Lethostole Jordan and Evermann, Fish. North and Mid. Amer., 1896, 792 (*Chirostoma estor* Jordan).

Eslopsarum Jordan, Rept. U. S. Fish Com., XIX, 1895, 330 (*Chirostoma jordani* Woolman).

Subgenus ESLOPSARUM Jordan.**1956. *Chirostoma jordani* Woolman.**

Basin of Rio Lerma and Valley of Mexico.

Chirostoma jordani Woolman, Bull. U. S. Fish Com., XIV, 1894, 62, pl. 2, canals at Salamanca and City of Mexico.

Atherinichthys brevis Therese von Baiern, Anz. Kais. Akad. Wiss. Wien, 1894, 149, Lake Cuitzeo, Mexico.

1957. *Chirostoma mezquital* Meek.

Durango.

Chirostoma mezquital Meek, Field Col. Mus., pub. 93, V, 1904, 170, Durango, Mexico.

1958. *Chirostoma labarcæ* Meek.

Rio Lerma, Mexico.

Chirostoma labarcæ Meek, Field Col. Mus., Zool., III, 1902, 113, Rio Lerma, Mexico.

1959. *Chirostoma arge* (Jordan and Snyder).

Rio Verde, Mexico.

Eslopsarum arge Jordan and Snyder, Bull. U. S. Fish Com., XIV, 1899, 133, fig. 12, Rio Verde, Mexico.

Subgenus CHIROSTOMA Swainson.**1960. *Chirostoma bartoni* Jordan and Evermann.**

Tributary of the Rio Lerma, near Guanajuato, Mexico; Lake Patzcuaro.

Chirostoma bartoni Jordan and Evermann, Fish. North and Mid. Amer., 1896, 793, tributary of the Rio Lerma, near Guanajuato, Mexico.

Chirostoma attenuatum Meek, Field Col. Mus., Zool., III, 1902, 112, Lake Patzcuaro, Mexico.

1961. *Chirostoma zirahuen* Meek.

Lago de Zirahuen, Mexico.

Chirostoma zirahuen Meek, Field Col. Mus., Zool., III, 1902, 114, Lago de Zirahuen, Mexico.

1962. *Chirostoma patzcuaro* Meek.

Lago de Patzcuaro, Mexico.

Chirostoma patzcuaro Meek, Field Col. Mus., Zool., III, 1902, 113, Lago de Patzcuaro, Mexico.

1963. *Chirostoma humboldtianum* (Cuvier and Valenciennes). *Pescado blanco de Chalco.*

Lakes near the City of Mexico; La Laguna, Jalisco, Mexico.

Atherina humboldtiana Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1835, 355 (479), Mexico.

Atherina vomerina Cuvier and Valenciennes, loc. cit., 357 (481), lake near City of Mexico.

1964. *Chirostoma regani* Jordan and Hubbs.

Valley of Mexico; Lake Patzcuaro.

Chirostoma regani Jordan and Hubbs, Leland Stanford Univ. Pub., Univ. Ser., 1919, 74, Xocho milco, Morelos, Mexico. After *Chirostoma breve* Regan; not of Steindachner.

Subgenus LETHOSTOLE Jordan and Evermann.

1965. *Chirostoma chapalæ* Jordan and Snyder.
Laguna de Chapala, Mexico.
Chirostoma chapalæ Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 135, fig. 13, Laguna de Chapala, Mexico, at Ocotlan.
1966. *Chirostoma consocium* Jordan and Hubbs.
Basin of Rio Lerma, Mexico.
Chirostoma consocium Jordan and Hubbs, Rev. Silversides, in Leland Stanford Univ. Pub. Univ. Ser., 1919, 76, Lake Chapala at La Palma, Mexico.
1967. *Chirostoma grandocule* (Therese von Baiern).
Lake Patzcuaro, Mexico.
Atherinichthys grandoculis Therese von Baiern, Anz. Kais. Akad. Wiss. Wien, 1894, Lake Patzcuaro, Mexico.
1968. *Chirostoma promelas* Jordan and Snyder.
Laguna de Chapala; Guadalajara, Mexico.
Chirostoma promelas Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 136, market of Guadalajara, Mexico, from Laguna de Chapala.
1969. *Chirostoma lucius* Boulenger.
Lake Chapala, Mexico.
Chirostoma lucius Boulenger, Ann. and Mag. Nat. Hist., ser. 7, V, January, 1900, 54, Lake Chapala, Mexico.
Chirostoma crystallinum Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1900), 139, fig. 16, Laguna de Chapala, Mexico, at Ocotlan.
1970. *Chirostoma diazi* Jordan and Snyder.
Lake Chapala, Mexico.
Chirostoma diazi Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1900), 137, fig. 15, Lake Chapala, Mexico.
1971. *Chirostoma sphyraena* Boulenger.
Lake Chapala, Mexico.
Chirostoma sphyraena Boulenger, Ann. Mag. Nat. Hist., ser. 7, V, 1900, 54, Lake Chapala, Mexico.
Chirostoma lermæ Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1900), 142, fig. 19.
1972. *Chirostoma ocotlane* Jordan and Snyder.
Lago de Chapala, Mexico.
Chirostoma ocotlane Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1900), 140, Lago de Chapala, Mexico.
1973. *Chirostoma estor* Jordan. *Pescado blanco de Chapala*.
Lake Chapala, Lake Patzcuaro, Lake Zirahuen, and Lake Xochimilco, Mexico.
Chirostoma estor Jordan, Proc. U. S. Nat. Mus., II, 1879, 298, Lake Chapala (an error for Lake Patzcuaro).
Atherinichthys albus Therese von Baiern, Anz. Kais. Akad. Wiss. Wien, 1894, 148, Lake Patzcuaro, Mexico.
- Genus 660. LABIDESTHES Cope.
Labidesthes Cope, Proc. Amer. Phil. Soc., XI, 1870, 455 (*Chirostoma sicculum* Cope).
1974. *Labidesthes sicculus* (Cope). *Brook silverside; Skipjack; Glassfish*.
Lake Ontario and southern Michigan to Iowa, Florida, and Texas, clear streams and lakes.
Chirostoma sicculum Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 81, Grosse Isle, Detroit River, Mich.
- Genus 661. LEURESTHES Jordan and Gilbert. *Grunion*.
Leuresthes Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 29 (*Atherinopsis tenuis* Ayres).
1975. *Leuresthes tenuis* (Ayres). *Grunion*.
Sandy coasts of California from San Francisco to Lower California; abundant southward.
Atherinopsis tenuis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 75, San Francisco.
Leuresthes crameri Jordan and Evermann, Fishes North and Mid. Amer., 1896, 802, Ballenas Bay, Lower California.

- Genus 662. **MUGILOPS** Meek and Hildebrand.
Mugilops Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 271 (*M. cyanellus* Meek and Hildebrand).
1976. **Mugilops cyanella** Meek and Hildebrand.
 Panama Bay.
Mugilops cyanellus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 271, Pl. XXII, fig. 1, Balboa, Panama Bay.
1977. **Mugilops marina** Meek and Hildebrand.
 Panama Bay.
Mugilops marinus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 272, Pl. XXII, fig. 2, Porto Bello, Panama Bay.
- Genus 663. **COLPICHTHYS** Hubbs.
Colpichthys Hubbs, Proc. Acad. Nat. Sci. Phila., LXVIX, 1917 (1918), 67 (*Atherinops regis* Jenkins and Evermann).
1978. **Colpichthys regis** (Jenkins and Evermann). *Pez del rey*.
 Gulf of California.
Atherinops regis Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 138, Bay of Guaymas, Sonora, Mexico.
- Genus 664. **ATHERINOPS** Steindachner. *Pascadillos del rey*.
Atherinops Steindachner, Ichth. Beitr., III, 1875, 61 (*Atherinopsis affinis* Ayres).
1979. **Atherinops affinis** (Ayres). *Little smelt; Pescadillo del rey*.
 Coast of California, and north to Oregon.
Atherinopsis affinis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 73, San Francisco.
Atherinops oregonia Jordan and Snyder, Proc. U. S. Nat. Mus., XLV, 1913, 575, pl. 46, Yachats River, Oreg.
1980. **Atherinops insularum** Gilbert.
 San Clemente and San Nicolas Islands of the Santa Barbara Group; Guadalupe Island.
Atherinops insularum Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 549, San Clemente, San Nicolas, and Guadalupe Islands.
1981. **Atherinops insularum cedroscensis** Hubbs.
 Shores of Cedros and San Benito Islands, Lower California.
Atherinops insularum cedroscensis Hubbs, Bull. Amer. Mus. Nat. Hist., XXXVIII, 1918, 434, southeast side of Cedros Island, Lower California.
1982. **Atherinops insularum guadalupæ** Hubbs.
 Guadalupe Island, off Lower California.
Atherinops insularum guadalupæ Hubbs, Bull. Amer. Mus. Nat. Hist., XXXVIII, 1918, 435, Guadalupe Island.
- Genus 665. **ATHERINOPSIS** Girard. *Pescado del rey*.
Atherinopsis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 134 (*A. californiensis* Girard).
1983. **Atherinopsis californiensis** Girard. *California blue smelt; Pescado del rey; Peize rey; Pesce rey*.
 Coast of California, from Cape Mendocino to Ballenas Bay, Lower California; abundant.
Atherinopsis californiensis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 134, San Francisco, Calif.
1984. **Atherinopsis sonoræ** Osburn and Nichols.
 Cedros Island, Lower California.
Atherinopsis sonoræ Osborn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 156, fig. 8, Cedros Island, Lower California.

Family 156. MUGILIDÆ. Mulletts

- Genus 666. **MUGIL** Linnæus. *Striped mullets* (soft dorsal and anal not scaled).
Mugil Artedi, in Linnæus, Syst. Nat., Ed. X, 1758, 316 (*M. cephalus* Linnæus).
Arnion Gistel, Thierreich, 1848, X (*M. cephalus* L.).

1985. *Mugil brasiliensis* Agassiz. *Liza*; *Lebranco*; *Queriman*; *Blueback mullet*; *Fantail mullet*.
Cuba to Patagonia; West Indies; Brazil; Habana, probably north to Virginia.
Mugil brasiliensis Agassiz, in Spix, Pisc. Brasil, 134 (234 by mispaging), 1829, pl. 72, Atlantic Ocean off Brazil.
Mugil liza Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 61 (83), Brazil; Porto Rico; Maracaibo; Surinam; Martinique.
Mugil lebranchius Poey, Memorias, II, 1861, 260, plate 18, fig. 3 Cuba.
Querimana gyraus Jordan and Gilbert, Proc., U. S. Nat. Mus., VII, 1884, 26, Key West; "the whirligig mullet" a little fish, with two anal spines, now regarded as the young of *Mugil*, but if so of *M. brasiliensis* as the scales are 28 to 29 (41 in *Mugil cephalus*). It is recorded from Woods Hole and Fortress Monroe; in revolving schools, perhaps mixed with the young of *M. cephalus*.
1986. *Mugil cephalus* Linnæus. *Common mullet*; *Striped mullet*; *Céfalo*; *Macho*; *Machuto*; *Lisa cabezuda*; *Callyfaver mullet*.
Cosmopolitan; coasts of southern Europe and northern Africa; Atlantic coast of America from Cape Cod to Brazil; Pacific coast of America from Monterey to Chile; Hawaii; Japan; entering rivers. Perhaps more than one species included, but we fail to detect differences. If different, the American species must be known as *Mugil albula* of the Atlantic and *Mugil mexicanus* of the Pacific.
Mugil cephalus Linnæus, Syst. Nat., Ed. X, 1758, 316, Europe; based on Artedi.
Mugil albula Linnæus, Syst. Nat., Ed. XII, 1766, 520, Charleston, S. C.
? *Mugil tang* Bloch, Ichthyologia, 1794, 171, Africa.
? *Mugil plumieri* Bloch, loc. cit., 173, St. Vincent; on a drawing by Plumier.
Mugil lineatus Mitchill, in Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 71 (96), New York.
Mugil rammelsbergii Tschudi, Fauna Peruana, Ichth., 1845, 20, Peru.
Mugil berlandieri Girard, U. S. and Mex. Bound. Surv., 1859, 20, pl. 10, figs. 1 to 4, St. Josephs Island, Indianola, Brazos Santiago; Brazos; Galveston; all on coast of Texas.
Mugil guentheri Gill, Proc. Acad. Nat. Sci. Phila., XXI, 1863, 169, western coast of Central America; not of Steindachner.
Mugil mexicanus Steindachner, Ichth. Beitr., III, 1875, 58, Acapulco, Mexico.
- Genus 667. *QUERIMANA* Jordan and Gilbert. *White mullets*. (Soft dorsal and anal closely scaled.)
Querimana Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 588 (*Myxus harengus* Günther, now regarded as the young of *Mugil curema*).
1987. *Querimana curema* (Cuvier and Valenciennes). *White mullet*; *Blueback mullet*; *Lisa*; *Lisa blanca*.
Both coasts of America, Cape Cod to Brazil and Mazatlan to Chile.
Mugil curema Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 64 (87), Brazil; Martinique; Cuba.
Mugil petrosus Cuvier and Valenciennes, loc. cit., 65 (89), Brazil; Surinam; Gulf of Mexico; Cuba.
Myxus harengus Günther, Cat., III, 1861, 467, Pacific coast of Central America; young examples with two anal spines.
1988. *Querimana incilis* (Hancock). *Trench mullet*.
Waters of Central America; Rio Chagres to Pará and Bahia.
Mugil incilis Hancock, Quart. Jour. Sci., 1830, 127, Guiana.
Mugil guentheri Steindachner, Ichth. Notizen, I, 1864, 12, British Guiana.
1989. *Querimana thoburni* (Jordan and Starks).
Pacific coast of tropical America, from Guatemala to Galapagos Islands.
Mugil thoburni Jordan and Starks, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 812, Galapagos Islands.

1990. *Querimana hospes* (Jordan and Culver). *Lisita*.
West coast of Mexico to Panama Bay.
Mugil hospes Jordan and Culver, in Jordan, Proc. Calif. Acad. Sci., V,
1895, 422, pl. 31, Mazatlan, Mexico.
1991. *Querimana gaimardiana* (Desmarest). *Redeye mullet; Liza ojo de Perdiz*.
Florida Keys to Cuba; Key West.
Mugil gaimardianus Desmarest, Dict. Class., 1831, pl. 109, Cuba.
1992. *Querimana setosus* (Gilbert).
Clarion Island of the Revillagigedo Group.
Mugil setosus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 549, Clarion Island.
1993. *Querimana trichodon* (Poey). *Fantail mullet*.
Florida Keys to Brazil; Key West.
Mugil trichodon Poey, Ann. Lyc. Nat. Hist. N. Y., XI, 1875, 66, pl. 8, figs. 4-8, Cuba.
- Genus 668. *CHÆNOMUGIL* Gill.
Chænomugil Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 169 (*Mugil proboscideus* Gunther).
1994. *Chænomugil proboscideus* (Günther).
Pacific coast of tropical America; Mazatlan; Cordova; Panama.
Mugil proboscideus Günther, Cat., III, 1861, 459, Island of Cordova (Cardon), west coast of Central America.
- Genus 669. *DAJAUS* Cuvier and Valenciennes. *Mountain mullets*.
? *Agonostomus* Bennet, Proc. Zool. Soc. Lond., I, 1830, 105 (*A. telfairii* Bennett).
Dajaus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 122 (164) (*Mugil monticola* Bancroft).
Neomugil Vaillant, Bull. Soc. Philom., Paris, 1894, 73 (*N. digueti* Vaillant).
1995. *Dajaus monticola* (Bancroft). *Trucha; Mountain mullet*.
Fresh waters of the West Indies and the Atlantic and Pacific streams of Mexico and Central America.
Mugil monticola Bancroft, in Griffith's ed. Cuvier's Animal Kingdom, Fishes, 1836, 367, pl. 36, Jamaica.
Mugil irretitus Gosse, Nat. Sojourn Jamaica, 1851, 84, Jamaica.
Agonostoma nasutum Günther, Cat., III, 1861, 463, Rio Geronimo, Guatemala.
Neomugil digueti Vaillant, Bull. Soc. Philom., IV, 1894, 73, Lower California.
Agonostomus salvini Regan, Ann. Mag. Nat. Hist., ser. XIX, 1907, 7, and Biol. Cent. Amer., Pisces., 1907, 68, Pl. XI, fig. 2, Nacasil, Guatemala.
1996. *Dajaus percoides* (Gunther).
San Domingo.
Agonostoma percoides Gunther, Cat., III, 1861, 464, San Domingo; perhaps not distinct from *Dajaus monticola*.
1997. *Dajaus microps* (Gunther).
West Indies and Central America.
Agonostoma microps Günther, Cat., III, 1861, 462; probably West Indies.
- Genus 670. *JOTURUS* Poey.
Joturus Poey, Memorias, II, 1861, 263 (*J. pichardi* Poey).
Xenorhynchichthys Regan, Ann. Mag. Nat. Hist., II, 1908, 461 (*Joturus stipes* Jordan and Gilbert).
1998. *Joturus pichardi* Poey. *Joturo*.
Rivers of Cuba; at the foot of waterfalls.
Joturus pichardi Poey, Memorias, II, 1861, 263, Rio Almendares, near Habana.

1999. *Joturus globiceps* (Günther). *Bobo*.
Rivers and cataracts of eastern Mexico, from Jalapa to Costa Rica and Panama.
Agonostoma globiceps Gunther, Ann. Mag. Nat. Hist., ser. 4, XIV, 1874, 370, Myzantia, Vera Cruz.
Joturus stipes Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 373, Rio Bayano, near Panama.

Family 157. SPHYRÆNIDÆ. Barracudas

Genus 671. SPHYRÆNA Röse.

- Sphyræna* Röse, in Artedi Piscium, 1793, 52 (*Esox sphyræna* Linnæus).
Sphyræna Artedi, in Bloch and Schneider, Syst. Ichth., 1801, 109 (*Esox sphyræna* Linnæus).
Agriosphyræna Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 749 (*Esox barracuda* Walbaum).

Subgenus SPHYRÆNA Röse.

2000. *Sphyræna sphyræna* (Linnæus). *European barracuda; Spet; Sennet*.
Coasts of southern Europe and neighboring islands, west to the Bermudas.
Esox sphyræna Linnæus, Syst. Nat., Ed. X, 1758, 313, Mediterranean Sea.
Esox spet Hatuy, Encycl. Metod. Poiss., 1787; after Linnæus.
Sphyræna vulgaris Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 242 (327), Mediterranean.
Sphyræna viridensis Cuvier and Valenciennes, loc. cit., 251 (339), St. Jago, Cape Verde Islands.
- 2001. *Sphyræna argentea* Girard. *California barracuda; Barracouta*.
Pacific coast, from San Francisco southward to Cape San Lucas.
Sphyræna argentea Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 144 San Diego, Calif.
Sphyræna lucasana Gill, loc. cit., XV, 1863, 86, Cape San Lucas.
2002. *Sphyræna ensis* Jordan and Gilbert. *Vicuda*.
Gulf of California to Panama.
Sphyræna ensis Jordan and Gilbert, Bull. U. S. Fish Com., II, 1882, 106, Mazatlan, Mexico.
2003. *Sphyræna borealis* DeKay. *Northern barracuda; Sennet*.
Atlantic coast of the United States, from Cape Cod to North Carolina.
Sphyræna borealis DeKay, N. Y. Fauna: Fishes, 1842, 39, pl. 60, New York.
2004. *Sphyræna guachancho* Cuvier and Valenciennes. *Guaguanche pelón; Guanuanche*.
West Indies north to Pensacola; Gulf Stream to Woods Hole.
Sphyræna guachancho Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 252 (342), Habana.
Sphyræna guntheri Haly, Ann. Mag. Nat. Hist., XV, 1875, 270, Colon.
2005. *Sphyræna picudilla* Poey. *Picudilla; Baracuda*.
West Indies, on the coasts of Cuba; Bahía.
Sphyræna picudilla Poey, Memorias, II, 1861, 162; Habana.
- Subgenus AGRIOSPHYRÆNA Fowler.
2006. *Sphyræna barracuda* (Walbaum). *Great barracuda; Picuda; Becuna; Short barracuda*.
West Indies and Brazil, north to Pensacola, Charleston, and the Bermudas; accidental to Woods Hole, Mass.
Umbra minor, etc. (the barracuda) Catesby, Fishes of Carolina, etc., II, 1743, 1, pl. 1, Bahamas.
Esox barracuda Walbaum, Artedi Piscium, III, 1792, 94; after Catesby.
Sphyræna sphyræna var. *picuda* Bloch and Schneider, ed. Bloch, Syst. Ichth., 1801, 110, Habana; after Parra.
Sphyræna becuna Lacépède, Hist. Nat. Poiss., V, 1803, pl. 9, fig. 3; after drawing by Plumier at Martinique.

Suborder RHEGNOPTERI

Family 158. POLYNEMIDÆ. Threadfins

Genus 672. POLYNEMUS Linnæus. *Barbudos*.

Polynemus (Gronow) Linnæus, Syst. Nat., Ed. X, 1758, 317, (*P. quinquarius*, *virginicus*, and *paradiseus* Linnæus); properly restricted by Gunther and by general usage to *Polynemus paradiseus* Linnæus.

Polydactylus Lacépède, Hist. Nat. Poiss., V, 1803, 419 (*P. plumieri* Lacépède = *Polynemus virginicus* Linnæus).

Trichidion Klein, in Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 274 (*Polynemus virginicus* Linnæus).

2007. *Polynemus approximans* Lav and Bennett. *Raton*

Pacific coast of tropical America; Guaymas to Panama; Santa Catalina and San Diego.

Polynemus approximans Lay and Bennett, Beechey's Voyage, Zool., Fish., 1839, 57, Mazatlan, Mexico.

Polynemus californiensis Thominot, Bull. de la Soc. Philom. de Paris, Séance, 27 Juin, 1886, (7), X, 161, California.

2008. *Polynemus virginicus* Linnæus *Barbudo; Barbu; Catfish*.

West Indies, north to the Florida Keys.

Polynemus virginicus Linnæus, Syst. Nat., Ed. X, 1758, 317, America.

Polydactylus plumieri Lacépède, Hist. Nat. Poiss., V, 1803, 419, Martinique; from a drawing by Plumier.

Polynemus mango Lacépède, loc. cit., 413, America, after Linnæus.

Polynemus americanus Cuvier and Valenciennes, loc. cit., III, 1829, 291 (393), Cayenne, San Domingo, Martinique.

Polynemus oligodon Günther, Cat., II, 1860, 322, Rio Janeiro.

2009. *Polydactylus opercularis* (Gill).

Pacific coast of tropical America, from Cape San Lucas to Panama.

Trichidion opercularis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 168, Cape San Lucas.

Polynemus melanopoma Günther, Trans. Zool. Soc. Lond., VI, 1868, 421, San José de Guatemala.

2010. *Polynemus octonemus* Girard. *Threadfin; Eight-threaded threadfish*.

New York to the Rio Grande.

Polynemus octonemus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 167, Brazos Santiago; Galveston.

Trichidion octofilis Gill, loc. cit., XIII, 1861, 280, New York.

Genus 673. PENTANEMUS Günther.

Pentanemus Günther, Cat., II, 1860, 331 (*Polynemus quinquarius* Linnæus).

2011. *Pentanemus quinquarius* (Linnæus).

Atlantic Ocean; West Indies to coast of Africa; rare.

Polynemus quinquarius Linnæus, Syst. Nat., Ed. X, 1758, 317, America; after Gronow.

Polynemus artedii Bennett, Proc. Zool. Soc. Lond., 1831, 146, Africa.

Polynemus macronemus Pel, Bydrage tot de Dierk., 1851, 9, Africa.

Series SCOMBRIFORMES

Family 159. ACANTHOCYBIIDÆ

Genus 674. ACANTHOCYBIUM Gill. *Petos*.

Acanthocybium Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 125 (*Cybium sara* Bennett).

2012. *Acanthocybium petus* (Poey). *Peto*.

West Indies north to Key West; recorded from Italy.

Cybium petus Poey, Memorias, II, 1861, 234, pl. 16, fig. 1, Habana.

Cybium verany Döderlein, Giorn. di Sci. Nat., ed. Econ., VIII, 1872, Palermo.

2013. *Acanthocybium solandri* (Cuvier and Valenciennes). *Ono; Peto*.

Warmer parts of the Pacific; Hawaii; Pacific coast of Mexico; and South Seas.

Cybium solandri Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 192, open seas; exact locality unknown; after Solander.

Family 160. CYBIIDÆ⁴⁴

Genus 675. SIERRA Fowler.

Sierra Fowler, Proc. Acad. Nat. Sci. Phila., LVI, 1904 (1905), 766
(*Cybiium cavalla* Cuvier).

2014. *Sierra cavalla* (Cuvier) *Sierra*; *Kingfish*; *Cavalla*.

Tropical Atlantic; Florida Keys and Charleston; Cape Cod; Africa;
Brazil.

Cybiium cavalla Cuvier, Règne Animal, ed. 2, II, 1829, 200; after *Guarapuca* of Maregrave.

Cybiium immaculatum Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 140 (191), no locality.

Cybiium caballa Cuvier and Valenciennes, loc. cit., 187, Brazil.

Genus 676. SCOMBEROMORUS Lacépède.

Scomberomorus Lacépède, Hist. Nat. Poiss., III, 1802, 292 (*S. plumieri* Lacépède=*Scomber regalis* Bloch).

Polypterus Rafinesque, Anal. de la Nat., 1815, 89; substitute for *Scomberomorus*.

Chromitra Lockington, Proc. Acad. Nat. Sci. Phila., XXXI, 1879, 133
(*C. concolor* Lockington).

2015. *Scomberomorus maculatus* (Mitchill). *Spanish mackerel*; *Spaniard*;
Spotted cybium; *Bay mackerel*; *Spotted mackerel*

East coast of America from Cape Ann to Brazil; common in Gulf of
Mexico but not recorded from Cuba

Scomber maculatus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 426,
New York; not of Couch.

2016. *Scomberomorus sierra* Jordan and Starks. *Sierra*.

Pacific coast of tropical America.

Scomberomorus sierra Jordan and Starks, in Jordan, Proc. Calif. Acad.
Sci., V, 1895, 428, Mazatlan, Mexico.

2017. *Scomberomorus regalis* (Bloch). *Sierra*; *Pintado*; *Cero*; *Sier*; *Siering*;
Searer; *Spotted cero*; *Kingfish*; *Black-spotted Spanish mackerel*.

Cape Cod to Brazil; Cuba.

Scomber regalis Bloch, Naturgesch. ausl. Fische, VII, 1793, 38, Marti-
nique; after a drawing by Plumier.

Scomberomorus plumieri Lacépède, Hist. Nat. Poiss., III, 1802, 292,
Martinique.

Cybiium acervum Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831,
136 (186), Martinique; Cuba; St. Domingo.

2018. *Scomberomorus concolor* (Lockington). *Monterey Spanish mackerel*.
Coast of California.

Chromitra concolor Lockington, Proc. Acad. Nat. Sci. Phila., XXXI,
1879, 133, Monterey, Calif.

Family 161. SCOMBRIDÆ

Genus 677. SCOMBER Linnæus.

Scomber Artedi, in Linnæus, Syst. Nat., ed. X, 1758, 297 (*S. scombrus*
Linnæus).

Cordylus Gronow, Cat. Fish., 1854, 163 (*Scomber scombrus* L.).

2019. *Scomber scombrus* Linnæus. *Common mackerel*; *Boston mackerel*;
Wawwhunnekesuog; *Caballa*.

North Atlantic; Norway and Labrador, south to Spain and Cape
Hatteras.

Scomber scombrus Linnæus, Syst. Nat., Ed. X, 1758, 297, Atlantic.

Scomber vernalis Mitchill, Trans. Lit. and Phil. Soc. N. Y., I, 1815,
423, Sandy Hook, N. J.

Genus 678. PNEUMATOPHORUS Jordan and Gilbert.

Pneumatophorus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
593 (*Scomber pneumatophorus* De la Roche=*Scomber colias* Gmelin).

⁴⁴ The family name *Cybiidæ* may be applied to the allies of *Scomberomorus* and *Cybiium*. The name may be preferred to one derived from *Scomberomorus*, as the genus *Cybiium* is also valid. *Cybiium comersoni*, the Asiatic type of *Cybiium* Cuvier, has but two gillrakers, while in *Scomberomorus* there are 8 to 11 below the angle. *Acanthocybiium* (giant fishes with the reticulated gills of a swordfish) should constitute a separate family. The earliest of the many fossils of this type are mainly allies of *Scomberomorus*.

2020. *Pneumatophorus grex* (Mitchill). *Chub mackerel*; *Tinker mackerel*; *Easter mackerel*; *Thimble-eyed mackerel*; *Spanish mackerel of England*. The small mackerel called by the above names is very abundant on our Atlantic coast. It is very close to the similar species of Europe, *Scomber colias* Gmelin.
?Scomber colias Gmelin, Syst. Nat., 1788, 1329, Sardinia; after Cetti.
?Scomber lacertus Walbaum, Artedi Pisc., 1792, 209, Sardinia; after Cetti.
?Scomber pneumatophorus De la Roche, Ann. Mus. Nat. Hist., XIII, 1809, 315, Balearic Islands.
?Scomber macrophthalmus Rafinesque, Indice, 1810, 20 and 53, Palermo.
Scomber grex Mitchill, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 422, New York.
?Scomber maculatus Couch, Mag. Nat. Hist., V, 1832, 22, England.
?Scomber undulatus Swainson, Nat. Hist. Fishes, Amphib., etc., II, 1839, 409, Sicily.
?Scomber gracilis Swainson, loc. cit., 410, Sicily.
Scomber dekaysi Storer, Fishes Mass., 1867, 130, Massachusetts.
2021. *Pneumatophorus diego* (Ayres).
 Coast of California; very abundant southward. Perhaps not separable from the Japanese species, *Scomber japonicus*; some small differences apparently permanent.
?Scomber japonicus Houttuyn, Verhand. Holl. Maatsch. Haarl., XX, 1782, 331, Japan.
?Scomber auratus Houttuyn, loc. cit., 333, Japan.
Scomber diego Ayres, Proc., Calif. Acad. Nat. Sci., I, 1856, 101, Santa Barbara, Calif.
2022. *Pneumatophorus australasicus* (Cuvier and Valenciennes).
 North coast of Australia, from Moreton Bay, Queensland to Lord Howe Island, Hawaii and Socorro Island off the west coast of Mexico.
Scomber australasicus Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 49, Australia.

Family 162. KATSUWONIDÆ. The victor fishes

- Genus 679. *KATSUWONUS* Kishinouye. *Oceanic bonito*.
Katsuwonus Kishinouye, Scombroid Fish. Japan, Coll. Agric. Tokyo, VIII, 1923, 452 (*Scomber pelamis* L.).
2023. *Katsuwonus pelamis* (Linnæus). *Oceanic bonito*.
 Open Atlantic; rarely north to Cape Cod and Bermudas on the Atlantic coast.
Scomber pelamis Linnæus, Syst. Nat., ed. X, 1758, 297, "in Pelago inter Tropicos."
Scomber pelamides Lacépède, Hist. Nat. Poiss., III, 1802, 14; after Linnæus.
2024. *Katsuwonus vagans* (Lesson). *Aku*; *Pacific bonito*.
 Pacific Ocean, Hawaii, southern California, and Japan; inordinately abundant in open sea.
Thynnus vagans Lesson, Voy. Coquille, Zool., II, 1828, 162, pl. 32.
Thynnus affinis Cantor, Cat. Malay Fishes, 1849, 106.
- Genus 680. *EUTHYNNUS* Lütken.
Euthynnus Lütken, Jordan and Evermann, Fishes North and Mid. Amer., 1896, 868 (*Thynnus thunnina* Cuvier and Valenciennes = *Scomber alletteratus* Rafinesque).
 This genus proves to be quite distinct from *Gymnosarda*, with which it has been confused. *Gymnosarda* is an Asiatic ally of *Sarda*.
2025. *Euthynnus alletteratus* (Rafinesque). *Little tunny*; *Bonito*.
 Warm seas; north in Gulf Stream to Cape Cod; rather common at New York in summer: West Indies; Mediterranean; Japan.
Scomber alletteratus Rafinesque, Caratteri, 1810, 46, Palermo.
Thynnus leachianus Risso, Eur. Mérid., III, 1826, 414, Nice.
Scomber quadripundatus Geoffrey St. Hilaire, Descr. Egypt. Poiss., 1827, pl. 24, fig. 3, Egypt.
Thynnus thunnina Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 76, (104), pl. 212, Mediterranean.
Thynnus brasiliensis Cuvier and Valenciennes, loc. cit., 80 (110), Brazil.
Thynnus brevipinnis Cuvier and Valenciennes, loc. cit., 81 (112), Mediterranean.

2026. *Euthynnus yaito* Kishinouye. *Kawakawa; Yaito*.
Japan; Hawaii; Pacific coast of Mexico.
Euthynnus yaito Kishinouye, Scombroïd Fishes, 1923, 420, Japan.
2027. *Euthynnus lineatus* Kishinouye.
Pacific coast of Mexico.
Euthynnus lineatus Kishinouye, Sici. Gakko, No. 111, 1920, 113; and
in Scombroïd Fishes, 1923, 430, fig 2, Manzanillo, Mexico.
- Genus 681. *AUXIS* Cuvier. *Frigate mackerels*.
Auxis Cuvier, Règne Anim., ed. 2, II, 1829, 199 (*Scomber rochei* Risso =
Scomber thazard Lacépède).
2028. *Auxis rochei* (Risso). *Frigate mackerel*.
Open Atlantic, at times northward to Cape Cod; perhaps not separable
from *A. thazard* (Lacépède) of the South Seas.
Scomber rochei Risso, Ichth. Nice, 1810, 165, Nice.
Scomber bisus Rafinesque, Caratteri, 1810, 45, Palermo.

Family 163. THUNNIDÆ. Tunnies and albacores

- Genus 682. *SARDA* Cuvier. *Bonitos*.
Sarda Cuvier, Règne Animal, ed. 2, II, 1829, 199 (*Scomber pelamys*
Brunnich; not of Linnæus = *Scomber sarda* Bloch).
Pelamys Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 149
(*Scomber sarda* L.); name preoccupied
Palamita Bonaparte, Ittiologia Generale, 1831, 107, substitute for
Pelamys.
Creotroctes Gistel, Thierreich, X, 1848 (*Scomber sarda* L.); substitute
for *Pelamys*.
2029. *Sarda sarda* (Bloch). *Bonito; Bone jack; Bone-eater; Skipjack*.
Atlantic ocean, on both coasts, north to Cape Cod; very abundant.
Scomber sarda Bloch, Naturgesch. Ausl. Fische, VII, 1793, 44, Europe.
Scomber mediterraneus Bloch and Schneider, Syst. Ichth., 1801, 23,
Marseilles; after Brunnich.
Scomber pelamitus Rafinesque, Caratteri, 1810, 44, pl. 2, Palermo.
2030. *Sarda lineolata* ⁴⁵ (Girard). *California bonito; Skipjack*.
California, Patagonia and Japan; if *S. orientalis* is not distinct.
? *Pelamys chiliensis* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII,
1831, 118 (163), Valparaiso.
? *Pelamys orientalis* Temminck and Schlegel, Fauna Japon., Poiss., 1844,
99, pl. 52, Japan.
Pelamys lineolata Girard, Pac. R. R. Surv., X, 1858, 106, San Diego,
Calif.
2031. *Sarda velox* Meek and Hildebrand.
Panama.
Sarda velox Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV,
1923, 320, pl. 24, Panama market. Reported by Nichols and
Breder from Block Island, R. I.; probably an error.
- Genus 683. *THUNNUS* South. *Great tunnies*.
Thynnus Cuvier, Règne Anim., ed. 1, 1817, 313 (*Scomber thynnus* L.);
name preoccupied in Lepidoptera.
Thunnus South, Encyclop. Metropol., V, 1845, 620 (*Scomber thynnus*
Linnæus); substitute for *Thynnus*.
Albacora Jordan, Man. Vert., ed. V, 1888, 106 (*Scomber thynnus* L.).
Orcynus Cooper, Proc., Calif. Acad. Nat. Sci., III, 1863, 77 (*Scomber*
thynnus L.); not *Orcynus* Gill, 1861, the name originally a misprint
for *Orcynus*.
2032. *Thunnus saliens* Jordan and Evermann. *Leaping tuna*.
Eastern Pacific; lower California, north to mouth of Columbia River;
the type locality Santa Catalina Island.
Thunnus saliens Jordan and Evermann, Occ. Papers, Calif. Acad. Sci.,
XII, Sept. 30, 1926, 10, Santa Catalina Island, Calif.

⁴⁵ This species is well separated from *Sarda sarda* of the Atlantic. *Sarda orientalis* of Japan and Hawaii is nearly related, and both may be inseparable from *S. chilensis* of Chile, as has been assumed by Jordan and Evermann (1896).

2033. *Thunnus secundodorsalis* Storer.
North Atlantic; New York to Labrador.
Thunnus secundodorsalis Storer, Hist. Fish. Mass., 1875, 163, Pl. XXII,
fig 1, Provincetown, Mass.
2034. *Thunnus subulatus* (Poey).
West Indies; Cuba.
Orcynus subulatus Poey, Enumeratio, 1875, 71, Cuba; perhaps same
as *T. coretta*.
2035. *Thunnus coretta* (Cuvier and Valenciennes) *Bonita of Martinique*.
West Indies.
Tynnus coretta Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831,
102, a young fish 11 inches long from Martinique.
- Genus 684. GERMO Jordan. *Long-finned albacores*.
Orcynus Cuvier, Règne Anim., Ed. I, 1817, 314 (*Scomber germo*
Lacépède = *Scomber alalunga* Gmelin; not *Orcynus* Rafinesque, 1810,
which is *Scomberoides*)
Germo Jordan, Proc Acad. Nat. Sci. Phila., XL, 1888, 180 (*Scombrus*
alalunga Gmelin); substitute for *Orcynus*, preoccupied.
2036. *Germo alalunga* (Gmelin). *Albacore; Alalunga*
Mediterranean; Atlantic; all warm seas.
Scomber alalunga Gmelin, Syst. Nat., I, 1788, 1330, Sardinia; mis-
printed *Alatunga*.
Thynnus atlanticus Lesson, Voy. Coquille Zool., II, 1830, 165, Atlantic
ocean.
Thynnus balteatus Cuvier and Valenciennes, Hist. Nat. Poiss., VIII,
1831, 98 (136), coast of Brazil ("vis-à-vis la Trinité du Brésil, par les
20° de latitude Australe").
2037. *Germo germo* (Lacépède).
Pacific Ocean; California; Hawaii, Japan, and southward.
Scomber germo Lacépède, Hist. Nat. Poiss., II, 1800, 598, lat. 17° S.,
long 103° W.
Thynnus pacificus Cuvier and Valenciennes, op. cit., VIII, 1831, 96
(133), lat. 26° S; long. 103° W.
- Genus 685. NEOTHUNNUS Kishinouye. *Yellow-finned albacores*
Neothunnus Kishinouye, Coll. Agric. Tokyo, VIII, 1923, 445 (*Thynnus*
macropterus Temminck and Schlegel).
2038. *Neothunnus catalinae* Jordan and Evermann. *Yellow-finned albacore*.
Open Pacific; southern California, about Santa Catalina Island.
Neothunnus catalinae Jordan and Evermann, Occ. Paper No. 12, Calif.
Acad. Sci., Sept. 30, 1926, 19, pl. 4, Santa Catalina Island, Calif.
2039. *Neothunnus albacores* (Bonnaterre). *West Indies long-finned albacore*.
West Indies.
Scomber albacores Bonnaterre, Encyc. Ichth., 1788, 140, Jamaica; based
on Sloane; not *albacora* Lowe.
Scomber sloani Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831,
107 (148), Jamaica; after Sloane.
2040. *Neothunnus allisoni* (Mowbray). *Allison's tuna*.
Coast of Florida.
Thunnus allisoni Mowbray, Copeia, No. 78, 1920, 9, with text fig.,
Biscayne Bay, east coast of Florida.

Family 164. GEMPYLIDÆ. Snake mackerel; Escolares

Genus 686. ESCOLAR Jordan and Evermann.

Escolar Jordan and Evermann, in Goode and Bean, Oceanic Ichth.,
1896, 519 (*Thyrstitops violaceus* Bean).

Bipinnula Jordan and Evermann, Fishes North and Mid. Amer., 1896,
878 (*Thyrstitops violaceus* Bean); duplication of names accidental
through noncorrection of proofs.

2041. *Escolar violaceus* (Bean).

Atlantic coast of America in deep water; LaHave Bank.

Thyrstitops violaceus Bean, Proc. U. S. Nat. Mus., X, 1887, 513, LaHave
Bank; perhaps the same as *Escolar niger*.

2042. *Escolar niger* (Poey).

Cuba.

Thyrsites niger Poey, Enumeratio, 1875, 74, deep water off Cuba.Genus 687. *RUVETTUS* Cocco. *Escolares*.*Ruvettus* Cocco, Giorn. Sci. Sicilia, XLII, 1829, 2 (*R. pretiosus* Cocco).*Aplurus* Lowe, Trans. Zool. Soc. London, II, 1841, 180 (*A. simplex* Lowe).*Acanthoderma* Cantraine, Jour. Acad. Sci. Belles-Lettres Bruxelles, X, 1837 (*A. temminckii*).2043. *Ruvettus pretiosus*⁴⁶ Cocco. *Escolar*; *Rovetto*; *Ruvetto*; *Chicolar*; *Oilfish*; *Scounfish*; *Plavntail*.

Mediterranean and westward to Cuba in Atlantic.

Ruvettus pretiosus Cocco, in Giornale di Scienze per la Sicilia, XLII, 1829, 21, Messina.*Tetragonurus simplex* Lowe, Proc. Zool. Soc. Lond., 1833, 143, Madeira.*Ruvettus temminckii* Cantraine, Giorn. Sci. Ft. Lit. Pisa, 1833, Malta; *vide* Poey.*Thyrsites acanthoderma* Lowe, Proc. Zool. Soc. Lond., 1839, 78, Madeira.*Thyrsites scholaris* Poey, Memorias, I, 1854, 372, pl. 32, fig. 1, Cuba.Genus 688. *EPINNULA* Poey.*Epinnula* Poey, Memorias, I, 1854, 369 (*E. magistralis* Poey).2044. *Epinnula magistralis* Poey. *Dómine*.

Habana.

Epinnula magistralis Poey, Memorias, I, 1854, 369, Habana.Genus 689. *NEALOTUS* Johnson.*Nealotus* Johnson, Proc. Zool. Soc. Lond., 1865, 434 (*N. tripes* Johnson).2045. *Nealotus tripes* Johnson.

About Madeira in deep water; between the Bahamas and Bermudas.

Nealotus tripes Johnson, Proc. Zool. Soc. Lond., 1865, 434, Madeira.Genus 690. *PROMETHICHTHYS* Gill. *Conejos*.*Prometheus* Lowe, Trans. Zool. Soc. Lond., II, 1841, 181, Madeira (*Psempylus prometheus* Cuvier and Valenciennes); name preoccupied.*Promethichthys* Gill, Mem. Nat. Acad. Sci., VI, 1893, 115 and 123 (*Prometheus atlanticus* Lowe=*Gempylus prometheus* Cuvier and Valenciennes).2046. *Promethichthys prometheus* (Cuvier and Valenciennes). *Rabbitfish*; *Coelho*; *Conejo*; *Bermuda catfish*.

Tropical Atlantic; Cuba; Bermuda.

Gempylus prometheus Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 156 (213), pl. 222, St. Helena.Genus 691. *PTAX*⁴⁷ Jordan and Evermann.*Ptax* Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 504 (*Dicrotus parvipinnus* Goode and Bean).2047. *Ptax parvipinnis* (Goode and Bean).

Western Atlantic in deep water.

Dicrotus parvipinnis Goode and Bean, Oceanic Ichth., 1895, 201, fig. 212, Gulf Stream at Albatross stations 2537, 2542, and 2601, at about 40° N., 70° W.Genus 692. *GEMPYLUS* Cuvier and Valenciennes. *Snake mackerels*.*Gempylus* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 152 (207) (*G. serpens* Cuvier and Valenciennes).*Lemnisoma* Lesson, Voyage Coquille, Poiss., 1826 (1831), 160, the date later than that of *Gempylus*, as the work of subsequent Cuvier and Valenciennes is quoted by Lesson (*Lemnisoma thyrsioides* Lesson).*Zypothyca* Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 239 (*Gempylus coluber* Cuvier and Valenciennes).⁴⁶ A species very similar, regarded by Doctor Gudger as the same, is *Ruvettus pretiosus* Weber=*R. pacificus* Jordan and Jordan, from the tropical Pacific north to Hawaii.⁴⁷ From *tax*, a rabbit, conejo, a Spanish name of *Promethichthys*. The genus is characterized by the number of fin rays—D. XXI, 11, A. II, 8—while in *Promethichthys* the rays are D. XVIII, 19-II; A. 16-II. *Dicrotus* has two lateral lines, the fin rays about as in *Promethichthys*.

2048. *Gempylus serpens* Cuvier and Valenciennes.

Deep seas of the Atlantic.

Gempylus serpens Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 152 (207), Martinique*Gempylus ophidianus* Poey, Memorias, II, 1861, 246, Cuba.**Family 165. TRICHIURIDÆ. Hairtails; Cutlassfishes****Genus 693. TRICHIURUS Linnæus. Hairtails.***Trichiurus* Linnæus, Syst. Nat., Ed. X, 1758, 246 (*T. lepturus* Linnæus); after Artedi.*Gymnogaster* Gronow, Mus. Ichth., I, 1754, 17, No. 47, and Zooph., I, 1763, 136 (*Trichiurus lepturus* Linnæus).*Enchelyopus* Klein, in Bleeker, Versl. Akad. Amsterdam, XIV, 1862, 109 (*Trichiurus lepturus* Linnæus); after Klein; not of Schneider nor of Gronow.*Lepturus* (Artedi) Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 126 (*Trichiurus lepturus* Linnæus, a revival of a pre-Linnæan name).**2049. *Trichiurus lepturus* Linnæus. Cutlassfish; Scabbardfish; Silverfish; Sable; Savola; Hairtail; Ribbonfish; Silvery hairtail.**

Western Atlantic north to New Jersey; West Indies.

Trichiurus lepturus Linnæus, Syst. Nat., Ed. X, 1758, 246, America; after *Lepturus* of Artedi.*Trichiurus argenteus* Shaw, Gen. Zool., IV, 1803, 90, pl. 12; after Linnæus.**2050. *Trichiurus nitens* Garman.**

Off Galapagos Islands. This or a similar species in the Gulf of California.

Trichiurus nitens Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 69, off Galapagos Islands, Albatross stations 3354, 3389.**Genus 694. APHANOPUS Lowe.***Aphanopus* Lowe, Proc. Zool. Soc. Lond., Pt. VI, 1839, 79 (*A. carbo* Lowe).**2051. *Aphanopus minor* Collett.**

East coast of Greenland.

Aphanopus minor Collett, Vidensk. Selsk. Forhandl. Christiania, No. 19, 1836, 3, east coast of Greenland at 65° N., 31° W.**Genus 695. EVOXYMETOPON Poey. Tirantes.***Evoxymetopon* Poey, in Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 227 (*E. tæniatus* Poey).**2052. *Evoxymetopon tæniatus* Poey. Tirante; Tyrantfish.**

Cuba.

Evoxymetopon tæniatus Poey, in Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 228, Cuba.**Genus 696. LEPIDOPUS Gouan. Frostfishes.***Lepidopus* Gouan, Hist. Nat. Poiss., 1770, 107 and 185 (*gouani*).*Vandellius* Shaw, Gen. Zool., IV, 1803, 199 (needless substitute for *Lepidopus* Gouan).*Scarcina* Rafinesque, Caratteri, 1810, 20 (*S. argyrea* Rafinesque).*Xiphotheca* Montagu, Werner. Mus., I, 1812, 82 (*Lepidopus caudatus* Euphrasen).**2053. *Lepidopus xantusi* Goode and Bean. Scabbardfish.**

Avalon, Santa Catalina; Lower California north to Santa Catalina, Calif.; San José del Cabo, Cape San Lucas.

Lepidopus xantusi Goode and Bean, Oceanic Ichth., 1895, 519, Cape San Lucas.**Genus 697. BENTHODESMUS Goode and Bean.***Benthodesmus* Goode and Bean, Proc. U. S. Nat. Mus., IV, 1881, 380 (*B. elongatus* Goode and Bean, not *Lepidopus elongatus* Clarke = *Benthodesmus atlanticus* Goode and Bean).**2054. *Benthodesmus atlanticus* Goode and Bean.**

Western edge of Grand Banks of Newfoundland in 80 fathoms.

Benthodesmus elongatus Goode and Bean, Proc. U. S. Nat. Mus., IV, 1881, 380-383; not of Clarke.*Benthodesmus atlanticus* Goode and Bean, Oceanic Ichth., 1895, 205, Grand Banks.

Series XIPHIIFORMES

Family 166. ISTIOPHORIDÆ. Sailfishes; Spearfishes

Genus 698. ISTIOPHORUS Lacépède. *Sailfishes*.

Istiophorus Lacépède, Hist. Nat. Poiss., III, 1803, 374 (*I. gladiifer* Lacépède = *Scomber gladius* Broussonet).

Notistium Herrmann, Obs. Zool., 1804, 305 (*N. gladius* Herrmann).

Zanclurus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 239 (*Histiophorus indicus* Cuvier and Valenciennes = *H. gladius*).

Histiophorus of authors, revised spelling.

2055. *Istiophorus americanus* (Cuvier and Valenciennes).

Brazil, and West Indies, north to Woods Hole.

Histiophorus americanus Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 303, Brazil; after Guebucú of Marcgrave.

2056. *Istiophorus wrighti* Jordan and Evermann.

West Indies northward to Florida.

Istiophorus wrighti Jordan and Evermann, Occ. Paper No. 12, Calif. Acad. Sci., Sept. 30, 1926, 43, pl. 12, fig. 2, coast of Florida.

2057. *Istiophorus greyi* Jordan and Hill. *Mexican sailfish; Pacific volador*.

Pacific coast of Mexico, north to Cape San Lucas.

Istiophorus greyi Jordan and Hill, in Jordan and Evermann, Occ. Paper No. 12, Proc. Calif. Acad. Sci., Sept. 30, 1926, 41, pl. 14, near Cape San Lucas, Lower California.

2058. *Istiophorus volador* Jordan and Evermann. *Volador; Florida sailfish*
About Florida Keys.

Istiophorus volador Jordan and Evermann, Occ. Paper No. 12, Calif. Acad. Sci., Sept. 30, 1926, 46, pl. 12, fig. 1, Long Key, Fla.

Istiophorus maguirei Jordan and Evermann, loc. cit., 45, pl. 13, West Indies; apparently young of *volador*.

Genus 699. MAKAIRA Lacépède. *Marlinspike fishes; Spearfishes*.

Makaira Lacépède, Hist. Nat. Poiss., IV, 1803, 688 (*M. nigricans* Lac.).

2059. *Makaira mazara* (Jordan and Snyder). *Black marlin; Kurokajiki*
(*Black spearfish*); *Mazaara* (*real swordfish*); *Kuroka* (*black marlin*).

Pacific Ocean, from Japan to Hawaiian Islands and the Channel Islands, Calif.

Tetrapturus mazara Jordan and Snyder, Jour. Coll. Sci. Imp. Univ. Tokyo, XV, May 20, 1901, 4, 305, Misaki, Japan.

2060. *Makaira ampla* (Poey). *Aguja de Casta*.

West Indies.

Tetrapturus amplus Poey, Memorias, II, 1861, 243, Habana.

2061. *Makaira marlina* Jordan and Evermann. *Giant black marlin*.

Southern California, south to Cape San Lucas.

Makaira marlina Jordan and Evermann, Occ. Paper No. 12, Calif. Acad. Sci., Sept. 30, 1926, 59, pl. 17, off Cape San Lucas, Lower California.

2062. *Makaira mitsukurii* (Jordan and Snyder). *Marlinspike fish; Barred marlin; Makajiki or true spearfish of Japan*.

Coasts of Japan; Hawaii; southern California; abundant off Santa Catalina.

?*Tetrapturus indicus* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 286, Sumatra, on a figure by Banks.

?*Histiophorus audax* Philippi.

Tetrapturus mitsukurii Jordan and Snyder, Jour., Coll. Sci. Imp. Univ. Tokyo, XV, pt. 2, 1901, 304, Pl. XVI, fig. 5, fishwell at Misaki.

2063. *Makaira holei* Jordan and Evermann. *Striped marlin*.

Off Pacific coast of Mexico, north to Cape San Lucas.

Makaira holei Jordan and Evermann, Occ. Papers No. 12, Calif. Acad. Sci., Sept. 30, 1926, 63, pl. 19, fig. 1, off Cape San Lucas, Lower California.

Family 167. XIPHIIDÆ. Swordfishes

Genus 700. XIPHIAS Linnæus. *Swordfishes*.*Xiphias* Linnæus, Syst. Nat., Ed. X, 1758, 248 (*X. gladius* Linnæus).*Phæthomichthys* Nichols, Amer. Mus. Nov., XIX, I, 1923 (*Phæthomichthys tuberculatus* Nichols; a young swordfish).2064. *Xiphias gladius* Linnæus. *Common swordfish; Broadbill; Espada; Espadón emperador.*

Open sea, Atlantic and Pacific Oceans, on both coasts; Cuba to Cape Breton and Newfoundland Banks; Pacific off Santa Barbara Islands, Hawaii, Japan, San Diego.

Xiphias gladius Linnæus, Syst. Nat., Ed. X, 1758, 248, Europe.*Xiphias rondeletii* Leach, in Wern. Mem., II, 1818, 58, pl. 2, fig. 1, Firth of Forth.*Xiphias imperator* Schneider, ed. Bloch, Syst. Ichth., 1801, 93, Pl. XXI, Mediterranean, after an erroneous figure of Duhamal.*Phæthomichthys tuberculatus* Nichols, Amer. Mus. Nat. Hist. Novitates, XIX, pl. 1, Oct. 19, 1923; very young of *X. gladius* L.

Series LUVARIFORMES

Family 168. LUVARIDÆ

Genus 701. LUVARUS Rafinesque.

Luvarus Rafinesque, Caratteri, 1810, 22 (*L. imperialis* Rafinesque).*Diana* Risso, Europe Méridionale, 1826, 267 (*D. semilunata* Risso).*Ausonia* Risso, op. cit., 341 (*A. cuvieri* Risso).*Proctostegus* Nardo, Diaro, Chem. et Hist. Nat. Ticino, I, 1827, 18, 42 (*P. proctostegus* Nardo).*Astrodermus* (Bonelli) Cuvier, Règne Anim., ed. 2, II, 1829, about 215 (*A. guttatus* Bonelli MS).*Scafaria* Gistel, Thierreich, VIII, 1848 (*Ausonia cuvieri* Risso).2065. *Luvarus imperialis* Rafinesque.

Open sea; once taken at San Pedro, Calif. (W. F. Thompson).

Luvarus imperialis Rafinesque, Caratteri, 1810, 22.*Diana semilunata* Risso, Europe Mérid., 1826, 267.*Ansonia cuvieri* Risso, loc. cit., 341.*Proctostegus proctostegus* Nardo, Diaro, Chem. et Hist. Nat. Ticino, I, 1827, 18, 42.*Astrodermus guttatus* Bonelli, in Cuvier, Règne Anim., ed. 2, II, 1829, 215 ca.*Scafaria* Gistel, Thierreich, VIII, 1848.

Series CORYPHÆNIFORMES

Family 169. CORYPHÆNIDÆ. Dolphins; Dorados

Genus 702. CORYPHÆNA Linnæus. *Dorados*.*Coryphæna* Linnæus, Syst. Nat., Ed. X, 1758, 261 (*C. hippurus* L.).*Lampugus* Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 317 (*Scomber pelagicus* L.; female).2066. *Coryphæna hippurus* Linnæus. *Common dolphin; Dorado; Dourade.*

Pelagic to Cape Cod and northward, South Carolina to Texas; also throughout the mid-Pacific.

Coryphæna hippurus Linnæus, Syst. Nat., ed. X, 1758, 261, open seas.*Scomber pelagicus* Linnæus, loc. cit., 299, no locality.*Coryphæna fasciolata* Pallas, Spicil. Zool., VIII, 1770, 23, Amboina.*Coryphæna imperialis* Rafinesque, Caratteri, 1810, 33, Sicily.*Coryphæna immaculata* Agassiz in Spix, Pisc. Brasil., 1829, 111, pl. 56, Atlantic, off Brazil.*Coryphæna marcgravi* Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 223 (301), America.*Coryphæna suerii* Cuvier and Valenciennes, loc. cit., 224 (302), Philadelphia.*Coryphæna dorado* Cuvier and Valenciennes, loc. cit., 225 (303), Brazil.*Coryphæna dolfyn* Cuvier and Valenciennes, loc. cit., 226 (305), Antilles.

- Coryphæna virgata* Cuvier and Valenciennes, loc. cit., 228 (308), Martinique.
Coryphæna argyrurus Cuvier and Valenciennes, loc. cit., 232 (314), Sea of Coromandel.
Coryphæna blainvillæ Cuvier and Valenciennes, loc. cit., 233 (315), seas of India.
Lampugus sculus Cuvier and Valenciennes, loc. cit., 240 (323), Sicily.
Coryphæna scomberoides Cuvier and Valenciennes, loc. cit., 234 (315), South Sea.
Ectenias brunneus Jordan and Thompson, Mem. Carnegie Mus., VI, 1914, 241, Japan; young.

2067. *Coryphæna equisetis* Linnæus. *Small dolphin*.
 Open Atlantic and Pacific; rare in the West Indies. Recently taken in some numbers off coast of southern California.
Coryphæna equisetis Linnæus, Syst. Nat., Ed. X, 1758, 261 (misprinted *equiselis*), high seas; after Osbeck.
Coryphæna aurata Rafinesque, Caratteri, 1810, 33, Sicily.
Coryphæna lessona Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 228 (307), India.
Lampugus punctulatus Cuvier and Valenciennes, loc. cit., 242 (429), Atlantic at the Equator.
Lampugus neapolitanus Cuvier and Valenciennes, loc. cit., 241 (325), Naples; after Bloch.
Coryphæna azorica Cuvier and Valenciennes, loc. cit., 227 (306), Azores.

Series BRAMIFORMES

Family 170. BRAMIDÆ. Sea breams

- Genus 703. *BRAMA* Bloch and Schneider. *Pomfret*.
Brama Bloch and Schneider, Ed. Bloch, Syst. Ichth., 1801, 98 (*Sparus rai* Bloch).
Lepodus Rafinesque, Caratteri, 1810, 54 (*L. saragus* Rafinesque = *Sparus rai* Bloch).

2068. *Brama rai* (Bloch). *Sea bream; Pomfret; Catagnole; Rondanin*.
 Open seas; Europe; Faroe Islands; Bermuda; Grand Banks; Pacific coast from Santa Catalina to Puget Sound; Japan; reported recently from Cape Flattery.
Brama marina cauda forcipata Ray, Syn. Meth. Piscium, 1713, 115, Middelburg.
Sparus rai Bloch, Naturgesch. Ausl. Fische, V, 1791, 95; after Ray, etc.
Sparus castaneola Shaw, Gen. Zool., IV, 1803, 424; after Lacépède.
Sparus niger Turton, British Fauna, 1807, 98, Swansea.
Lepodus saragus Rafinesque, Caratteri, etc., 1810, 54, Palermo.
Brama dussumieri Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 219 (294), Indian Sea, long. 85° E.
 ?*Brama oroni* Cuvier and Valenciennes, loc. cit., 220 (295), Indian Sea, long. 85° E.
 ?*Brama chilensis* Gay, Hist. Chile, Peces, 1843, 218, Chile.

2069. *Brama agassizii* Poey.
 Cuba; rare.
Brama agassizii Poey, Memorias, II, 1861, 204, Cuba.

2070. *Brama brevoortii* Poey.
 Cuba; rare.
Brama brevoortii Poey, Memorias, II, 1861, 206, Cuba.

- Genus 704. *TARACTES* Lowe.
Taractes Lowe, Proc. Zool. Soc. Lond., 1843, 82 (*T. asper* Lowe; probably young of *Taractes longipinnis*).
Argo Doderlein, Fische Japans, II, Denkschr. Acad. Wiss. Wien, XLVIII, 1883, 34 (*Argo steindachneri* Doderlein); name preoccupied.

2071. *Taractes saussurii* (Lunel).
 Habana.
Brama saussurii Lunel, Rêvue du Genre Brama, in Mém. Soc. Phys. Hist. Nat. Genève, XVIII, 1865, 185, tab. 2, Cuba.

Family 171. STEINEGERIIDÆ

Genus 705. STEINEGERIA Jordan and Evermann.

Steinegeria Jordan and Evermann, Proc. U. S. Nat. Mus., IX, 1886, 467 (*S. rubescens* Jordan and Evermann).2072. *Steinegeria rubescens* Jordan and Evermann.

Gulf of Mexico in deep water; one specimen known.

Steinegeria rubescens Jordan and Evermann, Proc. U. S. Nat. Mus., IX, 1886, 467, snapper banks off Pensacola, Fla.

Family 172. PTERACLIDÆ

Genus 706. PTERACLIS Gronow.

Pteraclis Gronow, Acta Helvetica, VII, 1772, 43 (*Coryphæna velifera* Pallas).*Pteridium* Scopoli, Introd. Hist. Nat., 1777, 454 (*Coryphæna velifera* Pallas).*Oligopodus* Lacépède, Hist. Nat. Poiss., II, 1800, 511 (*Coryphæna velifera* Pallas).*Pteraclidus* Rafinesque, Analyse de la Nat., 1815, 82, substitute for *Oligopodus* Lacépède.*Oligopodes* Risso, in Cuvier, Règne Anim., ed. 1, 1817, 328 (*Oligopode noir* Risso).*Oligopus* Risso, Europe Mérid., 1826, 105, substitute for *Oligopodus*.2073. *Pteraclis carolinus* Cuvier and Valenciennes.

Coast of South Carolina.

Pteraclis carolinus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 272 (368), off coast South Carolina.? *Pteraclis trichipterus* Cuvier and Valenciennes, loc. cit., 272 (367), locality unknown.

Genus 707. CENTROPHOLIS Hilgendorf.

Centropholis Hilgendorf, Sitzber. Naturf. Freunde, Berlin, 1878, 135 (*C. petersi* Hilgendorf).2074. *Centropholis goodei* Jordan.

Gulf Stream.

Pteraclis carolinus Goode and Bean, Oceanic Ichth., 212, Pl. LIX, fig. 218, Gulf Stream; not of Linnæus.*Centropholis goodei* Jordan, Ann. Carnegie Mus., IX, 1915, 333, Gulf Stream.

Series STROMATEIFORMES

Family 173. STROMATEIDÆ. Fiatolas

Genus 708. PEPRILUS Cuvier.

Rhombus Lacépède, Hist. Nat. Poiss., II, 1800, 321 (*Chætodon alepidotus* Linnæus); preoccupied in mollusks.*Peprilus* Cuvier, Règne Anim., ed. 2, II, 1829, 213 (*Stromateus longipinnis* Mitchell = *Stromateus paru* Linnæus).2075. *Peprilus paru* (Linnæus). Harvestfish; Poppyfish; Rudderfish.

South Atlantic coast of United States; West Indies; Cape Cod to Jamaica and Brazil.

Stromateus paru Linnæus, Syst. Nat., Ed. X, 1758, 248, Jamaica; based on Sloane.*Chætodon alepidotus* Linnæus, op. cit., Ed. XII, 1766, 460, Charleston, S. C.? *Sternoptyx gardenii* Bloch and Schneider, Syst. Ichth., 1801, 494, Carolinas; after Linnæus.*Stromateus longipinnis* Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 366, New York Bay.

Genus 709. PALOMETA Jordan and Evermann.

Palometa Jordan and Evermann, Fishes North and Mid. Amer., 1896, 966 (*Stromateus palometa* Jordan and Bollman).2076. *Palometa simillima* (Ayres). California pampano.

Pacific coast of United States, Puget Sound to San Diego.

Poronotus simillimus Ayres, Proc. Calif. Acad. Nat. Sci., 1st ser., II, 1860, 84, San Francisco.

2077. *Palometa media* (Peters). *Palometa*.
Pacific coast of North America, Mazatlan to Panama.
Stromateus medius Peters, Berl. Monatsb., 1869, 707, Mazatlan, Mexico.
2078. *Palometa palometa* (Jordan and Bollman).
Pacific Ocean, off Colombia.
Stromateus palometa Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 156, off coast of Colombia, lat. 8° 16' 30'' N., long. 79° 37' 45'' W., at Albatross station 2804.
- * 2079. *Palometa snyderi* (Gilbert and Starks).
Panama.
Peprilus snyderi Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 87, Panama.
- Genus 710. *PORONOTUS* Gill.
Poronotus Gill, Cat. Fish. E. Coast N. Amer., Sup. to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 35 (*Stromateus triacanthus* Peck).
2080. *Poronotus triacanthus* (Peck). *Dollarfish*; *Harvestfish*; *Butterfish*; *Lafayette*.
Nova Scotia to Florida; common.
Stromateus triacanthus Peck, Mem. Amer. Acad., II, 1800, pt. 2, 48, pl. 2, fig. 2, Piscataqua River, N. H.
Stromateus cryptosus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 365, New York Bay.
- Genus 711. *EUCROTUS* Bean (B. A.).
Eucrotus Bean, Proc. Biol. Soc. Wash., XXV, 1912, 123 (*E. ventralis* Bean).
2081. *Eucrotus ventralis* Bean.
Bermudas.
Eucrotus ventralis Bean, Proc. Biol. Soc. Wash., XXV, 1912, 125, Bermudas.

Family 174. CENTROLOPHIDÆ. Rudderfishes

- Genus 712. *CENTROLOPHUS* Lacépède. *Black ruffs*.
Centrolophus Lacépède, Hist. Nat. Poiss., IV, 1803, 441 (*Perca nigra* Gmelin=*Coryphæna pompilus* L.).
Acentrolophus Nardo, Prodromus. Adriat. Ichth. Giorn. Fisica di Pavia, sp. 62, 1827, (*A. maculatus* Nardo=*Centrolophus niger* Lacépède, the name *Centrolophus* deemed inappropriate).
Gymnocephalus Cocco, Nuov. Ann. Sci. Nat., II, 1838, 26 (*G. messinensis* Cocco); name preoccupied.
Pompilus Lowe, Suppl. Fish Madeira, Proc. Zool. Soc. London, VII, 1839, 81 (*Coryphæna pompilus* L.).
2082. *Centrolophus niger* (Gmelin). *Blackfish*; *Black ruffe*; *Borlase*.
Coasts of southern Europe; once taken at Dennis, Mass.
Perca nigra Gmelin, Syst. Nat., 1788, 1321, Cornwall.
Centrolophus liparis Risso, Eur. Mérid., III, 1826, 337, Nice, Italy.
Centrolophus morio Lacépède in Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 254 (342), Mediterranean.
Centrolophus pompilus Cuvier and Valenciennes, loc. cit., 247 (334), pl. 269; not of Linnaeus.
Acentrolophus maculosus Nardo, Prodr. Ichth. Adriat., No. 62, Isis, 1827, Adriatic.
Gymnocephalus messinensis Cocco, Nuovi Ann. Sci. Nat., II, 1838, 26, Messina.
- Genus 713. *PALINURICHTHYS* Bleeker. *Black rudderfishes*.
Palinurus DeKay, N. Y. Fauna: Fishes, 1842 (*Coryphæna perciformis* Mitchell); name preoccupied in Crustacea.
Palinurichthys Bleeker, Enum. Spec. Pisc. Arch. Ind., in Acta Soc. S. Indo-Neerl., VII, November, 1859, 22 (*Coryphæna perciformis*, Mitchell).
Palinurichthys Gill, Proc. Acad. Nat. Sci. Phila., XII, January, 1860, 20 (*Coryphæna perciformis* Mitchell).
Pammelas Günther, Cat., II, June, 1860, 485 (*Coryphæna perciformis* Mitchell).

2083. *Palinurichthys perciformis* (Mitchill). *Rudderfish, Logfish; Snipnose mullet; Black rudderfish*
Atlantic coast of North America from Cape Hatteras to Maine; once taken in Cornwall.
Coryphæna perciformis Mitchill, Amer. Month. Mag., II, 1818, 244, New York Harbor.
Pimalepterus cornubiensis Cornish Zoologist, IX, 1874, 4255, Penzance, Cornwall (a stray specimen floated across in a fish box).

Genus 714. *SCHEDOPHILUS* Cocco.

Schedophilus Cocco, Giorn. Innom. Messina, Ann., III, 1834, No. 7, 57, (*S. medusophagus* Cocco).

Crius Valenciennes, Ichth. Îles Canaries, II, 1844, 43 (*Crius bertheloti* Valenciennes = *Schedophilus medusophagus* Cocco).

2084. *Schedophilus heathi* Gilbert.

Monterey Bay.

Schedophilus heathii Gilbert, Proc. Calif. Acad. Sci., ser. 3, Zool., III, 1904, 260, Pl. XXVI, Pacific Grove, Calif.

Genus 715. *ICICHTHYS* Jordan and Gilbert.

Ichthys Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 305 (*I. lockingtoni* Jordan and Gilbert).

2085. *Ichthys lockingtoni* Jordan and Gilbert.

Off California, in deep water; a surface swimmer, hovering under jelly-fishes; known from off Monterey and San Francisco.

Ichthys lockingtoni Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 305, off San Francisco.

Family 175. *NOMEIDÆ*. Man-of-war fishes

Genus 716. *NOMEUS* Cuvier.

Nomeus Cuvier, Règne Anim., ed. 1, II, 1817, 315 (*Gobius gronovii* Gmelin).

2086. *Nomeus gronovii* (Gmelin). *Portuguese man-of-war fish; Harder; Pastor.*

Tropical parts of the Atlantic and Indian Oceans; Sargasso Sea; Florida; Bermuda; Woods Hole, Mass.; rocky pools at Panama.

Gobius gronovii Gmelin, Syst. Nat., Ed. XIII, 1788, 1205, tropical America; after Gronow.

Eleotris mauritii Bloch and Schneider, Syst. Ichth., 1801, 66, Brazil.

Nomeus maculosus Bennett, Proc. Comm. Zool. Soc., I, 1831, 146, East Indies.

Nomeus oxyurus Poey, Memorias, II, 1861, 236, Habana.

Genus 717. *PSENES* Cuvier and Valenciennes.

Psenes Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 193 (259), (*P. cyanophrys* Cuvier and Valenciennes).

2087. *Psenes cyanophrys* Cuvier and Valenciennes.

Open sea; Atlantic, Pacific, and Indian Oceans; Jamaica and Martinique.

Psenes cyanophrys Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 193 260, pl. 265, New Ireland.

Psenes javanicus Cuvier and Valenciennes, loc. cit., 196 (264), Batavia.

Psenes auratus Cuvier and Valenciennes, loc. cit., 197 (264), Gulf of Bengal.

Psenes fuscus Guichenot, Mem. Soc. Sci. Nat. Cherbourg, XII, 1866, 138, Madagascar.

Cubiceps multiradiatus Günther, Proc. Zool. Soc. Lond., 1871, 661, pl. 61, Manado.

2088. *Psenes maculatus* Lütken.

Open Atlantic.

Psenes maculatus Lütken, Spolia Atlantica, 1880, 110, open Atlantic, lat. 39° N., long. 25° S. and between 34° and 27° W. in 600 to 700 fathoms.

2089. *Psenes regulus* Poey.

Coasts of Cuba.

Psenes regulus Poey, Synopsis, 1868, 375, Cuba.

2090. *Psenes pellucidus* Lutken.
Gulf Stream at 32° 24' N., 76° 55' W., in 528 fathoms.
Psenes pellucidus Lutken, Spolia Atlantica, 1880, 516 (108), with fig., Strait of Surabaja.
2091. *Psenes edwardsii* Eigenmann.
Off Rhode Island.
Psenes edwardsii Eigenmann, Bull. U. S. Fish Com., XXI, 1901 (1902), 35, fig.; taken from under a medusa, 30 miles south of Newport, R. I.
2092. *Psenes chapmani* Fowler.
Open Atlantic, Cape Verde Islands, northwest of the Lesser Antilles.
Psenes chapmani Fowler, Proc. Acad. Nat. Sci. Phila., LVIII, 1906, 120; open Atlantic, lat. 24° 21' N, long. 34° 32' W.
2093. *Psenes pacificus* Meek and Hildebrand.
Panama.
Psenes pacificus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, 409, Chame Point, Panama.

Family 176. TETRAGONURIDÆ. Squaretails

Genus 718. TETRAGONURUS Risso.

Tetragonurus Risso, Ichth. Nice, 1810, 247 (*T. cuvieri* Risso).
Ctenodax Macleay, Proc. Linn. Soc. New South Wales, 1885, 716 (*C. williamsoni* Macleay).

2094. *Tetragonurus cuvieri* Risso. *Squaretail*; *Escolar de natura*; *Courpata*; *Sea raven*.
Open seas; Nice; Toulon and Marseilles; Madeira; once taken off Santa Catalina Island, Calif. (W. F. Thompson), and once at Woods Hole, Mass.; these possibly different species.
Tetragonurus cuvieri Risso, Ichth. Nice, 1810, 347, Nice.
Tetragonurus atlanticus Lowe, Fishes Madeira, pt. 5, 1860 (1843), 129, Pl. XIX, Madeira.

Family 177. ICOSTEIDÆ. Ragfishes

Genus 719. ICOSTEUS Lockington.

Icosteus Lockington, Proc. U. S. Nat. Mus., III, 1880, 63 (*I. ænigmaticus* Lockington).
Schedophilopsis Steindachner, Ichth. Beitr., XI, 1881, 4 (*S. spinosus* Steindachner=*Icosteus ænigmaticus* Lockington).

2095. *Icosteus ænigmaticus* Lockington. *Ragfish*.
Open sea off the coasts of California, Oregon, and Washington; scarce.
Icosteus ænigmaticus Lockington, Proc. U. S. Nat. Mus., III, 1880, 63, off San Francisco.
Schedophilopsis spinosus Steindachner, Ichth. Beitr., XI, 1881, 4, with figure, off San Francisco.

Family 178. ACROTIDÆ

Genus 720. ACROTUS Bean.

Acrotus Bean, Proc. U. S. Nat. Mus., X, 1887, 631 (*A. willoughbyi* Bean).

2096. *Acrotus willoughbyi* Bean.
Off coast of Washington, Whitby Island, Puget Sound, south to San Pedro, Calif. (W. F. Thompson), in rather deep water.
Acrotus willoughbyi Bean, Proc. U. S. Nat. Mus., X, 1887, 631, Quinault Agency, Wash.

Family 179. CARANGIDÆ. Cavallas; Pampanos

The *Carangidæ*, as heretofore understood, may be naturally divided into two families, the *Seriolidæ* (*Seriola*, *Elagates*, *Naucrates*, etc.) differing strongly in lacking the bony arch behind the abdominal cavity, this being made up of enlarged and ankylosed hæmals, its character somewhat different in different genera but developed in all *Carangidæ* in the narrow sense.

Genus 721. *DECAPTERUS* Bleeker. *Mackerel scads.*

Decapterus Bleeker, Naturk. Tijdsch., VIII, 1855, 417 (*Caranx kurra* Bleeker).

Eustomalodus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 261 (*Decapterus kurroides* Bleeker).

Gymneipignathus Gill, loc. cit., 261 (*Decapterus macrosoma* Bleeker).

Euepigymnus Gill, loc. cit., 261 (*Decapterus hypodus* Gill).

2097. *Decapterus punctatus* (Agassiz). *Scad; Round robin; Cigarfish; Quanaqua; Dotted scad.*

Cape Cod to Brazil, coasts of Florida and the West Indies.

Caranx punctatus Agassiz, in Spix, Pisc. Brasil., 1829, 108, pl. 56a, fig. 2, Brazil.

Caranx aureus Risso, in Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 25 (33), Mediterranean.

2098. *Decapterus sanctæ-helenæ* (Cuvier and Valenciennes).

Tropical Atlantic; Cuba.

Caranx sanctæ-helenæ Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 28 (37), St. Helena.

2099. *Decapterus hypodus* Gill.

Cape San Lucas, Lower California.

Decapterus hypodus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 262, Cape San Lucas, Lower California.

2100. *Decapterus macarellus* (Cuvier and Valenciennes). *Mackerel scad; Antonino; Cigarfish.*

West Indies, northward to Cape Cod.

Caranx macarellus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 30 (40), Martinique.

Genus 722. *TRACHURUS* Rafinesque. *Saurels.*

Trachurus Rafinesque, Caratteri, 1810, 41 (*Scomber trachurus* Linnæus).

2101. *Trachurus symmetricus* (Ayres). *Horse mackerel.*

Coast of California.

Caranx symmetricus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 62, Bay of San Francisco.

2102. *Trachurus trachurus* (Linnæus). *Sauwel; Gascon.*

North Atlantic, chiefly on the northern coast of Europe, south to Spain and Naples; Newport, R. I.; Pensacola.

Scomber trachurus Linnæus, Syst. Nat., Ed. X, 1758, 298, Mediterranean.

?*Caranxomorus plumieiranus* Lacépède, Hist. Nat. Poiss., III, 64, pl. 11, scarcely identifiable, from a figure by Plumier, supposed to have been made in Martinique.

Caranx semispinosus Nilsson, Prodr. Ichthyol. Scand., 1832, 84, Scandinavia; the northern form usually called *trachurus*, the valid name for the European form.

Trachurus europæus Gronow, Cat., Ed. Gray, 1854, 125, seas of Europe.

Trachurus linnæi Malm, Bohusläns Fauna, 421, Bohuslän, Sweden.

2103. *Trachurus japonicus* (Schlegel).

Coast of Japan, east of Lower California, the Mexican specimens few and not surely identified; may be *T. nova-zealandica* Richardson.

Caranx trachurus japonicus Schlegel, Faun. Japon., Poiss., 1850, 109, pl. 59, fig. 1, Japan.

2104. *Trachurus lathamii* Nichols.

Coast of Long Island (a stray from the Gulf Stream).

Trachurus lathamii Nichols, Bull. Amer. Mus. Nat. Hist., XLII, 1920, 479, Long Island.

Genus 723. *TRACHUROPS*⁴⁸ Gill.

Trachurops Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 238, 261, 431 (*Scomber crumenophthalmus* Bloch).

⁴⁸ The genus *Selar* Bleeker was based on four species, the first named being *Caranx hasselti*, which is a species of the modern genus *Atule* Jordan and Jordan. It was first restricted to *Scomber boops* Bleeker, a species of *Trachurops*; but this restriction can not stand as *boops* was not mentioned by Bleeker in connection with *Selar*; *Caranx hasselti* = *affinis* of the south Pacific must therefore be the type of *Selar*.

2105. *Trachurops crumenophthalma* (Bloch). *Goggler*; *Big-eyed scad*;
Goggle-eye jack; *Chacharro*; *Horse-eyed jack*.
Atlantic coast of the United States, Central America, South America,
and Africa; West Indies; probably not in the Pacific.
Scomber crumenophthalmus Bloch, *Naturgeschich. Ausl. Fische*, VII,
1793, 77, Acara in Guinea.
Scomber plumieri Bloch, loc. cit., 79, Antilles.
Scomber balantophthalmus Bloch and Schneider, *Syst. Ichth.*, 1801, 29,
Guinea.
Caranx macrophthalmus Agassiz, Spix, *Pisc. Brasil.*, 1829, 107, Brazil.
2106. *Trachurops brachychira* Gill. *Akule*; *Atule*.
Pacific Ocean, north to Panama, and the Gulf of California and Hawaii.
Trachurops brachychirus Gill, *Proc. Acad. Nat. Sci. Phila.*, XIV, 1862,
261, Cape San Lucas.
- Genus 724. **HEMICARANX** Bleeker.
Alepes Swainson, *Glassn. Anim.*, II, 1839, 248 (*Alepes melanoptera*
Swainson); identification uncertain.
Hemicaranx Bleeker, *Vesl. Kon. Ak. Wet.*, XIV, 1862, 135 (*H. mar-*
ginatus Bleeker).
Carangops Gill, *Proc. Acad. Nat. Sci. Phila.*, XIV, 1862, 238 (*Caranx*
heteropygus Poey).
2107. *Hemicaranx amblyrhynchus* (Cuvier and Valenciennes).
Cape Hatteras to Brazil; West Indies; Carolina and Florida.
Caranx amblyrhynchus Cuvier and Valenciennes, *Hist. Nat. Poiss.*, IX,
1833, 76 (100), pl. 248, Brazil.
Caranx heteropygus Poey, *Memorias*, II, 1861, 344, Cuba.
2108. *Hemicaranx falcatus* (Holbrook).
Charleston, S. C.
Caranx falcatus Holbrook, *Ichth. South Carolina*, 1855, 92, pl. 13, fig. 2,
Charleston, S. C.
2109. *Hemicaranx atrimanus* (Jordan and Gilbert).
Pacific coast of tropical America; Panama.
Caranx atrimanus Jordan and Gilbert, *Bull. U. S. Fish Com.*, I, 1881,
308, Panama.
2110. *Hemicaranx furthii* (Steindachner).
Panama.
Caranx fürthii Steindachner, *Ichth. Beitr.*, IV, 1875, 12, Panama.
2111. *Hemicaranx leucurus* (Günther).
Pacific coast of tropical America; Panama.
Caranx leucurus Günther, *Proc. Zool. Soc. Lond.*, 1864, 24, Panama.
2112. *Hemicaranx zelotes* Gilbert.
Panama.
Hemicaranx zelotes Gilbert, in Jordan and Evermann, *Fishes North*
and Mid. Amer., 1898, 2845, Panama.
2113. *Hemicaranx rhomboides* Meek and Hildebrand.
Atlantic coast of Isthmus of Panama.
Hemicaranx rhomboides Meek and Hildebrand, *Field Mus. Nat. Hist.*,
Zool., XV, 1925, 345, Fox Bay, Colon, Panama.
- Genus 725. **ELAPHOTOXON** ⁴⁸ Fowler.
Elaphotoxon Fowler, *Proc. Acad. Nat. Sci. Phila.*, LVII, 1905, 76
(*Scomber ruber* Bloch).
2114. *Elaphotoxon ruber* (Bloch). *Cibi mancho*; *Carbonero*; *Cibi*; *Green jack*.
West Indies and coast of the Carolinas.
Scomber ruber Bloch, *Naturgeschich. Ausl. Fische*, VII, 1793, 75, St.
Croix.
Caranx blochii Cuvier and Valenciennes, *Hist. Nat. Poiss.*, IX, 1833,
52 (69), St. Croix.
Caranx iridinus Poey, *Memorias*, II, 1861, 226, Cuba.

⁴⁸ This genus is near *Caranx*, the body rather elongate, the teeth in few series, without canines, the arch of the lateral line moderate, more than half the straight part; the dorsal and anal fins anteriorly but little elevated, maxillary not narrowed.

2115. *Elaphotoxon bartholomæi* (Cuvier and Valenciennes). *Yellow jack*;
Cibi amarilla.
West Indies, northward to Florida and North Carolina; Cuba.
Caranx bartholomæi Cuvier and Valenciennes, Hist. Nat. Poiss., IX,
1833, 75 (100), St. Bartholomew.
Caranx cibi Poey, Memorias, II, 1861, 224, Cuba.
Caranx beani Jordan, Proc. U. S. Nat. Mus., III, 1880, 486, Beaufort,
N. C.
- Genus 726. *PARATRACTUS* Gill.
Paratractus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 432 (*Caranx*
pisquetus Cuvier and Valenciennes = *Scomber crysos* Mitchell).
2116. *Paratractus crysos* (Mitchill). *Hardtail*; *Runner*; *Blue runner*; *Jurel*;
Yellow mackerel, *Crevallé*; *Cojnera*
Cape Cod to Brazil
Scomber crysos Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 424, New
York.
Trachurus squamosus Gronow, Cat. Fishes, ed. Gray, 1854, 125, Carolina.
Caranx pisquetus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833,
73, Santo Domingo; Cuba; Brazil.
2117. *Paratractus caballus* (Günther) *Cocinero*; *Cocinero dorado*; *Jurel*.
Pacific coast of tropical America, San Diego to Panama, recently taken
at San Pedro.
Caranx caballus Günther, Trans. Zool. Soc. Lond., VI, 1868, 431 (Nov.
27), Panama.
Trachurus boops Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 108,
San Diego; not *Caranx boops* of Cuvier and Valenciennes.
Caranx girardi Steindachner, Ichthyol. Notizen, IX, 1869, 25, Mazatlan.
- Genus 727. *XUREL* Jordan and Evermann.
Xurel Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI,
Apr. 27, 1927, 505 (*Caranx vinctus* Jordan and Gilbert).
2118. *Xurel vincta* (Jordan and Gilbert). *Cocinera*.
Pacific coast of Mexico, Mazatlan to Punta Arenas; Gulf of California
to Panama.
Caranx vinctus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
349, Mazatlan, Mexico.
2119. *Xurel fasciatus* (Cuvier and Valenciennes). *Segundo*; *Volantin*.
Cuba; Atlantic coast of Mexico.
? *Caranx fasciatus* Cuvier and Valenciennes, Hist. Nat. Poiss., IX,
1833, 53 (70), east coast of Mexico.
Caranx secundus Poey, Memorias, II, 1861, 223, Cuba.
2120. *Xurel lata* (Agassiz). *Jurel*; *Xurel*, *Horse-eye jack*.
Tropical Atlantic, north to Virginia; probably not in the Pacific, where
it is represented by the near related *Caranx melampygus*.
Caranx latus Agassiz, in Spix, Pisc. Bras., 1829, 105, Brazil.
Caranx lepturus Agassiz, loc. cit., 106, Brazil.
Caranx fallax Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 71
(95), Antilles; Brazil.
Caranx richardi Holbrook, Ichthyol. South Carolina, 1855, 94, pl. 13,
fig. 1, South Carolina.
Carangus aureus Poey, Enumeratio, 1875, 76, Cuba.
2121. *Xurel medusicola* (Jordan and Starks).
Pacific coast of Mexico, south to Panama.
Caranx medusicola Jordan and Starks, in Jordan, Proc. Calif. Acad.
Sci., V, 1895, 430, pl. 34, Mazatlan.
2122. *Xurel marginatus* (Gill).
Pacific Ocean; Mazatlan, Panama, and Hawaii.
Caranx marginatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 166,
Panama.
2123. *Xurel melampygus* (Cuvier and Valenciennes). *Ulua*.
Tropical Pacific, east to off-shore islands of Mexico; Hawaii. A very
important food fish.
Caranx melampygus Cuvier and Valenciennes, Hist. Nat. Poiss., IX,
1833, 103, East Indies.

2124. *Xurel stellatus* (Eydoux and Souleyet). *Omilu*.
Pacific Ocean generally; Hawaii; Revillagigedo Islands; with the "Ulua" one of the staple food fishes of the Pacific islands.
Caranx punctatus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 38; name preoccupied.
Caranx stellatus Eydoux and Souleyet, Voy. *Bonite*, Zool., I, Poiss., 1840, 167, Hawaiian Islands.
2125. *Xurel lugubris* (Poey). *Tñosa*.
Tropical Atlantic; West Indies.
Caranx lugubris Poey, Memorias, II, 1861, 222, Cuba.
Caranx frontalis Poey, loc. cit., 222, Cuba.
2126. *Xurel tenebrosus* Jordan and Evermann.
South seas; Revillagigedos; Hawaii.
Caranx lugubris Jordan and Evermann, Fishes North and Mid. Amer., 1896, 924, in part; not of Poey, based on examples so listed by Jordan and McGregor, Report U. S. Fish Com., XXIV, 1899, 227, from Clarion Island of the Revillagigedos Group.
Caranx tenebrosus Jordan and Evermann, Proc. Calif. Acad. Sci., 4th Ser., XVI, Nov. 14, 1927, 656, Clarion Island.
- Genus 728. *CARANX* Lacépède.
Caranx Lacépède, Hist. Nat. Poiss., III, 1802, 57 (*Scomber carangus* Bloch = *Scomber hippos* L., as now usually restricted) Species with the breast naked except for a central patch of scales.
Tricopterus Rafinesque, Caratteri, 1810, 41 (*Scomber carangus* Bloch = *Scomber hippos* Linnaeus), as restricted.
Carangichthys Bleeker, Nat. Tijdschr. Neder. Indie, III, 1852, 760 (*C. typus* Bleeker); immature; fish with breast naked; strong teeth present.
2127. *Caranx hippos* (Linnaeus). *Crevallé*; *Toro*; *Horse crevallé*; *Cavalla*; *Jack*; *Jaguagua*; *Yellow caranx*; *Yellow mackerel*.
East coast of tropical America, north to Cape Cod; apparently represented in the Pacific by the closely related *Caranx ignobilis*.
Scomber hippos Linnaeus, Syst. Nat., Ed. XII, 1766, 494, Charleston, S. C.
Scomber carangus Bloch, Naturgeschichte. Ausl. Fische, VII, 1793, 69, Antilles.
Caranx erithrurus Lacépède, Hist. Nat. Poiss., III, 1802, 68, South Carolina.
Caranx daubentonii Lacépède, loc. cit., 72, Martinique.
Caranx antillarum Bennett, Whaling Voyage, II, 1840, 282, West Indies.
Caranx defensor DeKay, N. Y. Fauna: Fishes, 1842, 120, New York.
Carangus esculentus Girard, U. S. Mex. Bound. Surv., 1859, 23, pl. XI, figs. 1-3, Brazos Santiago, Tex.
2128. *Caranx ignobilis* Forskål.
Tropical Pacific, east to offshore islands of Mexico; Hawaii.
Scomber ignobilis Forskål, Descr. Anim., 1775, Red Sea.
Carangus hippoides Jenkins, Bull. U. S. Fish Com., XXII, 1902, 443, fig. 15, Honolulu.
- Genus 729. *GNATHANODON* Bleeker.
Gnathanodon Bleeker, Verh. Batav. Genootsch., XXIV, Makreele, art. 5, 1851 (1852), pp. 6, 18, 30, 72, etc. (*Scomber speciosus* Forskål).
Hypocaranx Klunzinger, Fische Rothen Meeres, 1884, 92 (*Scomber speciosus* Forskål).
2129. *Gnathanodon speciosus* (Forskål). *Mojarra dorada*.
Tropical parts of the Pacific Ocean; East Indies; Mazatlan to Panama; Hawaii.
Scomber speciosus Forskål, Descr. Anim., 1775, 54, Red Sea.
Scomber rim Forskål, loc. cit., 54, Djidda.
Caranx panamensis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 166, Panama.
- Genus 730. *CITULA* Cuvier.
Citula Cuvier, Règne Anim., ed. 1, 1817, 315 (*Scizena armata* Forskål).

2130. *Citula orthogramma* (Jordan and Gilbert).
Revillagigedo Islands; Pacific Islands.
Caranx orthogrammus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 226, Sulphur Bay, Clarion Island, Revillagigedo Archipelago.
2131. *Citula dorsalis* (Gill). *Pampano*.
Pacific coast of tropical America; Mazatlan; Panama.
Carangoides dorsalis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 166, Panama.
Caranx otrynter Jordan and Gilbert, Proc. U. S. Nat. Mus., 1883, 202; new name substituted for *dorsalis* preoccupied in *Caranx*, but not in *Citula*.
- Genus 731. *ALECTIS*⁵⁰ Rafinesque. *Cobblerfishes*; *Cockfishes*.
Gallus Lacépède, Hist. Nat. Poiss., IV, 1803 (*Zeus gallus* L.); preoccupied in birds.
Alectis Rafinesque, Anal. Nature, 1815, (*Zeus gallus* L.); substitute for *Gallus*, preoccupied.
Blepharis Cuvier, Règne Animal, 1817, 322 (*Zeus ciliaris* Bloch).
Gallichthys Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 264 (*Zeus gallus* L.); substitute for *Gallus*, preoccupied.
Bleparichthys Gill, Cat. Fish. East Coast U. S., suppl. to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 36; substitute for *Blepharis*.
2132. *Alectis crinitus* (Mitchill). *Threadfish*; *Cobblerfish*; *Sunfish*; *Shoemaker*.
West Indies north to Cape Cod, Florida Keys and Cuba.
Zeus crinitus Mitchill, Amer. Jour. Sci. Arts, XI, 1826, 144, Shoreham, Mass.
Blepharis sutor Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 120 (161), pl. 253, Caribbean Sea.
Blepharis major Cuvier and Valenciennes, loc. cit., 121 (163), West Indies; Martinique; not *Gallichthys major*, an East Indian species.
Carangoides blepharis Bleeker, Verhand. Batav. Genootsch., XXIV, 1852, Makr. 667, East Indies.
Carangoides gallichthys Bleeker, loc. cit., 668, East Indies.
- Genus 732. *SCYRIS* Cuvier.
Scyris Cuvier, Règne Anim., ed. 2, II, 1829, 209 (*S. indica* Cuvier and Valenciennes).
2133. *Scyris analis* Poey.
Cuba.
Scyris analis Poey, Synopsis, 1868, 369, Cuba.
- Genus 733. *HYNNIS* Cuvier and Valenciennes.
Hynnys Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 145 (195), pl. 257 (*H. gorensis* Cuvier and Valenciennes).
2134. *Hynnys cubensis* Poey.
Cuba.
Hynnys cubensis Poey, Memorias, II, 1861, 235, Habana.
2135. *Hynnys hopkinsi* Jordan and Starks. *Pampano*.
Puerto Viejo, near Mazatlan, Mexico.
Hynnys hopkinsi Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 435, pl. 35, Mazatlan.
- Genus 734. *VOMER* Cuvier and Valenciennes.
Vomer Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 141 (189), pl. 256 (*V. brownii* Cuvier and Valenciennes=*Zeus setapinnis* Mitchell).
Platysomus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 250 (*Vomer browni* Cuvier and Valenciennes).

⁵⁰ The genera and species allied to *Alectis* have been much mixed from the first. While other possibilities exist, and may be defended, it seems best to adhere to current nomenclature, which, in fact, accords best with the Code.

- 2136. *Vomer setapinnis* (Mitchill).** *Jorobado; Moonfish; Horsefish; Blunt-nosed shiner; Sunfish; Pug-nosed shiner.*
Atlantic coast of America; from Maine to Florida.
Zeus setapinnis Mitchill, Trans. Lit. Phil. Soc. N. Y., 1815, 384, New York.
Vomer brownii Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 141 (189), pl. 256, New York.
Argyreosus unimaculatus Batchelder, Proc. Bost. Soc. Nat. Hist., II, 1845, 78, Maine.
Vomer curtus Cope, Proc. Acad. Nat. Sci. Phila., XXII, 1870, 119, Rhode Island.
- 2137. *Vomer setapinnis cubensis* Nichols.**
Coast of Cuba.
Vomer setapinnis cubensis Nichols, Bull. Amer. Mus. Nat. Hist., XXXVIII, 1918, 672, Cuba.
- 2138. *Vomer dorsalis* Gill.**
West coast of Africa, recorded from the West Indies.
Vomer dorsalis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 436; after Günther.
Argyreosus setipinnis var. B. Günther, Cat., II, 1860, 459, St. Vincent; Cape Verde Islands; West Indies.
Vomer senegalensis Guichenot, Ann. Soc. Linn. Maine et Loire, 1865, 35, Senegal.
Vomer goreensis Guichenot, loc. cit., 37, Gorea.
- 2139. *Vomer spixii* (Swainson).**
West Indies south to Brazil.
Platysomus spixii Swainson, Class. Fishes, etc., II, 1839, 250 and 406, Brazil; after Spix and Agassiz, pl. 37.
Platysomus micropteryx Swainson, loc. cit., 406, Pernambuco.
Vomer sanctæ-marthæ, columbianus, martinicensis, dominicensis, novæ-boracensis, sanctæ-petri, brasiliensis, cayennensis, and cubæ Guichenot, Ann. Soc. Linn. Maine et Loire, 38 to 44, 1865, localities indicated by names.
- 2140. *Vomer declivifrons* Meek and Hildebrand.**
Cape San Lucas to Peru.
Vomer declivifrons Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, 367, Pl. XXIX, fig. 2, Panama.
- 2141. *Vomer gabonensis* Guichenot.**
West Indies and west coast of Africa.
Vomer gabonensis Guichenot, Ann. Soc. Linn. Maine et Loire, 1865, 42, Gaboon.
- Genus 735. ARGYREIOSUS Lacépède.** *Moonfishes.*
Selene Lacépède, Hist. Nat. Poiss., IV, 1803, 560 (*S. argentea* Lacépède = *Zeus vomer* L.).
Argyreiosus Lacépède, loc. cit., IV, 1803, 566 (*Zeus vomer* Linnæus).
- 2142. *Argyreiosus vomer* (Linnæus).** *Moonfish; Jorobado; Lookdown; Horse-head; Hair-finned dory; Hair-finned silverfish; Dollarfish.*
Tropical America, on both coasts; Cape Cod to Brazil; Lower California to Peru.
Zeus vomer Linnæus, Syst. Nat., Ed. X, 1758, 266, America.
Zeus gallus Linnæus, loc. cit., America; after Artedi.
Zeus capillaris Mitchill, Trans. Lit. and Phil. Soc., I, 383, 1815, New York.
Zeus rostratus Mitchill, loc. cit., 384, New York.
Zeus geometricus Mitchill, Amer. Month. Mag., II, 245, 1818, New York.
Argyriosus triacanthus Swainson, Nat. Hist. Classn. Fish., 1839, 250, Brazil.
Argyriosus filamentosus Swainson, loc. cit., 250; after Cuvier and Valenciennes.
Argyriosus setifer Swainson, loc. cit., 407; after Cuvier and Valenciennes.
Argyriosus mauriceii Swainson, loc. cit., 408, Brazil.
Argyriosus mitchilli DeKay, N. Y. Fauna: Fishes, 1842, 126, New York.
- 2143. *Argyreiosus ærstedii* (Lütken).**
Pacific coast, Mazatlan to Panama.
Selene ærstedii Lütken, Spolia Atlantica, 1880, 144, Punta Arenas.

2144. *Argyreusos brevoorti* Gill.

Lower California to Peru.

Argyriosus brevoorti Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 83, Lower California.*Argyreusos pacificus* Lockington, Proc. Calif. Acad. Sci., first ser., VII, 1876, 84, Magdalena Bay, Lower California.Genus 736. *CHLOROSCOMBRUS* Girard. *Casabe*.*Chloroscombrus* Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 168 (*Micropteryx cosmopolita* Agassiz = *Scomber chrysurus* Linnaeus).*Micropteryx* Gunther, not of Agassiz, which is *Seriola*2145. *Chloroscombrus chrysurus* (Linnaeus). *Casabe*; *Bumper*; *Yellowtail*. Cape Cod to Brazil; common on our South Atlantic coast and in Cuba. *Scomber chrysurus* Linnaeus, Syst. Nat., ed. XII, 1766, 494, Charleston, S. C.*Micropteryx cosmopolita* Agassiz, in Spix, Pisc. Bras., 1829, 104, Tab. LIX, Atlantic Ocean.*Scomber latus* Gronow, Cat. Fishes, ed. Gray, 1854, 127, Carolina.*Chloroscombrus caribbaeus* Girard, Mex. Bound. Surv., Zool., 1859, 21, Pl. XI, fig. 6, St. Joseph Island, Tex.2146. *Chloroscombrus orqueta* Jordan and Gilbert. *Orqueta*; *Xurel de Castilla*.

Pacific coast of tropical America, Magdalena Bay to Panama.

Chloroscombrus orqueta Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1862, 646, Panama.*Chloroscombrus stirurus* Jordan and Gilbert, loc. cit., VI, 1883, 206, Panama; slip for *C. orqueta* imperfectly corrected proof.Genus 737. *TRACHINOTUS* Lacépède. *Pámpanos*.*Trachinotus* Lacépède, Hist. Nat. Poiss., III, 1802, 79 (*Labrus falcatus* Linnaeus).*Cæsiomorus* Lacépède, loc. cit., 92 (*C. bailloni* Lacépède).*Acanthion* Lacépède, loc. cit., IV, 1803, 499 (*Chætodon rhomboides* Linnaeus).*Baillonus* Rafinesque, Analyse Nature, X, 1815 (*Trachinotus bailloni* Lacépède); substitute for *Cæsiomorus*.*Bothrolæmus* Holbrook, Ichth. South Carolina, 1855, 80 (*Trachinotus pampanus* Cuvier and Valenciennes).*Dohodon* Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 168 (*Gasterosteus carolinus* Linnaeus).2147. *Trachinotus palometa* Regan. *Gafftopsail*; *Pámpano*; *Oldwife*; *Palometa*; *Banner pompano*.

Tropical America, from Virginia to the Caribbean Sea.

Chætodon glaucus Bloch, Ichth., 1787, pl. 210, Martinique; on a figure by Plumier (name preoccupied by *Lichia glauca* L., also a *Trachinotus*).*Trachynotus palometa* Regan, Ann. Mag. Nat. Hist., (7), XII, 1903, 349; substitute for *glaucus* Bloch).2148. *Trachinotus rhodopus* (Gill). *Pampanito*.

Pacific coast of tropical America; south to Panama.

Trachynotus rhodopus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 85, Cape San Lucas.? *Trachynotus nasutus* Gill, loc. cit., 85, Cape San Lucas.*Trachynotus fasciatus* Gill, loc. cit., 86, Panama.*Trachynotus glacioides* Günther, Proc. Zool. Soc. Lond., 1864, 150, San Jose de Guatemala.2149. *Trachinotus falcatus* (Linnaeus). *Round pampano*; *Palometa*; *Permit* (Indian River) *Spinous dory*; *Ovate pampano*.

East coast of United States, Cape Cod to Florida.

Labrus falcatus Linnaeus, Syst. Nat., ed. X, 1758, 284, America.*Trachinotus spinosus* DeKay, N. Y. Fauna: Fishes, 1842, 117, pl. 19, fig. 53, New York Harbor.*Trachinotus fuscus* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 302 (410), Brazil.

2150. *Trachinotus rhomboides* (Bloch). *Round pampano*.
West Indies to Brazil.
Chaetodon rhomboides Bloch, Nat. Aûsl. Fische, III, 1787, 100, Martinique.
2151. *Trachinotus culveri* Jordan and Starks.
Pacific coast of Mexico.
Trachinotus culveri Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 439, pl. 36, Astillero at Mazatlan, Mexico.
2152. *Trachinotus kennedyi* Steindachner. *Palometa*.
Magdalena Bay to Panama.
Trachinotus kennedyi Steindachner, Ichth. Beitr., III, 1875, 47, Pl. VII, Magdalena Bay.
2153. *Trachinotus goodei* Jordan and Evermann. *Permit*; *Palometa*; *Great pampano*.
West Indies, north to southern Florida.
Trachinotus goodei Jordan and Evermann, Fishes North and Mid. Amer., 1896, 943, Key West, Fla.
2154. *Trachinotus carolinus* (Linnæus). *Common pampano*; *Pámpano*; *Cobblerfish*; *Butterfish*, *Carolina pampano*.
South Atlantic and Gulf coasts of United States; Cape Cod; West Indies; Brazil. One of our best food fishes.
Gasterosteus carolinus Linnæus, Syst. Nat., ed. XII, 1766, 490, Carolina.
? *Trachinotus argenteus* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 304 (410), New York; Rio Janeiro.
? *Trachinotus cupreus* Cuvier and Valenciennes, loc. cit., 305 (414), Martinique.
Trachinotus pampanus Cuvier and Valenciennes, loc. cit., 305 (415), Brazil; Charleston, S. C.
2155. *Trachinotus paloma* Jordan and Starks.
Cape San Lucas, Mazatlan, San Juan Lagoon.
Trachinotus paloma Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 437, Mazatlan, Mexico.
- Genus 738. *ZALOCYS* Jordan and McGregor.
Zalocys Jordan and McGregor (R. C.), Rept. U. S. Fish. Com., XXIV, 1898 (1899), 276 (*Zalocys stilbe* Jordan and McGregor).
- 2156 *Zalocys stilbe* Jordan and McGregor.
Off Pacific coast of Mexico, Revillagigedo Islands.
Zalocys stilbe Jordan and McGregor, Rept. U. S. Fish. Com., XXIV, 1898 (1899), 277, pl. 5, Clarion Island.
- Genus 739. *OLIGOPLITES* Gill. *Leatherjackets*.
Oligoplites Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 166 (*O. occidentalis* Gill).
2157. *Oligoplites saurus* (Bloch and Schneider). *Leatherjacket*; *Leathercoat*; *Runner*; *Zapatero*; *Quiebra*; *Sauteur*.
Both coasts of Central America, and West Indies, north to New York and Lower California
Scomber saurus Bloch and Schneider, ed. Bloch, Syst. Ichth., 1801, 321, Jamaica; after *Saurus argenteus* of Browne.
Centronotus argenteus Lacépède, Hist. Nat. Poiss., III, 1802, 316, equatorial America.
Lichia quiebra Quoy and Gaimard, Voy. Freycinet, Zool., 1824, 365, equatorial America.
Chorinemus saltans Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 289 (393), Martinique, Brazil, and San Domingo.
2158. *Oligoplites inornatus* Gill. *Plain leatherjacket*.
Panama.
Oligoplites inornatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 166 Panama.
2159. *Oligoplites saliens* (Bloch). *Sauteur*.
West Indies.
Scomber saliens Bloch, Naturgeschichte. Aûsl. Fische, VII, 1793, 49, Martinique; on a figure by Plumier
Scomberoides saltator Lacépède, Hist. Nat. Poiss., II, 1800, pl. 19, fig. 2.

2160. *Oligoplites palometa* (Cuvier and Valenciennes).
Lake Maracaibo, Venezuela.
Chorinemus palometa Cuvier and Valenciennes, Hist. Nat. Poiss., VIII,
1831, 288 (392), Lake Maracaibo, Venezuela.
2161. *Oligoplites altus* (Günther). *Jurel*; *Monda*.
Pacific coast of tropical America; Panama.
Chorinemus altus Gunther, Trans. Zool. Soc. London, VI, 1869, 433,
Panama.
2162. *Oligoplites mundus* Jordan and Starks.
West coast of Mexico; Mazatlan.
Oligoplites mundus Jordan and Starks, in Jordan and Evermann,
Check List, 1896, 344, Mazatlan, Mexico (name only); Jordan
and Evermann, Fishes North and Mid. Amer., 1898, 2844, San
Juan Lagoon, Mexico.
2163. *Oligoplites refulgens* Gilbert and Starks.
Panama.
Oligoplites refulgens Gilbert and Starks, Mem. Calif. Acad. Sci., IV,
1904, 73, Panama.

Family 180. SERIOLIDÆ

Genus 740. NAUCRATES Rafinesque.

- Naucrates* Rafinesque, Caratteri 1810, 44 (*Centronotus conductor*
Lacépède=*Gasterosteus ductor* Linnæus).
Centronotus Lacépède, Hist. Nat. Poiss., III, 1802, 311, 51 (*C. conductor*
Lacépède).
Xystophorus Richardson, Voy. *Erebus* and *Terror*, 1844, 52, pl. 30,
fig. 22 (larval form, no species named).
Naucleus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 247
(*N. compressus*, Cuvier and Valenciennes; young of *Naucrates*).
2164. *Naucrates ductor*. (Linnæus). *Pilotfish*; *Romero*.
Atlantic coast from Cape Cod to West Indies.
Gasterosteus ductor Linnæus, Syst. Nat., ed. X, 1758, 295, "in Pelago."
Centronotus conductor Lacépède, Hist. Nat. Poiss., III, 1802, 311, no
locality.
Naucrates fanfarus Rafinesque, Caratteri, 1810, 45, Sicily.
Naucrates noveboracensis Cuvier and Valenciennes, Hist. Nat. Poiss.,
VIII, 1831, 239 (325), New York.
Seriola dussumieri Cuvier and Valenciennes, loc. cit., IX, 1833, 162
(217), Gulf of Bengal.
Seriola succincta Cuvier and Valenciennes, loc. cit., 163 (218), south of
St. Helena.
Naucleus compressus Cuvier and Valenciennes, loc. cit., 185 (249),
pl. 263, fig. 1, Molucca.
Naucleus abbreviatus Cuvier and Valenciennes, loc. cit., 187 (251),
Atlantic Ocean.
Naucleus brachycentrus Cuvier and Valenciennes, loc. cit., 187 (253),
Indian Ocean.
Naucleus triacanthus Cuvier and Valenciennes, loc. cit., 188 (253),
Atlantic Ocean.
Naucleus annularis Cuvier and Valenciennes, loc. cit., 189 (254), south
of St. Helena.
Naucleus lecurus Cuvier and Valenciennes, loc. cit., 189 (255), south
of St. Helena.
Naucrates cyanophrys Swainson, Classn. Fishes, II, 1839, 412, Palermo.
Naucrates serratus Swainson, loc. cit., 413, Palermo.
Thynnus pompilus Gronow, Cat. Fishes, 1854, 123, Mediterranean;
seas of America and Spain.
Xystophorus Richardson, Voy. *Erebus* and *Terror*, 1844, 52, Palermo.
- 2165 *Naucrates polysarcus* Fowler.
Pacific coast of Mexico.
Naucrates polysarcus Fowler, Proc. Acad. Nat. Sci. Phila., LVII, 1905,
62, west coast of Mexico.

Genus 741. *ELAGATIS* Bennett.

Elagatis Bennett, Narrative of Whaling Voyage, II, 1840, 283 (*Seriola bipinnulata* Quoy and Gaimard).

Seriolichthys Bleeker, Nat. Tijdschr. Neder.-Indië, VI, 1854, 196 (*Seriola bipinnulata* Quoy and Gaimard).

Decaptus Poey, Memorias, II, 1861, 391 (*Seriola pinnulata* Poey).

2166. *Elagatis bipinnulatus* (Quoy and Gaimard). *Runner; Yellowtail; Skippack; Shoemaker.*

Tropical seas; West Indies; Long Island; Hawaii.

Seriola bipinnulata Quoy and Gaimard, Voy. *Uranie*, Zool., I, 1824, 363, pl. 61, fig. 3, Keeling Islands.

Seriola pinnulata Poey, Memorias, II, 1861, 233, Habana, Cuba.

Genus 742. *SERIOLA* Cuvier.

Seriola Cuvier, Règne Animal, ed. 2, II, 1829, 205 (*Caranx dumerili* Risso).

Micropteryx Agassiz, Pisc. Brasil., 1829, 102, type (not named) *Seriola dumerili* Cuvier, substitute for *Seriola*, used in botany.

Zonichthys Swainson, Nat. Hist. Fishes, Amphib., etc., II, 1839, 248 (*Scomber fasciatus* Bloch).

Halatractus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 442 (*Seriola dumerili*, substitute for *Seriola*, used in botany).

Lepidomegas Thominot, Bull. Soc. Philom. Paris, ser. 7, IV, 1880, 173 (*L. muelleri* Thominot).

2167. *Seriola dorsalis* (Gill). *Yellowtail; White salmon.*

Pacific coast of America from Monterey southward to Mazatlan; abundant about the Santa Barbara Islands.

Halatractus dorsalis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 84, Cape San Lucas.

2168. *Seriola zonata* (Mitchill). *Rudderfish; Shark pilot; Pilotfish; Banded rudderfish; Jackfish; Banded mackerel.*

Atlantic coast of the United States from Cape Cod to Cape Hatteras.

Scomber zonatus Mitchill, Trans. Lit. Phil. Soc. N. Y., 1815, 427, New York Bay.

2169. *Seriola carolinensis* Holbrook.

Gulf of Mexico, and on Atlantic coast north to Cape Hatteras.

Seriola carolinensis Holbrook, Ichthyol. South Carolina, 1855, 70, Charleston, S. C.

Seriola stearnsi Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 48, Pensacola, Fla.

2170. *Seriola lalandi* Cuvier and Valenciennes. *Great amberfish; Amber jack; Coronado; Jenny Lind.*

New Jersey to Brazil; occasionally to New York.

Seriola lalandi Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 155 (208), Brazil.

Seriola gigas Poey, Memorias, II, 1861, 227, Cuba.

2171. *Seriola dumerili* (Risso). *Amber jack; Coronado.*

Mediterranean to West Indies, north to Key West and Pensacola.

Caranx dumerili Risso, Ichthyol. Nice, 1810, 175, pl. 6, fig. 20, Nice.

Trachurus aliciolus Rafinesque, Caratteri, 1810, 42, Pl. XI, fig. 2, Sicily.

Trachurus fasciatus Rafinesque, Indice, No. 108, 21 and 53, Sicily; not *Scomber fasciatus* of Bloch.

Seriola proxima Poey, Memorias, II, 1861, 229, Cuba.

Genus 743. *ZONICHTHYS* Swainson.

Zonichthys Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 248 (*Scomber fasciatus* Bloch).

2172. *Zonichthys mazatlanus* (Steindachner).

West coast of Mexico, Mazatlan.

Seriola mazatlanana Steindachner, Ichth. Beitr., V, 1876, 8, Mazatlan, Mexico.

2173. *Zonichthys fasciatus* (Bloch). *Madregal; Bonito.*

West Indies, north to Charleston, S. C.

Scomber fasciatus Bloch, Nat. Ausl. Fische, VII, 1793, 73, pl. 341, no locality.

Seriola semicoronata Poey, Memorias, II, 1861, 232, Habana.

2174. *Zonichthys rivolianus* Cuvier and Valenciennes.
Mediterranean to Brazil, West Indies, and South Carolina; Florida coast.
Seriola rivoliana Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 154 (207), the Greek Archipelago.
Seriola boscai Cuvier and Valenciennes, loc. cit., 156 (208), waters of Carolina.
Seriola bonariensis Cuvier and Valenciennes, loc. cit., 157 (211), Buenos Aires.
Seriola coronata Poey, Memorias, II, 1861, 232, Cuba.
2175. *Zonichthys falcatus* (Cuvier and Valenciennes). *Madregal*; *Rock salmon*.
West Indies, north to Florida and the Carolinas.
Seriola falcata Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 156 (210), Gulf of Mexico.
Seriola dubia Lowe, Proc. Zool. Soc. London, VII, 1839, 80, Madeira.
Seriola declivis Poey, Memorias, II, 1861, 230, Habana.
Seriola ligulata Poey, loc. cit., 231, Cuba.

Family 181. NEMATISTIDÆ. Peacockfishes; Papagallos

Genus 744. NEMATISTIUS Gill.

- Nematistius* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 258 (*N. pectoralis* Gill).
Seriolophus Guichenot, Mém. Soc. Nat. Sci. Cherbourg, XIII, 1868, 90 (*Seriolophus carangoides* Guichenot).
2176. *Nematistius pectoralis* Gill. *Per de gallo*; *Papagallo*; *Roosterfish*.
Gulf of California to Panama; generally common.
Nematistius pectoralis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 259, Cape San Lucas.
Seriolophus carangoides Guichenot, Mém. Soc. Nat. Sci. Cherbourg, XIII, 1868, 90, locality unknown.

Family 182. POMATOMIDÆ. Bluefishes

Genus 745. POMATOMUS Lacépède. Bluefishes.

- Pomatomus* Lacépède, Hist. Nat. Poiss., IV, 1802, 436 (*P. skib* Lacépède).
Lopharis Rafinesque, Caratteri, 1810, 52 (*Perca lophar* Forskål).
Gonenion Rafinesque, loc. cit., 53 (*G. serra* Rafinesque).
Temnodon Cuvier, Règne Animal, ed. I, II, 1817, 346 (*Cheilodipterus heptacanthus* Lacépède).
Sypterus Eichwald, Fauna Caspio-Caucas., 1841 (*Perca lophar* Forskål).
Chromis Gronow, Cat. Fishes, 1854, 149 (*C. epicurorum* Gronow = *Gasterosteus saltatrix* Linnæus); name preoccupied.
Sparactodon Rochebrune, Bull. Soc. Philom. Paris, ser. 7, IV, 1880, 162 (*S. nainal* Rochebrune).
2177. *Pomatomus saltatrix* (Linnæus). *Bluefish*; *Snap mackerel*; *Skipjack*; *Faiback*; *Snapping mackerel*; *Tailor*; *Snapper*; *Horse mackerel*; *Greenfish*; *Skip mackerel*.
Atlantic coast of United States; eastern part of the Mediterranean and South to Australia and Madagascar; locally and irregularly abundant.
Perca saltatrix Linnæus, Syst. Nat., ed. X, I, 1758, 293, Carolinas.
Perca lophar Forskål, Descr. Anim., 1775, 38, Constantinople.
Cheilodipterus heptacanthus Lacépède, Hist. Nat. Poiss., III, 1802, 542, Fort Dauphin, Madagascar.
Pomatomus skib Lacépède, loc. cit., IV, 1802, 436, Carolinas.
Lopharis mediterraneus Rafinesque, Indice, 1810, 17, after Forskål.
Gonenion serra Rafinesque, Caratteri, 1810, 53, Sicily.
Chromis epicurorum Gronow, Cat., ed. Gray, 1854, 149, Carolinas.
Sparactodon nainal De Rochebrune, Bull. Soc. Philom. Paris, 1880, 159, Senegambia.

Family 183. RACHYCENTRIDÆ. The Sergeantfishes

Genus 746. RACHYCENTRON Kaup. *Crab-eaters*.*Rachycentron* Kaup, Isis, XIX, 1826, col 89 (*R. typus* Kaup).*Elacate* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 328 (*E. malabarica* Cuvier and Valenciennes).*Mccladerma* Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 243(*M. nigerrima* Swainson = *Elacate pondicerriana* Cuvier and Valenciennes).2178. *Rachycentron canadus* (Linnæus). *Sergeantfish; Crab-eater; Bonito; Cobia*.

In all warm seas; Atlantic coast north to Cape Cod; also in Japan and the East Indies, if all members of the genus belong to one species.

Gasterosteus canadus Linnæus, Syst. Nat., ed. XII, 1766, 491, Carolinas.*Centronotus gardenii* Lacépède, Hist. Nat. Poiss., III, 1802, 357, Carolinas.*Centronotus spinosus* Mitchill, Trans. Lit. and Phil. Soc., I, 1815, 490, New York.? *Elacate pondicerriana* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 241 (329), Pondicherry.? *Elacate motta* Cuvier and Valenciennes, loc. cit., 244 (332), Orixá.? *Elacate malabarica* Cuvier and Valenciennes, loc. cit., 244 (332), Malabar.*Elacate atlantica* Cuvier and Valenciennes, loc. cit., 245 (334), Brazil.? *Elacate bivitata* Cuvier and Valenciennes, loc. cit., 248 (338), Molucca.*Elacata falcipinnis* Gosse, Nat. Sojourn Jamaica, 1851, 208, Jamaica.

Family 184. BATHYCLUPEIDÆ

Genus 747. BATHYCLUPEA Alcock.

Bathyclupea Alcock, Ann. Mag. Nat. Hist., VIII, 1891, 130 (*B. hoskynii* Alcock).2179. *Bathyclupea argentea* Goode and Bean.

West Indies, in deep water.

Bathyclupea argentea Goode and Bean, Oceanic Ichth., 1895, 190, off Nevis Island, West Indies, at Blake station 37 in 365 fathoms.

Family 185. PERCIDÆ. Perches

Genus 748. PERCA Linnæus. *River perches*.*Perca* Linnæus, Syst. Nat., ed. X, 1758, 289 (*P. fluviatilis* Linnæus).*Epirachys* Schulze, Jahrsber. Naturw. Ver. Magdeburg, 1889, 209 (*Perca fluviatilis* Linnæus).2180. *Perca flavescens* (Mitchill). *Yellow perch; American perch; Ringed perch; Raccoon perch; Yellow Ned; River perch; Lake perch*.

Fresh waters of the eastern United States, chiefly northward and eastward; abundant in the Great Lakes and in coastwise streams from Nova Scotia to North Carolina; common in the tributaries of the upper Mississippi, especially in lakes of Indiana, Michigan, Wisconsin, Iowa, and Minnesota; unknown from central Ohio southward; not known from the Ohio River or the lower Missouri.

Perca americana Schranck, Abh. Privatges. Oberdeutschland, I, 1792, 100, America; name preoccupied in *Morone*.*Morone flavescens* Mitchill, Rept. Fish. N. Y., 1814, 18, near New York City.*Centropomus luteus* Rafinesque, Précis des Découvertes Somnologiques, 1814, 19, Pennsylvania.*Perca notata* Rafinesque, Amer. Month. Mag., II, 1818, 205, Lake Erie.*Perca serrato-granulata* Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 34 (48), New York.*Perca granulata* Cuvier and Valenciennes, loc. cit., 35 (48), pl. 9, New York.*Perca acuta* Cuvier and Valenciennes, loc. cit., 36 (49), pl. 10, Lake Ontario.*Perca gracilis* Cuvier and Valenciennes, loc. cit., 36 (50), Skaneateles Lake, N. Y.

Genus 749. STIZOSTEDION⁵¹ Rafinesque. *American pike perches*.
Stizostedion Rafinesque, Ichth. Ohiensis, 1820, 23 (*Perca salmonea*
 Rafinesque=*Perca vitrea* Mitchill).
Stizostethum, attempted correction.

2181. *Stizostedion vitreum* (Mitchill). *Walleye; Wall-eyed pike; Pike perch; Dory; Glass-eye; Yellow pike; Blue pike; Jack salmon; White-eye; Spike* (small specimens about 8 inches long, at Maumee Bay, Sandusky Bay, Port Clinton, etc.); *Pickarel; River trout*.
 Great Lakes region, upper Mississippi, north to Assiniboia; east to Vermont and Pennsylvania; south to Georgia and Alabama; especially common northward.
Perca vitrea Mitchill, Am. Month. Mag., II, 1818, 247, Cayuga Lake, N. Y.
Lucioperca americana Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 90 (122), pl. 16, New York.
Perca salmonea Rafinesque, Amer. Month. Mag. III, 1818, 354, Ohio River.

2182. *Stizostedion glaucum* Hubbs. *Blue pike*.
 Great Lake region.
Stizostedion glaucum Hubbs, Univ. Mich. Mus. Zool., Misc. Publ. No. 15, July 7, 1926, 58, Pl. IV, fig. 1, Lake Erie.

Genus 750. CYNOPERCA Gill and Jordan.
Cynoperca Gill and Jordan, Bull. U. S. Nat. Mus., X, 1877, 44 (*Lucio-perca canadensis* C. H. Smith, in Griffith's edition of Cuvier's Règne Animal).

2183. *Cynoperca grisea* (DeKay). *Sauger; Sand pike; Gray pike; Pickering*.
 Great Lakes region and westward to Montana, Tennessee, and Arkansas; the common species which varies northward into the next.
Lucioperca grisea DeKay, New York Fauna: Fishes, 1842, 19, New York.
Lucioperca borea Girard, Proc. Acad. Nat. Sci. Phila., 1857, 201, Fort Sarpi, Nebr.
Lucioperca pepinus Estes, in Hallock's Sportsman's Gazetteer, 1827, 322, Lake Pepin.
2184. *Cynoperca canadensis* (Smith). *Sauger; Sand pike; Gray pike*.
 Northeastern North America, from Vermont to the Great Lakes.
Lucioperca canadensis C. H. Smith, in Griffith's edition of Cuvier's Règne Anim., Fishes, 1836, 275, Canada.

Family 186. ETHEOSTOMIDÆ Darters

We begin this series with the species most nearly related to the *Percidæ*, from which the darters are apparently descended. *Percina* is the most primitive genus, and in most respects the highest, the other forms being progressively specialized to their life in swift brooks. From *Percina* two lines diverge in different directions; the allies of *Crystallaria* and *Vigil* to *Ammocrypta* toward a quiescent life, buried in sand; the other through *Pacilichthys*, *Nothonotus* and *Oligocephalus* for life in weedy brooks, culminating in *Microperca* and *Alvarius*, the lowest forms.

Genus 751. PERCINA Haldeman. *Log perches*.
Percina Haldeman, Jour. Acad. Nat. Sci. Phila., VIII, 1842, 330 (*P. nebulosa* Haldeman=*Sciæna caprodes* Rafinesque).
Pileoma DeKay, New York Fauna: Fishes, 1842, 16 (*P. semifasciata* DeKay=*Etheostoma caprodes* Rafinesque).

⁵¹ The European genus *Sander* Oken (*Lucioperca* Cuvier and Valenciennes) differs from *Stizostedion* in the close-set ventrals and the less close approximation of the anal spines.

- 2185 *Percina caprodes* (Rafinesque). *Log perch; Rockfish; Hogmolly; Hogfish*. Great Lakes and streams of the South and West from Quebec to Lake Superior, and Iowa.
Sciæna caprodes Rafinesque, Amer. Month Mag., Vol. II, 1818, 354, Ohio River.
Perca (Percina) nebulosa Haldeman, Jour. Acad. Nat. Sci. Phila., VIII, 1842, 330, Susquehanna River.
Pileoma semifasciatum DeKay, New York Fauna: Fishes, 1842, 16, pl. 50, fig. 162, Lake Champlain, Westport, N. Y.
Percina bimaculata Haldeman, Proc. Bost. Soc. Nat. Hist., 1844, 157, Susquehanna River.
2186. *Percina carbonaria* (Baird and Girard).
 Streams of Mississippi to the Rio Grande.
Pileoma carbonaria Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 387, Rio Salada; Rio Medina; San Pedro Creek, Tex.
2187. *Percina zebra* (Agassiz). *Manitou darter*.
 Lakes of Michigan, northern Indiana, Wisconsin, and northward to Lake Superior; Cayuga Lake Basin. Probably varying into *P. caprodes*.
Pileoma zebra Agassiz, Lake Superior, 1850, 308, Lake Superior.
Percina manitou Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 53, Lake Manitou, Ind.
2188. *Percina rex* (Jordan and Evermann).
 Roanoke River, Va. One of the largest and most imposing of the darters.
Etheostoma rex Jordan and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 357, Pl. XLV, fig. 9, Roanoke River near Roanoke, Va.
- Genus 752. *HADROPTERUS* Agassiz. *Black-sided darters*.
Hadropterus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 305 (*H. nigrofasciatus* Agassiz).
2189. *Hadropterus nigrofasciatus* Agassiz. *Crawl-a-bottom*.
 South Carolina to Louisiana.
Hadropterus nigrofasciatus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 305, Mobile, Ala.
Alvordius spillmani Hay, Proc. U. S. Nat. Mus., III, 1880, 491, Chickasawha River, Miss.
- Genus 753. *ALVORDIUS* Girard.
Alvordius Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 68 (*A. maculatus* Girard).
2190. *Alvordius maculatus* Girard. *Black-sided darter*.
 Great Lakes region to the middle Missouri and north to Minnesota, southward through Missouri, Indiana, and Kentucky to Arkansas; especially common in the Ohio Valley.
Alvordius maculatus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 68, Fort Gratiot, Mich.
Alvordius aspro Cope and Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 51, White River, Indianapolis, Ind.; substitute for *Etheostoma blennioides* of Kirtland, not of Rafinesque; perhaps distinct from *A. maculatus* which, if correctly described, has not been found later.
2191. *Alvordius guntheri* (Eigenmann and Eigenmann). *Günther's darter*.
 Winnipeg south to Iowa.
Etheostoma guntheri Eigenmann and Eigenmann, Amer. Nat., November, 1892, 962, Souris River, Winnipeg.
2192. *Alvordius peltatus* (Stauffer). *Shielded darter*.
 Southeastern Pennsylvania, southward to South Carolina in coastwise streams.
Hadropterus maculatus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 100, tributary of Potomac River, Md.; name preoccupied.
Etheostoma peltatum Stauffer, in Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 233, Conestoga Creek, near Lancaster, Pa.
Etheostoma nevinsense Cope, Proc. Amer. Phil. Soc., XI, 1870, 261, falls of Neuse River, N. C.
Alvordius crassus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 12, near Greenville, S. C.

2193. *Alvordius sellaris* (Radcliffe and Welch). *Maryland darter*.
Streams of Maryland.
Hadropterus sellaris Radcliffe and Welch, Bull. U. S. Fish Com., XXXII, 1912, 31, Swan Creek, Havre de Grace, Md.
2194. *Alvordius ouachitæ* (Jordan and Gilbert). *Ouachita darter*.
Southern Indiana, western Kentucky, southwest to Arkansas, not rare; known from Patoka River, Ind.; lower Green and Obion Rivers, Kentucky and Tennessee; Saline River, Ark.
Etheostoma (*Hadropterus*) *ouachitæ* Jordan and Gilbert, Proc. U. S. Nat. Mus., X, 1887, 49, Saline River, Benton, Ark., a tributary of the Washita.
2195. *Alvordius maxinkuckiensis* (Evermann). *Maxinkuckee darter*.
Lake Maxinkuckee, Ind.
Hadropterus maxinkuckiensis Evermann, Rept. U. S. Fish Com., XXV, 1899 (1900), 366, pl. 17, fig. 2, Aubeenabee Creek, an eastern tributary to Lake Maxinkuckee, Ind.
2196. *Alvordius evermanni* (Moenkhaus). *Evermann's darter*.
Lake Tippecanoe, Ind.
Hadropterus evermanni Moenkhaus, Bull. U. S. Fish Com., XXII, 1902 (1904), 397, with figure on p. 398, Tippecanoe Lake, Ind.
2197. *Alvordius roanoka* (Jordan and Jenkins)
Roanoke River, Va.
Hadropterus roanoka Jordan and Jenkins, Proc. U. S. Nat. Mus., XI, 1883, 353, Roanoke River at Roanoke, Salem, and Allegheny Springs, Va.
2198. *Alvordius macrocephalus* (Cope). *Big-headed darter*.
West slope of the Alleghenies from Pennsylvania southward in mountain streams; known from Youghiogheny River, Pa.; North Fork of Holston River, Saltville, Va.; Middle Fork of Holston River, Glade Spring, Va.; Big Sandy, upper Green, and Cumberland Rivers, Ky.; Clinch River, Tenn.
Etheostoma macrocephalum Cope, Trans. Amer. Phil. Soc., XIII, 1869, 400, Youghiogheny River, Pa.
2199. *Alvordius phoxocephalus* (Nelson).
Ohio to Iowa, south to Kentucky and Oklahoma, in sandy rivers; locally common.
Etheostoma phoxocephalum Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 35, Illinois River and its tributaries.
- Genus 754. *SERRARIA* Gilbert.
Serraria Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 205 (*Hadropterus scierus* Swain).
2200. *Serraria sciera* (Swain). Northern Indiana to Tennessee and Texas.
Hadropterus scierus Swain, Proc. U. S. Nat. Mus., VI, 1883, 252, Bean Blossom Creek, Monroe County, Ind.
2201. *Serraria serrula* (Jordan and Gilbert).
From southern Arkansas southward through eastern Texas.
Hadropterus scierus serrula Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1886, 16, Red River, Fulton, Ark.
- Genus 755. *ERICOSMA* Jordan and Copeland.
Ericosma Jordan and Copeland, Bull. U. S. Nat. Mus., X, 1877, 8 (*Alvordius evdes* Jordan and Copeland).
2202. *Ericosma evdes* (Jordan and Copeland).
Indiana in the Wabash Basin, west to Cedar River, Iowa, and southward in Arkansas, Kentucky, and Tennessee; French Broad River and the Ozark region.
Alvordius evdes Jordan and Copeland, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 51, White River near Indianapolis, Ind.
- Genus 756. *HYPOHOMUS* Cope.
Hypohomus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 449 (*Cottogaster aurantiacus* Cope).

2203. *Hypohomus aurantiacus* (Cope).
Upper Tennessee Basin, North Fork of Holston, Clinch, Watauga,
and French Broad Rivers.
Cottogaster aurantiacus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1869,
211, North Fork of Holston River, Saltville, Va.
2204. *Hypohomus cymatotænia* (Gilbert and Meek).
Western Kentucky and southern Missouri.
Etheostoma (Hadropterus) cymatotænia Gilbert and Meek, Proc. U. S.
Nat. Mus., X, 1887, 51, Niangua River and Osage Fork of the Gas-
conade near Marshfield, Mo.; Sac River near Greenfield, Mo.
2205. *Hypohomus nianguæ* (Gilbert and Meek). *Niangua darter*.
Niangua River, Ozark region of Missouri.
Etheostoma (Hadropterus) nianguæ Gilbert and Meek, Proc. U. S. Nat.
Mus., X, 1887, 52, Niangua River near Marshfield, Mo.
2206. *Hypohomus spilotos* (Gilbert).
Kentucky River.
Etheostoma nianguæ spilotos Gilbert, Proc. U. S. Nat. Mus., X, 1887,
53, Sturgeon Creek, a tributary of Kentucky River near Travelers
Rest, Owsley County, Ky.
- Genus 757. *SWAINIA* Jordan and Evermann.
Swainia Jordan and Evermann, Fishes North and Mid. Amer., 1896,
1040 (*Etheostoma (Hadropterus) squamatus* Gilbert and Swain).
2207. *Swainia squamata* (Gilbert and Swain).
Upper Tennessee River Basin; known from Watauga and French
Broad Rivers.
Etheostoma (Hadropterus) squamatus Gilbert and Swain, Proc. U. S.
Nat. Mus., X, 1887, 50, French Broad River at mouth of Wolf Creek,
Tenn.
- Genus 758. *COTTOGASTER* Putnam.
Cottogaster Putnam, Bull. Mus. Comp. Zool., I, 1863, 4 (*Boleosoma
tessellatum* Thompson; not of DeKay = *Cottogaster putnami* Jordan
and Gilbert).
Rheocrypta Jordan, Bull. U. S. Nat. Mus., X, 1877, 9 (*R. copelandi*
Jordan).
2208. *Cottogaster copelandi* (Jordan). *Copeland's darter*.
Great Lakes region, south to the Black Warrior, westward to Missouri
and through the Ozark region, where it is abundant; also about
Indianapolis in clear brooks.
Rheocrypta copelandi Jordan, Bull. U. S. Nat. Mus., X, 1877, 9, White
River near Indianapolis, Ind.
2209. *Cottogaster putnami* (Jordan and Gilbert). *Putnam's darter*.
Lake Champlain.
Cottogaster putnami Jordan and Gilbert, Synopsis, 1883, 498, Westport
Brook, Lake Champlain, N. Y. Probably intergrades with *C.
copelandi*.
2210. *Cottogaster uranidea* (Jordan and Gilbert).
Lower Wabash Basin to southern Missouri, south through Arkansas
and Alabama to the Escambia; mostly in sandy lowland streams.
Etheostoma (Cottogaster) uranidea Jordan and Gilbert, Proc. U. S. Nat.
Mus., X, 1887, 48, Washita River, Arkadelphia, Ark.
- Genus 759. *IMOSTOMA* Jordan.
Imostoma Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 49 (*Ha-
dropterus shumardi* Girard).
2211. *Imostoma shumardi* (Girard). *Shumard's darter*.
Michigan, Ohio, Indiana, and Illinois southward to Kentucky and
Arkansas.
Hadropterus shumardi Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859,
100, Arkansas River near Fort Smith, Ark.
Etheostoma wrighti McCormick, Bull. Oberlin College Lab., No. 2,
1891, 30, near Oberlin, Ohio.

2212. *Imostoma gilberti* (Evermann and Thoburn). *Gilbert's darter*.
Clinch River, Tenn.
Ulocentra gilberti Evermann and Thoburn in Jordan and Evermann,
Fishes North and Mid. Amer., 1896, 1049, Clinch River at Walker's
ford near Tazewell, Tenn.
2213. *Imostoma meadiæ* (Jordan and Evermann). *Meadie's darter*.
Indian Creek, Powell River basin, eastern Tennessee.
Ulocentra meadiæ Jordan and Evermann, Fishes North and Mid. Amer.,
Nov. 26, 1898, 2852, Indian Creek, Cumberland Gap, Tenn.
2214. *Imostoma longimane* (Jordan).
Basin of James River, Va.
Etheostoma longimana Jordan, Proc. Acad. Nat. Sci. Phila., XL, 1888,
179, tributary of James River, Va.
2215. *Imostoma cheneyi* (Evermann and Kendall). *Cheney's darter*.
Racket River near Norfolk, St. Lawrence County, N. Y.
Cottogaster cheneyi Evermann and Kendall, Bull. U. S. Fish Com.,
XVII, 1897 (Feb. 9, 1898), 129, pl. 8, fig. 8, Racket River, near
Norfolk, N. Y.
2216. *Imostoma podostemone* (Jordan and Jenkins).
Roanoke River Basin, Va.
Etheostoma podostemone Jordan and Jenkins, Proc. U. S. Nat. Mus.,
XI, 1888, 359, Roanoke River, at Roanoke, Salem, and Allegheny
Springs, Va.
- Genus 760. *ETHEOSTOMA* Rafinesque.
Etheostoma Rafinesque, Jour. Phys., etc., 1819, 419 (*E. blennioides*
Rafinesque, as restricted by code).
Dipleston Rafinesque, Ichth. Ohiensis, 1820, 37 (*Etheostoma blennioides*
Rafinesque).
Hyostoma Agassiz, Amer. Jour. Sci., XVII, 1854, 305 (*H. newmani*
Agassiz = *Etheostoma blennioides* Rafinesque).
2217. *Etheostoma blennioides* Rafinesque. *Green-sided darter*.
New York to South Dakota and Kansas, south to lower Alabama
Basin; a beautiful fish.
Etheostoma blennioides Rafinesque, Jour. de Physique, LXXXVIII,
1819, 419, Ohio River.
Hyostoma newmani Agassiz, Amer. Jour. Sci. and Arts, X, VII, 1854,
305, vicinity of Huntsville, Ala.
Pileoma cymatogramma Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860,
327, no locality.
Hyostoma blennioperca Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868,
215, tributaries of Kanawha and Holston Rivers.
- Genus 761. *ULOCENTRA* Jordan.
Ulocentra Jordan, Man. Vertebrates, ed. 2, 1878, 223 (*Arlina atripinnis*
Jordan = *Hyostoma simotermum* Cope).
2218. *Ulocentra simotera* (Cope).
West Virginia, eastern Kentucky, and Tennessee, in the basins of the
Green, Cumberland, and Tennessee Rivers, southward through
Alabama to the Escambia River.
Hyostoma simotermum Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 215,
Holston River and its tributaries, near Nashville, Tenn.
Arlina atripinnis Jordan, Bull. U. S. Nat. Mus., X, 1877, 10, tributary
of Cumberland River near Nashville, Tenn.
? *Etheostoma duryi* Henshall, Jour. Cincinnati Soc. Hist., 1889, 32,
tributary of Tennessee River.
2219. *Ulocentra histrio* (Jordan and Gilbert).
Southern Indiana, southwestward to Arkansas; known from the lower
Wabash and tributaries; also from Green River, Ky., and Black,
Poteau, and Washita Rivers, Ark.
Etheostoma (Ulocentra) histrio Jordan and Gilbert, Proc. U. S. Nat.
Mus., X, 1887, 47, Poteau River near Hackett City, Ark.; Saline
River at Benton, and Washita River at Arkadelphia, Ark.

2220. *Ulocentra blennius* (Gilbert and Swain).
Northern Alabama, in streams tributary to the Tennessee River.
Etheostoma (Rhotheca) blennius Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 55, Cox Creek and Shoal Creek, tributary to Tennessee River, Florence, Ala.
2221. *Ulocentra verecunda* (Jordan and Evermann).
Tributaries of Holston River, Va.; rare.
Etheostoma verecundum Jordan and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 360, Pl. XLV, fig. 12, Middle Fork of the Holston River, about 5 miles south of Glade Spring, Va.
2222. *Ulocentra phlox* (Cope).
Trinity River, Tex.
Boleosoma phlox Cope, Bull. U. S. Nat. Mus., XVII, 1880, 30, Trinity River at Fort Worth, Tex.
2223. *Ulocentra stigmæa* (Jordan). *Speck*
Tennessee and Arkansas to Georgia and Louisiana.
Boleosoma stigmæum Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876 (1877), 311, small tributaries of Etowah and Oostanaula Rivers near Rome, Ga.
Pacilichthys saxatilis Hay, Proc. U. S. Nat. Mus., III, 1880, 495, stream at Enterprise, Miss.
- Genus 762. **BOLEOSOMA** DeKay. *Tessellated darters*.
Boleosoma DeKay, New York Fauna: Fishes, 1842, 20 (*B. tessellatum* DeKay = *Etheostoma olmstedii* Storer).
Arlina Girard, Proc. Acad. Nat. Sci. Phila., 1859, 64 (*A. effulgens* Girard).
Estrella Girard, loc. cit., 65 (*E. atromaculata* Girard).
2224. *Boleosoma olmstedii* (Storer). *Tessellated darter; Grand Orankee darter; Sand darter*.
Lake Ontario to Massachusetts and south to North Carolina, east of the Alleghenies.
Etheostoma olmstedii Storer, Jour., Bost. Soc. Nat. Hist., IV, 1842, 61, pl. 5, fig. 2, Hartford, Conn.
Perca minima Haldeman, Jour. Acad. Nat. Sci. Phila., VIII, 1842, 330, Susquehanna River.
Boleosoma tassellatum DeKay, New York Fauna: Fishes, 1842, 20, pl. 20, fig. 57; New York fresh-water streams.
Estrella atromaculata Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 66, Potomac River, vicinity of Washington, D. C.
? *Boleosoma æsopus* Cope, Proc. Amer. Phil. Soc., XI, 1870, 270, Loyalsoc Creek, Pa.
2225. *Boleosoma nigrum* (Rafinesque). *Johnny darter; Blind simon*.
Eastern United States, Ohio Valley, Great Lakes region, and upper Mississippi, west to Colorado and north to Manitoba.
Etheostoma nigra Rafinesque, Ichth. Ohienis, 1820, 37, Green River, Ky.
Boleosoma maculatum Agassiz, Lake Superior, 1850, 305, Lake Superior.
Boleosoma olmstedii brevipinnis Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 214, Kiskiminitas River, Pa.
Pacilichthys beani Jordan, Proc. U. S. Nat. Mus., VII, 1884, 479, Tabo Creek, Mo.
2226. *Boleosoma effulgens* (Girard). *Tassellated darter*.
Maryland to North Carolina.
Arlina effulgens Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 65, brooks and streams flowing into Potomac River.
2227. *Boleosoma mesæum* (Cope).
Platte River near Fort Kearney, Nebr.
Pacilichthys mesæus Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 232, Platte River near Fort Kearney, Nebr.
2228. *Boleosoma maculatiiceps* Cope.
Upper waters of Catawba River, N. C.
Boleosoma maculatiiceps Cope, Proc. Amer. Phil. Soc., XI, 1870, 269, upper waters of Catawba River, N. C.

- 2229. *Boleosoma vexillare* Jordan.**
Tributaries of the James, Roanoke, and Rappahannock Rivers.
Boleosoma vexillare Jordan, Proc. U. S. Nat. Mus., II, 1879, 235, Rappahannock River at Warrenton, Va.
- 2230. *Boleosoma susanæ* Jordan and Swain. *Susan's darter*.**
Basin of upper Cumberland River, Ky.
Boleosoma susanæ Jordan and Swain, Proc. U. S. Nat. Mus., VI., 1883, 249, Wolf Creek and other tributaries of the Clear Fork of Cumberland River near Pleasant View, Whitley County, Ky.
- Genus 763. VAILLANTIA Jordan.**
Vaillantia Jordan, Bull. U. S. Nat. Mus., XII, 1878, 89 (*Boleosoma camurum* Forbes).
- 2231. *Vaillantia camura* (Forbes).**
Indiana to Iowa and Mississippi, southwest to Houston, Tex.
Boleosoma camurum Forbes, Bull. Ill. Lab. Nat. Hist., I, 1878, 40, Cache River and Clear Creek, Union County, Johnston County, and Pekin, Ill.; name preoccupied in *Nothonotus* but not in *Boleosoma* or *Vaillantia*,
Vaillantia chlorosoma Hay, Proc. U. S. Nat. Mus., III, 1880, 495, Mississippi.
- Genus 764. CRYSTALLARIA Jordan and Gilbert.**
Crystallaria Jordan and Gilbert, in Jordan, Cat. Fishes N. Amer., 1885, 78 (*Pleurolepis asprellus* Jordan).
- 2232. *Crystallaria asprella* (Jordan)**
Southern Indiana and Illinois to Arkansas and Alabama, chiefly in the larger, clearer streams. Recorded from Ohio River at Rising Sun, Ind.; Wabash River at New Harmony, Vincennes, and Terre Haute; Green River, Ky.; Choccolo Creek, Ala.; and Washita River, Ark.; besides the original locality in Illinois.
Pleurolepis asprellus Jordan, Bull. Ill. Lab. Nat. Hist., I, 1878, No. 2, 38, 1878, rocky tributary of Mississippi River in Hancock County, Ill.
- Genus 765. VIGIL Jordan.**
Pleurolepis (Agassiz) Putnam, Bull. Mus. Comp. Zool., I, 1863, 5 (*Etheostoma pellucidum* (Baird) Agassiz; name preoccupied by Egerton for a genus of fossil fishes.
Vigil Jordan, Proc. Acad. Nat. Sci. Phila., LXX, 1918 (1919), 344 (*Pleurolepis pellucidus* Agassiz).
- 2233. *Vigil pellucidus* (Baird). *Sand darter*.**
Lake Erie to Minnesota, Kentucky, and Texas.
Pleurolepis pellucidus Baird, in Agassiz, Bull. Mus. Comp. Zool., I, 1863, 5, no locality; the specimens from Black River, Ohio, the types of *Etheostoma pellucidum* Baird (MS., 1853).
- 2234. *Vigil clarus* (Jordan and Meek).**
Mississippi Valley; Wabash River, west to central Iowa and Minnesota, and south to Arkansas and northern Texas.
Ammocrypta clara Jordan and Meek, Proc. U. S. Nat. Mus., VIII, 1885, 8, Des Moines River, Ottumwa, Iowa.
- 2235. *Vigil vivax* (Hay).**
Mississippi and northwest through Arkansas and south to Texas.
Ammocrypta vivax Hay, Bull. U. S. Fish Com., II, 1882, (1883), 58, Pearl River, Jackson, Miss.
- Genus 766. IOA Jordan and Brayton.**
Ioa Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 88 (*Pæclichthys vitreus* Cope).
- 2236. *Ioa vitrea* (Cope).**
Southeastern Virginia and western North Carolina.
Pæclichthys vitreus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 263, Walnut Creek, a tributary of Neuse River, Wake County, N. C.
- 2237. *Ioa vigil* Hay.**
Pearl River, Jackson, Miss.
Ioa vigil Hay, Bull. U. S. Fish Com., II, 1882 (1883), 59, Pearl River, Jackson, Miss.

- Genus 767. AMMOCRYPTA** Jordan. *Sand darters.*
Ammocrypta Jordan, Bull. U. S. Nat. Mus., X, 1877, 5 (*A. beanii* Jordan).
- 2238. *Ammocrypta beanii* Jordan.**
 Gulf States, Alabama to Louisiana.
Ammocrypta beanii Jordan, Bull. U. S. Nat. Mus., X, 1877, 5, Notalbany River, Tickfaw, La.
Ammocrypta gelida Hay, Proc. U. S. Nat. Mus., III, 1880, 490, Enterprise, Miss.
- Genus 768. PÆCILICHTHYS** Agassiz.
Pæciosoma Agassiz, Lake Superior, 1850, 299 (*Etheostoma variata* Kirtland); name preoccupied.
Pæcilichthys Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 305 (*Etheostoma variata* Kirtland).
- 2239. *Pæcilichthys variatus* (Kirtland).**
 Ohio River Basin, from western Pennsylvania to Kentucky; scarce. Long confused with *Oligocephalus cæruleus*.
Etheostoma variata Kirtland, Zoology of Ohio, 1838, 168, 192, Mahoning River, Ohio
Hadropterus tessellatus Jordan, Bull. U. S. Nat. Mus., X, 1877, 7, Allegheny River, Foxburg, Pa.
- Genus 769. NANOSTOMA** Putnam.
Nanostoma Putnam, in Jordan, Bull. U. S. Nat. Mus., X, 1877, 6 (*Pæcilichthys zonalis* Cope); not *Nannostomus* Gunther).
Rhothæa Jordan, Rept. U. S. Fish Com., XIII, 1885 (1887), 868, p. 80 of reprint issued 1885; substitute for *Nanostoma* Putnam, regarded as preoccupied by *Nannostomus* Günther.
- 2240. *Nanostoma zonale* (Cope). *Banded darter.***
 Mississippi Valley from Ohio and northern Indiana (Kankakee River) west through Iowa and south through Kentucky and Tennessee into Alabama, Louisiana, and Arkansas.
Pæcilichthys zonalis Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 212, Holston River, Va.
Nanostoma vincentipes Jordan, Proc. U. S. Nat. Mus., II, 1879, 236, tributary of Illinois River at Naperville, Ill.
- 2241. *Nanostoma arcansanum* (Jordan and Gilbert).**
 Ozark region of Missouri and Arkansas.
Etheostoma zonale arcansanum Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1886, 5, Spring River, Carthage, Mo.; James River, Marshfield, Mo.; Poteau River, Hackett City, Ark.; Washita River, Arkadelphia, Ark., and Saline River, Benton, Ark.
- 2242. *Nanostoma swannanoa* (Jordan and Evermann).**
 Upper waters of the Tennessee River.
Etheostoma swannanoa Jordan and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 360, Pl. XLV, fig. 13, South Fork of Holston River, Holstein Mills, Va.; Middle Fork of Holston River, Marion, Va. and South Fork of Swannanoa River, Black Mountain, N. C.
- 2243. *Nanostoma elegans* (Hay).**
 Chickasawha River, Miss.
Nanostoma elegans Hay, Proc. U. S. Nat. Mus., III, 1880, 493, shallow rocky branch of Chickasawha River, Enterprise, Miss.
- 2244. *Nanostoma thalassinum* (Jordan and Brayton).**
 Santee River Basin in North and South Carolina.
Nothnotus thalassinus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 13, Reedy River, Greenville, S. C.; Catawba River and tributaries in North Carolina; Ennoree River, Chick Springs, S. C.; and Saluda River, Farr's Mill, S. C.
- 2245. *Nanostoma inscriptum* (Jordan and Brayton).**
 Oconee River, Ga.
Nothnotus inscriptus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 34, Oconee River, Sulphur Springs, Hall County, Ga.

2246. *Nanostoma rupestre* (Gilbert and Swain).
North River, a tributary of Black Warrior River, Ala.
Etheostoma (Rhotheca) rupestre Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 57, North River near Tuscaloosa, Ala.
- Genus 770. *NOTHONOTUS* Agassiz.
Nothonotus Agassiz, Bull. Mus. Comp. Zool., I, 1863, 3 (*Etheostoma maculata* Kirtland).
2247. *Nothonotus maculatus* Kirtland.
Ohio and Indiana, southward through Kentucky and Tennessee to northern Alabama, in tributaries of the Wabash, Ohio, Cumberland, and Tennessee Rivers.
Etheostoma maculata Kirtland, Jour. Bost. Soc. Nat. Hist., 1840, 276, Mahoning River, Ohio.
Pacilichthys sanguifluus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 264, headwaters of the South Fork of the Cumberland River in Tennessee.
2248. *Nothonotus camurus* (Cope). *Blue-breasted darter*.
Lake Erie to Tennessee, in clear, swift waters, not common; the most brilliantly colored of fresh-water fishes
Pacilichthys camurus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 265, headwaters of Cumberland River in Tennessee.
2249. *Nothonotus vulneratus* (Cope).
French Broad River
Pacilichthys vulneratus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 266, tributary of French Broad River, Warm Springs, N. C.
2250. *Nothonotus cinereus* (Storer)
Tennessee and Cumberland Rivers; Tennessee River at Florence, Ala.; Obeys River at Olympus, Tenn.; Rock Creek near Whitley Station, Ky.
Etheostoma cinerea Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 49, Florence, Ala.
2251. *Nothonotus tessellatus* (Storer). *Tessellated darter*.
Florence, Ala.
Etheostoma tessellata Storer, Proc., Bost. Soc. Nat. Hist., II, 1845, 48, Tennessee River at Florence, Ala. (Not *Boleosoma tessellatum* DeKay).
2252. *Nothonotus rufilineatus* (Cope).
Upper tributaries of the Tennessee, Cumberland, and Green Rivers.
Pacilichthys rufilineatus Cope, Proc. Amer. Phil. Soc., XI, 1870, 267, Warm Springs Creek, French Broad River, Madison County, N. C.
2253. *Nothonotus jordani* (Gilbert). *Jordan's darter*.
Tributaries of Coosa River in the Alabama River Basin.
Etheostoma (Nothonotus) jordani Gilbert, Bull. U. S. Fish Com., I, 1889 (1891), 156, pl. 43, fig. 3, Choccolo Creek, Oxford, Ala., and Chestnut Creek, Verbena, Ala.
- Genus 771. *RAFINESQUIELLUS* Jordan and Evermann.
Aplesion Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 102 (*A. pottsi* Girard); not of Rafinesque.
Rafinesquiellus Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1082 (*Aplesion pottsi* Girard).
2254. *Rafinesquiellus pottsi* (Girard).
Streams of Chihuahua, Mexico, in Rio Grande Basin.
Aplesion pottsi (misprinted *potsii*) Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 102, tributaries of Chihuahua River, Mexico.
Etheostoma macropterus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 289, Chihuahua, Mexico.
- Genus 772. *OLIGOCEPHALUS* Girard.
Oligocephalus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 67 (*Boleosoma lepida* Baird and Girard).

- 2255. *Oligocephalus caeruleus* (Storer).** *Blue darter; Rainbow darter; Soldierfish, Blue johnny.*
Upper Mississippi Valley, from Kentucky to Michigan and east to New York.
Etheostoma caerulea Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 47, Fox River, Ill.
Pecilosoma erythrogastrum Kirtland, Cleveland Annals Sci., 1854, 4, near Cleveland, Ohio
Pacilichthys versicolor Agassiz, Amer. Jour. Sci. Arts, 1854, 304, Quincy, Ill.
Pacilichthys spectabilis Agassiz, Amer. Jour. Sci. Arts, 1854, 304, Osage River, Mo.
Pecilosoma transversum Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 326, Lake Superior.
- 2256. *Oligocephalus lepidus* (Baird and Girard).**
Streams of Arkansas, Texas, and Chihuahua; perhaps more than one species included.
Boleosoma lepidus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 388, upper tributaries of the Rio Nueces, Tex.
Oligocephalus grahmi Girard, loc. cit., XI, 1859, 102, Devil River, Tex.
Oligocephalus leonensis Girard, loc. cit., 102, Leon River, Tex.
Oligocephalus pulchellus Girard, loc. cit., 102, Gypsum Creek, a tributary to the Canadian River.
- 2257. *Oligocephalus fricksia* (Hildebrand).**
Tributaries of Savannah River.
Etheostoma fricksia Hildebrand, Bull. U. S. Bur. Fish., XXIX, 1924, 7, Brook near Augusta, Ga.
- 2258. *Oligocephalus iowæ* (Jordan and Meek).**
Upper Mississippi Valley, from Iowa, Nebraska, and Colorado north to Assiniboia; abundant and widely distributed.
Etheostoma iowæ Jordan and Meek, Proc. U. S. Nat. Mus., VIII, 1885, 10, Chariton River, Chariton, Iowa.
Etheostoma quappella Eigenmann and Eigenmann, Amer. Nat., 1892, 963, Qu'Appelle River, Manitoba.
- 2259. *Oligocephalus aubeenaubei* (Evermann).** *Aubeenaubee darter.*
Aubeenaubee Creek, an eastern inlet to Lake Maxinkuckee, Ind.
Etheostoma aubeenaubei Evermann, Rept., U. S. Fish Com., XXV, 1899 (1900), 367, Aubeenaubee Creek, Lake Maxinkuckee, Ind.
- 2260. *Oligocephalus hildebrandti* (Evermann and Clark).**
Fletcher Lake, Fulton Co., Ind.
Etheostoma hildebrandti Evermann and Clark, Proc. Biol. Soc. Wash., XXIII, May 4, 1910, 81-88, fig. 1, Fletcher Lake, Fulton Co., Ind.
- 2261. *Oligocephalus jessiae* (Jordan and Brayton).** *Jessie's darter.*
Southern Illinois to Georgia and Mississippi.
Pacilichthys jessiae Jordan and Brayton, in Jordan, Man. Vert., ed. 2, 1878, 227; creek near Ringgold, Ga.
Pacilichthys asprigenus Forbes, Bull. Ill. Lab. Nat. Hist., I, No. 2, 1877, 41, small creek at Pekin, Ill.
Pacilichthys swaini Jordan, Proc. U. S. Nat. Mus., VII, 1884, 479, tributary of Pearl River, Monticello, Miss.
- 2262. *Oligocephalus luteovinctus* (Gilbert and Swain).**
Stone River, Tenn.
Etheostoma luteovinctum Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 58, Stone River near Nashville, Tenn.
- Genus 773. *NIVICOLA* Jordan and Evermann.**
Nivicola Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1082 (*Pacilichthys borealis* Jordan).
- 2263. *Nivicola borealis* (Jordan).** *Northern darter.*
St. Lawrence River at Montreal, Canada, and Muskoka Lakes near Lake Huron.
Pacilichthys borealis Jordan, Proc. U. S. Nat. Mus., VII, 1884, 477, Montreal, Canada.

2264. *Nivicola cragini* (Gilbert). *Cragin's darter*.
Western portion of the Arkansas River Basin, from Garden City, Kans., to Canyon City, Colo.
Etheostoma cragini Gilbert, Bull. Washburn Coll. Lab., March and April, 1885, 99, small stream connecting the "Lake" at Garden City, Kans., with Arkansas River.
2265. *Nivicola punctulata* (Agassiz).
Ozark region of southwestern Missouri.
Paculichthys punctulatus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 304, Osage River, Mo.
2266. *Nivicola pagei* (Meek).
Tributary of Neosho River, Arkansas Basin.
Etheostoma pagei Meek, Amer. Nat., 1894, 957, Neosho, Mo.; only two specimens known.
2267. *Nivicola artesiae* (Hay).
Georgia to central Texas.
Paculichthys artesae Hay, Proc. U. S. Nat. Mus., III, 1880, 494, from a small branch of Catawba River, Artesia, Lowndes County, Miss.
2268. *Nivicola lepidogenys* (Evermann and Kendall).
Rio Comal, Tex.
Etheostoma lepidogenys Evermann and Kendall, Bull. U. S. Fish. Com., XII, 1892 (Feb. 6, 1894), 114, pl. 35, fig. 3, Rio Comal, New Braunfels, Tex.
2269. *Nivicola virgata* (Jordan).
Tributaries of Cumberland River in western Kentucky.
Paculichthys virgatus Jordan, Proc. U. S. Nat. Mus., II, 1879, 236, Rock Castle River, Livingston, Ky.
2270. *Nivicola obeyense* (Kirsch).
Tributaries of the Cumberland River in Clinton County, Ky.
Etheostoma obeyense Kirsch, Bull. U. S. Fish. Com., X, 1890 (1892), 292, with figure; all streams tributary to the Cumberland River in Clinton County, Ky.
2271. *Nivicola tippecanoe* (Jordan and Evermann).
Tippecanoe River, Ind.
Etheostoma tippecanoe Jordan and Evermann, Proc. U. S. Nat. Mus., XIII, 1890, 3 (with a figure of *Etheostoma camurum* by error of printer), Tippecanoe River, Marshland, Ind.
- Genus 774. **CLARICOLA** Jordan and Evermann.
Claricola Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1093 (*Etheostoma juliae* Meek).
2272. *Claricola juliae* (Meek). *Julia's darter*.
Basin of White River, Mo.
Etheostoma juliae Meek, Bull. U. S. Fish. Com., IX, 1889 (1891), 130, pl. 42, fig. 2, James River, near Springfield, Mo.
2273. *Claricola whipplii* (Girard). *Whipple's darter*.
Lower Arkansas Basin; locally abundant in clear tributaries of the Saline, Washita, etc.
Boleichthys whipplii Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 103, Coal Creek, Ark.
2274. *Claricola alabamæ* (Gilbert and Swain). *Alabama darter*.
Black Warrior and Big Catawba Rivers, Ala.
Etheostoma whipplei alabamæ Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 62, Black Warrior River near Morris and Tuscaloosa, Ala.
2275. *Claricola squamiceps* (Jordan).
Lower Wabash Valley, Ind., through western Kentucky and Tennessee to Georgia and western Florida. A peculiar species resembling *Catnotus*; perhaps not a *Claricola*.
Etheostoma squamiceps Jordan, Bull. U. S. Nat. Mus., X, 1877, 11, Russellville, Ky.
Etheostoma parvipinne Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 59, small spring branch, tributary to Black Warrior River, Tuscaloosa, Ala.

Genus 775. CATONOTUS Agassiz. *Fantail darters.*

Catonotus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 305 (*C. lineolatus* Agassiz).

Etheostoma Rafinesque, as restricted by Cope and Jordan.

Richia Coker, Bull. U. S. Bur. Fish., XLII, 1926, 105, 106; *R. brevispina* Coker, preoccupied by *Richia*, a genus of moths.

Richiella Coker, Copeia, No. 162, March, 1927, 17, substitute for *Richia*, preoccupied.

2276. *Catonotus flabellaris* (Rafinesque). *Fan-tailed darter.*

New York to Virginia, west to Iowa and south to South Carolina and northern Alabama.

Etheostoma flabellaris Rafinesque, Jour. de Physique, LXXXVIII, 1819, 419, tributaries of Ohio River.

Etheostoma flabellata Rafinesque, Ichth. Ohiensis, 1820, 36, Ohio River.

Etheostoma fontinalis Rafinesque, loc. cit., 85, Ohio River.

Etheostoma linsleyi Storer, Proc. Bost. Soc. Nat. Hist., IV, 1851, 37, Wolcott, Wayne County, N. Y.

Oligocephalus humeralis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 67, James River, Va.

Catonotus fasciatus Girard, loc. cit., 68, Madrid, N. Y., and Grass River.

Catonotus kennicottii Putnam, Bull. Mus. Comp. Zool., I, 1863, 3, southern Illinois.

Richia brevispina Coker, Bull. U. S. Bur. Fish., XLII, March, 1927, 106, fig. 1, Paddys Creek, N. C.

2277. *Catonotus flabellare lineolatum* (Agassiz).

Minnesota to northern Indiana and northern Missouri.

Catonotus lineolatus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 305, small creek near Quincy, Ill.

2278. *Catonotus flabellare cumberlandicum* (Jordan and Swain). *Cumberland darter*

Brooks in Cumberland Mountains tributary to Cumberland River.

Etheostoma cumberlandicum Jordan and Swain, Proc., U. S. Nat. Mus., VI, 1883, 251, small streams of the Cumberland Mountains; Wolf Creek and Briar Creek near Pleasant View, Whitley County, Ky.

Genus 776. TORRENTARIA Jordan and Evermann.

Torrentaria Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1080 (*Etheostoma sagitta* Jordan and Swain).

2279. *Torrentaria sagitta* (Jordan and Swain).

Cumberland River.

Pacilichthys sagitta Jordan and Swain, Proc. U. S. Nat. Mus., VI, 1883, 250, Wolf Creek, near Pleasant View, Whitley County, Ky.

2280. *Torrentaria fasciata* (Girard).

Chihuahua River, Mexico, in Rio Grande Basin.

Diplesion fasciatus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 101, Chihuahua River, Mexico; the name preoccupied in *Etheostoma*, not in *Torrentaria*.

Etheostoma australe Jordan, Proc. U. S. Nat. Mus., XI, 1888, 362, Chihuahua River, Mexico; based on the types of *Diplesion fasciatus* Girard.

Etheostoma scovellii Woolman, Amer. Nat., March, 1892, 260, Rio de las Conchas, Chihuahua, Mexico.

Genus 777. PSYCHROMASTER Jordan and Evermann.

Psychromaster Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1099 (*Etheostoma tuscumbia* Gilbert and Swain).

2281. *Psychromaster tuscumbia* (Gilbert and Swain). *Tuscumbia darter.*

Tennessee River Basin in northern Alabama

Etheostoma tuscumbia Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 63, stream flowing from the large spring at Tuscumbia, Ala.

Genus 778. BOLEICHTHYS Girard.

Boleichthys Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 103 (*Boleichthys exilis* Girard).

2282. *Boleichthys fusiformis* (Girard). *Fusiform darter*.
Massachusetts and south in coastwise streams.
Boleosoma fusiforme Girard, Proc. Bost. Soc. Nat. Hist., IV, 1854, 41,
Charles River, Mass.
Hololepis erochrous Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 233,
Brown's Mill, N. J., opposite Philadelphia
2283. *Boleichthys gracilis* (Girard). *Crimson darter*.
Southern Indiana to Georgia and Texas; perhaps more than one species
included; perhaps varying into the preceding.
Boleosoma gracile Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 103,
Rio Seco and Leona River, Tex.
Pacilichthys bullerianus Hay, Bull. U. S. Fish Com., II, 1882, 61, Big
Black River near Vaughns station, Yazoo County, Miss.
Pacilichthys palustris Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 209,
Switz City Swamp, Greene County, Ind.
2284. *Boleichthys eos* (Jordan and Copeland).
New York to Minnesota.
Boleichthys eos Jordan and Copeland, Proc. Acad. Nat. Sci. Phila.,
XXIX, 1877, 46, Wisconsin and Rock Rivers, Wis.; Fox River, Ill.;
northern Indiana.
2285. *Boleichthys exilis* Girard.
Upper Missouri River Basin and Red River of the North.
Boleichthys exilis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 103,
Little Muddy River, a tributary of the upper Missouri.
Boleichthys warreni Girard, loc. cit., 104, Cannonball River.
- Genus 779. *HOLELEPIS* Agassiz.
Hololepis Agassiz, in Putnam, Bull. Mus. Comp. Zool., I, 1863, 4
(*Etheostoma barrati* Holbrook).
Copelandellus Jordan and Evermann, Fishes North and Mid. Amer.,
1896, 1100 (*Pacilichthys quiescens* Jordan).
2286. *Hololepis barrati* Holbrook.
Swamps and streams of the lowlands from Virginia to Florida; known
from Blackwater River, Zuni, Va.; Allapaha River, Nashville, Ga.;
and various streams in Florida and South Carolina.
Hololepis barrati Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 56,
South Carolina.
Pacilichthys quiescens Jordan, Proc. U. S. Nat. Mus., VII, 1884, 478,
tributary of Allapaha River, Nashville, Ga.
- Genus 780. *MICROPERCA* Putnam.
Microperca Putnam, Bull. Mus. Comp. Zool., I, 1863, 4 (*M. punctulata*
Putnam).
2287. *Microperca punctulata* Putnam. *Least darter*.
Northwestern States; Indiana, Michigan, and Minnesota, south to
Arkansas; rare outside of tributaries of the Great Lakes.
Microperca punctulata Putnam, Bull. Mus. Comp. Zool., I, 1863, 4,
various points in Michigan, Wisconsin, Illinois, and Alabama; those
from Alabama probably *M. praeliarius* (Hay); name preoccupied in
Etheostoma.
Etheostoma microperca Jordan and Gilbert, in Jordan's Man. Vert.,
ed. V, 1888, 134, lakes of Laporte County, Ind., substitute name.
2288. *Microperca praeliarius* Hay.
Alabama and Mississippi; in lowland streams and ponds.
Microperca praeliarius Hay, Proc. U. S. Nat. Mus., III, 1880, 496, small
branch of the Tuscumbia River at Corinth, Miss.
- Genus 781. *ALVARIUS* Girard.
Alvarius Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 101 (*A. lateralis*
Girard).
This genus differs from *Microperca* in having but one anal spine. The
fins are short and low and the scales in the type are said to be small.
2289. *Alvarius lateralis* Girard.
Rio Grande about Brazos Santiago, Tex.
Alvarius lateralis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 101,
mouth of Rio Grande.

2290. *Alvarius fonticola* Jordan and Gilbert.
Arkansas and Texas, in clear rocky streams; known only from Washita River at Arkadelphia; San Marcos River at San Marcos, Tex.; Comal Creek at New Braunfels, Tex.
Alvarius fonticola Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1886, 23, San Marcos Spring at San Marcos, Tex.

Family 187. APOGONIDÆ. Cardinal fishes

- Genus 782. APOGON Lacépède.
Amia Gronow, Zoophyl., 1763, 80 (*Apogon moluccensis* Cuvier and Valenciennes); not *Amia* L., nonbinomial
Apogon Lacépède, Hist. Nat. Poiss., III, 1802, 411 (*A. ruber* Lacépède = *Mullus imberbis* L.).
Ostorhinchus Lacépède, loc. cit., IV, 1803, 23 (*O. fleureiu* Lacépède).
Dipterodon Lacépède, loc. cit., 165 (*D. hexacanthus* Lacépède).
Macrolepis Rafinesque, Analyse de la Nat., 1815, 86; a bare name, no type.
Monoprion Poey, Memorias, II, 1861, 123 (*M. maculatus* Poey).
2291. *Apogon imberbis* (Linnaeus). *King of the mullets*; *Alfoncino*; *Fucinita*.
Mediterranean and neighboring waters; one from Newport, R. I., and one from the island of Fernando de Noronha
Mullus imberbis Linnaeus, Syst. Nat., ed. X, 1758, 300, Malta; after Artedi.
Apogon ruber Lacépède, Hist. Nat. Poiss., III, 1802, 411, Malta.
Perca pusilla De la Roche, Ann. Mus. Nat. Hist., XIII, 1809, 318, Nice.
Apogon rex-mullorum Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 105 (143), Marseilles, Nice, Genoa, Naples, Palermo.
2292. *Apogon maculatus* (Poey).
Pensacola to Bahia; recorded from as far north as Woods Hole; common on the "snapper banks"; often found in the stomachs of snappers and groupers.
Monoprion maculatus Poey, Memorias, II, 1861, 123, Cuba.
2293. *Apogon pigmentarius* (Poey).
Cuba; common at Habana.
Monoprion pigmentarius Poey, Memorias, II, 1861, 123, Cuba.
2294. *Apogon sellicauda* Evermann and Marsh.
West Indies; Tortugas Archipelago; Porto Rico.
Apogon sellicauda Evermann and Marsh, Fish. Porto Rico, Bull. U. S. Fish Com., XX, 1900, 143, fig. 40, Culebra Island, Porto Rico.
2295. *Apogon americanus* Castelnau.
Brazil, north to the Tortugas.
Apogon americanum Castelnau, Anim. Amer. Sud, Poiss., 1855, 3, pl. III, fig. 2, Bahia.
2296. *Apogon binotatus* (Poey).
Cuba.
Amia binotata Poey, Repertorio, 1867, 234, Cuba.
2297. *Apogon retrosella* (Gill). *Cardenal*.
Pacific coast of Mexico.
Amia retrosella Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 251, Cape San Lucas.
2298. *Apogon dovii* Günther.
Mazatlan to Panama.
Apogon dovii Günther, Proc. Zool. Soc. Lond., 1861, 371, Panama.
2299. *Apogon atricaudus* (Jordan and McGregor).
Offshore islands, Pacific coast of Mexico.
Apogon atricaudus Jordan and McGregor, Rept. U. S. Fish Com., XXIV, 1898, 277, and in Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2853, Socorro, Clarion, and San Benedicto islands.
2300. *Apogon atrodorsatus* Heller and Snodgrass.
Galapagos Islands.
Apogon atrodorsatus Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 192, Pl. III, Charles Island.

2301. *Apogon guadalupensis* (Osburn and Nichols).
Guadalupe Island, off Lower California.
Ama guadalupensis Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 160, fig. 9, Guadalupe Island.
- Genus 783. *APOGONICHTHYS* Bleeker.
Apogonichthys Bleeker, Naturkundig Tijds. Nederl.-Indie, Floris, VI, 1854, 321 (*A. perdis* Bleeker).
2302. *Apogonichthys alutus* (Jordan and Gilbert).
Snapper banks off Pensacola and Tampa, Fla.; and at Porto Rico.
Apogon alutus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 279, and in Synopsis, 1883, 931, snapper banks off Pensacola, Fla.
2303. *Apogonichthys puncticulatus* Poey.
Cuba; Tortugas.
Apogonichthys puncticulatus Poey, Repertorio, II, 1867, 233, Cuba.
2304. *Apogonichthys melampodus* Blosser.
St. Croix Island, West Indies
Apogonichthys melampodus Blosser, Ann. Carnegie Mus., VI, 1909, No. 1, 296, St. Croix.
- Genus 784. *ASTRAPOGON* Fowler.
Astrapogon Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1906, 527 (*Apogonichthys stellatus* Cope).
2305. *Astrapogon stellatus* (Cope).
Bahamas.
Apogonichthys stellatus Cope, Trans. Amer. Phil. Soc., XIII, 1869, 400, Nassau, Bahama Islands.
Apogonichthys strombi Plate, Zool. Anzeiger, XXXIII, 1908, 393, with plate, Bahamas; living symbiotic in shells of *Strombus gigas*.
- Genus 785. *XYSTRAMIA* Jordan.
Xystramia Jordan, Copeia, May 24, 1917, 46 (*Glossamia pandionis* Goode and Bean).
2306. *Xystramia pandionis* (Goode and Bean).
Deep water off Chesapeake Bay.
Apogon pandionis Goode and Bean, Proc. U. S. Nat. Mus., IV, 1881, 160, deep water off Chesapeake Bay.
- Genus 786. *EPIGONUS* Rafinesque.
Epigonus Rafinesque, Indice, 1810, 64 (*E. macrophthalmus* Rafinesque).
Pomatomus Risso, in Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 171 (*P. telescopium* Risso); not of Lacépède.
Telescopus Bleeker, Arch. Néerl. Sci. Nat., XI, 1876, 261 (*Pomatomus telescopium* Risso); substitute for *Pomatomus* Cuvier; not of Lacépède.
Pomatomichthys Giglioli, Espos. Int. d. Pesca, Berlin, 1880, 20 (*P. constanciae* Giglioli=*Pomatomus telescopium* Risso).
2307. *Epigonus occidentalis* Goode and Bean.
Deep water off Barbados.
Epigonus occidentalis Goode and Bean, Oceanic Ichth., 1895, 233, off Barbados.
- Genus 787. *CHEILODIPTERUS* Lacépède.
Cheilodipterus Lacépède, Hist. Nat. Poiss., II, 1802, 539; restricted by Cuvier and Valenciennes, 1828, to *C. macrodon* Lacépède.
Clodipterus Rafinesque, Analyse Nat., 1815, 86, substitute for *Cheilodipterus* Lacépède; regarded as too long.
Paramia Bleeker, Neder. Tijdschr. Dierk., I, 1863, 233 (*Cheilodipterus lineatus* Lacépède); substitute for *Cheilodipterus* (Lacépède) Cuvier.
2308. *Cheilodipterus affinis* Poey.
Cuba; three specimens known.
Cheilodipterus affinis Poey, Ann. Acad. Nat. Sci. N. Y., XI, 1876, 58, Habana.

Family 188. SCOMBROPIDÆ

Genus 788. SPHYRÆNOPS Poey.

Sphyrænops Poey, *Memorias*, II, 1861, 249 (*S. bairdianus* Poey).

2309. *Sphyrænops bairdiana* Poey.

Deep water of Cuba; one specimen known

Sphyrænops bairdianus Poey, *Memorias*, II, 1861, 350, Cuba.

Genus 789. PARASPHYRÆNOPS Bean.

Parasphyrænops B. A. Bean, *Proc. Biol. Soc. Wash.*, XXV, 1912, 194

(*P. atrimanus* B. A. Bean).

2310. *Parasphyrænops atrimanus* B. A. Bean.

Bermudas.

Parasphyrænops atrimanus B. A. Bean, *Proc. Biol. Soc. Wash.*, XXV, 1912, 194, Bermudas.

Genus 790. LATEBRUS Poey.

Latebrus Poey, *Memorias*, II, 1860, 168 (*L. oculatus* Poey).

2311. *Latebrus oculatus* Poey. *Escolar chino*.

Coast of Cuba; in deep water.

Latebrus oculatus Poey, *Memorias*, II, 1861, 168, with plate, Cuba.

Genus 791. HYPOCLYDONIA Goode and Bean.

Hypoclydonia Goode and Bean, *Oceanic Ichth.*, 1895, 265 (*H. bella* Goode and Bean).

2312. *Hypoclydonia bella* Goode and Bean.

Gulf Stream.

Hypoclydonia bella Goode and Bean, *Oceanic Ichth.*, 1895, 236, Albatross stations 2314, 2397, 2401, 2417, etc., at depths from 90 to 159 fathoms.

Genus 792. AMIICHTHYS Poey.

Amiichthys Poey, in Jordan, *Proc. U. S. Nat. Mus.*, IX, 1886, 586 (*diapterus* Poey, 1861; only the specific name of this fish being published by Poey).

2313. *Amiichthys diapterus* Poey.

Coast of Cuba; two specimens known.

— *diapterus* Poey, *Synopsis*, 1861, 305, Cuba.

Family 189. CENTRARCHIDÆ. Black basses; Sunfishes and pumpkinseeds

Genus 793. HURO Cuvier and Valenciennes. *Large-mouth black bass*. *

?*Nemocampsis*⁵² Rafinesque, *Ichth. Ohiensis*, 1820, 30 (*Lepomis flexuolaris* Raf.).

Huro Cuvier and Valenciennes, *Hist. Nat.*, Poiss., II, 1828, 125 (*H. nigricans* Cuvier and Valenciennes = *Cichla floridana* LeSueur).

2314. *Huro floridana* (LeSueur). *Large-mouthed black bass*; *Oswego bass*; *Green bass*; *Bayou bass*; *Chub*; *Trout*; *Jumper*; *Welshman*; *Achigan grand bouche*; *Achigan noir*; *Big-mouthed black bass*; *Black bass of the Huron*; *Black huron*; *Bride perch*; *Cow bass*; *Dotted paintedtail*; *Fresh-water trout*; *Gray bass*; *Green perch*; *Green trout*; *Growler*; *Huron*; *Lake bass*; *Marsh bass*; *Mullpond chub*; *Moss bass*; *Mud bass*; *Paintedtail*; *Pale river bass*; *Perch*; *River bass*; *Rock bass*; *Salmon-formed growler*; *Slough bass*; *Southern chub*; *Speckled hen*; *Spotted bass*; *Straw bass*; *Striped bass*; *White bass*; *White salmon*; *White trout*; *Yellow bass*; *Yellow pondperch*.

Rivers of the United States from the Great Lakes and Red River of the North to Florida, Texas, and Mexico; west to the Dakotas, Nebraska and Kansas; introduced into lakes and streams of California.

Cichla floridana LeSueur, *Jour. Acad. Nat. Sci. Phila.*, II, 1822, 219, east Florida.

Huro nigricans Cuvier and Valenciennes, *Hist. Nat. Poiss.*, II, 1828, 93 (124), pl. 17, Lake Huron.

Micropterus salmoides of authors; not *Labrus salmoides* Lacépède.

⁵² Except for the phrase "Mouth large," Rafinesque gives no distinctive characters for the young black bass he calls "*Nemocampsis*"; but in *Micropterus* the mouth is decidedly large, though smaller than in *Huro*.

Genus 794. *MICROPTERUS* Lacépède. *Black bass.*

Micropterus Lacépède, Hist. Nat. Poiss., IV, 325, 1802 (*M. dolomieu* Lacépède).

Callurus Rafinesque, Jour. de Physique, 1819, 420 (*C. punctulatus* Raf.).

Aplites Rafinesque, Ichth. Ohiensis, 1820, 30 (*Lepomis pallida* Raf.).

?*Nemocampsis* Rafinesque, loc. cit., 30 (*Lepomis flexuolaris* Raf.).

Dioplites Rafinesque, loc. cit., 32 (*Lepomis notata* Raf.).

Aplesion Rafinesque, loc. cit., 36 (*Etheostoma callura* Raf.).

2315. *Micropterus dolomieu* Lacépède. *Small-mouthed black bass; Achigan noir; Achigan petite bouche; Bass; Bass hogfish; Black bass; Black fresh-water bass; Black perch; Bronze-backer; Brown bass; Brown trout; Dwarf bass; Gold bass; Green bass; Green perch; Green trout; Hog bass; Jumper, Little bass; Minny bass; Mountain trout, Redeye; River bass; Rock bass; Speckled hen; Spotted river bass; Streaked-cheeked river bass; Streaked lead; Swago bass; Swego; Trout bass; Trout perch; White trout; Yellow bass, Yellow perch.*

Lake Champlain to Manitoba and southward on both sides of the mountains to South Carolina and Arkansas.

Micropterus dolomieu Lacépède, Hist. Nat. Poiss., IV, 1802, 325, type locality uncertain, probably South Carolina.

Labrus salmoides Lacépède, loc. cit., IV, 1802, 716, South Carolina.

Bodianus achigan Rafinesque, Amer. Month Mag., II, 1817, 120, New York.

Callurus punctulatus Rafinesque, Jour. de Physique, 1819, 420, falls of the Ohio River.

Lepomis pallida Rafinesque, Ichth. Ohiensis, 1820, 30, Ohio, Miami, and Hocking Rivers.

Lepomis trifasciata Rafinesque, loc. cit., 31, Ohio River.

?*Lepomis flexuolaris* Rafinesque, loc. cit., 31, Ohio River.

Lepomis salmonea Rafinesque, loc. cit., 32, Kentucky, Green, Licking, and Ohio Rivers.

Lepomis notata Rafinesque, loc. cit., 32, Ohio River.

Etheostoma callura Rafinesque, loc. cit., 36, Ohio River; Salt River, Ky.

Cichla fasciata LeSueur, Jour., Acad. Nat. Sci. Phila., II, 1822, 216, Lake Erie.

Cichla ohiensis LeSueur, loc. cit., 218, Ohio River.

Cichla minima LeSueur, loc. cit., 220, Lake Erie.

Centrarchus obscurus DeKay, New York Fauna: Fishes, 1842, 30, pl. 17, fig. 48, Onondaga Creek, N. Y.

Dioplites variabilis Vaillant and Bocourt, Miss. Sci. Mex., Poissons, 1875-1883, 142, Wabash River, New Harmony, Ind.

2316. *Micropterus pseudoplites* Hubbs. *Kentucky black bass.*

Southeastern United States from western West Virginia northward to southern Ohio, westward to Kansas and central Texas.

Micropterus pseudaplites Hubbs, Proc. Okla. Acad. Sci., Part I, V, Jan 1, 1927, 137, nomen nudum.

Micropterus pseudoplites Hubbs, Occ. Papers, Mus. Zool. Univ. Mich. No. 184, June 29, 1927, 8, Pl. I, figs. 2, 3, and 5, Forbush Creek near Mill Springs, Ky.

Genus 795. *APOMOTIS* Rafinesque.

Apomotis Rafinesque, Jour. de Physique, LXXXVIII, 1819, 420 (*Lepomis cyanellus* Rafinesque).

Telipomis Rafinesque, Ichth. Ohiensis, 1820, 27 (*Lepomis cyanellus* Rafinesque).

Bryttus Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 454, 461 (*B. punctatus* Cuvier and Valenciennes).

2317. *Apomotis cyanellus* (Rafinesque). *Blue-spotted sunfish; Green sunfish; Little red-eye; Blackeyes; Blackeye sunfish; Blue-and-green sunfish; Blue bass; Bluefish; Blue sunfish; Bream; Buffalo sunfish; Perch; Red-eye; Sunfish; Blanco perch; Wood bass.*

Great Lakes region to Mexico; very abundant from Ohio southwestward to the Rio Grande.

Lepomis cyanellus Rafinesque, Jour. de Physique, LXXXVIII, 1819, 420, Ohio River.

Ichtheis melanops Rafinesque, Ichth. Ohiensis, 1820, 28, Ohio River.

- Pomotis longulus* Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 391, Otter Creek, Ark.
Calliurus diaphanus Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 200, Rio Blanco, Tex.
Calliurus formosus Girard, loc. cit., 1857, 200, Arkansas.
Calliurus microps Girard, loc. cit., 200, Rio Brazos, Tex.
Calliurus murinus Girard, loc. cit., 200, Texas.
Bryttus signifer Girard, loc. cit., 201, Rio Medina, Tex.
Bryttus mineopas Cope, loc. cit., XVII, 1865, 84, Whittlesey Lake, Minneapolis, Minn.
2318. *Apomotis euryorus* (McKay).
 Upper Great Lakes region. A form regarded as this occasional in the upper Mississippi.
Lepomis euryorus McKay, Proc. U. S. Nat. Mus., IV, 1881, 89, Lake Huron at Fort Gratiot, Mich. Regarded by Hubbs as a hybrid between *Apomotis cyanellus* and *Eupomotis gibbosus*.
2319. *Apomotis ischyurus* (Jordan and Nelson). *Big-nosed sunfish*.
 Upper Mississippi Valley.
Lepiopotis ischyurus Jordan and Nelson, Bull. U. S. Nat. Mus., X, 1877, 25, Illinois River, Ill. Rare; thought by Hubbs to be a hybrid.
2320. *Apomotis phenax* Cope and Jordan. *Deceptive sunfish*.
 Streams of New Jersey.
Apomotis phenax Cope and Jordan, Bull. U. S. Nat. Mus., X, 1877, 26, Beaseley's Point, N. J.; locality possibly erroneous; rare; thought to be a hybrid.
2321. *Apomotis punctatus* (Cuvier and Valenciennes). *Bream; Chinquapin perch; Spotted bream*.
 South Carolina to Florida.
Bryttus punctatus Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 347 (462), no locality, probably Charleston, S. C.
Bryttus reticulatus Cuvier and Valenciennes, loc. cit., 348 (463), Charleston, S. C.
Lepomis apiatus Cope, Proc. Amer. Phil. Soc., XVII, 1877, 65, Volusia, Fla.
2322. *Apomotis symmetricus* (Forbes)
 Mississippi Valley; Illinois to Louisiana and Texas.
Lepomis symmetricus Forbes, in Jordan and Gilbert, Synopsis, 1883, 473, Illinois River, Ill.
- Genus 796. *LEPOMIS* Rafinesque. *Sunfishes*.
Lepomis Rafinesque, Jour. de Physique, LXXXVIII, 1819, 420 (*Labrus auritus* Linnæus).
Pomotis Rafinesque, loc. cit., 420 (*Labrus auritus* Linnæus).
Ichthelis Rafinesque, Ichth. Ohiensis, 1820, 27 (*Labrus auritus* Linnæus).
2323. *Lepomis pallidus* (Mitchill).
 New York.
Labrus pallidus Mitchill, Trans. Lit. and Phil. Soc. N. Y., 1815, 407, New York. Misprint for *pallidus*.
Lepomis appendix Mitchill, Amer. Month. Mag., 1818, 247, New York. An uncertain species, perhaps based on *L. auritus*.
2324. *Lepomis auritus* (Linnæus). *Yellowbelly; Redbreast bream; Sunfish; Tobacobo; Long-eared sunfish; Sun perch; Bream; Redbreast; Black-eared pondfish; Flatfish; Flounder; Fresh-water sunfish; Kiver; Leather-eared; Leatherwing; Perch; Quiver; Redbelly; Red-bellied bream; Red-bellied robin perch; Red-headed bream; Red perch; Red sunfish; Red-tailed bream; Red-tailed pomotis; Roach; Robin; Robin perch; Yellow perch; River sunfish; Horned-eared sunfish*.
 Maine to Virginia; abundant in all streams east of the Alleghenies and south of New York.
Labrus auritus Linnæus, Syst. Nat., Ed. X, 1758, 283, Philadelphia.
Bryttus unicolor Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 348 (464), Philadelphia and Charleston.
Pomotis rubri-cauda Storer, Jour. Boston Soc. Nat. Hist., IV, 1842, 177, Concord, N. H.
Lepomis ophthalmicus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 223, 224, Roanoke River, Va.

2325. *Lepomis mystacalis* Cope.

East Florida

Lepomis mystacalis Cope, Proc. Amer. Phil. Soc., XVII, 1877, 66, east Florida.**2326. *Lepomis solis* (Cuvier and Valenciennes).**

Virginia to Louisiana, in coastwise streams.

Pomotis solis Cuvier and Valenciennes, Hist. Nat. Poiss., 1831, 352 (468), Lake Pontchartrain, La.*Pomotis elongatus* Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 47, pl. 5, fig. 1, St. Johns River, Fla.**Genus 797. *SCLEROTIS* Hubbs.***Sclerotis* Hubbs, Proc. Okla. Acad. Sci., Pt. I, VI, Jan. 1, 1927, 138 (*Lepomis minutus* Jordan)**2327. *Sclerotis miniatus* (Jordan). *Red perch.***

Mississippi Valley; southern Illinois to Louisiana and Texas.

Lepropomus minutus Jordan, Bull. U. S. Nat. Mus., X, 1877, 26, Tangipahoa River.*Lepomis garmani* Forbes, Bull. Ill. Lab. Nat. Hist., 1885, 135, Little Fox River at Phillipstown and Wabash River and Drew Pond at Carmi, Ill.**Genus 798. *ALLOTIS* Hubbs.***Allotis* Hubbs, Proc. Okla. Acad. Sci., Pt. I, VI, Jan. 1, 1927, 139 (*Bryttus humulus* Girard).**2328. *Allotis humilis* (Girard). *Red-spotted sunfish; Bream; Orange-spotted sunfish; Perch; Sunfish.***

Ohio and Kentucky to the Dakotas, Kansas, and Texas; locally abundant, especially in the lower Missouri Basin.

Bryttus humilis Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 201, Sugar Loaf Creek, Ark.*Bryttus oculatus* Cope, loc. cit., XVII, 1865, 83, Lake Whittlesey, Minn.*Lepomis anagallinus* Cope, Jour. Acad. Nat. Sci. Phila., 2d ser., VI, January, 1869, 221, Leavenworth, Kans.**Genus 799. *XENOTIS* Jordan.***Xenotis* Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 76 (*Pomotis fallax* Baird and Girard).**2329. *Xenotis megalotis* (Rafinesque.) *Long-eared sunfish; Big-eared sunfish; Blackears, Black-tailed sunfish; Bloody sunfish; Bream; Brilliant sunfish; Common sunfish; Eared sunfish; Long-scaled sunfish; Perch; Red-bellied bream; Redbelly; Red-eyed sunfish; Small green sunfish; Sun perch.***

Michigan to Minnesota, South Carolina, and southward to the Rio Grande

Ichthys megalotis Rafinesque, Ichth. Ohiensis, 1820, 29, Kentucky, Licking, and Sandy Rivers, Ky.*Pomotis nitida* Kirtland, Bost. Jour. Nat. Hist., III, 1840, 472, pl. 28, fig. 1, Mahoning River, Ohio*Pomotis breviceps* Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 390, Otter Creek, Ark.*Pomotis fallax* Baird and Girard, loc. cit., VII, 1854, 24, Elm Creek, Tex.*Pomotis convexifrons* Baird and Girard, loc. cit., 1854, 24, Rio Cibolo, Tex.*Pomotis nefastus* Baird and Girard, loc. cit., 24, Rio Cibolo and Salado Creek, Tex.*Pomotis sanguinolentus* Agassiz, Amer. Jour. Sci. and Arts, XVII, 1854, 301, Huntsville, Ala.*Pomotis inscriptus* Agassiz, loc. cit., 1854, 302, Huntsville, Ala.*Pomotis popei* Girard, Pacific R. R. Surv., X, 1858, 26, headwaters of Colorado River, Tex.*Lepomis peltastes* Cope, Proc. Amer. Phil. Soc., 1870, 454, Huron River, Mich.*Xenotis lythochloris* Jordan, Ann. N. Y. Lyc. Nat. Hist., XI, 1876 (1877), 376, White River.*Xenotis aureolus* Jordan, loc. cit., 376, White River.

2330. *Xenotis occidentalis* (Meek).

Rio Grande Basin in Mexico.

Lepomis occidentalis Meek, Field Col. Mus., Zool., III, 1902, 118, Rio Conchos, Mexico.**Genus 800. *HELIOPERCA* Jordan.***Heloperca* Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876 (1877), 355 (*Labrus pallidus* Cuvier and Valenciennes; not of Mitchill = *Pomotis incisor* Cuvier and Valenciennes).*Erichæta* Jordan, Klippart's Rept. Fish Com. Ohio, 1876 (1877), 48 (*Pomotis incisor* Cuvier and Valenciennes); uncorrected in proof, reading for *Heloperca*.**2331. *Heloperca incisor* (Cuvier and Valenciennes). *Bluegill; Blue bream; Blue sunfish; Copper-nosed bream; Dollardee; Black-eared pondfish; Blue Joe; Blue-mouthed sunfish; Blue perch; Bream; Copper-headed bream; Perch; Sunfish.***

Great Lakes to Florida and the Rio Grande.

Pomotis incisor Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 350 (466), New Orleans.*Pomotis gibbosus* Cuvier and Valenciennes, loc. cit., 1831, 351 (467), Charleston, S. C.; not of Linnæus.*Pomotis speciosus* Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 24, Brownsville, Tex.*Pomotis obscurus* Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 302, Huntsville, Ala.; specimens almost black.*Pomotis luna* Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 201, Fort Snelling, Minn.*Lepomis longispinus* Cope, loc. cit., 1865, 83, Leavenworth, Kans.*Lepomis ardesiacus* Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 222, Kiskiminitas River, Pa.*Lepomis purpurascens* Cope, Proc. Amer. Phil. Soc., XI, 1870, 454, Yadkin River, N. C.**2332. *Heloperca macrochira* (Rafinesque). *Bream; Chain-sided perch; Chain-side; Gilded sunfish; Gold perch; Perch; Sunfish.***

Ohio Valley and southwestward to Missouri and Kentucky.

Lepomis macrochirus Rafinesque, Jour. de Physique, LXXXVIII, 1819, 420, Ohio, Wabash, Green, and Licking Rivers**2333. *Heloperca nephela* (Cope).**

Western Pennsylvania.

Lepomis nephelus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 222, Kiskiminitas River, western Pennsylvania. Perhaps the same as *H. macrochira*.**2334. *Heloperca marginata* (Holbrook).**

Florida.

Pomotis marginatus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 49, Pl. VI, fig. 2, St. Johns River, Fla. This and the remaining species here referred to *Heloperca* are of uncertain value and of doubtful relationship.**2335. *Heloperca haplognatha* (Cope).**

Monterey, Nuevo Leon, Mexico.

Lepomis haplognathus Cope, Proc. Amer. Phil. Soc., XXII, 1884 (1885), 168, Monterey, Nuevo Leon, Mexico.**2336. *Heloperca bombifrons* (Agassiz).**

Tennessee River.

Pomotis bombifrons Agassiz, Amer. Jour. Sci. and Arts, XVII, 1854, 303, Huntsville, Ala.**Genus 801. *XYSTROPLITES* Jordan.***Xystroplites* Jordan, Bull. X, U. S. Nat. Mus., 1877, 24 (*X. gillii* Jordan = *Pomotis pallidus* Agassiz).

2337. *Xystroplites pallidus* (Agassiz).

Georgia to Texas.

Pomotis pallidus Agassiz, Amer. Jour. Sci. Arts, 1854, 303 (name pre-occupied in *Lepomis* by *Labrus palladus* Mitchill, but not in *Eupomotis*).

Bryttus albulus Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 200, Rio Blanco, Tex.

Xystroplites gilli Jordan, Bull. U. S. Nat. Mus., X, 1877, 24, "Garden Key, Florida"; evidently an error, as there is no fresh-water stream on Garden Key.

Lepomis lirus McKay, Proc. U. S. Nat. Mus., IV, 1881, 89; name a substitute for *pallidus*.

Genus 802. EUPOMOTIS Gill and Jordan.

Eupomotis Gill and Jordan, Field and Forest, II, 1877, 190 (*Sparus aureus* Walbaum = *Perca gibbosa* Linnæus).

2338. *Eupomotis heros* (Baird and Girard).

Southern Illinois, Indiana, and Ohio to Florida and the Rio Grande Basin.

Pomotis heros Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, March, 1854, 25, Rio Cibolo, Tex.

Pomotis notatus Agassiz, Amer. Jour. Sci. and Arts, XVII, 1854, 302, Huntsville, Ala.

2339. *Eupomotis holbrookii* (Cuvier and Valenciennes). Yellow bream; Shellcracker; Strawberry bass; Branch perch.

Virginia to Florida.

Pomotis holbrookii Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 350 (466), Charleston, S. C. Misprint for *holbrookii*.

Pomotis speciosus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 48, pl. 5, fig. 2, St. Johns River, Fla.; not of Baird and Girard.

Pomotis microlophus Gunther, Cat., I, 1859, 264, St. Johns River, Fla.

2340. *Eupomotis longimanus* (Cope).

Florida to Alabama.

Xystroplites longimanus Cope, Proc. Amer. Phil. Soc., XVII, 1877, 66, Volusia and Port Bay, Fla. Perhaps identical with *E. holbrookii*.

2341. *Eupomotis gibbosus* (Linnæus). Common sunfish; Bream; Pumpkinseed; Sunfish; Sunny; Crapet jaune; Female perch; Flatfish; Flounder; Harlequin roach; Kuer; Northern pomotis; Perch; Pond perch; Quiver; Redbelly; Roach; Robin; Robin perch; Ruff; Sand perch; Sun bass; Tobaccobox; Yellowbelly; Yellow perch.

Great Lakes region to Maine and southward to Florida, east of the Alleghenies.

Perca gibbosa Linnæus, Syst. Nat., Ed. X, 1758, 293, Carolinas; after *Perca fluviatilis gibbosa*, ventre luteo, of Catesby.

Sparus aureus Walbaum, Artedi Pisc., 1792, 290, lakes of New York.

Morone maculata Mitchill, Fishes of New York, 1814, 18, near New York City.

Pomotis vulgaris Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 67 (91), Lake Huron, New York, Virginia, and the Carolinas.

Pomotis ravenelii Cuvier and Valenciennes, loc. cit., VII, 1831, 349 (465), Philadelphia.

Pomotis catesbei Cuvier and Valenciennes, loc. cit., 352 (469), Philadelphia.

Genus 803. CHÆNOBRYTTUS Gill.

Chænobryttus Gill, Amer. Jour. Sci. Arts, XXXVII, 1864, 92 (*Calliurus melanops* Girard = *Pomotis gulosus* Cuvier and Valenciennes).

Glossoplites Jordan, Man. Vert., ed. 1, 1876, 223 (*Calliurus melanops* Girard).

2342. *Chænobryttus gulosus* (Cuvier and Valenciennes). Warmouth; Goggle-eye; Bigmouth; Big-mouthed sunfish; Black sunfish; Black warmouth; Bream; Buffalo bass; Chub; Jugmouth; Moremouth bream; Mud chub; Perch; Perchmouth bream; Redeye; Sac-a-lait, Sunfish; Sun trout; Warmouth perch; Warmouth bream; Wide-mouthed sunfish; Yawmouth perch; Indianfish.

Eastern United States, from the Great Lakes to Florida and Texas, west to Kansas and the Dakotas; chiefly west or south of the Alleghenies; common in South Carolina.

Pomotis gulosus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 367 (498), Lake Pontchartrain and lagoons about New Orleans.

Calliurus floridensis Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 53, Pl. VI, fig 1, St. Johns River, Fla.

Calliurus melanops Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 200, Texas.

Lepomis gillis Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 222, 225, Texas.

Lepomis charybdis Cope, loc. cit., 223, Texas.

Chænobryttus antistius McKay, Proc. U. S. Nat. Mus., IV, 1881, 88, Lake Michigan.

Genus 804. AMBLOPLITES Rafinesque. Rock basses.

Ambloplites Rafinesque, Ichth. Ohiensis, 1820, 33 (*Lepomis ichtheloides* Rafinesque = *Bodianus rupestris* Rafinesque).

2343. *Ambloplites rupestris* (Rafinesque). *Common rock bass; Redeye, Goggle-eye; Redeye perch; Bream; Bronzed centrarchus; Crapet mondoux; Crapet noir; Crapet vert; Fresh-water bass; Lake bass; Red-eyed bream; Red-eyed sunfish; Sunfish; Sunfish bass; Sun perch, White bass.*

Vermont to Great Lakes region and Manitoba, south to Louisiana; very abundant west of the Alleghenies.

Bodianus rupestris Rafinesque, Amer. Month Mag., II, 1817, 120, Lakes of New York, Vermont, and Canada

Ichelis erythrops Rafinesque, Ichth. Ohiensis, 1820, 29, Ohio River.

Lepomis ichteloides Rafinesque, loc. cit., 32, Ohio River.

Cichla xenea LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1822, 214, Lake Ontario.

Centrarchus pentacanthus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 65 (88), Wabash River.

2344. *Ambloplites cavifrons* Cope. *Southern rock bass.*

Roanoke River, Va.

Ambloplites cavifrons Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 217, Roanoke River, Va.

Genus 805. ARCHOPLITES Gill.

Archoplites Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 165 (*Centrarchus interruptus* Girard).

2345. *Archoplites interruptus* (Girard). *Sacramento perch.*

Sacramento and San Joaquin Rivers, Calif.

Centrarchus interruptus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 129, San Joaquin and Sacramento Rivers, Calif.

Centrarchus maculosus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 7, Sacramento River.

Genus 806. ACANTHARCHUS Gill.

Acantharchus Gill, Amer. Jour. Sci. Arts, XXXVI, 1864, 92 (*Centrarchus pomotis* Baird).

2346. *Acantharchus pomotis* (Baird). *Mud sunfish; Bass sunfish; Mud bass; Mud perch.*

Southern New York to South Carolina.

Centrarchus pomotis Baird, Ninth Smiths. Rept., 1855, 325, New Jersey, New York.

Genus 807. MESOGONISTIUS Gill.

Mesogonistius Gill, Amer. Jour. Sci. Arts, XXXVII, 1864, 92 (*Pomotis chætodon* Baird).

2347. *Mesogonistius chætodon* (Baird). *Black-banded sunfish.*

New Jersey to Maryland.

Pomotis chætodon Baird, Ninth Smiths. Rept., 1855, 324, Cedar Swamp Creek, N. J.

Genus 808. ENNEACANTHUS Gill.

Enneacanthus Gill, Amer. Jour. Sci. Arts, XXXVII, 1864, 92 (*Pomotis obesus* Baird).

Hemiplites Cope, Jour. Acad. Nat. Sci. Phila., 2d ser., VI, 1868, 217 (*H. simulans* Cope=*Enneacanthus gloriosus*; abnormal example with A. IV, the dorsal erroneously stated to be X.
Copelandia Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 56 (*C. eriarcha* Jordan, an abnormal example with D. X; A. IV, of *Enneacanthus gloriosus*).

2348. *Enneacanthus obesus* (Girard). *Banded sunfish; Little bream; Little sunfish; Spotted sunfish; Rock sunfish.*

Charles River, Mass., to Florida.

Pomotis obesus Girard, Proc. Bost. Soc. Nat. Hist., V, 1854, 40, waters about Hingham and in branch of Charles River at Holliston, Mass.

Bryttus fasciatus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 51, Pl. V, fig. 3, St. Johns River, Fla.

Pomotis guttatus Morris, Proc. Acad. Nat. Sci. Phila., XI, 1859, 3, Delaware River, Philadelphia.

2349. *Enneacanthus gloriosus* (Holbrook). *Blue-spotted sunfish; Little sunfish; Spotted-finned sunfish; Spotted sunfish.*

Atlantic States; New York to Florida.

Bryttus gloriosus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 52, Pl. V, fig. 4, Cooper River, S. C.

Hemiplites simulans Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 217, Tuckahoe Creek near Richmond, Va.; abnormal examples.

Enneacanthus pinniger Gill and Jordan, Bull. U. S. Nat. Mus., X, 1877, 27, Tar River, Kinston, N. C.

Enneacanthus margarotis Gill and Jordan, loc. cit., 28, Beaseleys Point, N. J.

Copelandia eriarcha Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 56, Menominee River, Wis.; evidently an error, an abnormal example; the locality for New Jersey.

Genus 809. CENTRARCHUS Cuvier and Valenciennes. *Round basses.*

Centrarchus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 62 (84) (*Labrus irideus* Lacépède=*Labrus macropterus* Lacépède).

2350. *Centrarchus macropterus* (Lacépède). *Round sunfish; Flier; Flying perch; Large-finned bass; Long-finned sunfish; Many-spined sunfish; Millpond perch; Perch; Round bass; Sac-a-lait; Shinning bass; Sunfish.*

Lowland streams from Virginia southward to Florida and Louisiana; northward in the Mississippi Valley to southern Illinois.

Labrus macropterus Lacépède, Hist. Nat. Poiss., III, 1802, 432, Charleston, S. C.

Labrus irideus Lacépède, loc. cit., IV, 716, Charleston, S. C.

Genus 810. POMOXIS Rafinesque. *Crappies.*

Pomoxis Rafinesque, Amer. Month. Mag., IV, 1818, 41 (*Pomoxis annularis* Rafinesque).

Hyperistius Gill, Amer. Jour. Sci. Arts, 1864, 92 (*Centrarchus hexacanthus* Cuvier and Valenciennes=*Labrus sparoides* Lacépède).

Subgenus POMOXIS Rafinesque.

2351. *Pomoxis annularis* Rafinesque. *Crappie; Bachelor; Sac-a-lait; New-light; Campbelite; Crapet; Bachelor perch; Bridge perch; Chinquapin perch; Goggle-eye; Goldring; John demon; Pale crappie; Papermouth; Shad; Silver perch; Speckled perch; Strawberry perch; Suckley perch; Timber crappie; Tinmouth; Tin perch; White crappie, White perch.*

Eastern United States, from the Great Lakes south to Texas and west to Kansas and Nebraska. Introduced into Columbia River and other waters on the Pacific coast.

Pomoxis annularis Rafinesque, Amer. Month. Mag., IV, 1818, 41, falls of the Ohio River.

Cichla storeria Kirtland, Rept., Zool. Ohio, 1838, 191, Ohio River.

Pomoxis nitidus Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 200, Houston River, Ky.

Pomoxys brevicauda Gill, loc. cit., XVII, 1865, 64, North Grand River, Mo.

Pomoxys intermedius Gill, loc. cit., 65, no locality.

Pomoxys prolaecanthus Gill, loc. cit., 66, Tarboro, N. C.

Subgenus **HYPERISTIUS** Gill.

- 2352. *Pomoxis sparoides* (Lacépède).** *Calico bass; Grass bass; Barfish; Strawberry bass; Bachelor; Banklick bass; Butterhead, Calico bream; Chunguapin perch; Crapet; Crappie, Dolly Varden; Goggle-eye; Lake bass; Lake Erie bass; Millpond fluer; Razorback; Roach; Rockfish; Sand perch; Silver bass, Silver perch; Speckled bass; Spotted perch; Spotted trout; Straw bass; Sun perch; Tanmouth; White perch; Black crappie.*

Great Lakes and upper Mississippi Valley to New Jersey and southward to Florida, Louisiana, and Texas. Introduced into Pacific coast streams. *Lábrus sparoides* Lacépède, Hist. Nat. Poiss., III, 1802, 517 and 518, South Carolina.

Cantharus nigro-maculatus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 65 (88), Wabash River.

Centrarchus hexacanthus Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 344 (459), Charleston, S. C.

Hyperistius carolinensis Gill, Amer. Jour. Sci. Arts, XXXVI, 1864, 93, South Carolina.

Family 190. **ELASSOMIDÆ**Genus 811. **ELASSOMA** Jordan.

Elassoma Jordan, Bull. U. S. Nat. Mus., X, 1877, 50 (*E. zonata* Jordan). *Elassosoma* Boulenger; attempted correction but the name is from *ἐλάσσωμα*, a diminution.

- 2353. *Elassoma zonatum* Jordan** *Pigmy sunfish.*

Southern Illinois to Texas and Alabama.

Elassoma zonata Jordan, Bull. U. S. Nat. Mus., X, 1877, 50, Little Red River, White County, Ark.

- 2354. *Elassoma evergladei* Jordan.** *Everglades pigmy sunfish.*

Swamps of southern Georgia and Florida, locally common in dark waters tributary to the Everglades.

Elassoma evergladei Jordan, Proc. U. S. Nat. Mus., VII, 1884, 323, St. Johns and Suwanee Rivers, Fla.

Family 191. **KUHLIDÆ.** Mountain bass; SeseleGenus 812. **SAFOLE** Jordan.

Safole Jordan, Proc. U. S. Nat. Mus., XLII, 1912, 655 (*Dules tæniurus* Cuvier and Valenciennes).

- 2355. *Safole tæniura* (Cuvier and Valenciennes).**

Galapagos and Revillagigedo archipelagoes; locally common; South Seas and Hawaii to Japan; throughout the warm parts of the Pacific, in the open sea, away from coral reefs; not in California.

Dules tæniurus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 85 (114), Java.

Kuhlia arge Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 159, Chatham Island, Galapagos.

Family 192. **CENTROPOMIDÆ.** RobalosGenus 813. **CENTROPOMUS** Lacépède. *Robalos.*

Centropomus Lacépède, Hist. Nat. Poiss., IV, 1802, 248 (*Lucioperca undecimalis*, etc.; restricted to *Scæna undecimalis* Bloch by Cuvier and Valenciennes).

Oxylabrax Bleeker, Archiv Neerl., XI, 1876, 264 (same type *Centropomus* being restricted to the first species named by Lacépède).

- 2356. *Centropomus undecimalis* (Bloch).** *Robalo; Sergeantfish; Snook; Brochet de mer.*

Coasts of Florida and Texas, southward among the West Indies to Surinam or beyond; Bay of Panama.

Scæna undecimalis Bloch, Naturgeschich. Ausl. Fische, VI, 1792, 60, Jamaica.

Centropomus undecim-radiatus Lacépède, Hist. Nat. Poiss., IV, 1802, 268; after Bloch.

Perca loubina Lacépède, loc. cit., 397, Cayenne.

Sphyræna aureoviridis Lacépède, loc. cit., V, 1803, 324, Martinique.

Centropomus appendiculatus Poey, Memorias, II, 1860, 119, Cuba.

2357. *Centropomus viridis* Lockington. *Robalo*.
Pacific coast of Mexico; common from Gulf of California to Panama.
Centropomus viridis Lockington, Proc. Calif. Acad. Sci., VII, 1876 (1877), 110, off Ascension Island, Lower California. Probably not separable from *C. undecimalis*.
2358. *Centropomus nigrescens* Gunther. *Robalo prieto*.
Pacific coast of tropical America; generally common from Mazatlan to Panama.
Centropomus nigrescens Gunther, Proc. Zool. Soc. Lond., 1864 (Mar. 22), 144, Chiapam.
2359. *Centropomus pedimacula* Poey. *Constantino de las aletas prietas*;
Robalito de las aletas prietas
Both coasts of tropical America; common in Cuba; Jamaica; south to Brazil.
Centropomus pedimacula Poey, Memorias, II, 1860, 122, Habana and Cienfuegos, Cuba.
Centropomus medius Gunther, Proc. Zool. Soc. Lond., 1864, 144, Chiapam.
Centropomus cuvieri Bocourt, Ann. Sci. Nat. Paris, IX, 1868, 91, Haiti.
Centropomus grandoculatus Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 139, Guaymas, Sonora. Probably distinct.
2360. *Centropomus parallelus* Poey.
Coasts of Cuba; San Domingo, Jamaica, Barbados, and Rio Chagres; Guiana, Pernambuco, and Bahia; Porto Rico.
Centropomus parallelus Poey, Memorias, II, 1861, 120, Habana and Cienfuegos, Cuba.
Centropomus mexicanus Bocourt, Ann. Sci. Nat. Paris, IX, 1868, 90, Gulf of Mexico.
2361. *Centropomus pectinatus* Poey.
Coasts of Cuba.
Centropomus undecimalis Cuvier and Valenciennes (part; not of Bloch), Hist. Nat. Poiss., II, 1828, 102 (*Centropomus pectinatus* Poey, Memorias, II, 1861, 121; Habana and Cienfuegos, Cuba).
2362. *Centropomus unionensis* Bocourt
Pacific coast of Central America; rather common at Panama.
Centropomus unionensis Bocourt, Ann. Sci. Nat. Paris, IX, 1868, 90, La Union, San Salvador.
2363. *Centropomus armatus* Gill.
Pacific coast of Central America; common from Chiapas to Panama.
Centropomus armatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 163, Panama.
2364. *Centropomus robalito* Jordan and Gilbert. *Constantino*; *Robalito las aletas amarillas*
Pacific coast of Mexico, from Mazatlan to Panama.
Centropomus robalito Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 462, Mazatlan; Acapulco.
2365. *Centropomus ensiferus* Poey.
West Indies; generally common from Cuba to Surinam.
Centropomus ensiferus Poey, Memorias, II, 1861, 122, Habana.
? *Centropomus brevis* Günther, Proc. Zool. Soc. Lond., 1864, 145, locality unknown, tropical America.
Centropomus affinis Steindachner, Ichth. Notizen, I, 1864, 1, pl. 1, fig. 1, Rio Janeiro and Cajutuba; Demerara, Guiana.
Centropomus scaber Bocourt, Ann. Sci. Nat., 5 ser., IX, 1868, 90, marshes of Belize.

Family 193. MORONIDÆ. White bass

Genus 814. *ROCCUS* Mitchill. *Striped bass*.

Roccus Mitchill, Fishes of New York, 1814, 24 (*R. striatus* Mitchill = *Scizena lineata* Bloch).

2366. *Roccus saxatilis*⁵³ (Walbaum) *Striped bass; Rockfish; Rock; Mis-suckee-kequoek*.

Atlantic coasts of North America from New Brunswick to Escambia River, Fla.; introduced by the U. S. Fish Commission into Sacramento River, Calif.

Perca saxatilis Walbaum, Artedi Gen. Pisc., 1792, 330, New York.

Perca septentrionalis Bloch and Schneider, Systema Ichthyol., 1801, 90, New York.

Roccus striatus Mitchill, Rep. Fishes N. Y., 1814, 24, New York.

Perca mitchilli Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 413, pl. 3, fig. 4, New York.

Perca mitchilli alternata Mitchill, loc. cit., 415, New York.

Perca mitchilli interrupta Mitchill, loc. cit., 415, New York.

Genus 815. *LEPIBEMA* Rafinesque.

Lepibema Rafinesque, Ichth. Ohiensis, 1820, 23 (*Perca chrysops* Rafinesque).

2367. *Lepibema chrysops* (Rafinesque). *White bass; White lake bass; sometimes silver bass at Sandusky, Ohio*.

Great Lakes region, upper Mississippi and Ohio Valleys, south to Washita River.

Perca chrysops Rafinesque, Ichth. Ohiensis, 1820, 22, falls of the Ohio.

Labrax multilineatus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1830, 360 (488), Wabash River.

Labrax notatus Richardson, Fauna Bor.-Amer., III, 1836, 8, lower St. Lawrence River.

Labrax abidus DeKay, N. Y. Fauna: Fishes, 1842, 13, pl. 51, fig. 165, Buffalo, N. Y.

Labrax osculati Filippi, Rev. et. Mag. de Zoologie, 2d series, V, 1853, 164, Canada.

Genus 816. *CHRYSOPERCA* Fowler.

Chrysoperca Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 250 (*Morone interrupta* Gill).

2368. *Chrysoperca interrupta* (Gill). *Yellow bass*.

Lower Mississippi Valley north to Cincinnati, Terre Haute, and St. Louis.

Morone interrupta Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 118, St. Louis; New Orleans.

Morone mississippiensis Jordan and Eigenmann, Proc. Acad. Nat. Sci. Phila., XXXIX, 1887, 295; substitute for *Morone interrupta*, preoccupied in *Roccus*, not in *Morone* nor in *Chrysoperca*.

Genus 817. *MORONE* Mitchill. *White perch*.

Morone Mitchill, Fishes of New York, 1814, 17 (*M. pallida* Mitchill = *Perca americana* Gmelin).

2369. *Morone americana* (Gmelin). *White perch; Silver perch; Sea perch*.

Atlantic coast of North America from Nova Scotia to South Carolina in brackish water, ascending streams, and frequently landlocked in ponds.

Perca americana Gmelin, Syst. Nat., I, III, 1789, 1308, New York; after Schöpf.

Perca immaculata Walbaum, Artedi Genera Piscium, 1792, 330, New York; after Schöpf.

Morone rufa Mitchill, Rep. Fishes N. Y., 1814, 18, New York.

Morone pallida Mitchill, loc. cit., 17, New York.

?*Centropomus albus* Rafinesque, Précis des Découvertes Somiolog., 1814, 19, Philadelphia.

⁵³ This species is usually called *Roccus lineatus* after *Scizena lineata* Bloch (Ausländische Fische, VI, 1792, 62), but it can not be the same. The form, serræ of the preopercle, and the stout spines of the fin, as well as the asserted locality ("Mediterranean") indicate that the species concerned is *Dicentrarchus lupus* of Europe. The only resemblance to *Roccus* is found in the striped color, but Bloch says that the stripes on the sides are yellow.

Perca mucronata Rafinesque, Amer. Month. Mag. and Crit. Rev., II, 1818, 204, Delaware, Schuylkill, and Susquehanna Rivers.
Labrax nigriscans DeKay, Nat. Hist. N. Y., Fishes, 1842, 12, pl. 50, fig. 160, Long Island.

Family 194. EPINEPHELIDÆ. Sea basses, groupers, garrupas, etc.

Genus 818. STEREOLEPIS Ayres. Jewfishes.

Stereolepis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 28 (*S. gigas* Ayres).

Megaperca Hilgendorf, Sitzb. Gesell. Naturf. Freunde, Berlin, 1878, 155 (*M. ishinnagi* Hilgendorf).

2370. *Stereolepis gigas* Ayres. California jewfish.

Coast of California, north to the Farallons.

Stereolepis gigas Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 28, southern California.

Stereolepis californicus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 330, California; substitute for *S. gigas*, supposed to be preoccupied in this genus.

Genus 819. POLYPRION Cuvier. Wreckfishes.

Polyprion Cuvier, Règne Animal, ed. 1, II, 1817, 282 (?*Amphiprion americanus* Bloch and Schneider).

2371. *Polyprion americanus* (Bloch and Schneider). Wreckfish; Stone bass; Cernier.

Gulf stream; Grand Banks and New Jersey; also in Europe if *P. massiliensis* is the same, which is not certain.

Amphiprion americanus Bloch and Schneider, Syst. Ichth., 1801, 205, pl. 47; based on a drawing sent by Latham to Schneider representing some fish called in America "Girom"; called *Amphiprion australis* on pl. 47.

?*Scorpaena massiliensis* Risso, Ichth. Nice, 1810, 184, Marseilles.

?*Polyprion cernium* Cuvier, Mém. du Museum, XI, 1824, 265, Pl. XVII, France.

?*Holocentrus gulo* Risso, Europe Mérid., III, 1826, 367, Nice.

?*Serranus couchus* Yarrell, British Fishes, ed. 1, I, 1836, 12, Great Britain.

Genus 820. GONIOPECTRUS Gill. Spanish flags.

Gonioplectrus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 236, 237 (*Plectropoma hispanum* Cuvier and Valenciennes).

2372. *Gonioplectrus hispanus* (Cuvier and Valenciennes). Spanish flag; Ouatilibi espagnol; Brajaiba de lo alto.

West Indies; Cuba.

Plectropoma hispanum Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 298 (396), Martinique.

Genus 821. PETROMETOPON Gill. Enjambre.

Petrometopon Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 105 (*Perca guttatus* Poey = *Sparus cruentatus* Lacépède).

2373. *Petrometopon cruentatus* (Lacépède). Enjambre; Coney; Rock hind.
 West Indian fauna, Brazil to Florida Keys; very common on Cuba coast.

Sparus cruentatus Lacépède, Hist. Nat. Poiss., IV, 1802, 156, Martinique; on a copy of Plumier's drawing.

?*Perca guttata* Bloch, Ichthyol., 1792, 312, Martinique, not of Linnæus.

Saranus apiarius Poey, Memorias, II, 1861, 143, Habana.

2374. *Petrometopon cruentatus coronatus* (Cuvier and Valenciennes). Brown hind; Petite nègre.

West Indies, north to Key West.

Serranus coronatus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 278 (371), Martinique.

2375. *Petrometopon panamensis* (Steindachner).

Panama.

Serranus panamensis Steindachner, Ichth. Beitr., IV, 1875, 1, with plate, Panama.

Genus 822. CEPHALOPHOLIS Bloch and Schneider.

Cephalopholis Bloch and Schneider, Syst. Ichth., 1801, 311 (*C. argus* Bloch and Schneider).

Urophæton Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 202 (*Serranus*⁵⁴ *phaeton* Cuvier and Valenciennes = *U. microleptes* Swainson).

Enneacentrus Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 105 (*Serranus ouatalibi* Cuvier and Valenciennes = *Labrus fulvus* Linnæus).

Bodianus Bloch, in part, restricted by Cuvier and Valenciennes to *Bodianus guttatus*, etc., later restricted to *Bodianus bodianus* = *Harpe rufa* through tautonymy.

- 2376. *Cephalopholis fulvus***⁵⁵ (Linnæus) *Guatwere*; *Niggerfish*; *Yellowfish*; *Butterfish*; *Lemon-yellow butterfish*; *Guatwere amarilla*.

West Indies; Bermuda and Florida Keys to Bahia.

Labrus fulvus Linnæus, Syst. Nat., Ed. X, 1758, 287, Bahamas; after Catesby.

Guatwere amarilla Parra, Descr. Dif. Piezas, Hist. Nat. Cuba, 8 lam., V, 1787, fig. 2, Cuba.

Holocentrus auratus Bloch, Naturgesch. Ausl. Fische, IV, 1790, 75, East Indies.

Bodianus guatwere Bloch and Schneider, Syst. Ichth., 1801, 336 (after Parra's figure).

- 2377. *Cephalopholis fulvus ruber*** (Bloch and Schneider). *Red guatwere*; *Ouatlibi*; *Rock hind*.

West Indies, etc.; very common.

Gymnocephalus ruber Bloch and Schneider, Syst. Ichth., 1801, 346, pl. 67, on *Carauna* of Marcgrave; not *Epinephelus ruber* Bloch.

Serranus ouatalibi Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 286 (381), Habana.

Serranus carauna Cuvier and Valenciennes, loc. cit., 288 (384), no locality given.

- 2378. *Cephalopholis fulvus punctatus*** (Linnæus). *Niggerfish*; *Negrofish*; *Black guatwere*.

West Indies, etc.; everywhere common.

Perca punctata Linnæus, Syst. Nat., ed. X, 1758, 291, Bahamas; based on Catesby.

- 2379. *Cephalopholis stellatus*** (Blosser).

St. Croix.

Bodianus stellatus Blosser, Ann. Carnegie Mus., VI, 1909, 297, St. Croix.

- 2380. *Cephalopholis tæniops*** (Cuvier and Valenciennes)

West coast of Africa, straying to Florida.

Serranus tæniops Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 277 (370), Cape Verde Islands.

Genus 823. MENEPHORUS Poey.

Menephorus Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1871, 50 (*Serranus dubius* Poey).

- 2381. *Menephorus dubius*** (Poey).

Cuba.

Serranus dubius Poey, Memorias, II, 1861, 142, Cuba.

- 2382. *Menephorus punctiferus*** (Poey).

Cuba.

Menephorus punctiferus Poey, Enumeratio, 1875, 21, Habana.

Genus 824. ENNEISTUS Jordan and Evermann.

Enneistus Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1147 (*Bodianus acanthistius* Gilbert).

- 2383. *Enneistus acanthistius*** (Gilbert).

Cape Lobos, eastern shore of Gulf of California.

Bodianus acanthistius Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 552, Cape Lobos, eastern shore of Gulf of California at *Albatross* station 3017, in 58 fathoms.

⁵⁴ A spurious species, *Cephalopolis nigripinnis* to which the tail of a *Fistularia* had been attached

⁵⁵ The various forms here treated as subspecies seem to inhabit varying depths of water and should be regarded rather as forms of *C. fulvus*.

Genus 825. EPINEPHELUS Bloch. Groupers.

- Epinephelus* Bloch, Nat. Ausl. Fische, VII, 1793, 11 (*E. marginalis*).
Cerna Bonaparte, Introd. Fauna Italica, III, 1833, pt. 1 (*Perca gigas* Brunnich).
Cromileptes Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 201 (*Serranus altivelis* Cuvier and Valenciennes).
Cymatichthys Swainson, loc. cit., 201 (*Perca flavopurpurea* Bennett = *Holocentrus flavoceruleus* Lacépède).
Hyporthodus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 98 (*Serranus niveatus* Cuvier and Valenciennes).
Labroperca Gill, loc. cit., XIV, 1862, 236 (*Serranus labriformis* Jenyns).
Schistorus Gill, loc. cit., 237 (*Serranus mystacinus* Poey).
Merus Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1871, 39 (*Serranus gigas* Cuvier and Valenciennes).
Pracanthichthys Day, Proc. Zool. Soc. Lond., 1868, 193 (*P. maderaspatensis* Day = *Epinephelus latifasciatus* Temminck and Schlegel).
Homalagrystes Alleyne and Macleay, Proc. Linn. Soc. New South Wales, 1877, 268 (*H. guentheri* Alleyne and Macleay).
Hyposerranus Klunzinger, Fische des Rothen Meeres, 1884, 3 (*Serranus morrhua* Cuvier and Valenciennes).

Subgenus SCHISTORUS Gill.**2384. *Epinephelus mystacinus* (Poey). *Cherna de lo alto*.**

West Indies south to Brazil.

Serranus mystacinus Poey, Memorias, I, 1852, 52, Tab. 10, fig. 1, Cuba.

Subgenus EPINEPHELUS Bloch.**2385. *Epinephelus analogus* Gill. *Cabrilla pinta*.**

Pacific coast of tropical America; common on coast of Mexico.

Epinephelus analogus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 163, Panama.

?*Serranus courtaderi* Bocourt, Ann. Sci. Nat. Paris, X, 1868, 222, La Union, San Salvador.

2386. *Epinephelus adscensionis* (Osbeck). *Rock hind; Cabra mora*.

West Indies; Florida Keys to Brazil; Ascension and St. Helena Islands; Woods Hole, Mass.

Perca tota Siba, Thesaurus, LII, 1758, Tab. 27.

Trachinus adscensionis Osbeck, Iter Chin., etc., 1757, and English ed., 1771, 96, Ascension Island.

Trachinus punctatus Bonnaterre, Tableau Encycl., 1788, 46; after Osbeck.

Perca stellio Walbaum, Artedi Pisc., 1792, 349; after Siba.

?*Perca maculata* Bloch, Nat. Ausl. Fische, 1792, 92, Martinique; on a figure by Plumier. Name preoccupied in *Epinephelus*.

Trachinus osbeck Lacépède, Hist. Nat. Poiss., II, 1800, 364; after Osbeck.

Sparus atlanticus Lacépède, loc. cit., IV, 1803, 158, pl. 5, fig. 1, Martinique; on a drawing by Plumier.

Serranus nigriculus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 381 (375); Martinique.

Serranus pizanga Cuvier and Valenciennes, loc. cit., 288 (383), after Pira-Pixanga or Gat-Vichof, Marcgrave, Hist. Brazil, 1648, 152.

Serranus aspersus Jenyns, Voy. Beagle, Fishes, 1842, 6, Porto Praya, St. Jago of the Cape Verde Islands.

Serranus impetiginosus Muller and Troschel, Schomburgk's Hist. Barb., 1848, 665, Barbados.

?*Serranus varius* Bocourt, Ann. Sci. Nat., X, 1868, 222, Gulf coast of Mexico.

2387. *Epinephelus guaza* (Linnæus). *Merou; Mérou; Guasa*.

Coasts of southern Europe and western Africa, ranging north to England, southward to Cape of Good Hope, and westward to Rio Janeiro and to Guiana. The American form, *E. menzeli* (Cuvier and Valenciennes), may prove different.

Labrus guaza Linnæus, Syst. Nat., ed. X, 1758, 285, "Habitat in Pelago."

Perca gigas Brunnich, Ichth. Massiliensis, No. 81, 1768, 65, Marseilles.

Holocentrus merou Lacépède, Hist. Nat. Poiss., IV, 1803, 377, Marseilles; after Brunnich.

- ?*Serranus menizelus* Cuvier and Valenciennes, loc. cit., II, 1828, 216 (291), coast of Brazil.
Perca robusta Couch, Mag. Nat. Hist., V, 1832, 21, fig. 7, Polperro, Cornwall.
Serranus marginatus Lowe, Proc. Zool. Soc. Lond., pt. 1, 1833, 142, Madeira.
Serranus fimbriatus Lowe, Trans. Cambr. Phil. Soc., 1836, 195, pl. 1, Madeira.
 ?*Epinephelus brachysoma* Cope, Trans. Amer. Phil. Soc., XIV, 1871, 466, Rio Janeiro.
2388. *Epinephelus labriformis* (Jenyns).
 Pacific coast of tropical America; Cape San Lucas to Galapagos Islands.
Serranus labriformis Jenyns, Voyage *Beagle*, Fishes, 1842, 8, pl. 3, Galapagos Islands.
Epinephelus selkcauda Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 250, Cape San Lucas.
Epinephelus ordinatus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 466, Panama.
2389. *Epinephelus flavolimbatus* Poey. *Yellow-finned grouper*.
 West Indies, north to Pensacola.
Epinephelus flavolimbatus Poey, Repertorio, I, 1866, 183, Cuba.
2390. *Epinephelus niveatus* (Cuvier and Valenciennes). *Spotted grouper*;
Snowy grouper.
 West Indies to Brazil, occasionally northward in the Gulf Stream as far as Newport, R. I., and Woods Hole, Mass.
Serranus niveatus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 285 (380), Brazil.
Serranus margaritifera Gunther, Cat., I, 1859, 131, South America.
Serranus conspersus Poey, Memorias, II, 1860, 139, Cuba.
Hyporthodus flavicauda Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 98, Newport, R. I.
2391. *Epinephelus striatus* (Bloch). *Nassau grouper*; *Hamlet*; *Cherna crotilla*; *Grouper*; *Rockfish*.
 West Indian fauna; Key West to Brazil.
Cherna Parra, Dif. Piezas, 1787, 50, Habana.
Anthias striatus Bloch, Naturgesch. Ausl. Fische, VI, 1792, 125, Martinique; on figure by Plumier.
Anthias cherna Bloch and Schneider, Syst. Ichth., 1801, 310, Cuba; after Parra.
Sparus chrysomelanurus Lacépède, Hist. Nat. Poiss., IV, 1803, 160, Martinique; on a copy of Plumier.
2392. *Epinephelus guttatus* (Linnaeus). *Cabrilla*; *Red hind*.
 West Indies; Carolina to Brazil, occasionally north to Charleston, the Florida Keys, and the Bermudas.
Perca guttata Linnaeus, Syst. Nat., ed. X, 1758, 292, Brazil; after Marcgrave, in part, but with a type specimen, examined by Lönnberg.
Cabrilla Parra, Dif. Piezas, Hist. Nat. Cuba, 1787, 93, pl. 36, Habana.
Lutjanus lunulatus Bloch and Schneider, Syst. Ichth., 1801, 329; after Parra, not *L. lunulatus* of Park.
Serranus catus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 250 (373), Martinique.
Serranus maculosus Cuvier and Valenciennes, loc. cit., 332, verified by Boulenger.
Serranus arara Cuvier and Valenciennes, loc. cit., 283 (377), Habana.
Epinephelus cubanus Poey, Repertorio, I, 1866, 202, Cuba.
Serranus stadthouderi Vaillant, Miss. Sci. Mex., Poiss., 1877, 69, substitute for *maculosus* regarded as preoccupied by *maculatus*.
2393. *Epinephelus angustifrons* (Steindachner).
 Coast of Cuba.
Serranus angustifrons Steindachner, Verh. Ges. Wien, XIV, 230, Pl. VII, figs. 2 and 3, 1864, Cuba.

2394. *Epinephelus drummond-hayi* Goode and Bean. *Speckled hind*; *John paw*.
Bermudas, south Atlantic and Gulf coast of United States; Charleston, S. C.
Epinephelus drummond-hayi Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 173, 174, Pensacola; Bermuda.
2395. *Epinephelus niphobles* Gilbert and Starks.
Magdalena Bay, Lower California.
Epinephelus niphobles Gilbert and Starks, Proc. U. S. Nat. Mus., XIX, 1896, 442, Magdalena Bay, Lower California.
- Subgenus **VIVERO**⁵⁶ Jordan and Evermann.
2396. *Epinephelus morio* (Cuvier and Valenciennes). *Red grouper*; *Cherna americana*; *Cherna de vivero*; *Nègre*.
Atlantic coast of America; Woods Hole to Rio Janeiro.
Serranus morio Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 211 (285), New York and San Domingo.
Serranus erythrogaster DeKay, New York Fauna: Fishes, 1842, 21, pl. 19, fig. 52, Florida.
Serranus remotus Poey, Memorias, II, 1860, 140, Cuba.
- Genus 826. **GARRUPA** Jordan. *Black groupers*.
Garrupa Jordan, Bull. U. S. Fish Com., VIII, 1888 (1890), 353 (*Serranus nigrinus* Holbrook).
2397. *Garrupa nigrata* (Holbrook). *Black jewfish*; *Black grouper*; *Mero de lo alto*.
South Atlantic and Gulf coast of the United States, Charleston and Pensacola to Cuba and Brazil; perhaps straying to Sicily.
Serranus nigratus Holbrook, Ichth. South Carolina, Ed. I, 1855, 173, pl. 25, fig. 11, Charleston, S. C.
Centropristis merus Poey, Synopsis Piscium Cubensium, 1868, 288, Cuba.
? *Cerna sicana* Doderlein, Rivista delle Specie del Genere *Epinephelus* o *Cerna*, 1882, 81, Palermo.
- Genus 827. **PROMICROPS** Gill. *Guasas*.
Promicrops Gill, in Poey, Synopsis, 1868, 287 (*Serranus guasa* Poey = *Itaiara* Lichtenstein).
Itaiara Vaillant and Bocourt, Miss. Sci. Mexique, 1875, 90 (*Serranus itaiara* Lichtenstein).
2398. *Promicrops itaiara* (Lichtenstein). *Guasa*; *Spotted jewfish*; *Méro*.
Atlantic coast of tropical America, from Florida to Brazil; not uncommon about rocks.
Cugupuguacu Catesby, Nat. Hist. Carolina, 1771, 14; nonbinomial.
Serranus itaiara Lichtenstein, Abhandl. Berlin Acad. Wiss., 1821 (1822), 278, Brazil; not *Itaiara* of Marcgrave.
Serranus galeus Müller and Troschel, in Schomburgk's Reise in Brit. Guiana, 1848, 621, Guiana.
Serranus guasa Poey, Memorias, II, 1861, 141; 1861, 354, Cuba.
2399. *Promicrops quinquefasciata* (Bocourt).
Pacific coast of Central America.
Serranus quinquefasciatus Bocourt, Ann. Sci. Nat., X, 1868, 223, Naguolate, Pacific coast of Guatemala; very close to *P. itaiara*.
- Genus 828. **ALPHESTES** Bloch and Schneider.
Alphestes Bloch and Schneider, Syst. Ichth., 1801, 236 (*Epinephelus afer* Bloch).
Prospinus Poey, in Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 287 (*Plectropoma chloropterus* Cuvier and Valenciennes = *Alphestes afer* Bloch).
2400. *Alphestes afer* (Bloch).
West Indies; Cuba to Brazil; recorded perhaps doubtfully from Guinea, if so, the American species, *A. chloropterus*, may prove different.
Epinephelus afer Bloch, Ichthyol., 1793, pl. 327, Acará in Guinea.
Plectropoma chloropterus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 300 (398), San Domingo, Martinique.

⁵⁶ The subgenus *Vivero* is based on *Epinephelus morio* (Cuv. & Val.) which differs from all other species of *Epinephelus* in the elevation of the second dorsal spine.

Plectropoma monacanthus Muller and Troschel, in Schomburgk's Hist. Barbados, 1847, 665, Barbados.

?*Serranus armatus* Osorio, Jour. Sci. Lisbon, III, 1894, 74.

2401. *Alphestes multiguttatus* (Gunther).

Pacific coast of tropical America; Mazatlan to Panama.

Plectropoma multiguttatum Gunther, Proc. Zool. Soc. Lond., 1866, 600, Panama.

2402. *Alpestes lightfooti* (Fowler).

San Domingo, West Indies.

Epinephelus lightfooti Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 258, fig 3, San Domingo.

Genus 829. *DERMATOLEPIS* Gill.

Dermatolepis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 54 (*Dermatolepis punctatus* Gill).

Lioperca Gill, loc. cit., XIV, 1862, 237 (*Serranus inermis* Cuvier and Valenciennes).

2403. *Dermatolepis punctatus* Gill.

West coast of Mexico; Cape San Lucas; the Venados; the Revillagigedos.

Dermatolepis punctatus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 54, Cape San Lucas.

2404. *Dermatolepis zancus* Evermann and Kendall.

Key West.

Dermatolepis zancus Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 129, pl. 8, fig. 9, Key West, Fla.

Subgenus *LIOPERCA* Gill.

2405. *Dermatolepis inermis* (Cuvier and Valenciennes).

West Indies.

Serranus inermis Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 322 (436), Antilles.

Genus 830. *TRISOTROPIS* Gill.

Trisotropis Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 104 (*Johnius guttatus* Bloch and Schneider = *Perca venenosa* Linnaeus).

Archoperca Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1171 (*Mycteroperca boulengeri* Jordan and Starks).

Subgenus *ARCHOPERCA* Jordan and Evermann.

2406. *Trisotropis boulengeri* (Jordan and Starks). *Cabrilla de raizero*; *Mangrove grouper*.

Mazatlan, Mexico.

Mycteroperca boulengeri Jordan and Starks, in Jordan, Fishes Sinaloa, Proc. Calif. Acad. Sci., V, 1895, 445, pl. 38, taken in the Astillero at Mazatlan, Mexico.

Subgenus *TRISOTROPIS* Gill.

2407. *Trisotropis venenosus* (Linnaeus). *Rockfish*; *Yellow-finned grouper*; *Bonaci de Piedra*.

Bahamas, Florida Keys, and southward.

Perca venenosa Linnaeus, Syst. Nat., Ed. X, 1758, 292, Bahamas; after Catesby.

Serranus petrosus Poey, Memorias, II, 1861, 136, Cuba.

2408. *Trisotropis venenosus apua* (Bloch). *Bonaci cardenal*.

West Indies, Florida Keys, and southward to Brazil. A red form of *T. venenosus*.

Bodianus apua Bloch, Nat. Ausl. Fische, IV, 1790, 50, Brazil; after a drawing by Prince Maurice—the same used by Marcgrave.

Pirati apia Marcgrave, Hist. Brasil., 1648, 158, Brazil.

Bonaci cardenal Parra, Piezas de Hist. Nat. Cuba, 1787, 29, Pl. XVI, Habana.

Johnius guttatus Bloch and Schneider, Syst. Ichth., 1801, 77; after Parra.

Bodianus marginatus Bloch and Schneider, loc. cit., 331; after Marcgrave.

Serranus rupestris Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 323 (437), San Domingo.

2409. *Trisotropis bonaci* (Poey). *Marbled rockfish; Bonaci arara; Black grouper; Aguaji; Bonaci*.
West Indies; Pensacola to Brazil; Woods Hole, Mass.
Serranus bonaci Poey, *Memorias*, II, 1861, 129, Cuba.
Serranus brunneus Poey, loc. cit., 131, Cuba.
Serranus decimatus Poey, loc. cit., 138, Cuba; specimen with ten dorsal spines.
Trisotropis aguaji Poey, *Repertorio*, II, 1868, 229, Habana.
Serranus cyclopomatus Poey, *Memorias*, 1861, 353, Cuba.
Serranus latepectus Poey, loc. cit., 353, Cuba.
2410. *Trisotropis xanthostictus* (Jordan and Swain).
Snapper Banks off Pensacola, Fla.
Mycteroperca bonaci xanthosticta Jordan and Swain, *Proc. U. S. Nat. Mus.*, VII, 1884, 371, Snapper Banks off Pensacola, Fla.
2411. *Trisotropis jordani* (Jenkins and Evermann). *Cabrilla de astillero; Baya*.
Pacific coast of Mexico from Gulf of California to Mazatlan.
Epinephelus jordani Jenkins and Evermann, *Proc. U. S. Nat. Mus.*, XI, 1888, 140, Guaymas, Sonora.
2412. *Trisotropis microlepis* Goode and Bean. *Gog; Aguaji*.
South Atlantic and Gulf coast of United States, north to Virginia, south to Pensacola and Key West.
Trisotropis microlepis Goode and Bean, *Proc. U. S. Nat. Mus.*, II, 1879, 141, west Florida.
Trisotropis stomias Goode and Bean, loc. cit., V, 1882, 427, Pensacola; Key West.
2413. *Trisotropis interstitialis* (Poey).
Coast of Cuba; Woods Hole, Mass.
Serranus interstitialis Poey, *Memorias*, II, 1861, 127, Cuba.
Trisotropis chlorostomus Poey, *Repertorio*, II, 1868, 231, Cuba.
2414. *Trisotropis dimidiatus* (Poey).
Coast of Cuba.
Serranus dimidiatus Poey, *Memorias*, II, 1861, 129, Cuba.
2415. *Trisotropis xenarchus* (Jordan).
Pacific coast of Mexico to the Venados Islands near Mazatlan and to the Galapagos Archipelago.
Mycteroperca xenarcha Jordan, *Proc. Acad. Nat. Sci. Phila.*, XXXIX, 1887, 387, Galapagos Islands; Payta, Peru.
- Genus 831. MYCTEROPERCA Gill.
Mycteroperca Gill, *Proc. Acad. Nat. Sci. Phila.*, XV, 1863, 80 (*Serranus olfax* Jenyns).
2416. *Mycteroperca venadorum* Jordan and Starks. *Garlopa*.
West coast of Mexico at Mazatlan.
Mycteroperca venadorum Jordan and Starks, in Jordan, *Proc. Calif. Acad. Sci.*, V, 1895, 446, Mazatlan, Mexico.
2417. *Mycteroperca bowersi* Evermann and Marsh.
Culebra and Vieques Islands, Porto Rico.
Mycteroperca bowersi Evermann and Marsh, *Bull. Bur. Fish.*, XX, 1900 (1902), 158, Culebra Island.
2418. *Mycteroperca hopkinsi* Jordan and Rutter.
Jamaica.
Mycteroperca hopkinsi Jordan and Rutter, *Proc. Acad. Nat. Sci. Phila.*, XLIX, 1897, 105, Jamaica.
2419. *Mycteroperca olfax* (Jenyns). *Yellow grouper; Gourami*.
James Island, Galapagos Archipelago; Panama.
Serranus olfax Jenyns, *Voyage Beagle*, Fishes, 1840, 9, pl. 4, Galapagos Islands.
2420. *Mycteroperca rosacea* (Streets). *Cabrilla calamaria*.
Gulf of California.
Epinephelus rosaceus Streets, *Bull. U. S. Nat. Mus.*, VII, 1877, 51, Angel Island, Gulf of California.

2421. *Mycteroperca falcata* (Poey). *Scamp; Bacalao; Abadejo*.
West Indies, north to Bermuda.
Serranus falcatus Poey, Memorias, II, 1861, 138, Habana.
2422. *Mycteroperca phenax* Jordan and Swain. *Scamp; Bacalao*.
Coasts of southern Florida; abundant about the keys.
Mycteroperca falcata phenax Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 363, Key West, Fla.
2423. *Mycteroperca calliura* Poey.
Coast of Cuba.
Mycteroperca calliura Poey, Repertorio, I, 1867, 181, Cuba.
2424. *Mycteroperca tigris* (Cuvier and Valenciennes). *Bonaci gato; Rockfish*.
West Indies, north to Bermuda.
Serranus tigris Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 325 (440), San Domingo.
Serranus felinus Poey, Memorias, II, 1861, 134, Habana
Serranus repandus Poey, loc. cit., 135, Habana
Trisotropis reticulatus Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 105, Barbados.
2425. *Mycteroperca camelopardalis* (Poey).
West Indies.
Serranus camelopardalis Poey, Memorias, II, 1861, 132, Cuba, a bright red form, otherwise similar to *M. tigris*.
- Genus 832. *XYSTROPERCA* Jordan and Evermann.
Xystroperca Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1181 (*Mycteroperca pardalis* Gilbert).
2426. *Xystroperca pardalis* (Gilbert). *Cabrilla pirutita*.
Gulf of California.
Mycteroperca pardalis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 551, La Paz Bay, Lower California.
- Genus 833. *PAREPINEPHELUS* Bleeker.
Parepinephelus Bleeker, Systema Percarum Revisum, in Archiv. Neer. des Sci. Exet. Nat., XI, 1875, 1876, 257 (*Serranus acutirostris* Cuvier and Valenciennes = *Epinephelus ruber* Bloch).
2427. *Parepinephelus acutirostris* (Cuvier and Valenciennes). *Scirenga*.
West Indies, Brazil, Mediterranean Sea, and islands of the eastern Atlantic.
*?Epinephelus ruber*⁵⁷ Bloch, Nat. Ausl. Fische, VII, 1793, 22, pl. 330, "Japan"; probably from the Mediterranean.
Serranus acutirostris Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 212 (286), Brazil.
Serranus undulosus Cuvier and Valenciennes, loc. cit., 219 (295), Brazil.
Serranus tinca Cantraine, "Nouv. Mem. Ac. Brux., XI, 1831," Naples.
Serranus fuscus Lowe, Trans. Cambr. Philos. Soc., VI, 1838, 196, Madeira.
Cerna nebulosa Cocco, "Indice Pesci. Messina, Gen. 45, sp. 2," 1844, Messina.
Cerna macrogenis Sassi, "Descr. Genovail Genovasato, I, 139," 1846, Genoa.
Serranus emarginatus Valenciennes, Ichth. Îles Canaries, 1839, 10, Canary Islands.
Epinephelus chalinus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 465, St. Martins.
Cerna acutirostris lata Doderlein, loc. cit., 74, Palermo.
- Genus 834. *CHORISTISTIUM* Gill.
Chorististium Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 15 (*Liopropoma rubra* Poey).
2428. *Chorististium rubrum* (Poey).
Coast of Cuba.
Liopropoma rubre Poey, Memorias, II, 1861, 418, Habana.

⁵⁷ The original type of *E. ruber* has 24 gill rakers; all others examined (Athem, Brazil) have 30 to 35; for these we retain the name *acutirostris* pending investigation.

Genus 835. LIOPROPOMA Gill.

Liopropoma Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 52 (*Perca aberrans* Poey).

2429. *Liopropoma aberrans* (Poey).

Coast of Cuba

Perca aberrans Poey, Memorias, II, 1861, 125, Cuba.

2430. *Liopropoma longilepis* Garman.

Eastern Pacific in deep water.

Liopropoma longilepis Garman, Mem. Mus. Comp. Zool., XXIV, 1889, 47, lat 7° 30' N., long 78° 34' W.

Genus 836. RYPTICUS Cuvier. Soapfishes.

Rypticus Cuvier, Règne Animal, ed. 2, II, 1829, 144 (*Anthias saponaceus* Bloch and Schneider).

Smeeticus Valenciennes, Ichth. Voy. *Vénus*, 1855, 305 (*S. bicolor* Valenciennes).

Eleutheractis Cope, Trans. Amer. Phil. Soc., XIV, 1871, 467 (*E. coriaceus* Cope).

Rhypticus authors, corrected spelling.

2431. *Rypticus saponaceus* (Bloch and Schneider). Soapfish; Jabon; Jabon-cillo.

West Indies; Pensacola to west Africa and Brazil.

Jaboncillo Parra, Difer. Piezas de Hist. Nat., 1787, 51, lam. 24, fig. 2, Habana.

Anthias saponaceus Bloch and Schneider, Syst. Ichth., 1801, 310, Habana.

2432. *Rypticus arenatus* Cuvier and Valenciennes.

West Indies and coast of Brazil; recorded from Jamaica, Trinity, Barbados, and St. Thomas.

Rypticus arenatus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 48 (65), pl. 46, Brazil

?*Rhypticus subbifrenatus* Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 53, St. Thomas

?*Rhypticus nigromaculatus* Steindachner, Ichth. Notiz., VI, 1867, 42, Barbados; dorsal with four spines instead of three.

2433. *Rypticus coriaceus* (Cope). Black soapfish.

West Indies; recorded from St. Martins and Jamaica.

Eleutheractis coriaceus Cope, Trans. Amer. Phil. Soc., XIV, 1870 (1871), 467, St. Martins.

2434. *Rypticus xanti* (Gill).

Pacific coast of Mexico, from Cape San Lucas to Colima and Mazatlan.

Rhypticus xanti Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 250, Cape San Lucas.

2435. *Rypticus bicolor* (Valenciennes).

Galapagos Archipelago.

Smeeticus bicolor Valenciennes, Voy. *Vénus*, Poissons, 1855, 307, pl. 2, fig. 2, Galapagos Archipelago.

Genus 837. PROMICROPTERUS Gill.

Promicropterus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 53 (*Rhypticus maculatus* Holbrook=*Bodianus bistrispinus* Mitchill).

2436. *Promicropterus bistrispinus* (Mitchill). Soapfish.

South Atlantic coast of United States; frequent off Charleston, Pensacola, and Key West, Porto Rico; occasional as far north as Newport, R. I.

Bodianus bistrispinus Mitchill, Amer. Month. Mag. and Crit. Review, II, February, 1818, 247, Straits of Bahama.

Rypticus maculatus Holbrook, Ichth. South Carolina, Ed. I, 1855, 39, Cape Romain, S. C.

Rhypticus pituitosus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 341, Key West, Fla.

2437. *Promicropterus nigripinnis* (Gill).
Pacific coast of tropical America, Cape San Lucas to Panama.
Rhypticus nigripinnis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 53, Panama.
Rhypticus maculatus Gill, loc. cit., XIV, 1862, 251, Cape San Lucas.
Promicropterus decoratus Gill, loc. cit., XV, 1863, 164, Panama.

Family 195. SERRANIDÆ. Sea bass

- Genus 838. *PARALABRAX* Girard. *Cabrillas verdes*.
Paralabrax Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 131 (*Labrax nebulifer* Girard).
2438. *Paralabrax nebulifer* (Girard). *Johnny verde*; *Kelp bass*.
Southern California, from Monterey to Magdalena Bay.
Labrax nebulifer Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 142, Monterey, Calif.
2439. *Paralabrax maculatofasciatus* (Steindachner). *Spotted cabrilla*.
Pacific coast of America from Lower California to San Pedro and Mazatlan.
Serranus maculatofasciatus Steindachner, Ichth. Notizen, VII, 1868, 5, Mazatlan, Mexico.
? *Serranus acanthophorus* Bocourt, Ann. Sci. Nat., X, 1870, 223, Pacific coast of Mexico.
2440. *Paralabrax humeralis* (Cuvier and Valenciennes).
Pacific coast of South America; Panama to Juan Fernandez.
Serranus humeralis Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 183 (246), Chile.
Serranus albomaculatus Jenyns, Zool. Beagle, Fishes, 1840, 3, pl. 2, Galapagos Archipelago.
? *Serranus semifasciatus* Guichenot, in Gay, Hist. Chile, II, 1847, 151, pl. 1, fig. 2, Juan Fernandez.
2441. *Paralabrax clathratus* (Girard). *Cabrilla*; *Rock bass*.
Coast of southern California, from San Francisco to Cedros Island.
Labrax clathratus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 143, San Diego.
- Genus 839. *CRATINUS* Steindachner.
Cratinus Steindachner, Ichth. Beitr., VII, 1878, 19 (*C. agassizii* Steindachner).
2442. *Cratinus agassizii* Steindachner.
Galapagos Archipelago.
Cratinus agassizii Steindachner, Ichth. Beitr., VII, 1878, 19, Galapagos Islands.
- Genus 840. *CENTROPRISTES* Cuvier and Valenciennes. *Black sea bass*.
Centropristes Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 27 (36) (*C. nigricans* Cuvier and Valenciennes = *Labrus striatus* Linnæus).
Centropristes of authors; variation in spelling.
2443. *Centropristes striatus* (Linnæus). *Black sea bass*; *Blackfish*; *Tallywag*; *Hannahill*; *Black Will*; *Black Harry*; *Black perch*; *Black bass*; *Bluefish*.
Atlantic coast of United States, Cape Ann to northern Florida.
Labrus striatus Linnæus, Syst. Nat., ed. X, 1758, 285, "America"; description very brief, but not to be referred to any other fish.
Perca atraria Linnæus, loc. cit., ed. XII, 1766, 485, Carolina.
Perca furva Walbaum, Artedi Pisc., 1792, 336, New York; after Schöpfung.
Coryphæna nigrescens Bloch and Schneider, Syst. Ichth., 1801, 297, New York.
Lutjanus trilobatus Lacépède, Hist. Nat. Poiss., IV, 1802, 246, locality unknown.
Perca varia Mitchill, Trans. Lit. & Phil. Soc. N. Y., I, 1815, 415, pl. 3, fig. 6, New York.

2444. *Centropristes ocyurus* (Jordan and Evermann) *Gulf sea bass*.
Gulf of Mexico at Snapper Banks off Pensacola, Fla
Serranus ocyurus Jordan and Evermann, Proc. U. S. Nat. Mus., IX,
1886, 468, Snapper Banks off Pensacola, Fla.

2445. *Centropristes rufus* Cuvier and Valenciennes. *Hogfish*.
Martinique
Centropristes rufus Cuvier and Valenciennes, Hist. Nat. Poiss., III,
1829, 35 (47), Martinique.

Genus 841. **TRILOBURUS** Gill.

Triloburus Gill, Cat. Fish. East Coast U. S., Supplement to Proc. Acad.
Nat. Sci. Phila., XIII, 1861, 30 (*Perca trifurca* Storer = *Perca philadel-*
phica Linnæus)

2446. *Triloburus philadelphicus* (Linnæus). *Rock sea bass, Redfish*.
Coast of South Carolina.
Perca philadelphica Linnæus, Syst. Nat., ed. X, 1758, 291, America
Perca trifurca Linnæus, loc. cit., ed. XII, 1766, 489, Carolinas
Lutjanus tridens Lacépède, Hist. Nat. Poiss., IV, 1802, 246, Carolinas.

Genus 842. **DIPLECTRUM** Holbrook. *Squirrelfishes*.

Diplectrum Holbrook, Ichth. South Carolina, ed. I, 1855, 32 (*Serranus fascicularis* Cuvier and Valenciennes = *Perca formosa* Linnæus).
Haliperca Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 236, (*Serranus bivittatus* Cuvier and Valenciennes = *Serranus radialis* Quoy and Gaimard, and other species).
Paraserranus Bleeker, Verh. Akad. Amst., XIV, 1873, 6 (*P. hasselti* Bleeker).

Subgenus **HALIPERCA** Gill.

2447. *Diplectrum radiale* (Quoy and Gaimard). *Aguavina*.
Cuba to Brazil
Serranus radialis Quoy and Gaimard, Voyage *Uranie*, 1824, 316, Rio Janeiro.
Serranus bivittatus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 179 (241), Martinique.
Centropristes ayresi Steindachner, Ichth. Notizen, VII, 1868, 1, pl. 1, fig. 1, Santos, Brazil.

2448. *Diplectrum pacificum* Meek and Hildebrand.

Panama.
Diplectrum radiale Gilbert and Starks, Memoir Calif. Acad. Sci., IV, 1904, 97 (not of Quoy and Gaimard).
Diplectrum pacificum Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, Apr. 15, 1925, 477, Pl. XLVI, fig. 2, Naos Island, Panama.

2449. *Diplectrum macropoma* (Günther).

Pacific coast of tropical America.
Centropristes macropoma Gunther, Proc. Zool. Soc. Lond., 1864, 145, Panama.

2450. *Diplectrum sciurus* Gilbert.

Gulf of California.
Diplectrum sciurus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 550, Gulf of California at Albatross stations Nos 3014, 3021, 3026, and 3033, all in shallow water.

2451. *Diplectrum euryplectrum* Jordan and Bollman.

Coast of Colombia, southwest of Panama
Diplectrum euryplectrum Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 157, Pacific Ocean, off coast of Colombia.

Subgenus **DIPLECTRUM** Holbrook.

2452. *Diplectrum formosus* (Linnæus). *Squirrelfish; Sandfish; Serrano*.
West Indies; common on the south Atlantic and Gulf coasts of the United States, from Charleston south to Montevideo.
Perca formosa Linnæus, Syst. Nat., ed. XII, 1766, 488, Carolinas.
Epinephelus striatus Bloch, Naturgesch. Ausl. Fische, VII, 1793, 20, Jamaica (not *Anthias striatus* Bloch).
Serranus radialis Quoy and Gaimard, Voy. *Uranie*, Poiss., 1824, 313, pl. 58, fig. 2, Rio de Janeiro.

Serranus irradians Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 181 (244), Montevideo.

Serranus fascicularis Cuvier and Valenciennes, loc. cit., 182 (245), pl. 30, Brazil.

Genus 843. PRIONODES Jenyns.

Prionodes Jenyns, Voy. *Beagle*, Fishes, 1842, 46 (*P. fasciatus* Jenyns).

2453. *Prionodes æquidens* (Gilbert).

Gulf of California.

Serranus æquidens Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 61, Gulf of California at Albatross station No. 2996, in 112 fathoms.

2454. *Prionodes fuscus* (Poey).

Cuba.

Centropristes fuscus Poey, Memorias, II, 1861, 342, Cuba.

2455. *Prionodes phæbe* (Poey). *Phæbe*.

West Indies, north to Pensacola, Fla.

Serranus phæbe Poey, Memorias, I, 1852, 55, pl. 2, fig. 3, Cuba.

2456. *Prionodes baldwini* Evermann and Marsh.

Porto Rico.

Prionodes baldwini Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (1900), 353, Culebra and Vieques Islands.

2457. *Prionodes fasciatus* Jenyns.

Pacific coast of Mexico, Cape San Lucas to Galapagos Islands.

Prionodes fasciatus Jenyns, Voy. *Beagle*, Fishes, 1840, 46, Chatham Island, Galapagos Archipelago.

Serranus psittacinus Valenciennes, Voy. *Vénus*, Poiss., 1855, 299, pl. 1, fig. 1, Galapagos Islands.

Serranus calopteryx Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 350, Mazatlan.

2458. *Prionodes bulleri* (Boulenger).

Coast of Jalisco, western Mexico.

Serranus bulleri Boulenger, Cat., I, 1895, 288, Las Peñas, Jalisco, Mexico.

2459. *Prionodes deweegeri* (Metzelaar).

Caribbean Sea.

Serranus (Paralabrax) deweegeri Metzelaar, Trop. Atl. Visschen, 1919, 52, fig. 20, Guanta, coast of Venezuela.

2460. *Prionodes tabacaria* (Cuvier and Valenciennes) *Jacome; Bout de tabac*.

West Indies.

Centropristes tabacarius Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 33 (44), Martinique.

Serranus jacome Poey, Memorias, I, 1852, 57, pl. 2, fig. 1, Cuba.

2461. *Prionodes tigrinus* (Bloch).

West Indies

Holocentrus tigrinus Bloch, Naturgesch. Ausl. Fische, IV, 1790, 77, East Indies; after Seba, Thesaurus, III, Pl. XXVII, fig. 5

Serranus prestigiator Poey, Memorias, I, 1852, 58, pl. 2, fig. 2, Habana.

?*Centropristes annularis* Gunther, Shore Fishes, *Challenger* Rept., Zool., I, pt. 6, 1880, 6, pl. 1, fig. C, Pernambuco.

2462. *Prionodes flavescens* (Cuvier and Valenciennes).

Martinique.

Serranus flavescens Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 380 (506), Martinique.

Genus 844. MENTIPERCA Gill.

Mentiperca Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 236 (*Serranus luciopercanus* Poey).

2463. *Mentiperca luciopercana* (Poey).

Coast of Cuba.

Serranus luciopercanus Poey, Memorias, I, 1852, 56, pl. 9, fig. 1, Cuba.

Genus 845. CENTRISTHMUS Garman.

Centristhmus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 47 (*Centristhmus signifer* Garman).

2464. *Centristhmus signifer* Garman.

Off Bay of Panama in deep water.

Centristhmus signifer Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 48, Pl. LXIX, off Bay of Panama.**Genus 846. *DULES* Cuvier.***Dules* Cuvier, Règne Animal, ed. 2, II, 1829, 147 (*Dules auriga* Cuvier and Valenciennes).*Eudulus* Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 264 (*Dules auriga* Cuvier and Valenciennes; *Dulus* being a genus of birds).**2465. *Dules auriga* Cuvier and Valenciennes. *Coachman*; *Charioteer*.**

Coasts of Brazil and Uruguay, and occasionally northward.

Dules auriga Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 83 (112), pl. 51, Brazil.*Centropristis brasiliensis* Brissot de Barneville, Révue Zoologique, 1847, 131, Bahia**Genus 847. *CALLIDULUS* Fowler.***Callidulus* Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 265 (*Centropristes subligarius* Cope).**2466. *Callidulus flaviventris* Cuvier and Valenciennes. *Grassy ground rock-fish*.**

Tropical America, Charleston to Brazil.

Dules flaviventris Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 85, Brazil.*Centropristes displurus* Gunther, Proc. Zool. Soc. Lond., 1867, 99, Trinidad.*Centropristes subligarius* Cope, Proc. Acad. Nat. Sci. Phila., XXII, 1870, 120, Pensacola, Fla.**Genus 848. *PARANTHIAS* Guichenot.***Brachyrhinus* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 286 (*Serranus creolus* Cuvier and Valenciennes).*Paranthias* Guichenot, Ann. Soc. Linn. Maine-et-Loire, X, 1868 (*Serranus furcifer* Cuvier and Valenciennes=*Serranus creolus* Cuvier and Valenciennes).*Creolus* Jordan and Gilbert, Synopsis, Bull. U. S. Nat. Mus., XVI, 1882, XXXVI (withdrawn in the text) (*Serranus furcifer* Cuvier and Valenciennes).**2467. *Paranthias furcifer* (Cuvier and Valenciennes). *Rabirubia de lo alto*; *Creolefish*.**

Both coasts of tropical America, Cuba to Brazil, Cape San Lucas to the Galapagos Archipelago.

Serranus furcifer Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 196 (264), Brazil.*Serranus creolus* Cuvier and Valenciennes, loc. cit., 197 (265), Martinique; San Domingo*Corvina oxyptera* DeKay, New York Fauna: Fishes, 1842, 77, Pl. XXX, fig. 96, locality unknown.*Serranus colonus* Valenciennes, Voy. *Vénus*, Zool., 1855, 300, pl. 2, fig. 1, Galapagos Islands.**Genus 849. *OCYANTHIAS* Jordan and Evermann.***Ocyanthias* Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1227 (*Aylopon martinicensis* Guichenot).**2468. *Ocyanthias martinicensis* (Guichenot).**

West Indies.

Aylopon martinicensis Guichenot, Anthiani, Ann. Linn. Soc., X, 1868, Martinique.**Genus 850. *HYPOPLECTRUS*⁸⁸ Gill. *Vacas*.***Hypoplectrus* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 236 (*Plectropoma puella* Cuvier and Valenciennes=variety of *Perca unicolor* Walbaum).

⁸⁸ The West Indian forms correctly referred to this species fall into several groups here provisionally designated as subspecies of the form first named (*H. unicolor*), apparently differing only in color. These range from mottled olive and orange to intense indigo blue.

2469. *Hypoplectrus unicolor* (Walbaum). *Vaca*; *Petitnègre*.
West Indies, north to Florida Keys
Perca unicolor Walbaum, *Artedi Piscium*, III, 1792, 352, locality unknown.
Plectropoma ephippium Cuvier and Valenciennes, *Hist. Nat. Poiss.*, II, 1828, 307, (408), locality unknown.
Hypoplectrus maculiferus Poey, *Ann. Lyc. Nat. Hist. N. Y.*, X, 1871, 78, Habana.
2470. *Hypoplectrus unicolor puella* (Cuvier and Valenciennes). *Vaca*.
Martinique.
Plectropoma puella Cuvier and Valenciennes, *Hist. Nat. Poiss.*, II, 1828, 305 (405), pl. 37, Martinique.
2471. *Hypoplectrus unicolor vitulinus* (Poey).
Cuba.
Plectropoma vitulinum Poey, *Memorias*, I, 1852, 68, Cuba.
2472. *Hypoplectrus unicolor pinnivarius* (Poey).
Habana.
Hypoplectrus pinnivaria Poey, *Synopsis*, 1868, 291, Habana.
2473. *Hypoplectrus unicolor guttavarius* (Poey).
Cuba.
Plectropoma guttavarium Poey, *Memorias*, I, 1852, 70, Cuba.
Plectropoma melanorhina Guichenot, *Poissons*, in *Ramon de la Sagra*, *Hist. Cuba*, 1853, 18, pl. 1, fig. 1, Cuba.
2474. *Hypoplectrus unicolor gummigutta* (Poey).
Cuba.
Plectropoma gummigutta Poey, *Memorias*, I, 1852, 70, Cuba.
2475. *Hypoplectrus unicolor crocotus* (Cope).
West Indies.
Plectropoma crocota Cope, *Trans. Amer. Phil. Soc.*, XIV, 1870 (1871), 466, St. Martins, West Indies.
2476. *Hypoplectrus unicolor aberrans* (Poey).
Cuba.
Hypoplectrus aberrans Poey, *Synopsis*, 1868, 291, Cuba.
2477. *Hypoplectrus unicolor accensus* (Poey).
Cuba.
Plectropoma accensum Poey, *Memorias*, I, 1852, 72, Cuba.
2478. *Hypoplectrus unicolor affinis* (Poey).
Cuba.
Plectropoma affine Poey, *Memorias*, II, 1861, 427, Cuba.
2479. *Hypoplectrus unicolor chlorurus* (Cuvier and Valenciennes).
Martinique.
Plectropoma chlorurum Cuvier and Valenciennes, *Hist. Nat. Poiss.*, II, 1828, 306 (406), Martinique.
2480. *Hypoplectrus unicolor nigricans* (Poey).
Cuba.
Plectropoma nigricans Poey, *Memorias*, I, 1852, 71, Cuba.
2481. *Hypoplectrus unicolor indigo* (Poey). *Añil*.
Cuba.
Plectropoma indigo Poey, *Memorias*, I, 1852, 69, pl. 3, fig. 1, Cuba.
2482. *Hypoplectrus unicolor bovinus* (Poey).
Cuba.
Plectropoma bovinum Poey, *Memorias*, I, 1852, 69, Cuba.
2483. *Hypoplectrus unicolor gemma* Goode and Bean.
Florida Keys; known from one specimen from Garden Key, Fla.
Hypoplectrus gemma Goode and Bean, *Proc. U. S. Nat. Mus.*, V, 1882, 428, Garden Key, Fla.
2484. *Hypoplectrus lamprurus* (Jordan and Gilbert).
Panama.
Serranus lamprurus Jordan and Gilbert, *Bull. U. S. Fish Com.*, I, 1881, 322, Panama.

Genus 851. ANTHIAS Bloch. *Barbiers*.*Anthias* Bloch, Nat. Ausl. Fische, VI, 1792, 97 (*Labrus anthias* L.).*Aylopon* Rafinesque, Caratteri, 1810, 52, (*Labrus anthias* L.; the name *Anthias* said to be preoccupied).**2485. *Anthias tenuis* Nichols.**

Bermuda.

Anthias tenuis Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 60, Bermudas.**Genus 852. HEMIANTHIAS Steindachner.***Hemianthias* Steindachner, Ichth. Beitr., I, 1874, 4 (*Anthias* (*Hemianthias*) *peruanus* Steindachner).**2486. *Hemianthias peruanus* (Steindachner).**

Coasts of Peru and Chile, occasionally northward; off the coast of Lower California.

Anthias (*Hemianthias*) *peruanus* Steindachner, Ichth. Beitr., I, 1874, 4, Payta; Trujillo.**2487. *Hemianthias vivanus* (Jordan and Swain).**

Gulf of Mexico; Snapper Banks between Pensacola and Tampa, Fla.

Anthias vivanus Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 544, from stomach of a red snapper taken at the Snapper Banks off Pensacola, Fla.**Genus 853. PRNOTOGRAMMUS Gill.***Pronotogrammus* Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 80 (*P. multifasciatus* Gill).**2488. *Pronotogrammus multifasciatus* Gill.**

Off Bay of Panama; Cape San Lucas, Lower California.

Pronotogrammus multifasciatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 81, Cape San Lucas, Lower California.**2489. *Pronotogrammus eos* Gilbert.**

Pacific coast of tropical America.

Pronotogrammus eos Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 62, west coast of Mexico, at Albatross station 2996, in 112 fathoms.**Family 196. PSEUDOCROMIDÆ****Genus 854. GRAMMA Poey.***Gramma* Poey, Synopsis, 1868, 296 (*G. loreto* Poey).**2490. *Gramma loreto* Poey.**

Matanzas, Cuba; only the type known.

Gramma loreto Poey, Synopsis, 1868, 296, Matanzas, Cuba.**Family 197. RHEGMATIDÆ****Genus 855. RHEGMA Gilbert.***Rhegma* Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 3169 (*Rhegma thaumasium* Gilbert).**2491. *Rhegma thaumasium* Gilbert.**

Panama.

Rhegma thaumasium Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 3170, Panama.**Family 198. PRIACANTHIDÆ. Big-eyes or catalufas****Genus 856. PRIACANTHUS Cuvier.***Priacanthus* Cuvier, Règne Animal, ed. 1, 1817, 281 (*P. macrophthalmus* Cuvier).**2492. *Priacanthus cruentatus* (Lacépède). *Big-eye; Catalufa*.**Tortugas; West Indies to St. Helena and the Canaries (*Priacanthus carolinus* Lesson), a species very similar or identical in Hawaii and the western Pacific.*Labrus cruentatus* Lacépède, Hist. Nat. Poiss., III, 1802, 522, Martinique; from a copy by Aubriet of a plate made by Plumier at Martinique.

- Priacanthus cepedianus* Desmarest, Prém. Dec. Ichthy., 1823, 9, pl. 1, Habana.
 ?*Priacanthus carolinus* Lesson, Voy. Coquille, 1826, 204, Caroline Islands.
2493. *Priacanthus arenatus* Cuvier and Valenciennes. *Catalufa*; *Big-eye*. West Indies, south to Brazil; occasionally northward in the Gulf Stream to Newport and Woods Hole, Mass.
Catalufa Parra, Dif. Piezas Hist. Nat., 1787, 20, Habana.
Priacanthus arenatus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 75 (101), Brazil.
Priacanthus fulgens Lowe, Trans. Zool. Soc. Lond., II, 1839, 174, Madeira.
Priacanthus catalufa Poey, Proc. Acad. Nat. Sci. Phila., XV, 1863, 182, Habana.
- Genus 857. **PSEUDOPRIACANTHUS** Bleeker.
Pseudopriacanthus Bleeker, Versl. Ak. Wet. Amsterdam, 1869 (2), 241 (*Priacanthus nipponius* Cuvier and Valenciennes).
2494. *Pseudopriacanthus altus* (Gill)
 West Indies and the Gulf Stream.
Priacanthus altus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 732, Narragansett Bay.
2495. *Pseudopriacanthus serrula* (Gilbert).
 Pacific coast of Colombia.
Priacanthus serrula Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 450, Albatross station, 2797, west coast of Colombia.

Family 199. **PEMPHERIDÆ**. Deep-water catalufas

- Genus 858. **PEMPHERIS** Cuvier.
Pempheris Cuvier, Règne Anim., ed. 2, II, 1829, 195 (*Kurtus argenteus* Bloch and Schneider = *Pempheris touea* Cuvier).
2496. *Pempheris schomburgkii* Müller and Troschel.
 Barbados and Cuba.
Pempheris schomburgkii Müller and Troschel, in Schomburgk's History of Barbados, 1848, 669, Barbados.
2497. *Pempheris mülleri* Poey. *Catalufa de lo alto*.
 West Indies to Brazil.
Pempheris mülleri Poey, Memorias, II, 1861, 203, Cuba.
2498. *Pempheris poeyi* Bean.
 Cuba.
Pempheris poeyi Bean, Proc. U. S. Nat. Mus., VIII, 1885, 229, Habana.
2499. *Pempheris mexicanus* Cuvier and Valenciennes.
 Pacific coast of Mexico, Acapulco.
Pempheris mexicana Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 230 (308), Acapulco, Mexico.

Family 200. **LOBOTIDÆ**. Flashers or tripletails

- Genus 859. **LOBOTES** Cuvier.
Lobotes Cuvier, Règne Anim., ed. 2, II, 1829, 177 (*L. erate* Cuvier and Valenciennes).
2500. *Lobotes surinamensis* (Bloch). *Flasher*; *Tripletail*; *Dormeur*; *Black perch*; *Black grunt*.
 Atlantic coast from Cape Cod to Surinam; Mediterranean; Japan.
Holocentrus surinamensis Bloch, Naturgesch. Ausl. Fische, IV, 1790, 98, Surinam.
Bodianus triourus Mitchell, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 418, Powells Hook, N. J.
Lobotes somnolentus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 244 (324), pl. 126, San Domingo.
Lobotes erate Cuvier and Valenciennes, loc. cit., 322, Pondicherry.
Lobotes auctorum Günther, Cat., I, 1859, 338, Cuba; Calcutta; China.
- Genus 860. **VERRUGATO** Jordan.
Verrugato Jordan, Classn. Fishes, 1923, 195 (*Lobotes pacificus* Gilbert).

2501. *Verrugato pacificus* (Gilbert). *Berrugate*.

Panama.

Lobotes pacificus Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2857, Panama.**Family 201. XENICHTHYIDÆ****Genus 861. XENICHTHYS Gill.***Xenichthys* Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 82 (*X. xanti* Gill).**2502. *Xenichthys xanti* Gill.**

Pacific coast of tropical America, from Cape San Lucas to Panama.

Xenichthys xanti Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 82, Cape San Lucas.*Xenichthys xenops* Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 325, Panama.**Genus 862. XENISTIUS Jordan and Gilbert.***Xenistius* Jordan and Gilbert, Synopsis, 1883, 920 (*Xenichthys californiensis* Steindachner).**2503. *Xenistius californiensis* (Steindachner).**

Pacific coast of America, from San Diego southward to Guaymas, Mexico.

Xenichthys californiensis Steindachner, Ichth. Beitr., III, 1875, 3, San Diego.**Family 202. HOPILOPAGRIDÆ****Genus 863. HOPILOPAGRUS Gill.***Hoplopagrus* Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 78; and 1862, 253 (*Hoplopagrus guntheri* Gill.)**2504. *Hoplopagrus guntheri* Gill. *Pargo*.**

Pacific coast of tropical America, from Guaymas to Panama.

Hoplopagrus guntheri Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 78, and 1862, 253, Cape San Lucas, Lower California.**Family 203. LUTIANIDÆ. Snappers****Genus 864. LUTIANUS Bloch.***Lutianus* Bloch, Nat. Ausl. Fische, IV, 1790, 105 (*L. lutianus* Bloch).*Diacope* Cuvier, Mém. Mus. Hist. Nat. Paris, I, 1816, 360 (*Holocentrus bengalensis* Bloch, preoccupied).*Mesoprion* Cuvier (Règne Animal, 143), Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 441 (*L. lutianus* Bloch).*Genyorange* Cantor, Jour. Roy. Asiat. Soc. Bengal, XVIII, 1849, 12 (*Diacope sebæ* Cuvier and Valenciennes).*Neomænis* Girard, Proc. Acad. Nat. Sci. Phila., X, 1858 (1859), 167 (*Lobotes emarginatus* Baird and Girard = *Labrus griseus* Linnaeus).*Hypoplites* Gill, loc. cit., XIV, 1862, 236 (*Mesoprion retrospinis* Cuvier and Valenciennes).*Evoplites* Gill, loc. cit., 236 (*Mesoprion pomacanthus* Bleeker, young of *L. kasmira*).*Lutjanus* Bloch and later writers, change of spelling; the word from the Indian "Lutjang."**Subgenus EVOPLITES Gill.****2505. *Lutianus viridis* (Valenciennes).**

Rocky islands of the eastern Pacific; known from Galapagos, Tres Marias, and Revillagigedo Islands.

Diacope viridis Valenciennes, Voy. *Vénus*, 1855, 303, pl. 1, fig. 2 (very bad), Galapagos Islands.**Subgenus NEOMÆNIS Girard.****2506. *Lutianus jordani* (Gilbert)**

Panama; Cocos Island.

Neomænis jordani Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1252, Panama.

- 2507. *Lutianus novemfasciatus* (Gill).** *Pargo prieto; Pargo moreno; Pargo negro.*
Pacific coast of tropical America; Cape San Lucas; Guaymas; Mazatlan; Punta Arenas; San Blas; Panama.
Lutjanus novemfasciatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 251, Cape San Lucas, Lower California.
Mesoprion pacificus Bocourt, Ann. Sci. Nat. Paris, X, 1868, 223, Tauesco, Pacific coast of Guatemala.
Lutjanus prieto Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 353, Mazatlan.
- 2508 *Lutianus cyanopterus* (Cuvier and Valenciennes).** *Cubera.*
West Indies to Brazil.
Mesoprion cyanopterus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 357 (472), Brazil.
? *Mesoprion pargus* Cuvier and Valenciennes, loc. cit., 358 (473), Porto Rico.
Lutjanus cubera Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1871, 75, Cuba.
Lutjanus dentatus A. Duméril in Vaillant and Bocourt, Misc. Sci. Mex. Poiss., 1881, 125, Brazil.
- 2509. *Lutianus griseus* (Linnæus).** *Gray snapper; Mangrove snapper; Cabellero; Lawyer.*
West Indies; ranging from New York to Brazil.
Turdus pinus branchialis carens Catesby, Hist. Carolinas, II, 1743, 5, pl. 9, Bahamas.
Labrus griseus Linnæus, Syst. Nat., ed. X, 1758, 283, after Catesby.
Caballero Parra, Descr. Dif. Piezas, Hist. Nat., 1787, 52, pl. 25, fig. 1, Habana.
Sparus tetracanthus Bloch, Nat. Ausl. Fische, 1791, 116, Martinique.
Anthias caballero Bloch and Schneider, Syst. Ichth., 1801, 310, Cuba; after Parra.
Bodianus vivanet Lacépède, Hist. Nat. Poiss., IV, 1802, pl. 4, fig. 3, Martinique.
Mesoprion griseus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 355 (469), San Domingo, not after Linnæus.
Lobotes emarginatus Baird and Girard, 9th Smith. Rept., 1854 (1855), 332, Beaseley Point, N. J.
Lutjanus stearns Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 179, Pensacola, Fla.
- 2510. *Lutianus jocú* (Bloch and Schneider).** *Dog snapper; Jocú.*
West Indies, north to Florida Keys, south to Bahia; one young recorded from Woods Hole, Mass. (Nichols and Breder).
Anthias jocú Bloch and Schneider, Syst. Ichth., 1801, 310, Cuba; after Parra.
Jocú Parra, Descr. Dif. Piezas, Hist. Nat., I, 1787, 53, pl. 25, fig. 2, Cuba.
Mesoprion litura Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 353 (467), Cayenne; St. Thomas.
- 2511. *Lutianus apodus* (Walbaum).** *Schoolmaster; Caji.*
West Indies, north to Key West, south to Bahia; young casual at Woods Hole, Mass. (Nichols and Breder).
Perca apoda Walbaum, Artedi Piscium, 1792, 351; based on the "Schoolmaster" of Catesby, which, in the very bad plate, lacks the pectoral fin.
Caxis Parra, Descr. Dif. Piezas, Hist. Nat., 1787, 14, pl. 8, fig. 2, Habana.
Sparus caxis Bloch and Schneider, Syst. Ichth., 1801, 284, Habana; after Parra.
Bodianus striatus Bloch and Schneider, loc. cit., 1801, 335, pl. 65, West Indies, misprinted *albostratus* and *fasciatus*.
Lutjanus acutirostris Desmarest, Prem. Dec. Ichth., 1823, 12, pl. 3, Cuba.
Mesoprion cynodon Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 351 (465), Martinique, San Domingo.
Mesoprion linea Cuvier and Valenciennes, loc. cit., 354 (468), Cuba; San Domingo.
Mesoprion flavescens Cuvier and Valenciennes, loc. cit., 357 (472), Martinique.

2512. *Lutianus lutjanoides* (Poey).
Cuba
Ocyurus lutjanoides Poey, Ann. Lyc. Nat. Hist., IX, 1870, 319, Cuba;
known from one specimen, suspected to be a hybrid between *Ocyurus*
chrysurus and *Lutianus apodus*.
2513. *Lutianus brachypterus* (Cope).
Bahama Islands.
One specimen known
Lutjanus brachypterus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 470,
New Providence, Bahama Islands; perhaps a hybrid of *Lutianus*
griseus and *L. synagris*.
2514. *Lutianus argentiventris* (Peters). *Pargo amarillo*
Pacific coast of tropical America; Galapagos and Cocos Islands.
Mesoprion argentiventris Peters, Berlin Monatsber., 1869, 704, Mazatlan.
2515. *Lutianus hastingsi* (Bean) *Bermuda silk snapper*.
Bermuda.
Neomænis hastingsi Bean, Bull. Amer. Mus. Nat. Hist., X, Art. III,
1898, 45, Bermuda.
2516. *Lutianus buccanella* (Cuvier and Valenciennes). *Sesú de lo alto*;
Orielle noire; *Boucanello*; *Blackfin snapper*.
West Indies; Habana
Mesoprion buccanella Cuvier and Valenciennes, Hist. Nat. Poiss., II,
1828, 344 (455), Martinique.
Mesoprion caudanotatus Poey, Memorias, I, 1854, pl. 3, fig. 3, 440, Cuba.
2517. *Lutianus vivanus* (Cuvier and Valenciennes). *Pargo de lo alto*; *Silk*
snapper; *West Indian snapper*.
West Indies; Habana.
Mesoprion vivanus Cuvier and Valenciennes, Hist. Nat. Poiss., II,
1828, 343 (454), Martinique
Mesoprion profundus Poey, Memorias, II, 1861, 150, Cuba.
Lutjanus torridus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 469, St.
Kitts.
Lutjanus purpureus Poey, Enumeratio, 1875, 29.
2518. *Lutianus campechanus* ⁵⁹ (Poey). *Red snapper*; *Mexican snapper*;
Pargo guachenarigo.
Long Island to Yucatan and Texas; a valued food fish.
Mesoprion campechanus Poey, Memorias, II, 1861, 149, Campeche.
2519. *Lutianus blackfordii* Goode and Bean. *Pensacola snapper*.
Lutjanus blackfordii Goode and Bean, Proc. U. S. Nat. Mus., I, 1878,
176, Pensacola, Fla.
2520. *Lutianus analis* (Cuvier and Valenciennes). *Muttonfish*; *Pargo*; *Pargo*
criollo.
West Indies; Pensacola to Brazil, rarely north to Woods Hole, common
and valued.
Anthias quartus rondeleti (the muttonfish) Catesby, Nat. Hist. Carolina,
II, 1743, 25, pl. 25, Bahamas.
Mesoprion analis Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828,
341 (452), San Domingo.
Mesoprion sobra Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828,
342 (453), Martinique.
Mesoprion isodon Cuvier and Valenciennes, loc. cit., IX, 1833, 328 (443),
San Domingo.
Mesoprion rosaceus Poey, Ann. Lyc. Nat. Hist. N. Y., IX, 1870, 317,
Cuba.
2521. *Lutianus colorado* (Jordan and Gilbert). *Pargo colorado*.
Pacific coast of Mexico, Guaymas to Panama.
Lutjanus colorado Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
338, name only, 351, 355, Mazatlan, Mexico.
2522. *Lutianus guttatus* (Steindachner). *Flamenco*.
Pacific coast of Mexico, from Guaymas to Panama.
Mesoprion guttatus Steindachner, Ichthyol. Notizen, IX, 1869, 18, pl. 8,
Mazatlan, Mexico.

* The common red snapper is here admitted provisionally as distinct from the Brazilian *Lutianus aya* from which it is said by Hildebrand to differ tangibly

- 2523. *Lutianus synagris* (Linnæus).** *Lane snapper; Bjaiaiba; Redtail snapper.*
West Indies; Florida Keys to Colon and Brazil
Sparus synagris Linnæus, Syst. Nat., Ed. X, 1758, 280, Carolinas; after Catesby.
Salpa purpurascens variegata (Lane snapper) Catesby, Nat. Hist. Carolina, II, 1743, 17, pl. 17, Bahamas
Sparus vermicularis Bloch and Schneider, Syst. Ichth., 1801, 275, Martinique, on a drawing by Plumier.
?Lutjanus aubrieti Desmarest, Prem. Dec. Ichth., 1823, 17, pl. 2, Cuba.
Mesoprion uninotatus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 339 (449), pl. 39, San Domingo; Martinique.

2524. *Lutianus ambiguus* (Poey).

Cuba.

Mesoprion ambiguus Poey, Memorias, II, 1861, 152, pl. 12, fig. 4; pl. 13, fig. 8; Habana; two specimens known; suspected to be a hybrid of *Lutianus synagris* and *Ocyurus chrysurus*)

2525. *Lutianus mahogoni* (Cuvier and Valenciennes). *Ojanco, Mahogany snapper.*

West Indies; Habana.

Mesoprion mahogoni Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 338 (447), Martinique.

Mesoprion ricardi Cuvier and Valenciennes, loc. cit., 339 (447), Martinique? No locality given.

Mesoprion ojanco Poey, Memorias, II, 1861, 150, pl. 13, fig. 10, Cuba.

2526. *Lutianus megalophthalmus* (Evermann and Marsh).

Porto Rico.

Neomænis megalophthalmus Evermann and Marsh, Bull. U. S. Fish Com., XX, 1900 (1902), 177, Puerto Real, Porto Rico.

Genus 865. RAIZERO Jordan and Fesler.

Raizero Jordan and Fesler, Rept. U. S. Fish. Com., XVII, 1889 (1893), 438 (*Mesoprion aratus* Günther).

2527. *Raizero aratus* (Günther). *Pargo de raizero.*

Pacific coast of tropical America

Mesoprion aratus Günther, Proc. Zool. Soc. Lond., 1864, 145, Panama; Chiapam.

Genus 866. RABIRUBIA Jordan and Fesler.

Rabirubia Jordan and Fesler, Rept. U. S. Fish. Com., XVII, 1889 (1893), 438 (*Mesoprion inermis* Peters).

2528. *Rabirubia inermis* (Peters).

Pacific coast of America from Mazatlan to Panama.

Mesoprion inermis Peters, Berliner Monatsber., 1869, 705, Mazatlan, Mexico.

Genus 867. OCYURUS Gill. *Rabirubias*.

Ocyurus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 236 (*Sparus chrysurus* Bloch).

2529. *Ocyurus chrysurus* (Bloch). *Yellowtail; Rabirubia.*

West Indies, southern Florida to Brazil; single specimen recorded at Woods Hole, Mass.

Sparus chrysurus Bloch, Nat. Ausl. Fische, V, 1791, 28, Brazil; after Marcgrave.

Acara pilamba Marcgrave, Hist. Brasil, 1648, 155, Brazil.

Rabirubia Parra, Descr. Dif. Piezas, Hist. Nat., 1787, 42, pl. 20, fig. 1, Cuba.

Anthias rabirubia Bloch and Schneider, Syst. Ichth., 1801, 309, Cuba; after Parra.

Sparus semiluna Lacépède, Hist. Nat. Poiss., IV, 1802, 141, Martinique.

Mesoprion aurovittatus Agassiz, Spix, Pisc. Brasil., 1829, 121, pl. 66, Brazil

Ocyurus riggersmai Cope, Trans. Amer. Phil. Soc., XIV, 1871, 468, fig. 4, St. Kitts.

Genus 868. RHOMBOPLITES Gill.

Rhomboplites Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 237 (*Centropomus aurorubens* Cuvier and Valenciennes).

2530. *Rhomboplites aurorubens* (Cuvier and Valenciennes). *Cagon de lo alto*.

West Indies, north to Charleston, south to Rio Janeiro.

Centropristis aurorubens Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 34 (45), Brazil, Martinique, and San Domingo.

Mesopriion elegans Poey, Memorias, II, 1861, 153, Cuba.

Aprion ariommus Jordan and Gilbert, Proc. U. S. Nat. Mus., VI, 1883, 142, Pensacola, Fla.

Genus 869. *TROPIDINIUS* Gill.

Tropidinus Gill, in Poey, Synopsis, 1868, 296 (*Mesopriion arnullo* Poey = *Apsilus dentatus* Guichenot).

2531. *Tropidinius dentatus* (Guichenot). *Arnullo*.

West Indies; Habana.

Apsilus dentatus Guichenot, in Ramon de la Sagra, Hist. Cuba, Poiss., 1853, 29, pl. 1, fig. 2, Habana.

Mesopriion arnullo Poey, Memorias, II, 1861, 154, Cuba.

Genus 870. *PRISTIPOMOIDES* Bleeker.

Pristipomoides Bleeker, Tijdschr. Ned. Ind., III, 1852, 574 (*Pristipomoides typus* Bleeker).

Platyinnus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 237 (*Mesopriion vorax* Poey = *Centropristis macrophthalmus* Muller and Troschel).

Bowersia Jordan and Evermann, Bull. U. S. Fish Com., XXII, 1902 (1903), 182 (*B. violescens* Jordan and Evermann).

Subgenus *PLATYINIUS* Gill.

2532. *Pristipomoides macrophthalmus* (Müller and Troschel). *Vorax*

West Indies, Habana.

Centropristis macrophthalmus Müller and Troschel, in Schomburgk, Hist. Barbados, 1848, 666, Barbados.

Mesopriion vorax Poey, Memorias, II, 1861, 151, Cuba.

Genus 871. *ETELIS* Cuvier and Valenciennes.

Etelis Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 93 (127) (*Etelis carbunculus* Cuvier and Valenciennes, an East Indian species).

Elastoma Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 202 (*Etelis oculatus* Cuvier and Valenciennes).

Hesperanthias Lowe, Fishes of Madeira, 1843, 14 (*Serranus oculatus* Cuvier and Valenciennes).

Macrops Duméril, Mém. Acad. Sci. Paris, XXVII, 1856, 279 (*Serranus aculeatus* (oculatus) Cuvier and Valenciennes).

2533. *Etelis oculatus* ⁹⁰ (Cuvier and Valenciennes). *Cachucho*.

Warm seas; West Indies to Madeira; Hawaii.

Etelis carbunculus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 94 (127), pl. 18, Seychelles Archipelago.

Serranus oculatus Cuvier and Valenciennes, loc. cit., II, 198 (266), pl. 32, Martinique.

?*Etelis evurus* Jordan and Evermann, Bull. U. S. Fish Com., XXII, 1902, 142, Hawaii.

Genus 872. *ETELIDES* Jordan and Thompson.

Etelides Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904, 241 (*Anthias aquilonaris* Goode and Bean).

2534. *Etelides aquilonaris* (Goode and Bean).

Gulf of Mexico; Gulf Stream near Carolina coast.

Anthias aquilonaris Goode and Bean, Oceanic Ichth., 1895, 238, Gulf of Mexico.

Family 204. *VERILIDÆ*

Genus 873. *VERILUS* Poey.

Verilus Poey, Memorias, II, 1861, 124 (*V. sordidus* Poey).

2535. *Verilus sordidus* Poey. *Escolar chino*.

West Indies.

Verilus sordidus Poey, Memorias, II, 1861, 125, pl. 12, fig. 6, Cuba.

⁹⁰ This widely distributed and brilliantly colored fish seems to be of a single species throughout the warm seas. We let the American species stand, pending investigation, though we can not detect any difference from the common *evurus*, Hawaii, and *carbunculus* of Japan.

Family 205. HÆMULIDÆ

Genus 874. HÆMULON Cuvier. *Roncos* or *grunts*.

- Diabasis* Desmarest, Prem. Décade Ichth., 1823, 34 (*D. parra* Desmarest; name preoccupied in beetles).
Hæmulon Cuvier, Règne Anim., ed. 2, II, 1829, 175 (*elegans*, etc., restricted later to *Hæmulon elegans* Cuvier = *Sparus sciurus* Shaw).
Anarmostus Scudder, in Putnam, Bull. Mus. Comp. Zool., I, 1863, 12; name preoccupied in insects.
2536. *Hæmulon sexfaciatum* Gill. *Mojarra almejero*.
 Pacific coast of tropical America, Guaymas to Panama.
Hæmulon sexfasciatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 254, Cape San Lucas.
Hæmulon maculosum Peters, Berliner Monatsber., 1869, 705, Mazatlan.
2537. *Hæmulon album* Cuvier and Valenciennes. *Margatefish*; *Jallao*; *Margaret grunt*.
 West Indies; Florida Keys to Brazil.
Perca marina gibbosa cinerea (margatefish) Catesby, Nat. Hist. Carolina, II, 1743, 2, pl. 2, Bahamas.
Perca gibbosa Walbaum, Artedi Pisc., 1792, 342, not of Linnæus, after Catesby.
Hæmulon album Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 179 (241), St. Thomas.
Hæmulon macrophthalmum Gunther, Cat., I, 1859, 306, America.
2538. *Hæmulon mowbrayi* Jordan and Evermann.
 Key West, Fla.
Hæmulon chrysopterum Mowbray, N. Y. Zool. Soc. Bull., XVIII, No. 6, 1915, 1298, with fig., Key West; not of Cuvier, which is *Bathystoma rimator* Jordan and Swain.
Hæmulon mowbrayi Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 505.
2539. *Hæmulon macrostomum* Günther. *Gray grunt*; *Striped grunt*.
 West Indies, north to Florida Keys.
Hæmulon macrostoma Gunther, Cat., I, 1859, 308, Jamaica.
Hæmulon fremebundum Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 340, Clearwater Harbor, Fla.
2540. *Hæmulon bonariense* Cuvier and Valenciennes. *Black grunt*; *Ponce prieto*.
 West Indies, south to Buenos Aires.
Hæmulon bonariense Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 174 (254), Buenos Aires.
Hæmulon canna Cuvier and Valenciennes, loc. cit., V, 173 (233), Martinique.
Hæmulon notatum Poey, Memorias, II, 1861, 179, Cuba.
Hæmulon retrocurrans Poey, Repertorio, II, 1868, 236, 461, Cuba.
Hæmulon continuum Poey, Enumeratio, 1875, 46, Cuba.
2541. *Hæmulon parra* (Desmarest). *Sailors' choice*; *Ronco blanco*; *Ronco prieto*; *Bastard margaret*.
 West Indies; southern Florida to Brazil; very common at Key West and Habana.
Diabasis parra Desmarest, Prem. Décade Ichth., 1823, 30, pl. 2, fig. 2, Habana.
Hæmulon caudimacula Cuvier, Règne Anim., ed. 2, II, 1829, 176, Brazil; Habana.
Hæmulon chromis Broussonet, in Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 180 (242), Jamaica.
Hæmulon acutum Poey, Memorias, II, 1861, 180, Cuba.
Hæmulon serratum Poey, loc. cit., 181, Cuba.
Hæmulon albidum Poey, loc. cit., 181, Cuba.
Hæmulon canna Agassiz in Spix, Pisc. Brasil., 1829, 130, pl. 69, Brazil; not of Cuvier and Valenciennes.

2542. *Hæmulon scudderi* Gill *Mojarra prieta*.
Pacific coast of tropical America, Guaymas to Panama
Hæmulon scudderi Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 253,
Cape San Lucas.
Hæmulon brevirostrum Gunther, Trans. Zool. Soc. Lond., VI, 1868, 418
(Nov. 27), Panama.
Hæmulon undecimale Steindachner, Ichth. Beitr., III, 1875, 11, Panama.
2543. *Hæmulon carbonarium* Poey *Ronco carbonero; Cæsar grunt*.
West Indies and Bermudas, south to Brazil; Habana.
Hæmulon carbonarium Poey, Memorias, II, 1861, 176, Cuba.
2544. *Hæmulon steindachneri* (Jordan and Gilbert) *Roncador raído*.
Both coasts of tropical America; Guaymas to Panama; a similar
species (unnamed) recorded from St. Lucia and Rio Janeiro.
Diabasis steindachneri Jordan and Gilbert, Bull. U. S. Fish Com., I,
1881 (1882), 322, Panama and Mazatlan.
2545. *Hæmulon melanurum* (Linnaeus). *Jeniguana; Cæsar grunt*.
West Indies; Habana and southward; Bush Key; Tortugas
Perca marina cauda nigra (blacktail) Catesby, Hist. Carolina, II, 1743,
7, pl. 7, fig. 2, Bahamas.
Perca melanura Linnaeus, Syst. Nat., ed. X, 1758, 292, and ed. XII,
1766, 486, Bahamas; based on Catesby.
Hæmulon schrankii Agassiz in Spix, Pisc. Brasil, 1829, 131, pl. 69a,
Brazil.
Hæmulon dorsale Poey, Memorias, II, 1861, 179, Cuba.
2546. *Hæmulon sciurus* (Shaw). *Yellow grunt; Ronco amarillo; Boar
grunt; Humpback grunt*.
West Indies; Florida Keys to Brazil.
Anthias formosus Bloch, Nat. Ausl. Fische, VI, 1792, 122, Antilles;
not *Perca formosa* of Linnaeus as identified.
Sparus sciurus Shaw, General Zoology, IV, 1803, 439, pl. 64, Antilles;
based on the description and figure of Bloch.
? *Diabasis obliquatus* Bennett, Zool. Jour. London, V, 1837, 90, Jamaica.
? *Hæmulon sumilis* Castelnau, Anim. Nouv. Rares, II, 1855, Bahia.
Hæmulon luteum Poey, Memorias, II, 1861, 174, Cuba.
Hæmulon multilineatum Poey, loc. cit., 178, Cuba.
? *Hæmulon hians* Haly, Ann. Nat. Hist., XV, 1875, 268, Aspinwall.
2547. *Hæmulon plumieri* (Lacépède). *Common grunt; Ronco ronco; Ronco
arará; Boar grunt; Squirrelfish; Ronco grande*.
West Indies; abundant from Cape Hatteras to Rio Janeiro.
Perca marina capite striato Catesby, Nat. Hist. Carolina, II, 1743, 6,
pl. 6, Bahamas.
Labrus plumieri Lacépède, Hist. Nat. Poiss., III, 1802, 480, pl. 2, fig. 2,
Martinique; on a copy of a drawing by Plumier, identified with this
species by Cuvier.
Guabicoara brasiliensibus Marcgrave, Hist. Brasil., 1648, 163, Brazil.
Hæmulon formosum Cuvier, Règne Anim., ed. 2, II, 1829, 175, Mar-
tinique.
Hæmulon arcuatum Cuvier and Valenciennes, Hist. Nat. Poiss., IX,
1833, 356 (481), Charleston, S. C.
Hæmulon arara Poey, Memorias, II, 1861, 177, Cuba.
Hæmulon subarcuatum Poey, loc. cit., 419, Cuba.
2548. *Hæmulon flavolineatum* (Desmarest). *French grunt; Open-mouthed
grunt; Ronco condenado*.
West Indies; Florida Keys and Bermudas to Brazil.
Diabasis flavolineatus Desmarest, Prem. Décade Ichth., 1823, 35, pl. 2,
fig. 1, Cuba.
Hæmulon heterodon Cuvier, Règne Anim., ed. 2, II, 1829, 176, Cuba,
based on Desmarest.
Hæmulon xanihoipteron Cuvier and Valenciennes, Hist. Nat. Poiss., V,
1830, 174 (254), Martinique.

Genus 875. BRACHYGENYS Scudder.

Brachygenys Scudder, in Poey, Enumeratio, 1875, 47 (*Hæmulon tænia-
tum* Poey = *Hæmulon chrysargyreum* Günther).

2549. *Brachygenys chrysargyreus* (Günther).
Key West, Habana, and Trinidad.
Hæmulon chrysargyreum Günther, Cat., 1859, 314, Trinidad.
Hæmulon læniatum Poey, Memorias, II, 1861, 182, Cuba.
- Genus 876. *BATHYSTOMA* Scudder. *Tomtates*.
Bathystoma Scudder, in Putnam, Bull. Mus. Comp. Zool., I, 1863, 12;
no definition. (*Hæmulon jeniguano* Poey = *Hæmulon aurolineatum*
Cuvier and Valenciennes); not *Bathystomus* Förster, 1862.
2550. *Bathystoma rimator* (Jordan and Swain). *Tomtate*; *Redmouth grunt*;
Cæsar; *Young grunt*.
West Indies; Cape Hatteras to Trinidad.
Hæmulon rimator Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884,
308, Charleston; Key West; Pensacola.
2551. *Bathystoma aurolineatum* (Cuvier and Valenciennes). *Jeniguano*;
Tomtate.
West Indies; Florida Keys to Brazil.
Hæmulon aurolineatum Cuvier and Valenciennes, Hist. Nat. Poiss., V,
1830, 176 (237), Brazil; San Domingo.
Hæmulon jeniguano Poey, Memorias, II, 1861, 183, Cuba.
2552. *Bathystoma striatum* (Linnæus) *White grunt*.
West Indies; Bermudas to Brazil.
Capeuna brasiliensis Marcgrave, Hist. Brasil., 1648, 155, Brazil
(nonbinomial).
Perca striata Linnæus, Syst. Nat., Ed. X, 1758, 293, North America.
Grammistes truttatus Bloch and Schneider, Syst. Ichth., 1801, 188,
Brazil, after Marcgrave.
Serranus capuena Lichtenstein, Abhandl. Berlin. Akad., 1820, 288,
Brazil; after Marcgrave.
Hæmulon quadrilineatum Cuvier and Valenciennes, Hist. Nat. Poiss.,
V, 1830, 177 (238), pl. 120, San Domingo.
Hæmulon quinquelineatum Poey, Memorias, II, 1861, 419, Cuba.
- Genus 877. *LYTHRULON* Jordan and Swain.
Lythrulon Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 287
(*Hæmulon flaviguttatus* Gill).
2553. *Lythrulon flaviguttatum* (Gill). *Peixe fonda*.
Pacific coast of tropical America, Guaymas to Panama.
Hæmulon flaviguttatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
254, Cape San Lucas, Lower California.
Hæmulon margaritifera Günther, Proc. Zool. Soc. Lond., 1864, 147,
Panama.
2554. *Lythrulon opalescens* Jordan and Starks.
Mazatlan, Mexico.
Lythrulon opalescens Jordan and Starks, in Jordan, Proc. Calif. Acad.
Sci., V, 1895, 459, pl. 40, Mazatlan, Mexico.
- Genus 878. *ORTHOSTÆCHUS* Gill. *Striped grunts*.
Orthostæchus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 255 (*O.*
maculicauda Gill).
2555. *Orthostæchus maculicauda* Gill. *Roncador raro*.
Pacific coast of tropical America; Guaymas to Panama.
Orthostæchus maculicauda Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
225, Cape San Lucas, Lower California.
Hæmulon mazatlanum Steindachner, Ichth. Notizen, VIII, 1869, 12,
Pl. VI, Mazatlan.
- Genus 879. *PARACONODON* Bleeker.
Paraconodon Bleeker, Archiv. Neerl., XI, 1876, 272 (*Conodon pacifici*
Günther).
2556. *Paraconodon pacifici* (Günther).
Pacific coast of tropical America.
Conodon pacifici Günther, Proc. Zool. Soc. Lond., 1864, 147, Chiapam.
2557. *Paraconodon cæsius* (Jordan and Gilbert).
Pacific coast of Mexico; Mazatlan and Acapulco; Bay of Panama.
Pomadasy cæsius Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
383, Mazatlan.

2558. *Paraconodon dovii* (Gunther).
Pacific coast of tropical America at Mazatlan and Panama.
Pristipoma dovii Gunther, Proc. Zool. Soc. Lond., 1864, 23, Panama.
- Genus 880. ANISOTREMUS Gill.
Anisotremus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 105 (*Sparus virginicus* Linnæus).
Gentyremus Gill, loc. cit., XIV, 1862, 256 (*Pristipoma bilineatum* Cuvier and Valenciennes = *Anisotremus surinamensis* Bloch).
- Subgenus GENTYREMUS Gill.
2559. *Anisotremus surinamensis* (Bloch). *Pompon*.
Tropical America, from Louisiana (New Orleans market) to Brazil.
?Lutjanus surinamensis Bloch, Naturgesch. Ausl. Fische, IV, 1790, 98, Surinam.
Pristipoma bilineatum Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 202 (271), pl. 122, Martinique.
Pristipoma melanopteryum Cuvier and Valenciennes, loc. cit., 203 (273), Brazil.
Hæmulon obtusum Poey, Memorias, II, 1861, 182, Habana.
Hæmulon labridum Poey, loc. cit., 1861, 419, Cuba.
2560. *Anisotremus interruptus* (Gill). *Mojarron*.
Pacific coast, Magdalena Bay to Panama and the Galapagos.
Gentyremus interruptus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1861, 256, Cape San Lucas, Lower California.
Pristipoma furthii Steindachner, Ichth. Beitr., V, 1876, 4, Panama.
2561. *Anisotremus bicolor* (Castelnau). *Maria prieta*.
Coast of Brazil; perhaps Cuba.
Pristipoma bicolor Castelnau, Anim. Amér. du Sud, Poiss., 1855, 8, pl. 2, fig. 2, Bahia, Brazil.
?Pristipoma trilineatum Poey, Memorias, II, 1861, 343, Habana, young.
Pristipoma brasiliense Steindachner, Sitzungs. k. Akad. Wiss. Wien, 1863, 1013, Bahia.
- Subgenus ANISOTREMUS Gill.
2562. *Anisotremus scapularis* (Tschudi).
Coast of Peru; said to have been once taken at Mazatlan, a doubtful record.
Pristipoma scapulare Tschudi, Fauna Peruana, 12, 1845, Huacho, Peru.
?Diagramma melanospilum Kner, Sitzungs. k. Akad. Wiss., 1807, 4, west coast of South America.
Pristipoma notatum Peters, Monatsb. Akad. Wiss. Berlin, 1869, 706, "angeblich aus Mazatlan."
2563. *Anisotremus davidsonii* (Steindachner). *Sargo ranado*.
Coast of southern California.
Pristipoma davidsonii Steindachner, Ichth. Beitr., III, 1875, 6, San Diego.
2564. *Anisotremus spleniatum* (Poey).
Habana.
Pristipoma spleniatum Poey, Memorias, II, 1861, 187, Habana.
2565. *Anisotremus tæniatus* Gill. *Catalina*; *Striped grunt*.
Pacific coast of tropical America; Magdalena Bay to Panama.
Anisotremus tæniatus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 107, Panama.
2566. *Anisotremus virginicus* (Linnæus). *Porkfish*; *Catalineta*.
West Indies; Florida Keys to Brazil.
Guatucupa juba Marcgrave, Hist. Brasil., 1648, 147, Brazil.
Acara pinima Marcgrave, loc. cit., 1648, 152, Brazil.
Sparus virginicus Linnæus, Syst. Nat., ed. X, 1758, 281, South America.
?Sparus vittatus Bloch, Naturgesch. Ausl. Fische, V, 1791, 33, fig. 2, Brazil; after *Acara pinima* Marcgrave.
Perca juba Bloch, loc. cit., VI, 1792, 77, Brazil, after Marcgrave.
Grammistes mauritii Bloch and Schneider, Syst. Ichthyol., 1801, 185, after Bloch.

- ?*Pristipoma catharinæ* Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 201, 269, St. Catharine Island, Brazil.
Pristipoma acara-pinna Castelnau, Anim. Nouv. Rares, 1856, 8, Brazil.
Pristipoma rodo Cuvier and Valenciennes, loc. cit., 205 (274), Martinique, Brazil, Porto Rico, and San Domingo.
2567. *Anisotremus serrula* (Cuvier and Valenciennes). *Tête de roche*; *Petite scie*.
 Martinique.
Pristipoma serrula Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 203 (272), Martinique.
Pristipoma auratum Cuvier and Valenciennes, loc. cit., 203 (272), Martinique.
- Genus 881. **CONODON** Cuvier and Valenciennes.
Conodon Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 156 (*C. antillanus* Cuvier and Valenciennes = *Perca nobilis* Linnæus).
Cheloniger Plumier, in Lacépède, Hist. Nat. Poiss., V, 1803 (*Perca nobilis* Linnæus); name not accepted by Lacépède.
2568. *Conodon nobilis* (Linnæus).
 West Indies; coast of Texas to Brazil.
Perca nobilis Linnæus, Syst. Nat., Ed. X, 1758, 291, North America.
Scæna plumieri Bloch, Nat. Ausl. Fische, VI, 1792, 66, taf. 306, Martinique.
Scæna coro Bloch, loc. cit., 1792, 70, pl. 307, fig. 2, Brazil; after *Coro coro* Marcgrave.
Conodon antillanus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 116 (156), Jamaica.
2569. *Conodon serrifer* Jordan and Gilbert.
 Lower California.
Conodon serrifer Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 351, Boca Soledad, Lower California.
- Genus 882. **BRACHYDEUTERUS** Gill. *Burrilos*.
Brachydeuterus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 17 (*B. auratus*, an African species).
Hæmulopsis Steindachner, Ichth. Notizen, VIII, 1869, 9, Mazatlan (*Hæmulon corvinæformis* Steindachner).
2570. *Brachydeuterus nitidus* (Steindachner).
 Pacific coast of tropical America; Panama, Mazatlan, and the Gulf of California.
Pristipoma (*Hæmulopsis*) *nitidum* Steindachner, Ichth. Notizen, VIII, 1869, 5, Mazatlan.
2571. *Brachydeuterus leuciscus* (Gunther). *Burrito*.
 Pacific coast of tropical America; not rare at Mazatlan and Panama; south to northern Peru.
Pristipoma leuciscus Gunther, Proc. Zool. Soc. Lond., 1864, 147, San Jose de Nicaragua; Chiapam.
Pomadasis elongatus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 352, Mazatlan, Mexico.
2572. *Brachydeuterus axillaris* (Steindachner). *Burro blanco*.
 Pacific coast of Mexico; Mazatlan and Guaymas
Pristipoma (*Hæmulopsis*) *axillare* Steindachner, Ichth. Notizen, VIII, 1869, 7, Mazatlan.
2573. *Brachydeuterus corvinæformis* (Steindachner).
 West Indies to Brazil; Jamaica.
Hæmulon corvinæforme Steindachner, Ichth. Notizen, VII, 1868, 16, Santos, Brazil.
- Genus 883. **RHENCUS** Jordan and Evermann.
Rhencus Jordan and Evermann, Check List, 1895, 387 (*Pristipoma panamense* Steindachner).

2574. *Rhencus panamensis* (Steindachner).
Pacific coast of tropical America; Panama and Mazatlan.
Pristipoma panamense Steindachner, Ichth. Beitr., III, 1875, 8, Panama.
- Genus 884. **POMADASYS** Lacépède. *Burros*.
Pomadasys Lacépède, Hist. Nat. Poiss., IV, 1803, 516 (*Sciæna argentea* Forskål).
Pristipomus Oken, Isis, 1817, 279 (*Lutjanus hasta* Bloch; after *les Pristipomes* Cuvier).
Pristipoma Cuvier and Valenciennes, changed spelling.
- Subgenus **PRISTOPOMUS** Cuvier.
2575. *Pomadasys bayanus* Jordan and Evermann.
Pacific coast of Panama, in fresh water.
Pristipoma humile Kner and Steindachner, Sitzgber. Akad. Wiss. Munch., 1863, 222, Rio Bayano, near Panama; name preoccupied by *Pristipoma humilis* Bowdich, Fishes of Madeira, 1825, 236.
Pomadasys bayanus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1331, Rio Bayano, near Panama.
2576. *Pomadasys productus* (Poey).
Cuba
Pristipoma productum Poey, Memorias, II, 1861, 186, Habana.
2577. *Pomadasys macracanthus* (Günther). *Burro*.
Pacific coast of tropical America; Panama; Mazatlan; Chiapas, and Punta Arenas.
Pristipoma macracanthum Günther, Proc. Zool. Soc. Lond., 1864, 146, Chiapam.
2578. *Pomadasys grandis* Meek.
Lake Nicaragua
Pomadasys grandis Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 116, Lake Nicaragua.
2579. *Pomadasys templei* Meek.
Valles, San Luis Potosí, Mexico.
Pomadasys templei Meek, Field Col. Mus., Zool., V, 1904, 201, Valles, San Luis Potosí, Mexico.
2580. *Pomadasys boschmæ* (Metzelaar).
West Indies.
Pristipoma boschmæ Metzelaar, Tropisch Atl. Vissch., 1919, 83, fig. 27; probably from Curaçao.
2581. *Pomadasys starri* Meek. *Burro*.
Perez, Vera Cruz, Mexico.
Pomadasys starri Meek, Field Col. Mus., Zool., V, 1904, 200, Perez, Vera Cruz, Mexico.
- Genus 885. **RHONCISCUS** Jordan and Evermann.
Rhonciscus Jordan and Evermann, Check List, 1895, 387 (*Pristipoma croco* Cuvier and Valenciennes).
2582. *Rhonciscus croco* (Cuvier and Valenciennes)
West Indies; Cuba to Brazil.
Pristipoma croco Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 197 (264), Martinique.
Pristipoma cultiferum Poey, Memorias, II, 1861, 185, Habana.
Pomadasys approximans Bean and Dresel, Proc. U. S. Nat. Mus., VII, 1884, 160, Jamaica.
2583. *Rhonciscus branicki* (Steindachner). *Burrito*.
Pacific coast of tropical America; Mazatlan to Peru.
Pristipoma branicki Steindachner, Denkschr. K. Akad. Wiss. Wien, XLI, 1879, 28, Tumbes, Peru.
2584. *Rhonciscus ramosus* (Poey).
West Indies; south to Brazil.
Pristipoma ramosum Poey, Memorias, II, 1861, 186, Habana.
Pristipoma boucardi Steindachner, Ichth. Notizen, VIII, 1869, 1, Cojimar River near Habana.

2585. *Rhonciscus sinuosus* (Eigenmann).
North coast of Colombia.
Pomadour sinuosus Eigenmann, Proc. Amer. Phil. Soc., LVI, 1917, 683,
Patia, between Magna and Telembi.
- Genus 886. ORTHOPRISTIS Girard. *Pufffishes*.
Orthoprists Girard, U. S. Mex. Bound. Surv., Zool., 1859, 15 (*O. duplex*
Girard=*Perca chrysopterus* Linnaeus).
Pristocantharus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 256
(*Pristipoma cantharinum* Jenyns).
2586. *Orthoprists chalcus* (Günther).
Pacific coast of tropical America, from Cape San Lucas to the Gala-
pagos.
Pristipoma chalcum Günther, Proc. Zool. Soc. Lond., 1864, 146,
Panama.
Pristipoma kneri Steindachner, Ichth. Notizen, VIII, 1869, 3, Mazatlan,
Mexico.
2587. *Orthoprists reddingi* Jordan and Richardson.
La Paz, Lower California
Orthoprists reddingi Jordan and Richardson, in Jordan, Proc. Calif.
Acad. Sci., V, 1895, 509, pl. 41, La Paz, Lower California.
2588. *Orthoprists chrysopterus* (Linnaeus). *Pufffish*; *Sailors' choice*, *Hogfish*;
Red-mouthed grunt.
South Atlantic and Gulf coasts of the United States; young common
as far north as New York.
Perca chrysoptera Linnaeus, Syst. Nat., Ed. XII, 1766, 485, Charleston,
S. C.
Labrus fulvo-maculatus Mitchill, Trans. Lit. and Phil. Soc. N. Y.,
1814, 406, New York.
Pristipoma fasciatum Cuvier and Valenciennes, Hist. Nat. Poiss., V,
1830, 213 (285), New York.
Orthoprists duplex Girard, U. S. Mex. Bound. Surv., Zool., Fishes, 1859,
15, pl. 9, figs. 1 to 4, Indianola and Brazos Santiago, Tex.
2589. *Orthoprists poeyi* Scudder.
West Indies; Habana
Orthoprists poeyi Scudder, in Poey, Synopsis, 1868, 312, Habana.
- Genus 887. MICROLEPIDOTUS Gill.
Microlepidotus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 255 (*M.*
inornatus Gill).
Isaciella Jordan and Fesler, Rept. U. S. Fish Com., XVII, 1889-1891
(1893), 497 (*Pristipoma brevipinna* Steindachner).
2590. *Microlepidotus brevipinnis* (Steindachner).
Pacific coast of Mexico.
Pristipoma (Hemulopsis) brevipinna Steindachner, Ichth. Notizen, VIII,
1869, 10, Mazatlan.
2591. *Microlepidotus inornatus* Gill. *Jopaton*.
Gulf of California; La Paz and Mazatlan.
Microlepidotus inornatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
256, Cape San Lucas, Lower California.
- Genus 888. GENYATREMUS Gill.
Genyatremus Gill, Proc., Acad. Nat. Sci. Phila., XIV, 1862, 256 (*Dia-*
gramma cavifrons Cuvier and Valenciennes=*Lutjanus luteus* Bloch).
2592. *Genyatremus luteus* (Bloch).
Lesser Antilles to Brazil.
Lutjanus luteus Bloch, Nat. Ausl. Fische, IV, 1795, 111, Martinique;
on a drawing by Plumier.
Grammistes hepatus Bloch and Schneider, Syst. Ichth., 1801, 187,
after Bloch.
Diagramma cavifrons Cuvier and Valenciennes, Hist. Nat. Poiss., I,
1830, 217 (290), pl. 123, Rio Janeiro.

Family 206. SPARIDÆ. Porgies; Pargos

Genus 889. OTRYNTER Jordan and Evermann.

Otrynter Jordan and Evermann, Check List, 1895, 388 (*Stenotomus caprinus* Bean).2593. *Otrynter caprinus* (Bean).

Banks off the west coast of Florida.

Stenotomus caprinus Bean, Proc. U. S. Nat. Mus., V, 1882, 426, Snapper Banks off Pensacola, Fla.

Genus 890. STENOTOMUS Gill.

Stenotomus Gill, Canadian Nat., 1865, 266 (*Sparus argyrops* Linnæus = *Sparus chrysops* Linnæus).*Stenesthes* Jordan, Copeia, Oct. 4, 1917 (same type); substitute name, there being an earlier genus *Stenotoma*.2594. *Stenotomus versicolor* (Mitchill). *Common scup; Porgy; Scuppaug; Mishcuppaug; Bream; Fair maid.*

Atlantic coast of United States, from Cape Cod to North Carolina; common northward.

Labrus versicolor Mitchill, Trans. Lit. and Phil. Soc., I, 1815, 404, New York.*Sargus ambassis* Gunther, Cat., I, 1859, 449, New York.2595. *Stenotomus chrysops* (Linnæus). *Southern porgy.*

Southern Atlantic and Gulf coast of United States, Cape Hatteras to Texas.

Sparus argyrops Linnæus, Syst. Nat., Ed. XII, 1766, 471, Charleston; young.*Sparus chrysops* Linnæus, loc. cit., 1766, 471, Charleston, S. C.*Sparus xanthurus* Lacépède, Hist. Nat. Poiss., IV, 1803, 120, Charleston; after *argyrops* L.*Chrysophrys aculeata* Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1830, 137, Charleston, S. C.

Genus 891. CALAMUS Swainson.

Calamus Swainson, Nat. Hist. Fishes, etc., II, 1839, 171 and 221 (*Pagellus calamus* Cuvier and Valenciennes).*Grammateus* Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1874 (*Pagellus microps* Guichenot = *Pagellus penna* Cuvier and Valenciennes).2596. *Calamus calamus* (Cuvier and Valenciennes). *Saucereye porgy; Pez de pluma.*

West Indies, north to Florida Keys.

Pagellus calamus Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 152 (206), pl. 152, Martinique; San Domingo.*Calamus megacephalus* Swainson, Nat. Hist. Fishes, II, 1839, 222, after Cuvier and Valenciennes.*Pagellus orbitarius* Poey, Memorias, II, 1861, 201, Cuba.2597. *Calamus bajonado* (Bloch and Schneider). *Jolt-head porgy; Bajonado.*

West Indies, north to Florida Keys.

Sparus bajonado Bloch and Schneider, ed. Bloch, Syst. Ichth., 1801, 284, West Indies; after Parra.*Bajonado* Parra, Dif. Piezas Hist. Nat. Cuba, 1787, 13, lam. 8, Habana.*Pagellus caninus* Poey, Memorias, II, 1861, 199, Habana.*Calamus plumatula* Guichenot, Rev. Pagels in Mém. Soc. Sci. Nat. Cherbourg, XIV, 1868, 119, Martinique.2598. *Calamus proridens* Jordan and Gilbert. *Littlehead porgy; Pez de pluma.*

West Indies, north to the Florida Keys.

Calamus proridens Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 150, Key West, Fla.2599. *Calamus pennatula* Guichenot. *Littlehead porgy.*

West Indies.

Calamus pennatula Guichenot, Révision des Pagels in Mém. Soc. Sci. Nat. Cherbourg, XIV, 1868, 116, Martinique.

2600. *Calamus brachysomus* (Lockington). *Mojarra garabata*.
Gulf of California and neighboring waters, locally abundant from
Magdalena Bay to Panama.
Sparus brachysomus Lockington, Proc. U. S. Nat. Mus., III, 1880, 284,
Magdalena Bay, Lower California.
2601. *Calamus leucosteus* Jordan and Gilbert. *Whitebone porgy*.
South Atlantic coast of United States; known only from the markets
of Charleston, S. C.
Calamus leucosteus Jordan and Gilbert, in Jordan, Cat. Fish. N. Amer.,
in Rept. U. S. Fish Com., XIII, 1885, 879, Charleston, S. C.
2602. *Calamus macrops* Poey.
Cuba.
Calamus macrops Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1872, 181,
Pl. VII, fig. 3, Habana.
2603. *Calamus penna* (Cuvier and Valenciennes). *Littlemouth porgy*; *Sheeps-
head porgy*; *Speckled porgy*.
Southern Florida to Brazil; known from southern Florida, Key West,
Rio Janeiro, St. Thomas, Habana, Camaru, and Rio Grande do Sul.
Pagellus penna Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830,
154 (209), Brazil.
Pagellus microps Guichenot, in Ramon de la Sagra, Hist. Ile Cuba,
1853, 83, pl. 3, fig. 1, Habana.
Pagellus humilis Poey, Synopsis, 1868, 308, Habana.
Pagellus milneri Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 134,
Charlotte Harbor, Fla.
2604. *Calamus arctifrons* Goode and Bean. *Grass porgy*; *Shad porgy*.
Gulf of Mexico, from Pensacola to Key West; West Indies; Porto Rico.
Calamus arctifrons Goode and Bean, Proc. U. S. Nat. Mus., V, 1882,
425, Pensacola, Fla.
2605. *Calamus medius* (Poey).
West Indies; Habana.
Grammateus medius Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1872, 183,
pl. 7, fig. 4, Habana.
2606. *Calamus kendalli* Evermann and Marsh. *Pluma*.
Porto Rico.
Calamus kendalli Evermann and Marsh, Rept. U. S. Fish Com., XXV,
1899, 354, Mayaguez, Porto Rico.
- Genus 892. *PAGRUS* Cuvier.
Pagrus Cuvier, Règne Animal, ed. 1, 1817, 272 (*Sparus argenteus* Bloch
and Schneider = *Sparus pagrus* Linnæus).
2607. *Pagrus pagrus* (Linnæus). *Red porgy*; *Besugo*; *Pargo colorado*.
Southern Europe and south Atlantic and Gulf coasts of the United
States; south to Uruguay; less common in America.
Sparus pagrus Linnæus, Syst. Nat., Ed. X, 1758, 279, southern Europe.
Sparus argenteus Bloch and Schneider, Syst. Ichth., 1801, 271, Europe.
Pagrus vulgaris Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830,
104 (142), pl. 148, southern Europe.
- Genus 893. *LAGODON* Holbrook. *Chopa spina*; *Sailors' choice*.
Lagodon Holbrook, Ichth. South Carolina, 1855, 56 (*Sparus rhomboides*
Linnæus).
2608. *Lagodon rhomboides* (Linnæus). *Pinfish*; *Bream*; *Sailors' choice*;
Chopa spina; *Sargo*.
Atlantic and Gulf coasts of United States; Cape Cod to Cuba.
Sparus rhomboides Linnæus, Syst. Nat., Ed. XII, 1766, 470, Charleston,
S. C.
- Genus 894. *SALEMA* Jordan and Evermann.
Salema ⁶¹ Jordan and Evermann, Check List, 1895, 390 (*Perca uni-
maculata* Bloch).

⁶¹ Not *Salema* Bowdich 1825 = *Kyphosus*.

2609. *Salema rhomboidalis* (Linnæus). *Salema*.
West Indies, north to Key West, south to Rio Janeiro.
Perca rhomboidalis Linnæus, Syst. Nat., Ed. X, 1758, Jamaica; after
Bream, Browne, Jamaica, 1756, 446.
Perca unimaculata Bloch, Nat. Ausl. Fische, VI, 1792, 75, Brazil.
Sparus salin Lacépède, Hist. Nat. Poiss., IV, 1802, 136; based on
Bloch.
Sargus humerimaculatus Quoy and Gaimard, Voy. Freycinet, Zool,
1824, 297, Rio Janeiro.
Sargus flavolineatus Cuvier and Valenciennes, Hist. Nat. Poiss., VI,
1830, 44 (60), Cuba.
Cynædus brama Gronow, Cat. Fishes, ed Gray, 1854, 56, South Carolina.
Sargus caribæus Poey, Memorias, II, 1861, 197, Cuba.
2610. *Salema tridens* (Poey).
Cuba.
Sargus tridens Poey, Enumeratio, 1875, 57, Cuba.
- Genus 895. **ARCHOSARGUS** Gill. *Sheepsheads*.
Archosargus Gill, Canadian Nat., 1865, 266 (*Sparus probatocephalus*
Walbaum).
2611. *Archosargus probatocephalus* (Walbaum). *Sheepshead*; *Sargo raiado*.
Atlantic and Gulf coasts of the United States; Cape Cod to Florida
Keys and Texas.
Sparus probatocephalus Walbaum, Artedi Pisc., 1792, 295, New York;
based on *Sparus* of Schopf.
Sparus ovinephalus Bloch and Schneider, Syst. Ichth., 1801, 280, New
York.
Sargus ovis Mitchell, Trans. Lit. and Phil. Soc. N. Y., I, 1814, 392,
pl. 2, fig. 5, New York.
2612. *Archosargus aries* (Cuvier and Valenciennes).
Honduras to Brazil; Rio Janeiro; Maracaibo; Belize
Sargus aries Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 42
(58), Rio Janeiro; Maracaibo.
- Genus 896. **DIPLODUS** Rafinesque.
Diplodus Rafinesque, Indice, 1810, 54 (*Sparus annularis* L.).
Sargus Cuvier, Règne Anim., I, 1817, 272 (*Sparus sargus* Linnæus);
preoccupied in insects.
2613. *Diplodus holbrookii* (Bean).
South Atlantic and Gulf coasts of the United States; Cape Hatteras to
Cedar Keys.
Sargus holbrookii Bean, Forest and Stream, June 13, 1878, 359, Charles-
ton, S. C.
2614. *Diplodus argenteus* (Cuvier and Valenciennes). *Sargo*.
West Indies; Florida and Bermudas, south to Argentina.
Sargus argenteus Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830,
44 (60), Brazil.
Sargus caudimacula Poey, Memorias, II, 1861, 198, Cuba.
2615. *Diplodus sargus* (Linnæus). *Sargo*.
Coasts of southern Europe; Bermudas.
Sargus sargus Linnæus, Syst. Nat., Ed. X, 1758, 278, Mediterranean.
Sargus variegatus Lacépède, Hist. Nat. Poiss., IV, 1803, 207, Medi-
terranean.
Sargus raucus Geoffrey St. Hilaire, Descr. de l'Égypte, Poiss., 1813,
Pl. XVIII, fig. 1, coast of Egypt.
Sargus rondeletii Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830,
10 (14), pl. 141, Mediterranean.
Sargus vitula Cuvier and Valenciennes, loc. cit., 35 (48), Naples.

Family 207. GIRELLIDÆ

Genus 897. **GIRELLA** ⁶² Gray.

- Girella* Gray, Illus. Indian Zool., II, 1833, 98 (*G. punctata* Gray).
Camarna Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860 (1863), 81 (*C.*
nigricans Ayres).

⁶² *Melanichthys* Schlegel, 1844 = *Incisidens* Gill, 1862; differs from *Girella* in having two to four rows of
incisors and is probably a valid genus.

2616. *Girella nigricans* (Ayres). *Greenfish; California bluefish; White spot.*
Coast of southern California from Monterey to Cape San Lucas.
Camarina nigricans Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 81, fig. 22, California.
Girella dorsomacula Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 244, Cape San Lucas.
- Genus 898. **INCISIDENS** Gill.
Incidens Gill, Proc. Acad. Sci. Phila., XIV, 1862, 244 (*Crenidens simplex* ⁶³ Richardson).
Aplodon Duméril, in Thominot, Bull. Soc. Philom. Paris, 1883, 141 (*Aplodon margaritifera* Duméril); name preoccupied.
2617. *Incidens simplicidens* (Osburn and Nichols).
Gulf of California.
Girella simplicidens Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 165, fig. 10, San Francisco.

Family 208. **KYPHOSIDÆ**. Pilotfishes

- Genus 899. **KYPHOSUS** Lacépède.
Kyphosus Lacépède, Hist. Nat. Poiss., III, 1802, 114 (*K. bigibbus* Lacépède=*fuscus* Lacépède).
Pimelepterus Lacépède, Hist. Nat. Poiss., IV, 1803, 429 (*P. bosci* Lacépède).
Dorsuarius Lacépède, loc. cit., V, 1803, 482 (*D. nigrescens* Lacépède).
Xyster Lacépède, loc. cit., X, 484 (*X. fuscus* Lacépède).
Xysterus Rafinesque, Anal. Nature, 1815, 95; substitute for *Xyster*.
Selerma Bowdich, Fishes Madeira, 1825, 238 (*S. aurata* Bowdich).
Opisthistrus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 245 (*Sciæna tahmel* Forskål).
Cyphosus of authors, corrected spelling.
2618. *Kyphosus analogus* (Gill). *Salema*.
Pacific coast of tropical America, Gulf of California to Panama.
Pimelepterus analogus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 245, Cape San Lucas.
2619. *Kyphosus incisor* (Cuvier and Valenciennes). *Chopa amarilla*.
West Indies to Brazil and the Canaries; Tortugas.
Pimelepterus incisor Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 198 (266), Brazil.
? *Salema aurata* Bowdich, Excursion Madeira, 1825, 238, Bona Vista Island.
Pimelepterus flavo-lineatus Poey, Repertorio, I, 1866, 319, Habana.
2620. *Kyphosus elegans* (Peters). *Chopa*.
Pacific coast of tropical America, from Guaymas to Mazatlan; a species apparently the same in Hawaii.
Pimelepterus elegans Peters, Berlin. Monatsb., 1869, 707, Mazatlan, Mexico.
Pimelepterus sandvicensis Sauvage, Bull. Soc. Philom., July, 1880 (7), IV, 221, Sandwich Islands.
2621. *Kyphosus metzelaari* Jordan and Evermann.
Caribbean Sea.
Cyphosus elegans Metzelaar, Trop. Atl. Vissch., 1919, 44, Curaçao; Venezuela; not *Pimelepterus elegans* Peters.
This species, according to Metzelaar's account, is extremely close to *Kyphosus elegans* of the Pacific, but the scales are larger; 52 in the lateral line (67 in *elegans*), the teeth are also fewer, "15 in upper jaw"; in *elegans* they are "86 in each jaw."
Kyphosus metzelaari Jordan and Evermann, Proc., Calif. Acad. Sci., 4th Ser., XVI, Apr. 27, 1927, 506, Curaçao, Venezuela.

⁶³ This species with entire teeth is said to be the female of *Girella tricuspidata*. If this is true the genus *Incidens* may not be tenable.

- 2622. *Kyphosus sectatrix* (Linnæus).** *Rudderfish; Bermuda chub; Chub; Chopá blanca.*
West Indies, ranging from Cape Cod to Brazil, crossing the ocean to the Canary Islands; accidental in the Mediterranean; one taken at Palermo.
Perca marina sectatrix Catesby, Nat. Hist. Carolina, II, 1743, 8, Carolinas.
Perca sectatrix Linnæus, Syst. Nat., Ed. XII, 1766, 486 (correction), misprinted for *saltatrix*, after Catesby.
Chætodon cyprinaceus Gmelin, Syst. Nat., I, 1789, 1269, tropical America.
Pimelepterus bosqui Lacépède, Hist. Nat. Poiss., IV, 1803, 429, South Carolina.
Pimelepterus oblongior Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 197 (264), locality unknown.
Pimelepterus bosci sacula Doderlein, Nat. Siciliano, Giornale Scienze Nat., III, 1883, 83, Gulf of Palermo.
- 2623. *Kyphosus lutescens* (Jordan and Gilbert).**
Revillagigedo Archipelago.
Pimelepterus lutescens Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 229, Braithwaite Bay, Socorro Island.
- Genus 900. SECTATOR** Jordan and Fesler.
Sectator Jordan and Fesler, Rept. U. S. Fish Com., XVII, 1889-91, 534 (*Pimelepterus ocyurus* Jordan and Gilbert).
- 2624. *Sectator ocyurus* (Jordan and Gilbert).**
Panama.
Pimelepterus ocyurus Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 327, 328, Bay of Panama.
- Genus 901. HERMOSILLA** Jenkins and Evermann.
Hermosilla Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 144 (*H. azurea* Jenkins and Evermann).
- 2625. *Hermosilla azurea* Jenkins and Evermann.**
Gulf of California.
Hermosilla azurea Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 144, Guaymas, Mexico.
- 2626. *Hermosilla robusta* Osburn and Nichols.**
Gulf of California.
Hermosilla robusta Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 166, fig. 11, Tiburon Island.

Family 209. INERMIDÆ

- Genus 902. INERMIA** Poey.
Inermia Poey, Memorias, II, 1861, 193 (*I. vittata* Poey).
- 2627. *Inermia vittata* (Poey).** *Boga.*
Caribbean Sea, north to Cuba; often in large schools, off Venezuela.
Inermia vittata Poey, Memorias, II, 1861, 193, Habana.

Family 210. MÆNIDÆ

- Genus 903. SPICARA** Rafinesque.
Spicara Rafinesque, Caratteri, 1810, 51 (*S. flexuosa* Rafinesque = *Sparus smar* Linnæus).
Smaris Cuvier, Règne Animal, ed. 1, 1817, 269 (*Sparus smar* Linnæus).
- 2628. *Spicara martinica* (Cuvier and Valenciennes).**
West Indies; probably an error.
Smaris martinicus Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 319 (424), Martinique; most likely an error, type probably from the Mediterranean.

Family 211. GERRIDÆ. Mojarras; Silver perch

- Genus 904. EUCINOSTOMUS** Baird and Girard. *Mojarritas.*
Eucinostomus Baird and Girard, Ninth Smithsonian Rept., 1854, 344 (*E. argenteus* Baird and Girard = *Gerres gula* Cuvier and Valenciennes).

2629. *Eucinostomus gula* (Cuvier and Valenciennes). *Silver jenny*; *Mojarra de ley*; *Petite queue*; *Shad*.
Gerres gula Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 349 (464), Martinique.
Eucinostomus argenteus Baird and Girard, Ninth Smith. Rept., 1854, 335, Bessley Point, N. J.
Eucinostomus gulula Poey, Enumeratio, 1875, 54, pl. 2, Habana.
Diapterus homonymus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 340, Clearwater Harbor, Fla.
2630. *Eucinostomus pseudogula* Poey.
 Tortugas; West Indies to Brazil; Bermudas; St. Lucia; Bahia.
Eucinostomus pseudo-gula Poey, Enumeratio, 1875, 53, pl. 1, Cuba.
Gerres jonesi Günther, Ann. Mag. Nat. Hist., 1879, 150, Bermudas.
Eucinostomus harengulus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 132, west Florida.
2631. *Eucinostomus californiensis* (Gill). *Mojarra cantilena*.
 Pacific coast of Mexico; Guaymas and Cape San Lucas to Panama.
Diapterus californiensis Gill, Proc. Acad. Nat. Sci., Phila., XIV, 1862, 245, Cape San Lucas, Lower California.
Diapterus gracilis Gill, loc. cit., 245, Cape San Lucas.
Diapterus dowii Gill, loc. cit., XV, 1863, 162, Panama.
2632. *Eucinostomus elongatus* Meek and Hildebrand.
 Panama.
Eucinostomus californiensis elongatus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, 586, Sobago Island, Panama.
- Genus 905. *ULÆMA* Jordan and Evermann.
Ulæma Jordan and Evermann, Proc. Calif. Acad. Sci., V, 1895, 471 (*Diapterus lefroyi* Goode).
2633. *Ulæma lefroyi* (Goode).
 West Indies, on sandy shores north to Cedar Keys; Bermudas; Cuba; Key West; recorded north at Beaufort, N. C.
Diapterus lefroyi Goode, Amer. Jour. Sci. Arts, 3d ser., VIII, 1874, 123, Bermudas.
2634. *Ulæma producta* (Poey).
 Cuba.
Eucinostomus productus Poey, Enumeratio, 1875, 55, Habana.
2635. *Ulæma meeki* (Eigenmann).
 Rio San Juan, in western Cuba.
Eucinostomus meeki Eigenmann, Bull. U. S. Fish Com., XXII, 1902, 229, fig. 10, San Juan River, just below its first ford, west Cuba; apparently a *Ulæma* having but two anal spines, the second hæmal spine not described.
- Genus 906. *GERRES* (Cuvier) Quoy and Gaimard. *Mojarras blancas*.
Gerres Cuvier in Quoy and Gaimard, Voyage Uranie, 1824, 293, restricted by Jordan 1919 to *Gerres vaignensis* Quoy and Gaimard = *Sciæna argyrea* Forskål, an ally of *Mugil cinereus* Walbaum. The only other species named in this earlier paper is *Gerres gula*, belonging to *Eucinostomus*. This use of the name *Gerres* from Cuvier's MS. antedates that in the Règne Animal, 1829, and unfortunately must stand.
Podager Gistel, Thierreich, IX, 1848, a substitute for *Gerres* regarded as preoccupied by *Gerris*; *Podager* is actually so, however.
Catochænum Cantor, Malayan Fishes, 1849, 55, same type, being a substitute for *Gerres*, deemed preoccupied by *Gerris*, a genus of insects.
Xystæma Jordan and Evermann, Proc. Calif. Acad. Sci., V, 1895, 471 (*Mugil cinereus* Walbaum).
2636. *Gerres cinereus* (Walbaum). *Mojarra de Casta*; *Mojarra blanca*; *Broad shad*.
 Coasts of tropical America; West Indies, north to Florida; Habana; Jamaica; Martinique; Bahamas; Barbados; Florida Keys.
Turdus cinereus peltatus Catesby, Nat. Hist. Carolinas, II, 1743, 11, Bahamas.
Mugil cinereus Walbaum, Arted. Pisc., 1792, 228, Bahamas; after Catesby

- Gerres aprion* Cuvier, Règne Animal, ed. 2, II, 1829, 188; based on Catesby.
Gerres zebra Muller and Troschel, in Schomburgk's Hist. Barbados, 1848, 668, Barbados.
Gerres squamipinnus Gunther, Cat., I, 1859, 349, Jamaica; Guatemala.
2637. *Gerres simillimus* Regan.
 Pacific coast of tropical America, Mazatlan to Panama.
Gerres simillimus Regan, Biol. Central-Amér., Pisces, 1906, 38, Panama.
2638. *Gerres havana* (Nichols).
 Miami, Florida; Cuba.
Xystæma havana Nichols, Bull. Amer. Mus. Nat. Hist., XXXI, 1912, 189, fig. 2, Marianas Beach, Habana.
- Genus 907. **MOHARRA** Poey.
Moharra Poey, Enumeratio, 1875, 50 (*Gerres rhombeus* Cuvier and Valenciennes).
2639. *Moharra rhombea* (Cuvier and Valenciennes).
 West Indies and Atlantic coast of tropical America; Jamaica; San Domingo; Martinique; Puerto Cabello; Habana; Aspinwall; Río Magdalena; Santa Lucia; Bahia.
Gerres rhombeus Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 345 (459), Martinique; San Domingo.
2640. *Moharra evermanni* (Meek and Hildebrand).
 Atlantic waters of Panama.
Diapterus evermanni Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, 594, Pl. LXII, Mindi River near Mindi, Panama Canal Zone.
- Genus 908. **DIAPTERUS** Ranzani.
Diapterus Ranzani, Nov. Comment. Bonon., V, 1842, 340 (*Diapterus auratus* Ranzani, a species near *D. alsthostomus*).
2641. *Diapterus aureolus* (Jordan and Gilbert).
 Panama; only the type known.
Gerres aureolus Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 328, Panama.
2642. *Diapterus axillaris* (Gunther).
 Pacific coast of Mexico.
Gerres axillaris Günther, Proc. Zool. Soc. Lond., 1864, 152, Chiapas.
2643. *Diapterus peruvianus* (Cuvier and Valenciennes). *Mojarra de las aletas amarillas*.
 West coast of tropical America; Mazatlan; Salina Cruz; Chiapas; Panama; Peru.
Gerres peruvianus Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 351 (467), Payta, northern Peru.
Gerres brevirostris Sauvage, Bull. Soc. Philom. Paris, III, 1879, 208, Río Guayas, Ecuador.
2644. *Diapterus olisthostomus* (Goode and Bean). *Irish pompano*; *Mutton-fish*.
 West Indies, north to southern Florida.
Gerres olisthostoma Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 423, Indian River, Fla.
- Genus 909. **EUGERRES** Jordan and Evermann.
Eugerres Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 506 (*Gerres plumieri* Cuvier and Valenciennes).
2645. *Eugerres lineatus* (Humboldt). *Mojarra china*.
 Pacific coast of Mexico; Acapulco; Mazatlan; San Blas; Chiapas.
Smaris lineatus Humboldt, Observ. Zool., II, 1811, 185, pl. 46, Acapulco.
2646. *Eugerres brasiliensis* (Cuvier and Valenciennes). *Pataó*.
 West Indies to Brazil.
Gerres brasiliensis Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 344, 458, Brazil.
Gerres patao Poey, Synopsis, VI, 1868, 320, Habana; scarcely distinguishable from *G. lineatus* of the Pacific coast.
Gerres embryx Jordan and Starks, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1379, Charleston, S. C.; large example.

2647. *Eugerres brevimanus* (Gunther).
Pacific coast of tropical America and Panama Bay.
Gerres brevimanus Gunther, Proc. Zool. Soc. Lond., 1864, 152, Chiapam.
2648. *Eugerres plumieri* (Cuvier and Valenciennes). *Mojarra*.
Atlantic coast of tropical America and West Indies; Habana; Porto Rico; San Domingo; Jamaica; Pernambuco; Bahia; Aspinwall; Guatemala.
Gerres plumieri Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 340 (452), pl. 467, Antilles; Porto Rico.
2649. *Eugerres mexicanus* (Steindachner). *Mexican mojarra*.
Rio Teapa, Mexico.
Gerres mexicanus Steindachner, Ueber eine Neue Gerres-Art aus Mexico, Verh. K. K. Ges. Wien, XIII, 1863, 383, Rio Teapa, Mexico.

Family 212. MULLIDÆ. Surmulletts

- Genus 910. *MULLUS* Linnæus. *Surmulletts*.
Mullus Linnæus, Syst. Nat., Ed. X, 1758, 299 (*M. barbatus* Linnæus).
2650. *Mullus auratus* Jordan and Gilbert. *Goatfish; Red mullet*.
Eastern coast of North America, Cape Cod to Pensacola; rare.
Mullus barbatus auratus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 280, Pensacola, Fla.
- Genus 911. *UPENEUS* Cuvier. *Goatfishes*.
Upeneus Cuvier, Règne Animal, Ed. II, 1829, 157 (*uttatus; russelii; bifasciatus; trifasciatus*; first restricted by Bleeker to *bifasciatus*, an ally to *U. maculatus*).
Pseudupeneus Bleeker, Versl. Akad. Amsterdam, XIV, 1862, 134 (*P. prayensis* Bleeker).
Parupeneus Bleeker, Neder. Tijdschr. Dierk., I, 1863, 342 (*Mullus bifasciatus* Lacépède).
Mullhypeneus Poey, Repertorio, II, 1867, 160 (*Upeneus flavovittatus* Poey).
Brachymullus Bleeker, Arch. Neerl. Sci. Nat., XI, 1876, 333 (*Upeneus tetraspilus* Gunther).
Hypeneus Agassiz, corrected spelling.
2651. *Upeneus maculatus* (Bloch). *Red goatfish; Salmonete*.
West Indies and Brazil, Key West to Rio Janeiro; generally common.
Mullus maculatus Bloch, Naturgesch. Ausl. Fische, VII, 1893, 95, Brazil.
Upeneus punctatus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 355, 482, Martinique.
2652. *Upeneus dentatus* Gill.
Pacific coast of Mexico, Cape San Lucas, La Paz, and the Tres Marias Islands.
Upeneus dentatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 256, Cape San Lucas.
2653. *Upeneus parvus* Poey.
Cuba.
Upeneus parvus Poey, Memorias, I, 1853, 226, Cuba.
2654. *Upeneus martinicus* Cuvier and Valenciennes. *Yellow goatfish; Salmonete amarilla; King mullet*.
West Indies, north to Key West.
Upeneus martinicus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 356 (483), Martinique.
Upeneus balteatus Cuvier and Valenciennes, loc. cit., 356 (483), Cuba.
Upeneus flavovittatus Poey, Memorias, I, 1853, 224, Cuba.
2655. *Upeneus xanthogrammus* Gilbert.
Coast of Lower California.
Upeneus xanthogrammus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 553, La Paz, Lower California.

2656. *Upeneus grandisquamis* Gill. *Chivo*.
Pacific coast of Mexico and Central America, Guaymas to Panama.
Upeneus grandisquamis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 168, western coast of Central America.
Upeneus tetraspilus Gunther, Proc. Zool. Soc. Lond., 1864, 148, Panama.
- Genus 912. **MULLOIDES** Bleeker.
Mulloidies Bleeker, Verh. Bat. Gen., XXII, 1849, 12 (*Mullus flavolineatus* Lacépède).
2657. *Mulloidies rathbuni* (Evermann and Jenkins).
Gulf of California.
Upeneus rathbuni Evermann and Jenkins, Proc. U. S. Nat. Mus., XIV, 1891, 158, pl. 2, fig. 4, Bay of Guaymas, Sonora.
- Family 213. **SCIÆNIDÆ**. Croakers; Roncadores
- Genus 913. **LARIMUS** Cuvier and Valenciennes.
Larimus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 108 (145), (*L. breviceps* Cuvier and Valenciennes).
Amblyscion Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 165 (*A. argenteus* Gill).
Monosira Poey, Anales de l'Hist. Nat. Esp., 1881, 326 (*M. stahli* Poey).
- Subgenus **LARIMUS**.
2658. *Larimus effulgens* Gilbert
Pacific coast of Mexico, Sonora to Panama.
Larimus effulgens Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1421, San Juan Lagoon, Sonora
- Subgenus **AMBLYSCION** Gill.
2659. *Larimus argenteus* (Gill).
Panama.
Amblyscion argenteus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 165, western coast of Central America, the extreme form of the series.
2660. *Larimus acclivis* Jordan and Bristol.
Pacific coast of Mexico and Central America, from Sonora to Panama.
Larimus acclivis Jordan and Bristol (S. B.), in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1422, San Juan Lagoon, Sonora.
2661. *Larimus breviceps* Cuvier and Valenciennes. *Cabezon*.
West Indies, south to Brazil.
Larimus breviceps Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 108 (145), pl. 140, Brazil; San Domingo.
Monosira stahli Poey, Anal. Soc. Española Hist. Nat., X, 1881, 326, Pl. VI, Porto Rico.
2662. *Larimus pacificus* Jordan and Bollman.
Pacific Ocean, off coast of Colombia.
Larimus pacificus Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 161, Albatross station 2802, 8° 38' N., 79° 31' 30" W., between Galapagos Islands and Panama.
2663. *Larimus fasciatus* Holbrook. *Banded larimus*.
South Atlantic coast of the United States, from Chesapeake Bay to Galveston, Tex.; not common.
Larimus fasciatus Holbrook, Ichth. South Carolina, 1855, 153, pl. 22, fig. 1, Charleston, S. C.
- Genus 914. **ODONTOSCION** Gill.
Odontoscion Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 18 (*Corvina dentex* Cuvier and Valenciennes).
2664. *Odontoscion dentex* (Cuvier and Valenciennes). *Corvina*.
West Indies, north to the Tortugas.
Corvina dentex Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 103 (139), pl. 109, San Domingo.
2665. *Odontoscion xanthops* Gilbert.
Panama.
Odontoscion xanthops Gilbert in Jordan and Evermann, Check List, 1895, 396, Panama.

- Genus 915. CORVULA** Jordan and Eigenmann.
Corvula Jordan and Eigenmann, Review Sciænida Europe and America, in Rept. U. S. Fish Com., XIV, 1886, 377 (35), (*Johnius batabanus* Poey).
2666. *Corvula batabana* (Poey).
 Cuba and Porto Rico.
Johnius batabanus Poey, Memorias, II, 1861, 184, Batabano, south coast of Cuba.
- Genus 916. VACUOQUA** Jordan and Evermann.
Vacuogua Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 506 (*Corvina macrops* Steindachner).
 This genus differs from the aberrant *Corvula batabana* in having the deeper, symmetrical body and the silvery coloration of *Bairdiella* and related species.
2667. *Vacuogua macrops* (Steindachner). *Vacuocua*.
 Pacific coast of tropical America; Mazatlan to Panama.
Corvina macrops Steindachner, Ichth. Beitr., III, 1875, 24, fig. 2. Panama.
2668. *Vacuogua sialis* (Jordan and Eigenmann).
 Florida Keys.
Corvula sialis Jordan and Eigenmann, Rept. U. S. Fish Com., XIV, 1886, 379, Key West, Fla.
2669. *Vacuogua subæqualis* (Poey).
 West Indies.
Corvina subæqualis Poey, Ann. Lyc. Nat. Hist. N. Y., XI, 1875, 58, Cuba.
2670. *Vacuogua sanctæ-luciæ* (Jordan).
 West Indies.
Corvula sanctæ-luciæ Jordan, Proc. U. S. Nat. Mus., XII, 1889, 649, Port Castries, St. Lucia.
- Genus 917. ELATTARCHUS** Jordan and Evermann.
Elattarchus Jordan and Evermann, Check List, 1895, 397 (*Odontoscion archidum* Jordan and Gilbert).
2671. *Elattarchus archidum* (Jordan and Gilbert).
 Panama.
Odontoscion archidum Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 317, Panama.
- Genus 918. BAIRDIELLA** Gill. *Mademoiselles*
Bairdiella Gill, Cat. Fish. E. Coast N. Amer., Suppl. to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 33 (*Bodianus argyroleucos* Mitchell = *Dipterodon chrysurus* Lacépède).
2672. *Bairdiella chrysura* (Lacépède). *Mademoiselle*; *Yellowtail*; *Silver perch*.
 South Atlantic and Gulf coasts of United States, north to New York; very abundant on our sandy shores from Long Island to Texas.
Caranx chrysurus Lacépède, Hist. Nat. Poiss., III, 1802, 64, South Carolina; after Linnæus.
Perca punctata Linnæus, Syst. Nat., Ed. XII, 1766, 482, in part, South Carolina.
Bodianus argyro-leucos et *exiguus* Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 417, pl. 6, fig. 9, New York.
Bodianus pallidus Mitchell, loc. cit., 420, New York.
Homoprion subtruncatus Gill, Cat. Fish. E. Coast N. Amer., Suppl. to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 33, South Carolina; after Holbrook.
2673. *Bairdiella ensifera* (Jordan and Gilbert).
 Panama; common.
Sciæna ensifera Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 313, Bay of Panama; Punta Arenas.
Corvina fulgens Vaillant, Miss. Sci. au Mexique, 1883, 164, Pacific coast of Mexico.

2674. *Bairdiella icistia* (Jordan and Gilbert). *Corbineta*
Pacific coast of Mexico; rather common about Mazatlan.
Sciæna icistia Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 356, Mazatlan.
2675. *Bairdiella ronchus* (Cuvier and Valenciennes). *Ronco*, Ground drummer.
Atlantic coasts of tropical America; generally common in the West Indies and along the coast of Brazil.
Corvina ronchus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 79 (107), Maracaibo; Surinam.
2676. *Bairdiella bedoti* (Regan).
Cuba.
Sciæna (Bairdiella) bedoti Regan, Rev. Suisse Zool., XIII, 1905, 391, Pl. VI, Cuba.
- Genus 919. *NECTOR* Jordan and Evermann.
Nector Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1432 (*Corvina chrysoleuca* Gunther).
2677. *Nector armatus* (Gill).
Both coasts of tropical America; not uncommon on the Pacific coast about Panama, and equally abundant on the Atlantic coast.
Bairdiella armata Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 164, west coast of Central America.
Corvina acutirostris Steindachner, Ichth. Beitr., III, 1875, 28, pl. 4, Panama.
2678. *Nector veræ-crucis* (Jordan and Dickerson).
Gulf of Mexico, Vera Cruz.
Bairdiella veræ-crucis Jordan and Dickerson, Proc. U. S. Nat. Mus., XXXIV, 1908, 16, fig. 1, Vera Cruz, Mexico.
2679. *Nector alutus* (Jordan and Gilbert).
Pacific coast of Central America.
Sciæna aluta Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 232, La Union, Salvador.
2680. *Nector chrysoleucus* (Gunther).
Panama.
Corvina chrysoleuca Gunther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 427, pl. 67, fig. 1, Panama.
- Genus 920. *STELLIFER* (Cuvier) Oken.
Les Stellifères Cuvier, Règne Anim., Ed. I, 1817, 283 (*Bodianus stellifer* Bloch).
Stellifer (Cuvier) Oken, Isis, I, 1817, 1182 (*Bodianus stellifer* Bloch).
Stelliferus Stark, Elem. Nat. Hist., 1, 1828, 459 (*Bodianus stellifer* Bloch).
Homoprion Holbrook, Ichth. South Car., 1st ed. (mostly destroyed by fire), 1855, 168 (*Homoprion lanceolatus* Holbrook).
2681. *Stellifer stellifer* (Bloch).
Coasts of Guiana and Brazil.
Bodianus stellifer Bloch, Naturgesch. Ausl. Fische, IV, 1790, 55, Cape of Good Hope.
Corvina trispinosa Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 80 (109), Brazil; Cayenne.
2682. *Stellifer lanceolatus* (Holbrook).
South Atlantic and Gulf coasts of United States, Charleston to Texas.
Homoprion lanceolatus Holbrook, Ichth. South Carolina, Ed. I, 1855, 168, pl. 23, Port Royal Sound, S. C.
2683. *Stellifer ericymba* (Jordan and Gilbert).
Panama.
Sciæna ericymba Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 311, Bay of Panama.
2684. *Stellifer microps* (Steindachner).
Coast of Brazil and Guiana.
Corvina microps Steindachner, Ichth. Notizen, I, 1864, 6, pl. 2, fig. 1, Guiana.
Corvina stellifera Günther, Cat., II, 1860, 299, West Indies (not *Bodianus stellifer* Bloch).

2685. *Stellifer mindii* Meek and Hildebrand.
Atlantic coast of Panama.
Stellifer mindii Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, 626, Pl. LXVI, fig. 2, Mindi Reef, near Colon.
- Genus 921. ZESTIS Gilbert.
Zestis Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1439 (*Stellifer oscitans* Jordan and Gilbert).
2686. *Zestis oscitans* (Jordan and Gilbert).
Panama.
Sciæna oscitans Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 312, Bay of Panama.
2687. *Zestis furthii* (Steindacher).
Panama.
Corvina (*Homoprion*) *furthii* Steindachner, Ichth. Beitr., III, 1875, 26, fig. 3, Panama.
2688. *Zestis melanochir* (Eigenmann).
North coast of Colombia.
Stellifer melanochir Eigenmann, Proc. Am. Phil. Soc., LVI, 1917, 682, Tumneo.
- Genus 922. ZESTIDIUM Gilbert.
Zestidium Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1439 (*Stellifer illecebrosus* Gilbert).
2689. *Zestidium illecebrosus* (Gilbert).
Panama.
Stelliferus illecebrosus Gilbert, in Jordan and Evermann, Check List, 1895, 393, and Fishes North and Mid. Amer., 1896, 1443, Panama.
2690. *Zestidium colonense* (Meek and Hildebrand).
Coral reefs near Colon, Atlantic side of Isthmus of Panama.
Stellifer colonensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, 623, Mindi Reef, Colon.
- Genus 923. STELLICARENS Gilbert.
Stellicarens Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1445 (*Stellifer zestocarus* Gilbert).
2691. *Stellicarens zestocarus* (Gilbert).
Panama.
Stelliferus zestocarus Gilbert, in Jordan and Evermann, Check List, 1895, 398, and Fishes North and Mid. Amer., 1896, 1445, Panama.
- Genus 924. OPHIOSCION Gill.
Ophioscion Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 164 (*Ophioson typicus* Gill).
2692. *Ophioscion adustus* (Agassiz).
West Indies to coast of Brazil.
Corvina adusta Agassiz, in Spix, Pisc. Brasil., 1829, 126, pl. 70, Brazil.
2693. *Ophioscion punctatissimus* Meek and Hildebrand.
Panama.
Ophioscion punctatissimus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, pt. 2, 644, Pl. LXVIII, Panama Canal at Cristobal.
2694. *Ophioscion typicus* Gill.
Panama.
Ophioscion typicus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 165, west coast Central America.
Corvina ophioscion Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 428, Panama.
2695. *Ophioscion strabo* Gilbert.
Panama.
Ophioscion strabo Gilbert, Proc. U. S. Nat. Mus., XIX, 1896, 444, San Juan Lagoön, south of Guaymas, Mexico.

2696. *Ophioscion simulus* Gilbert.

Panama.

Ophioscion simulus Gilbert, in Jordan and Evermann, Check List, 1895, 398, and Fishes North and Mid. Amer., 1896, 1449, Panama.2697. *Ophioscion imiceps* (Jordan and Gilbert).

Panama.

Sciæna imiceps Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 309, Bay of Panama.2698. *Ophioscion scierus* (Jordan and Gilbert).

Pacific coast of tropical America, from Mazatlan to Panama.

Sciæna sciera Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 480, Panama.Genus 925. *SIGMURUS* Gilbert.*Sigmurus* Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1446 (*Corvina vermicularis* Günther).2699. *Sigmurus vermicularis* (Günther).

Panama.

Corvina vermicularis Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 427, pl. 67, fig. 2, Panama.Genus 926. *SCIÆNOPS* Gill. *Red drums.**Sciænops* Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 30 (*Perca ocellata* Linnæus).2700. *Sciænops ocellata* (Linnæus). *Red drum; Channel bass; Redfish; Pescado colorado; Bull redfish; Bass; Red bass; Sea bass; Spotted bass; Spottail.*

South Atlantic and Gulf coasts of the United States, New York to Texas; very abundant south.

Perca ocellata Linnæus, Syst. Nat., Ed. XII, 1766, 483, South Carolina. *Lutjanus triangulum* Lacépède, Hist. Nat. Poiss., IV, 1802, 181, pl. 24, fig. 3, Sumatra.*Sciæna imberbis* Mitchill, Trans. Lit. and Phil. Soc. New York, 1815, 411, New York.Genus 927. *CALLAUS* Jordan.*Callaus* Jordan, Rept. U. S. Fish Com., XIV, 1886, 401, 406 (59) (*Corvina deliciosa* Tschudi).2701. *Callaus deliciosa* (Tschudi).

Pacific coast of South America, from Panama to Peru.

Corvina deliciosa Tschudi, Fauna Peruana, Ichthyol., 1845, 8, Peru.Genus 928. *RHINOSCION* ⁶⁴Gill. *Red roncadors.**Rhinoscion* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 17 (*Amblodon saturnus* Girard).2702. *Rhinoscion saturnus* (Girard). *Red roncador; Black croaker.*

Coast of southern California, from Santa Barbara to Cedros Island.

Amblodon saturnus Girard, U. S. Pac. R. R. Surv., 1858, 98, San Diego, Calif.*Corvina* (*Johnius*) *jacobi* Steindachner, Ichth. Beitr., VIII, 1879, 3, San Diego; young.Genus 929. *RONCADOR* Jordan and Gilbert.*Roncador* Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 28 (*Corvina stearnsi* Steindachner).2703. *Roncador stearnsi* (Steindachner). *Roncador.*

Coast of southern California, north to Santa Barbara.

Corvina stearnsi Steindachner, Ichth. Beitr., III, 1875, 22, San Diego, Calif.Genus 930. *LEIOSTOMUS* Lacépède. *Goodies.**Leistomus* Lacépède, Hist. Nat. Poiss., IV, 1803, 439 (*L. xanthurus* Lacépède).*Leistomus* authors, revised spelling.

⁶⁴ This genus is very close to the European genus *Corvina* on the one hand and to the Peruvian *Chelotrema* on the other, but probably deserves to stand alone. There is no genuine species in America of *Sciæna* as now restricted.

2704. *Leiostomus xanthurus* Lacépède. *Spot; Goody; Postcroaker; Oldwife; Lafayette; Cape May goody; Porgy; Yellowtail.*
South Atlantic and Gulf coasts of United States; Cape Cod to Texas; once doubtfully recorded from Martinique.
Leiostomus xanthurus Lacépède, Hist. Nat. Poiss., IV, 1802, 439, pl. 10, fig. 1, Carolina
Mugil obliquus Mitchill, Trans. Lit. Phil. Soc. New York, 1815, 405, New York.
Sciaena multifasciata LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 255, east Florida
Leiostomus humeralis Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 105 (141), pl. 110, New York.
- Genus 931. **PACHYPOPS** Gill.
Pachypops Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 87 (*Micropogon trifilis* Muller and Troschel).
2705. *Pachypops fuscus* (Lacépède).
Rivers and estuaries of Guiana.
Perca fuscus Lacépède, Hist. Nat. Poiss., IV, 1802, 398, 424, Surinam.
Corvina biloba Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 83 (112), habitat unknown.
- Genus 932. **GENYONEMUS** Gill.
Genyonemus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 87 (*Leiostomus lineatus* Ayres).
2706. *Genyonemus lineatus* (Ayres). *Little roncadore; Kingfish; Croaker.*
Coast of southern California, San Francisco to Cedros Island.
Leiostomus lineatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 24, San Francisco.
- Genus 933. **MICROPOGON** Cuvier and Valenciennes. *Croaker.*
Micropogon Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 158 (213) (*M. lineatus* Cuvier and Valenciennes = *Perca undulata* Linnaeus).
Micropogonias Bonaparte, Ittiologia Generale, 1831, 104 (*Perca undulata* Linnaeus; substitute for *Micropogon* said to be preoccupied; but the genus of birds so named dates from 1838).
2707. *Micropogon undulatus* (Linnaeus). *Croaker; Roncodina; Corvina; Hardhead.*
South Atlantic and Gulf coasts of United States; Cape Cod to Texas.
Perca undulata Linnaeus, Syst. Nat., Ed. XII, 1766, 483, South Carolina.
Sciaena croker Lacépède, Hist. Nat. Poiss., IV, 1802, 309, Carolinas.
Bodianus costatus Mitchill, Trans. Lit. Phil. Soc. New York, 1815, 417, New York.
Micropogon lineatus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 160 (215), pl. 119, New York.
2708. *Micropogon furnieri* (Desmarest). *Verrugato; Whitemouth drummer.*
West Indies and coasts of South America.
Umbrina furnieri Desmarest, Première Décade Ichth., 22, pl. 2, fig. 3, 1823, 182, Habana.
? *Micropogon argenteus* Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 162 (218), Surinam.
2709. *Micropogon megalops* Gilbert.
Gulf of California.
Micropogon megalops Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 64, Gulf of California in 14 fathoms at Albatross station 3021.
2710. *Micropogon ectenes* Jordan and Gilbert. *Verrugate.*
Pacific coast of Mexico, Mazatlan.
Micropogon ectenes Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 355, Mazatlan.
2711. *Micropogon altipinnis* Günther.
Panama.
Micropogon altipinnis Günther, Proc. Zool. Soc. Lond., 1864, 149, San Jose; Panama; Chiapam.

Genus 934. **UMBRINA** Cuvier.

Umbina Cuvier, Règne Anim., Ed. I, II, 1817, 297 (*Sciæna cirrhosa* Linnaeus).

Atilus Gistel, Naturgesch. d. Thierreichs, 1848, 109 (*Sciæna cirrhosa* Linnaeus).

2712. ***Umbina broussonnetii*** Cuvier and Valenciennes.

West Indies; Florida to Brazil.

Umbina broussonnetii Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 139 (187), Jamaica.

2713. ***Umbina coroides*** Cuvier and Valenciennes.

Southern Florida to the West Indies and Brazil.

Umbina coroides Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 139 (187), Brazil.

2714. ***Umbina gracilicirrhus*** Metzelaar.

Coast of Venezuela.

Umbina gracilicirrhus Metzelaar, Tropisch Atlantisch Vissch., 1919, 72, coast of Venezuela.

- 2715. ***Umbina roncadior*** Jordan and Gilbert. *Yellow-finned roncadior*;
Yellow-tailed croaker

Coast of southern California, from Point Conception to Guaymas.

Umbina roncadior Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 277, Pequena Bay, west coast of Lower California.

2716. ***Umbina xanti*** Gill *Codorniz*.

Pacific coast of tropical America, Cape San Lucas to Panama.

Umbina xanti Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 257, Cape San Lucas.

Umbina analis Gunther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 426. Panama.

2717. ***Umbina sinaloæ*** Scofield.

Pacific coast of Mexico; Mazatlan.

Umbina sinaloæ Scofield, Proc. Calif. Acad. Sci., VI, 1896, 220, Pl. XXV, Mazatlan, Mexico.

2718. ***Umbina dorsalis*** Gill.

Pacific coast of Mexico.

Umbina dorsalis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 257, Cape San Lucas

Genus 935. **MENTICIRRHUS** Gill. *Kingfishes*

Menticirrhus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 86 (*Perca alburnus* Linnaeus = *Cyprinus americanus* Linnaeus).

- 2719. ***Menticirrhus americanus*** (Linnaeus). *Carolina whiting*; *Sand whiting*;
Kingfish; *Sea smelt*; *Sea mullet*.

South Atlantic and Gulf coasts of United States; Chesapeake Bay to Texas.

Cyprinus americanus Linnaeus, Syst. Nat., Ed. X, 1758, 321, Carolinas; (based on *Alburnus americanus* the whiting of Catesby, nonbinomial; not *Cyprinus americanus* of 12th edition of Systema Naturæ, which is a cyprinoid, *Abramis bosci* Cuvier and Valenciennes).

Perca alburnus Linnaeus, Syst. Nat., Ed. XII, 1766, 482. Charleston, S. C.

Umbina phalæna Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 167, Indianola and Brazos Santiago, Tex.

2720. ***Menticirrhus saxatilis*** (Bloch and Schneider). *Kingfish*; *Sea mink*;
Northern whiting; *Hake*; *Barb*.

Atlantic and Gulf coasts of the United States; Cape Ann to Key West and Pensacola; most common northward.

Johnius saxatilis Schneider, Ed. Bloch, Syst. Ichth., 1801, 75, New York.

2721. ***Menticirrhus undulatus*** (Girard). *California whiting*; *Sand sucker*.

Southern California, north to Santa Barbara.

Umbina undulata Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 148, San Diego, Calif.

- 2722. *Menticirrhus martinicensis*** (Cuvier and Valenciennes). *Jewsharp drummer*.
West Indies; Martinique; Jamaica to Patagonia; common in Brazil.
Umbrina martinicensis Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 186, Martinique.
Umbrina gracilis Cuvier and Valenciennes, loc. cit., 1830, 186, Brazil.
Umbrina arenata Cuvier and Valenciennes, loc. cit., 141 (190), Brazil.
Umbrina januaria Steindachner, Ichth. Beitr., V, 1876, 122, Rio Janeiro.
- 2723. *Menticirrhus nasus*** (Gunther).
Panama.
Umbrina nasus Gunther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 426, Panama.
Menticirrhus simus Jordan and Eigenmann, Review Sciænidæ, Rept. U. S. Fish Com., XIV, 1886, 427, Mazatlan and Panama.
- 2724. *Menticirrhus panamensis*** (Steindachner).
Pacific coast of tropical America; Mazatlan to Panama
Umbrina panamensis Steindachner, Ichth. Beitr., IV, 1875, 9, Panama.
- Genus 936. UMBRULA** Jordan and Eigenmann.
Umbrula Jordan and Eigenmann, Review Sciænidæ, Rept. U. S. Fish Com., XIV, 1886, 424 (*Umbrina littoralis* Holbrook).
- 2725. *Umbrula littoralis*** (Holbrook). *Surf whiting; Silver whiting; Shore whiting*.
South Atlantic and Gulf coasts of United States, North Carolina to Texas.
Umbrina littoralis Holbrook, Ichth. South Carolina, Ed. I, 1855, 142, pl. 20, fig. 1, South Carolina.
- 2726. *Umbrula elongata*** (Günther). *Verrugata*.
Pacific coast of tropical America, Mazatlan to Panama.
Umbrina elongata Gunther, Proc. Zool. Soc. Lond., 1864, 148, Chiapam.
- Genus 937. PARALONCHURUS** Bocourt.
Paralanchurus Bocourt, Nouv. Arch. Mus. d'Hist. Nat., IV, 1869, 21 (*P. petersi* Bocourt).
- 2727. *Paralanchurus petersi*** Bocourt.
Coast of Central America; rare at Panama.
Paralanchurus petersi Bocourt, Nouv. Archiv. du Muséum, IV, 1869, 22, La Union, San Salvador.
- Genus 938. POLYCLEMUS** Berg. *Corvalos*.
Polycirrhus Bocourt, Nouv. Arch. Mus. Paris, IV, 1869, 22 (*P. dumerili* Bocourt); name preoccupied in worms.
Polyclemus Berg, Ann. Mus. Nac. Buenos Aires, IV, 1895, 52 (*Polycirrhus dumerili* Bocourt).
- 2728. *Polyclemus dumerili*** (Bocourt).
Panama.
Polycirrhus dumerili Bocourt, Nouv. Arch. Mus. d'Hist. Nat., IV, 1869, 23, La Union, San Salvador.
Genyonemus fasciatus Steindachner, Ichth. Beitr., II, 1875, 31, Panama.
- Genus 939. ZONOSCION** Jordan and Evermann.
Zonoscion Jordan and Evermann, Check List, 1895, 401 (*Polycirrhus rathbuni* Jordan and Bollman).
- 2729. *Zonoscion rathbuni*** (Jordan and Bollman).
Panama.
Polycirrhus rathbuni Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 162, Panama.
- Genus 940. ZACLEMUS** Gilbert.
Zaclemus Gilbert, in Jordan and Evermann, Check List, 1895, 401 (*Paralanchurus goodei* Gilbert).
- 2730. *Zaclemus goodei*** (Gilbert).
Panama.
Paralanchurus goodei Gilbert, in Jordan and Evermann, Check List, 1895, 401, and Fishes North and Mid. Amer., 1896, 1480, Panama.

Genus 941. LONCHIURUS Bloch.

Lonchurus Bloch, Naturgesch. Ausl. Fische, VII, 1793, 144 (*Lonchurus barbat* Bloch=*Perca lanceolata* Bloch).

Lonchurus authors, preferred spelling.

2731. *Lonchurus lanceolatus* (Bloch).

West Indies to Guiana.

Perca lanceolata Bloch, Nov. Act. Sc., Copenh., III, 1788, 383, India.

Lonchurus depressus Bloch, Naturgesch. Ausl. Fische, VII, 1793, 144, Surinam.

Genus 942. POGONIAS Lacépède. Sea drums.

Pogonias Lacépède, Hist. Nat. Poiss., III, 1802, 138 (*P. fasciatus* Lacépède=*Labrus cromis* Linnæus).

Pogonathus Lacépède, loc. cit., V, 1803, 121 (*P. courbina* Lacépède).

Pogonathus Bonaparte, Ittiologia Generale, 1831 (*Pogonias fasciatus* Lacépède).

2732. *Pogonias cromis* (Linnæus). Gray drum; Drum; Black drum; Drumfish; Banded drum; Big drum.

Atlantic coast of America; Long Island to mouth of Rio Grande.

Labrus cromis Linnæus, Syst. Nat., Ed. XII, 1766, 479, Carolinas.

Pogonias fasciatus Lacépède, Hist. Nat. Poiss., III, 1802, 137.

Mugil grunniens Mitchell, Rept. in part, Fishes New York, 1814, 15, New York.

Sciaena fusca Mitchell, Trans. Lit. Phil. Soc. N. Y., 1815, 409, New York.

Sciaena gigas Mitchell, loc. cit., 412, New York.

2733. *Pogonias courbina* (Lacépède).

Guiana to Uruguay; rather common on coast of Brazil.

Pogonathus courbina Lacépède, Hist. Nat. Poiss., V, 1803, 121, Rio de la Plata.

Genus 943. APLODINOTUS Rafinesque. River drums; Thunderpumpers.

Aplodinotus Rafinesque, Jour. de Phys., LXXXVIII, 1819, 418 (*A. grunniens* Rafinesque).

Ambodon Rafinesque, Jour. de Phys., 1819, 421, based on teeth supposed to belong to *Ictiobus*, later restricted by Rafinesque to *Ambodon grunniens*.

Eutychelithus Jordan, Man. Vert., Ed. I, 1876, 242 (*Corvina richardsoni* Cuvier and Valenciennes, an injured or deformed example of *A. grunniens*).

Haplodonotus Gill, revised spelling.

2734. *Aplodinotus grunniens* Rafinesque. Fresh-water drum; Gaspergou; Thunderpumper; Lake sheepshead; Croaker; Bubbler; White perch.

Great Lakes to Texas; abundant in lakes and large streams between the Alleghenies and the Great Plains.

Aplodinotus grunniens Rafinesque, Jour. de Phys., 1819, 88, Ohio River.

Sciaena oscula LeSueur, Jour. Acad. Nat. Sci. Phila., 1822, 252, pl. 13, Lake Ontario.

Sciaena grisea LeSueur, loc. cit., 254, Ohio River at Pittsburgh, Pa.

Corvina richardsoni Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 100, Lake Huron, based on an abnormal specimen.

Ambodon concinnus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 307, Tennessee River.

Ambodon lineatus Agassiz, loc. cit., 307, Osage River, Mo.

2735. *Aplodinotus neglectus* (Girard).

Streams of eastern Mexico from Rio Grande to Rio Usumacinta.

Ambodon neglectus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 167, mouth of the Rio Grande; Rio Usumacinta. This Mexican form apparently distinguished by larger scales (43 instead of 55 to 60).

Genus 944. EQUETUS Rafinesque.

Eques Bloch, Nat. Ausl. Fische, VII, 1793, 91 (*E. americanus* Bloch=*Chætodon lanceolatus* Linnæus); preoccupied.

Equetus Rafinesque, Anal. Nat., 1815, 89; substitute for *Eques* Bloch.

2736. *Equetus lanceolatus* (Linnæus). *Ribbonfish; Guapena; Serrana*.
West Indies, ranging northward to Pensacola.
Ribbon fish, Edwards, Gleanings, pl. 210, 1758, Guadeloupe; Carolina.
Chætodon lanceolatus Linnæus, Syst. Nat., Ed. X, 1758, 277, "Caribbean Islands"; based on Edwards, pl. 210.
? *Serrana* Parra, Piezas, d'Hist. de Cuba, 1787, pl. 2, Cuba.
Eques americanus Bloch, Nat. Ausl. Fische, VII, 1793, West Indies.
Eques balleatus Cuvier, Règne Anim., ed. 2, II, 1829, 175, pl. 29, fig. 2, Martinique, after Edwards.
Sciæna edwardsi Gronow, Cat. Fish., ed. Gray, 1854, 53, after Edwards.
- Genus 945. **PAREQUES** Gill.
Pareques Gill in Goode, Bull. U. S. Nat. Mus., V, 1875, 50 (*Grammistes acuminatus* Bloch and Schneider).
2737. *Pareques viola* (Gilbert).
Panama
Eques viola Gilbert in Jordan and Evermann, Check List, 1895, 402, Panama.
2738. *Pareques acuminatus* (Bloch and Schneider).
West Indies; South Carolina to Brazil.
Grammistes acuminatus Bloch and Schneider, ed. Bloch, Syst. Ichth., 1801, 184, no locality; after Seba.
Eques lineatus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 126 (169), Brazil.
2739. *Pareques umbrosus* (Jordan and Eigenmann).
Southeast coast of the United States, Charleston to Pensacola.
Eques acuminatus umbrosus Jordan and Eigenmann, Review *Sciænidæ*, Rept. U. S. Fish Com., XIV, 1886, 440, Charleston and Pensacola.
2740. *Pareques punctatus* (Bloch and Schneider). *Serrana hispana*.
West Indies.
Serrana hispana Parra, Piezas, Hist. Nat. Cuba, 1787, 2, pl. 2, lower figure, Cuba.
Eques punctatus Bloch and Schneider, Syst. Ichth., 1801, 106, Cuba; based on Parra, 2, pl. 2, fig. 2.
2741. *Pareques pulcher* (Steindachner).
Tortugas Archipelago; Barbados.
Eques pulcher Steindachner, Ichth. Notizen, VI, 1867, 43, Barbados.

Family 214. OTOLITHIDÆ. Weakfishes; Queenfishes

- Genus 946. **SERIPHUS** Ayres.
Seriphus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1858-1862 (1863; read Nov. 5, 1860), 80 (*S. politus* Ayres).
2742. *Seriphus politus* Ayres. *Queenfish; White croaker*.
Coast of southern California from Point Conception to Cedros Island; common on sandy shores.
Seriphus politus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1858-1862 (1863; read Nov. 5, 1860), 80, fig. 21, about San Diego.
- Genus 947. **ISOPISTHUS** Gill.
Isopisthus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 18 (*Ancylodon parvipinnis* Cuvier and Valenciennes).
2743. *Isopisthus parvipinnis* (Cuvier and Valenciennes).
Coast of Brazil, north to Cayenne and Guiana.
Ancylodon parvipinnis Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 62 (84), pl. 105, Cayenne
Isopisthus affinis Steindachner, Denksch. Mat. Nat. Kais. Acad. Wiss., XLI, Bd. I., Abth., 1879, 43, pl. 2, fig. 2, Porto Alegre, Brazil.
2744. *Isopisthus remifer* Jordan and Gilbert.
Panama, on sandy shores, rather common.
Isopisthus remifer Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 320, Panama.

- 2745. *Isopisthus harroweri* Fowler.**
Atlantic shore of Central America.
Isopisthus harroweri Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 402, Colon
- Genus 948. NEBRIS Cuvier and Valenciennes.**
Nebris Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 111 (149)
(*N. microps* Cuvier and Valenciennes)
- 2746. *Nebris microps* Cuvier and Valenciennes.**
Atlantic coast of northern South America; sandy shores.
Nebris microps Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 111 (149), pl. 112, Surinam
- 2747. *Nebris occidentalis* Vaillant. *Guavina*.**
Pacific coast of tropical America
Nebris occidentalis Vaillant, Bull. Mus. Paris, 1897, 124, Pacific coast of Central America.
Nebris zestus Jordan and Starks, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1417, Panama.
- Genus 949. MACRODON Schintz.**
Ancylodon Cuvier, in Oken, Isis, 1817, 1182 (*Lonchurus ancylodon* Schneider); name preoccupied in mammals.
Macrodon Schintz, Das Thierreich, II, 1822, 482 (*Lonchurus ancylodon* Schneider).
Nomalus Gistel, Thierreich, X, 1848, substitute for *Ancylodon*.
Sagenichthys Carlos Berg, Ann. Mus. Nat. Buenos Aires, 1895, 42 (*Lonchurus ancylodon* Schneider); substitute for *Ancylodon*.
- 2748. *Macrodon ancylodon* (Bloch and Schneider). *Pescadillo del rey*.**
West Indies, Guiana; Brazil; and Uruguay; rather scarce.
Lonchurus ancylodon Bloch and Schneider, ed. Bloch, Syst. Ichth., 1801, 102, pl. 25, Surinam
Ancyloden jaculidens Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 60 (81), Cayenne.
- 2749. *Macrodon mordax* (Gilbert and Starks).**
Panama.
Sagenichthys mordax Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 121, Pl. XVI, Panama.
- Genus 950. BUCCONE Jordan and Evermann.**
Buccone Jordan and Evermann, Check List, 1895, 394 (*Cestreus prædatorius* Jordan and Gilbert).
- 2750. *Buccone prædatoria* (Jordan and Gilbert). *Bocone*.**
Coast of Panama; scarce.
Cestreus prædatorius Jordan and Gilbert, Rept. U. S. Fish Com., XIV, 1886, 363, Panama.
- Genus 951. CYNOSCIION Gill.**
Cestreus Gronow, Cat. Fish., ed. Gray, 1854, 49 (*Cestreus carolinensis* Gronow=*Otolithus nebulosus* Cuvier and Valenciennes); preoccupied in gobies.
Cynoscion Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 81 (*Johnius regalis* Bloch and Schneider).
Apeudobranchus Gill, loc. cit., XIV, 1862, 18 (*Otolithus tæ-ræ* Cuvier and Valenciennes=*Cheilodipterus acoupa* Lacépède).
- Subgenus CYNOSCIION Gill.**
- 2751. *Cynoscion acoupa* (Lacépède). *Acoupa*; *Toeroe*.**
Atlantic coast of South America, Brazil north to Venezuela; generally common.
Cheilodipterus acoupa Lacépède, Hist. Nat. Poiss., III, 1802, 546, Cayenne.
Lutjanus cayanensis Lacépède, loc. cit., IV, 1802, 196, Cayenne.
Otolithus rhomboidalis Cuvier, Règne Anim., ed. 2, II, 1829, 173, Cayenne.
Otolithus toe-roë Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 54 (72), pl. 103, Cayenne.

2752. *Cynoscion squamipinnis* (Gunther).
Pacific coast of tropical America; known from a few specimens taken at La Union and Panama
Otolithus squamipinnis Gunther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 429, Panama.
2753. *Cynoscion othonopterus* Jordan and Gilbert.
Gulf of California
Cynoscion othonopterus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 274, Punta San Felipe, Mexico.
2754. *Cynoscion obliquatus* (Cuvier and Valenciennes).
Martinique
Otolithus obliquatus Cuvier and Valenciennes, in Sauvage, Bull. Soc. Philom. Paris, III, 1879, 209, Martinique.
2755. *Cynoscion jamaicensis* (Vallant and Bocourt). *Mongolar drummer*.
Jamaica and Porto Rico.
Otolithus jamaicensis Vallant and Bocourt, Miss. Sci. au Mexique, Poissons, 1874, 156, Jamaica.
2756. *Cynoscion regalis* (Bloch and Schneider). *Common weakfish; Squeteague; Sea trout; Squit; Shecuts; Chickwick; Salt-water trout, Gray trout*.
Atlantic and Gulf coasts of United States, Cape Cod southward to Mobile
Johmsus regalis Bloch and Schneider, Syst. Ichth., 1801, 75, New York.
Roccus comes Mitchill, Report, in part, Fishes New York, 1814, 25, New York.
Labrus squeteague Mitchill, Trans. Lit. Phil. Soc. New York, 1815, 396, pl. 2, fig. 1, New York
2757. *Cynoscion nothus* (Holbrook). *Bastard weakfish*.
South Atlantic and Gulf coasts of United States; rather rare at Charleston. This unspotted fish is now generally regarded as a pale variant of *Cynoscion regalis*.
Otolithus nothus Holbrook, Ichth. South Carolina, 1855, 134, pl. 19, fig. 1, South Carolina.
2758. *Cynoscion thalassinus* (Holbrook).
Pensacola; Pass Christian, Miss.; Hampton Roads, Va.
Otolithus thalassinus Holbrook, Ichth. South Carolina, 1855, 132, pl. 18, fig. 2, Charleston, S. C.
2759. *Cynoscion leiarchus* (Cuvier and Valenciennes).
Coasts of Brazil and Guiana
Otolithus leiarchus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 58 (78), Brazil; Cayenne.
2760. *Cynoscion microlepidotus* (Cuvier and Valenciennes).
Coast of Brazil and Guiana.
Otolithus microlepidotus Cuvier and Valenciennes, Hist. Nat. Poiss.; V, 1830, 59 (79), Surinam.
- Genus 952. *ERISCION* Jordan and Evermann.
Eriscion Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 506 (*Otolithus nebulosus* Cuvier and Valenciennes).
2761. *Eriscion reticulatus* (Günther). *Corvina*.
Pacific coast of tropical America, Mazatlan to Panama.
Otolithus reticulatus Günther, Proc. Zool. Soc. Lond., 1864, 149, San Jose de Guatemala; Chiapam
2762. *Eriscion nebulosus* (Cuvier and Valenciennes). *Spotted weakfish; Spotted sea trout; Spotted Squeteague*.
South Atlantic and Gulf coasts of the United States; New York to Texas; everywhere common on our southern coast; rare north of Virginia.
Labrus squeteague maculatus Mitchill, Trans. Lit. Phil. Soc., 1815, 396, New York; not *Labrus maculatus* Bloch.
Otolithus nebulosus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 59 (79), locality unknown.
Otolithus carolinensis Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 351, (475), South Carolina.
Otolithus drummondi Richardson, Fauna Bor.-Amer., Fish., 1836, 70, New Orleans.

2763. *Eriscion parvipinnis* (Ayres). *California bluefish*.
Coast of Lower California; Santa Barbara Islands to Guaymas;
common along the coasts of southern California as far north as
San Pedro.
Cynoscion parvipinnis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1858-1862
(read Dec. 1, 1861, published Sept., 1862), 156, fig. 48, coast of
Lower California
Otolithus magdalenæ Steindachner, Ichth. Beit., III, 1875, 34, Mag-
dalena Bay, Lower California.
2764. *Eriscion xanthulus* (Jordan and Gilbert). *Corvina de las aletas amaril-
las; Yellowmouth*.
Pacific coast of Mexico; not rare about Mazatlan.
Cynoscion xanthulum Jordan and Gilbert, Proc. U. S. Nat. Mus., IV,
1881, 460, Mazatlan.
2765. *Eriscion albus* (Günther).
Pacific coast of tropical America.
Otolithus albus Günther, Proc. Zool. Soc. Lond., 1864, 149, Chiapam;
Panama.
Otolithus cayennensis Vaillant and Bocourt, Miss. Sci. Mexique, Poiss.,
1874, 156, La Union, Salvador.
2766. *Eriscion stolzmanni* (Steindachner).
Pacific coast of tropical America, from Panama to Peru.
Otolithus stolzmanni Steindachner, Denkschr. Mat. Nat. Kais. Acad.
Wiss., XLI, Bd. I, Abth. 35, 1879, pl. 2, fig. 1, Tumbes, Peru.
2767. *Eriscion virescens* (Cuvier and Valenciennes).
Coasts of Guiana and Brazil.
Otolithus virescens Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830,
54 (72), Surinam.
Otolithus microps Steindachner, Denkschr. Mat. Nat. Kais. Acad.
Wiss., XLI, Bd. I, Abth. 38, 1879, pl. 8, fig. 2, Porto Alegre, Brazil.
2768. *Eriscion macdonaldi* (Gilbert). *Totuava*.
Gulf of California.
Cynoscion macdonaldi Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 64,
head of Gulf of California.
- Genus 953. *ATRACTOSCION* Gill.
Atractoscion Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 18 (*Otolithus
æquidens* Cuvier and Valenciennes).
2769. *Atractoscion nobilis* (Ayres). *White sea bass of California*.
Coast of California, north to San Francisco, occasionally straying as
far north as Boca de Quadra Inlet, Alaska.
Johnius nobilis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 77, San
Francisco.
Otolithus californiensis Steindachner, Ichth. Beitr., III, 1875, 31, San
Diego, Calif., and Magdalena Bay, Lower California.
2770. *Atractoscion phoxocephalus* (Jordan and Gilbert).
Pacific coast of tropical America; Panama.
Cynoscion phoxocephalum Jordan and Gilbert, Bull. U. S. Fish Com.,
I, 1881, 318, Panama.

Family 215. MALACANTHIDÆ. Matajuelos

Genus 954. MALACANTHUS Cuvier.

Malacanthus Cuvier, Règne Animal, ed. 2, II, 1829, 264 (*Coryphæna
plumieri* Bloch).

2771. *Malacanthus plumieri* (Bloch). *Matajuelo blanco*.
West Indies, generally common.
Coryphæna plumieri Bloch, Ichth., V, 1787, 119, pl. 175, Martinique;
from a drawing by Plumier.

Family 216. BRANCHIOSTEGIDÆ.⁶⁵ Blanquillos

Genus 955. CAULOLATILUS Gill.

- Caulolatilus* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 240 (?*Latilus chrysops* Cuvier and Valenciennes).
Dekaya Cooper, Proc. Calif. Acad. Nat. Sci., III, pt. 1, 1863 (Nov.), 70, fig. 17 (*D. anomala* Cooper).

2772. *Caulolatilus chrysops* (Cuvier and Valenciennes). *Blanquillo*.
 Gulf of Mexico and southward.
Latilus chrysops Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 496, Brazil.
2773. *Caulolatilus microps* Goode and Bean.
 Gulf of Mexico.
Caulolatilus microps Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 43, off Pensacola, Fla.
2774. *Caulolatilus princeps* (Jenyns). *Blanquillo*.
 Cape San Lucas to the Galapagos.
Latilus princeps Jenyns, Zool. Voy. *Beagle*, Fishes, 1842, 52, Chatham Island, Galapagos Archipelago.
Caulolatilus affinis Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 68, Cape San Lucas
2775. *Caulolatilus anomalus* (Cooper). *Blanquillo*; *Sea whitefish*.
 Coast of southern California, about rocky islands from Point Conception southward; common.
Dekaya anomala Cooper, Proc. Calif. Acad. Nat. Sci., III, pt. 1, 1863 (Nov.), 70, fig. 17, Santa Catalina Island off southern California.
2776. *Caulolatilus cyanops* Poey. *Blanquillo*.
 Coasts of Cuba and Porto Rico.
Caulolatilus cyanops Poey, Repertorio, I, 1866, 311-312, Cuba.
- Genus 956. LOPHOLATILUS Goode and Bean. *Tilefish*.
Lopholatilus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 205 (*L. chamæleonticeps* Goode and Bean).
- 2777-2778. *Lopholatilus chamæleonticeps* Goode and Bean. *Tilefish*.
 Deep waters of the western Atlantic; abundant at intervals but subject to calamities; a valued food fish.
Lopholatilus chamæleonticeps Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 205; 80 miles S. by E. of No-man's Land, 40° N., 70° W., in 84 fathoms.

Family 217. TRICHODONTIDÆ

Genus 957. TRICHODON Steller.

- Trichodon* Steller, in Tilesius, Mém. Acad. St. Petersb., IV, 1811, 468 (*Trachinus trichodon* Tilesius).
2779. *Trichodon trichodon* (Tilesius). *Sandfish*.
 Coasts of Kamchatka and eastward; Herendeen Bay and elsewhere in Bering Sea, and south to coast of Oregon.
Trachinus trichodon Tilesius, Mém. Acad. St. Petersb., IV, 1811, fig. 8, 73, Kamchatka.
Trachinus gasteropelecus Tilesius, loc. cit., 1811, 466, Kamchatka.
Trichodon lineatus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 60, San Francisco.
- Genus 958. ARCTOSCOPUS Jordan and Evermann.
Arctoscopus Jordan and Evermann, Check List, 1895, 464 (*Trichodon japonicus* Steindachner).
2780. *Arctoscopus japonicus* (Steindachner).
 Sea of Japan, eastward to Sitka, Alaska.
Trichodon japonicus Steindachner, Ichth. Beitr., X, 1881, 4 Strietok and Sitka.

⁶⁵ The name *branchistegus* Rafinesque (1815) has priority over *Latilus* Cuvier and Valenciennes (1830).

Family 218. CIRRHITIDÆ

Genus 959. CIRRHITUS Lacépède.

Cirrhutus Lacépède, Hist. Nat. Poiss., V, 1803, 3 (*C. maculatus* Lacépède).
Cirrhutichthys Gunther, Cat., II, 1860, 73; not of Bleeker.
Cirrhutes of authors, revised spelling.

2781. *Cirrhitus rivulatus* (Valenciennes).

Cape San Lucas to the Revillagigedos and Galapagos Islands.
Cirrhutes rivulatus Valenciennes, Voy. *Vénus*, Poiss., 1855, 309, pl. 3,
 fig. 1, Galapagos Islands
Cirrhutus betaurus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 259,
 Cape San Lucas; young.

Family 219. ANTIGONIIDÆ

Genus 960. ANTIGONIA Lowe.

Antigonia Lowe, Proc. Zool. Soc. Lond., 1843, 85 (*A. capros* Lowe).
Caprophonus Muller and Troschel, Horæ Ichth., III, 1845, 28, pl. 5,
 fig. 1 (*C. aurora* Muller and Troschel).
Hypsinothus Temminck and Schlegel, Faun. Japon., Poiss., 1844, 84
 (*H. rubescens* Gunther).

2782. *Antigonia capros* Lowe.

Tropical Atlantic, in rather deep water.
Antigonia capros Lowe, Proc. Zool. Soc. Lond., 1843, 85, Madeira.
Caprophonus aurora Muller and Troschel, Horæ Ichthyologicæ, III,
 1845, p. 28, pl. 5, fig. 1, Barbados.

Family 220. SCORPIDÆ. Halfmoon fishes

Genus 961. MEDIALUNA Jordan and Fesler.

Medialuna Jordan and Fesler, Rept. U. S. Fish Com., XVII, 1889-1891,
 536 (*Scorpius californiensis* Steindachner).

2783. *Medialuna californiensis* (Steindachner). *Medialuna*; Halfmoon.

Coast of southern California from Point Conception southward to
 Cedros Island
Scorpius californiensis Steindachner, Ichth. Beitr., III, 1875, 19, San
 Diego.

Suborder SQUAMIPENNES

Family 221. EPHIPPIDÆ

Genus 962. CHÆTODIPTERUS Lacépède.

Chætodipterus Lacépède, Hist. Nat. Poiss., IV, 1803, 503 (*Chætodon*
plumieri Bloch = *Chætodon faber* Broussonet).
Parephippus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 165 (*Ephip-*
pus zonatus Girard).

2784. *Chætodipterus faber* (Broussonet). *Angelfish*; *Spadefish*; *Moonfish*;
Tripletail; *Three-banded sheephead*; *Three-tailed porgy*.

West Indies, Cape Cod to Rio Janeiro, south Atlantic coast
Faber marinus fere quadratus (the pilotfish) Sloane, Jamaica, II, 1793,
 290, pl. 251, Jamaica.

Chætodon faber Broussonet, Ichth., V, Dec. 1, 1782, pl. 4, Jamaica.
Zeus quadratus Gmelin, Syst. Nat., I, 1788, 1225, Jamaica; after
 Sloane.

Chætodon plumieri Bloch, Naturgesch. Ausl. Fische, III, 1787, 104,
 Martinique.

Selene quadrangularis Lacépède, Hist. Nat. Poiss., IV, 1803, 564,
 Jamaica; after Sloane.

Chætodon oniformis Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 247,
 pl. 5, fig. 4, New York.

Ephippus gigas Cuvier, Règne Anim., ed. 2, II, 1829, 191, America.

2785. *Chætodipterus zonatus* (Girard).

Pacific coast of America, from San Diego to Panama; common south-
 ward.

Ephippus zonatus Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 110,
 San Diego, Calif.

Genus 963. PARAPSETTUS Steindachner.

Parapsettus Steindachner, Ichth. Beitr., III, 1875, 50 (*P. panamensis* Steindachner).

2786. Parapsettus panamensis Steindachner.

Panama.

Parapsettus panamensis Steindachner, Ichth. Beitr., III, 1875, 51, Panama.

Family 222. CHÆTODONTIDÆ. Butterflyfishes; Coralfishes**Genus 964. PROGNOTHODES Gill.**

Prognathodes Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 238 (*Chelmo pelta* Günther = *Chelmon aculeatus* Poey).

2787. Prognathodes aculeatus (Poey).

West Indies.

Chelmon aculeatus Poey, Memorias, II, 1861, 202, Cuba.

Chelmo pelta Günther, Cat., II, 1860, 38, locality unknown.

Genus 965. FORCIPIGER Jordan and McGregor.

Forcipiger Jordan and McGregor, Rept. U. S. Fish Com., XXIV, 1898, 279 (*Chætodon longirostris* Broussonet).

2788. Forcipiger longirostris (Broussonet).

South Seas east to Revillagigedo Islands.

Chætodon longirostris Broussonet, Desc. Ichth., I, 1782, 23, pl. 7, Society and Sandwich Islands.

Forcipiger flavissimus Jordan and McGregor, Rept. U. S. Fish Com., XXIV, 1898, 279, Clarion Island, Revillagigedo Archipelago.

Genus 966. CHÆTODON⁶⁶ Linnæus. Butterflyfishes.

Chætodon Linnæus, Syst. Nat., Ed. X, 1758, 272 (*C. capistratus* Linnæus).

Rabdophorus Swainson, Classn. Fishes, II, 1839, 21 (*Chætodon ephippium* Cuvier and Valenciennes).

Citrædus Kaup, Wiegmann's Archiv, XXVI, 1860, 136 (*Chætodon ornatus* Solander).

Linophora Kaup, loc. cit., 137 (*Chætodon auriga* Forskål).

Sarothrodus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 99 (*Chætodon capistratus* Gill).

Tholichthys Günther, Ann. Mag. Nat. Hist., I, 4th ser., 1868, 457 (*T. osseus* Günther; a larval form).

Chætodontops Bleeker, Arch. Neerl. Soc. Nat., XI, 1876, 304 (*Chætodon collaris* Bloch).

Tetragonopterus Klein, in Bleeker, loc. cit., 305 (*Chætodon striatus* L.); not *Tetragonopterus* of Cuvier.

Lepidochætodon Bleeker, loc. cit., 305 (*Chætodon unimaculatus* Bloch).

Gonochætodon Bleeker, loc. cit., 306 (*Chætodon triangulum* Kuhl and Van Hasselt).

Orychætodon Bleeker, loc. cit., 306 (*Chætodon lineolatus* Quoy and Gaimard).

Hemichætodon Bleeker, Versl. Ak. Amsterdam, XII, 1876, 38 (*Chætodon capistratus* Linnæus).

Antisochætodon Klunzinger, Fische Rothen Meeres, I, 1884, 3 (*Chætodon auriga* Cuvier and Valenciennes).

Subgenus CHÆTODON Linnæus.**2789. Chætodon capistratus Linnæus. Parché.**

West Indies, Habana, north to Woods Hole.

Chætodon capistratus Linnæus, Syst. Nat., Ed. X, 1758, 275, India.

Chætodon briceii Smith, Bull. U. S. Fish Com., XVII, 1897, 102, with text fig., Woods Hole, Mass.; young.

2790. Chætodon striatus Linnæus. Portuguese butterfly.

West Indies.

Chætodon striatus Linnæus, Syst. Nat., Ed. X, 1758, 275, India.

⁶⁶ These nominal genera, here included under *Chætodon*, are mostly imperfectly defined. The following may be valid: *Chætodon*, lines of scales running upward and backward from the lateral line, no produced rays, *Linophora*, similar but one dorsal ray much produced, *Orychætodon*, scales of sides enlarged, snout sharp; *Chætodontops*, snout blunt, etc. *Lepidochætodon*, scales of sides much enlarged, in horizontal rows, etc. *Rabdophorus*, scales horizontal, much reduced above lateral line (see Jordan and Jordan, Mem. Carnegie Mus. X, 55, 1922).

Subgenus CHÆTODONTOPS Bleeker.

2791. *Chætodon bimaculatus* Bloch⁶⁷ *Parché; Isabelita de la alto.*
West Indies, north in Gulf Stream to New Jersey and Rhode Island.
Chætodon ocellatus Bloch, Naturgesch. Ausl. Fische, III, pl. 211, 1787,
105, "also from the East Indies"
Chætodon bimaculatus Bloch, loc. cit., 1790, pl. 219, fig. 1
Sarothrodus maculo-cinctus Gill, Proc. Acad. Nat. Sci. Phila., XIII,
1861, 99, Newport, R. I.
Sarothrodus amplexicollis Poey, Enumeratio, 1875, 63, Cuba
2792. *Chætodon nigrirostris* (Gill)
Cape San Lucas, Lower California; Panama Bay.
Sarothrodus nigrirostris Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
243, Cape San Lucas, Lower California.
2793. *Chætodon humeralis* Gunther. *Muñeca; Dollfish.*
Pacific coast of America, from Guaymas to Panama
Chætodon humeralis Gunther, Cat., II, 1860, 19, "Sandwich Islands,"
by error for Central America.
2794. *Chætodon sedentarius* Poey.
West Indian fauna.
Chætodon sedentarius Poey, Memorias, II, 1860, 203, Cuba.
2795. *Chætodon gracilis* Gunther.
West Indies.
Chætodon gracilis Gunther, Cat., II, 1860, 20, Caribbean Sea, West
Indies.
2796. *Chætodon aya* Jordan.
Gulf of Mexico.
Chætodon aya Jordan, Proc. U. S. Nat. Mus., IX, 1886, 225, Snapper
Banks, south of Pensacola, Fla.
2797. *Chætodon atæniatus* (Poey).
Habana Cuba.
Sarothrodus atæniatus Poey, Synopsis, 1868, 353, Cuba.
? *Chætodon unicolor* Sauvage, Bull. Soc. Philom., IV, 1880, 222, Mar-
tinique.

Genus 967. POMACANTHUS Lacépède. *Chirivitas.*

- Pomacanthus* Lacépède, Hist. Nat. Poiss., IV, 1803, 517 (*Chætodon*
arcuatus Linnæus, as restricted by Cuvier).
2798. *Pomacanthus arcuatus* (Linnæus). *Black angel; Chirivita; Portugais.*
West Indies, north to New Jersey, south to Bahia.
Chætodon arcuatus Linnæus, Syst. Nat., Ed. X, 1758, 273, "India."
Chætodon aureus Bloch, Naturgesch. Ausl. Fische, III, 1787, 49, Mar-
tinique; on drawing by Plumier.
Chætodon lutescens Bonnaterre, Encycl. Meth., 1788, 82, Jamaica; after
Browne.
Chætodon littoricola Poey, Synopsis, 1868, 351, Cuba.
Pomacanthus balleatus Cuvier and Valenciennes, Hist. Nat. Poiss., VII,
1831, 156 (208), Porto Rico.
Pomacanthus cingulatus Cuvier and Valenciennes, loc. cit., 156 (209),
pl. 185, West Indies (probably Porto Rico).
Pomacanthus quinquecinctus Cuvier and Valenciennes, loc. cit., 157
(210), West Indies (probably Porto Rico).
2799. *Pomacanthus paru* (Bloch). *Paru; Indianfish; Flatfish; Chirivita.*
West Indies, south to Bahia; Tortugas.
Chætodon paru Bloch, Naturgesch. Ausl. Fische, III, 1787, 57, Brazil;
on a drawing.

Genus 968. POMACANTHODES Gill.

- Pomacanthodes* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 244 (*P.*
zonipectus Gill).

⁶⁷ Poey seems to have been the first to recognize the identity of *ocellatus* (which has page priority) with *bimaculatus*. He chose to use the name *bimaculatus*, which should therefore stand.

- 2800. *Pomacanthodes zonipectus* Gill.** *Mojarra de las piedras.*
West coast of tropical America, from Mazatlan to Panama.
Pomacanthodes zonipectus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 244, San Salvador.
Pomacanthus crescentalis Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 358, Mazatlan.
- Genus 969. CENTROPYGE Kaup.**
Centropyge Kaup, Wiegmann's Archiv, XXVI, 1860, 140 (*Holacanthus tubicen* Cuvier and Valenciennes).
- 2801. *Centropyge passer* (Valenciennes).**
West coast of tropical America; Cape San Lucas to Galapagos Islands.
Holacanthus passer Valenciennes, Voy. *Vénus*, 1846, 327, pl. 6, Galapagos Archipelago.
Holacanthus strigatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 243, Cape San Lucas.
- 2802. *Centropyge clarionensis* (Gilbert).**
Clarion, Socorro, and San Benedicto Islands of the Revillagigedo group.
Holacanthus clarionensis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 72, Revillagigedo Islands; Clarion, Socorro, and San Benedicto.
- 2803. *Centropyge lunata* (Blosser).**
St. Croix.
Holacanthus lunatus Blosser, Ann. Carnegie Mus., VI, 1909, 299, St. Croix.
This small species with low fins and rounded caudal if fairly mature can not be placed in *Holacanthus*. It seems to belong to the genus *Centropyge*, a genus thus far known only from the Pacific.
- Genus 970. HOLACANTHUS Lacépède.** *Catalineta*.
Holacanthus Lacépède, Hist. Nat. Poiss., IV, 1803, 525 (*Chætodon tricolor* Bloch).
- 2804. *Holacanthus tricolor* (Bloch).** *Rock beauty; Catalineta; Vaqueta de los colores.*
West Indies and Bermudas; south to Bahia.
Chætodon tricolor Bloch, Naturgesch. Ausl. Fische, IX, 1795, 103, Cuba.
Catalineta Parra, Descr. Dif. Piez., Hist. Nat. Cuba, 1787, 12, Pl. VII, fig. 2, Cuba.
- Genus 971. ANGELICHTHYS Jordan and Evermann.**
Angelichthys Jordan and Evermann, Check List, 1895, 420 (*Chætodon ciliaris* Linnaeus).
- 2805. *Angelichthys ciliaris* (Linnaeus).** *Angelfish; Isabelita; Yellow angel; Golden angel; Blue angel.*
West Indies north to Key West.
Chætodon ciliaris Linnaeus, Syst. Nat., Ed. X, 1758, 276, Indies, in part.
Isabelita Parra, Dif. Piezas, Hist. Nat. Cuba, 1787, 11, pl. 7, fig. 1, Cuba.
Chætodon squamulosus Shaw, Naturalist's Miscellany, VIII, 1796, 275; after the angelfish of Catesby.
Chætodon parrae Bloch and Schneider, Syst. Ichth., 1801, 235, Cuba; after *Isabelita* of Parra.
Holacanthus cornutus Desmarest, Décade Ichthyologique, 1823, 44, pl. 3, fig. 3, Cuba.
Holacanthus formosus Castelnau, Anim. Amer. Sud, Poiss., 1855, 19, Pl. XI, fig. 2, Bahia.
- 2806. *Angelichthys isabelita* Jordan and Rutter.** *Angelfish.*
West Indies; north to Key West, Fla.
Angelichthys isabelita Jordan and Rutter, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1685, Key West.
- 2807. *Angelichthys townsendi* Nichols and Mowbray.**
Florida.
Angelichthys townsendi Nichols and Mowbray, Bull. Amer. Mus. Nat. Hist., XXXIII, 1914, 581, Key West, Fla.

Family 223. ZANCLIDÆ. Moorish idols

Genus 972. ZANCLUS Cuvier and Valenciennes

Zanclus Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 70 (102) (*Chætodon cornutus* Linnæus).

Gonopterus Gronow, Cat. Fish., ed. Gray, 1854, 77 (*G. mærens* Gronow = *Chætodon cornutus* Linnæus).

Gnathocentrum Guichenot, Ann. Maine et Loire, IX, 1866, 4 (*Zanclus centrognathus* Cuvier and Valenciennes); young

2808. *Zanclus cornutus* (Linnæus). Moorish idol, Besan; Piquier; Porte enseigne.

Warm seas of the Pacific; Revillagigedo, Cocos, and Galapagos islands; Hawaii; Samoa; East Indies and islands of Polynesia; common and widely distributed throughout the Pacific.

Chætodon cornutus Linnæus, Syst. Nat., Ed. X, 1758, 273, Indies; after Artedi.

Zanclus centrognathus Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 397 (528), near Equator, 75° E.

Gonopterus mærens Gronow, Cat. Fishes, ed. Gray, 1854, 77, India.

Family 224. ACANTHURIDÆ. Surgeonfishes; Tangs

Genus 973. ACANTHURUS Forskål. Surgeonfishes.

Acanthurus Forskål, Descr. Anim., 1775, 59 (*Chætodon unicornis*), etc.

Harpurus (Forster) Gmelin, Enchiridion, 1778, 84 (*H. fasciatus* Forster).

Aspisurus Lacépède, Hist. Nat. Poiss., IV, 1803, 556 (*Chætodon sohar* Forskål).

Aconurus Gronow, Cat. Fishes, 1854, 190 (*Teuthis hepatus* Linnæus).

Rhombotides (Klein) Bleeker, Neder. Tijdschr. Dierk., I (*Chætodon triostegus* Linnæus).

Subgenus HARPURUS Forster.

2809. *Acanthurus triostegus* (Linnæus).

South Pacific, from New Zealand and Australia to the Revillagigedos.

Chætodon triostegus Linnæus, Syst. Nat., Ed. X, 1758, 274, India.

Harpurus fasciatus Forster, Enchiridion, 1778, 84, South Seas.

Acanthurus zebra Lacépède, Hist. Nat. Poiss., III, 1802, 548, fig. 3, no locality.

Chætodon couaga Lacépède, loc. cit. IV, 1802, 727, no locality.

Teuthis australis Gray, in King, Surv. Australia, II, 1826, 435.

Acanthurus hirudo Bennett, Ceylon Fishes, 1851, 11, pl. 11, Ceylon.

Acanthurus subarmatus Bennett, Whaling Voyage, II, 1840, 278, Raiatea, Society Islands.

Subgenus TEUTHIS.

2810. *Acanthurus cæruleus* Bloch and Schneider. Barbero; Blue surgeon; Blue tang.

West Indies, Key West, and Bermuda to Bahia, north to Woods Hole, Mass.

Acanthurus cæruleus Bloch and Schneider, Syst. Ichth., 1801, 214, Carolinas; Habana; Jamaica; after Catesby, Parra, and Browne.

Acanthurus broussoneti Desmarest, Prem. Dec. Ichth., 1823, 26, Cuba.

Acanthurus brevis Poey, Memorias, II, 1861, 207, Antilles and Habana.

Aconurus cæruleatus Poey, Enumeratio, 1875, 69, Cuba; young.

2811. *Acanthurus hepatus* (Linnæus). Common surgeon; Doctor fish; Lancel fish; Tang; Tangbarbero; Saigneur; Barbero.

West Indian fauna; Florida to Bahia; Key West and Cuba; Charleston; north to Woods Hole, Mass.

Teuthis hepatus Linnæus, Syst. Nat., Ed. XII, 1766, 507, Carolinas.

Chætodon chirurgus Bloch, Naturgesch. Ausl. Fische, III, 1787, 99, Martinique; on drawing by Plumier.

Acanthurus phlebotomus Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1835, 129 (176), pl. 287 Martinique, Brazil, Habana, New York.

Aconurus fuscus Gronow, Fish., ed. Gray, 119, 1854, West Indies; young.

Aconurus carneus Poey, Memorias, II, 1860, 207, Cuba.

2812. *Acanthurus bahianus* Castelnau. *Barbero; Ocean tang.*
West Indies to Brazil.
Acanthurus bahianus Castelnau, Anim. Amer. Sud, Poiss., 1855, 24,
Pl. XI, fig. 1, Bahia.
Acanthurus trachus Poey, Memorias, II, 1861, 208, Cuba.
Acronurus nigraculus Poey, Enumeratio, 1875, 69, Cuba.
2813. *Acanthurus crestonis* (Jordan and Starks). *Barbero negro.*
West coast of Mexico and Central America, Mazatlan to Panama.
Teuthus crestonis Jordan and Starks, Proc. Calif. Acad. Sci., ser. 2,
1895, V, 485, pl. 47, Mazatlan.
2814. *Acanthurus aliala* Lesson. *Philosophe.*
East Indies to the west coast of Mexico; abundant at Socorro and
Clarion Islands.
Acanthurus aliala Lesson, Voy. Coquille, Zool., II, 1830, 150, Oualan.
Acanthurus glaucopareus Cuvier and Valenciennes, Hist. Nat. Poiss.,
X, 1835, 139 (190), Oualan; Molucca.
- Genus 974. *XESURUS* Jordan and Evermann.
Xesurus Jordan and Evermann, Check List, 1895, 421 (*Prionurus*
punctatus Gill).
2815. *Xesurus punctatus* (Gill). *Cochinito.*
Cape San Lucas; Mazatlan; Creston Island.
Prionurus punctatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 242,
Cape San Lucas.
2816. *Xesurus clarionis* Gilbert and Starks.
Galapagos Islands.
Xesurus clarionis Gilbert and Starks, Proc. U. S. Nat. Mus., XIX,
1896, 437, 445, Clarion Island, Galapagos Archipelago.
- Genus 975. *CTENOCHÆTUS* Gill.
Ctenodon Bonaparte, in Klunzinger, Fische Rothen Meeres, 1884,
I Theil, 54 (*Acanthurus strigosus* Bennett); name preoccupied.
Ctenochætus Gill, Proc. U. S. Nat. Mus., VII, 1884, 277 (*Acanthurus*
strigosus Bennett).
2817. *Ctenochætus striatus* (Quoy and Gaimard). *Kale.*
South seas, recorded from Cocos Island.
Acanthurus striatus Quoy and Gaimard, Voyage l'Uranie, Zool., 1824,
373, pl. 63, fig. 3, Guam.
Acanthurus strigosus Bennett, Zool. Jour., IV, XIII, Art. III, 1828, 41,
Honolulu.
Acanthurus ctenodon Cuvier and Valenciennes, Hist. Nat. Poiss., X,
1835, 178 (241), pl. 289, the Carolines and New Guinea.

Order RR. CATAPHRACTI

Family 225. SCORPÆNIDÆ. Rockfishes and scorpionfishes

- Genus 976. *SEBASTES* Cuvier. *Rosefishes.*
Sebastes Cuvier, Règne Anim., ed. 2, II, 1829, 166 (*Perca norvegica*
Ascanius=*Perca marina* Linnaeus).
Eusebastes Sauvage, Nouv. Archives Mus. Paris, 1878, (2), 1, 1421
(*Perca norvegica* Ascanius).
2818. *Sebastes marinus* (Linnaeus). *Rosefish; Redfish, Snapper, Hemdurgan;*
Red seafish; Norway haddock.
North Atlantic, both coasts, south to Faroe Islands; Norway; Maine;
in deep water off coast of middle New Jersey.
Perca marina Linnaeus, Syst. Nat., Ed. X, 1758, 290, Norway.
Perca norvegica Ascanius, Icones Rev. Nat., II, 1772, 7, Norway.
Holocentrus sanguineus Faber, Natur. der Fische Islands, 1829, 126,
Iceland.
Sebastes fasciatus Storer, Proc. Bost. Soc. Nat. Hist., V, 1854, 31,
Provincetown, Mass.
Sebastes regulus Ekstrom, Skand. Fish., Ed. I, 197, plate only.

- 2819. *Sebastes viviparus* Krøyer.**
North Atlantic, in shallow bays, more southerly not certainly recorded from America.
Sebastes viviparus Krøyer, Naturh. Tidsskr., I, 1845, 275, Norway, in shallow water.
- Genus 977. SEBASTOLOBUS Gill.**
Sebastolobus Gill, Rept. Smithson. Inst., 1880, 375 (*Sebastes macrochir* Günther).
- 2820. *Sebastolobus altivelis* Gilbert.**
Aleutian Islands, Alaska, and in deep water off the coast of California.
Sebastolobus altivelis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 410, pl. 23, south of Alaska Peninsula, at Albatross station 3338, in 625 fathoms.
- 2821. *Sebastolobus alascanus* Bean.**
Bering Sea; Pacific coast of Alaska, Washington, Oregon, and California, in deep water; recently taken in San Pedro.
Sebastolobus alascanus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 44, off Trinity Islands, Alaska, at Albatross station 2853, 56° N., 154° W., in 159 fathoms.
- Genus 978. EMMELAS Jordan and Evermann.**
Emmelas Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1773 (*Sebastes glaucus* Hilgendorf).
- 2822. *Emmelas glaucus* (Hilgendorf).**
Hokkaido, north to Bering Island.
Sebastes glaucus Hilgendorf, Sitzber. Ges. Naturf. Freunde, Berlin, 1880, 172, Hokkaido, north Japan; no description given.
- Genus 979. SEBASTODES⁸⁸ Gill. Rockfishes.**
Sebastes Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 165 (*Sebastes paucispinis* Ayres).
- 2823. *Sebastes paucispinis* (Ayres). Bocaccio; Mérou; Jack.**
San Diego to Barclay Sound, British Columbia; common south. A valued food fish.
Sebastes paucispinis Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 6, San Francisco.
- 2824. *Sebastes jordani* Gilbert.**
Coast of California in deep water.
Sebastes jordani Gilbert, Rept. U. S. Fish Com., XIX, 1893, 466, coast of California, at Albatross station 2935, 3103, and 3114, in 64 to 124 fathoms.
- 2825. *Sebastes goodei* Eigenmann and Eigenmann.**
San Diego to San Francisco; not common.
Sebastes goodei Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., III, 1890, 12, San Diego, Calif.
- Genus 980. SEBASTOSOMUS Gill.**
Sebastosomus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 59 (*Sebastes melanops* Girard).
Primospina Eigenmann and Beeson, Am. Nat., XXVII, 669 (*Sebastichthys mystinus* Jordan and Gilbert).
- Subgenus PRIMOSPINA Eigenmann and Beeson.**
- 2826. *Sebastosomus flavidus* (Ayres). Yellowtail rockfish.**
San Diego to San Francisco; common northward.
Sebastes flavidus Ayres, Proc. Calif. Acad. Nat. Sci., ser. I, II, 1862, 209, fig. 64, San Francisco.

⁸⁸ The 13-spined allies of *Sebastes* constitute a most striking feature of the fauna of the north Pacific, the species being very numerous on the rocky coast of California, Alaska, and Japan, a few also in Chile. The species are in general well defined, showing little individual variation and no sex changes. It is extremely difficult, however, to divide them into genera, the various traits being intertangled to a remarkable degree. Yet such species as *paucispinis*, *rosaceus*, and *microcinctus* could hardly stand in one genus in a natural system. It may be noted that the group called *Sebastes* is the simplest in organization and also the oldest, four well-marked species related to *Sebastes paucispinis* occurring in the upper Miocene of California.

2827. *Sebastosomus serranoides* (Eigenmann and Eigenmann).
Cortez Banks, off San Diego.
Sebastodes serranoides Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., III, 1890, 36, Cortez Banks.
2828. *Sebastosomus melanops* (Girard).
Pacific coast of America, from Monterey to Sitka.
Sebastodes melanops Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 135, Astoria, Oreg.
Sebastosomus simulans Gill, loc. cit., XVI, 1864, 147, Cape Flattery.
2829. *Sebastosomus mystinus* (Jordan and Gilbert). *Pêche prêtre; Priestfish; Black rockfish*.
Pacific coast of America, from Aleutian Islands to San Diego.
Sebastichthys mystinus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 455, San Francisco.
2830. *Sebastosomus ciliatus* (Tilesius).
Coast of Alaska.
Epinephelus ciliatus Tilesius, Mem. Acad. Sci. St. Petersburg, IV, 1810, 474, Aleutian Islands.
Perca variabilis Pallas, Zoogr. Rosso-Asiat., III, 1811 (1831), 241, Aleutian Islands; including *S. aleuticus*.
- Genus 981. *ROSICOLA* Jordan and Evermann.
Rosicola Jordan and Evermann, Check List, 1895, 429 (*Sebastosomus pinniger* Gill).
2831. *Rosicola babcocki* (Thompson).
Coast of Alaska and British Columbia, common.
Sebastodes babcocki W. F. Thompson, Fisheries British Columbia, 1914, Middleton Island, Alaska.
2832. *Rosicola pinniger* (Gill). *Floma; Orange rockfish*.
Pacific coast of America, from San Diego to Puget Sound.
Sebastosomus pinniger Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 147, San Francisco.
2833. *Rosicola miniatus* (Jordan and Gilbert). *Rasciera; Rasher*.
Pacific coast of America, from San Francisco to San Diego.
Sebastichthys miniatus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 70, Monterey, Calif.
2834. *Rosicola alexandri* (Evermann and Goldsborough).
Monterey Bay.
Sebastodes alexandri Evermann and Goldsborough, Proc. U. S. Nat. Mus., XXXI, 1907, 651, Monterey Bay in 65 fathoms.
- Genus 982. *ZALOPYR* Jordan and Evermann.
Zalopyr Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1795 (*Sebastodes aleutianus* Jordan and Evermann).
2835. *Zalopyr aleutianus* (Jordan and Evermann).
Alaska; Aleutian Islands.
Sebastodes aleutianus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1795, Shelikof Straits, Alaska.
2836. *Zalopyr atrorubens* (Gilbert).
Monterey, Calif.
Sebastodes atrorubens Gilbert, in Jordan and Evermann, Check List, 1895, 429, and Fishes North and Mid. Amer., 1898, 1797, San Francisco.
2837. *Zalopyr atrovirens* (Jordan and Gilbert). *Garrupa*.
Coast of California, from San Diego, to San Francisco.
Sebastichthys atrovirens Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 289, Monterey, Calif.
- Genus 983. *ACUTOMENTUM*⁶⁹ Eigenmann and Beeson.
Acutomentum Eigenmann and Beeson, Amer. Nat., XXVII, 1893, 669 (*Sebastes ovalis* Ayres).

⁶⁹ As here arranged, this is far from a natural assemblage.

2838. *Acutomentum ovale* (Ayres). *Viuva*; *Widowfish*.
Pacific coast of United States from San Diego to San Francisco.
Sebastes ovalis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1862, 209-212,
fig. 65, San Francisco.
2839. *Acutomentum prorigerum* (Jordan and Gilbert).
San Diego to San Francisco.
Sebastichthys proriger Jordan and Gilbert, Proc. U. S. Nat. Mus., III,
1880, 327, Monterey, Calif.
2840. *Acutomentum brevispine* (Bean).
Coast of Alaska.
Sebastichthys proriger var. *brevispinis* Bean, Proc. U. S. Nat. Mus.,
VI, 1883, 359, Hassler Harbor, Alaska.
2841. *Acutomentum clavilatum* (Starks).
Puget Sound.
Sebastes clavilatus Starks, Ann. Carnegie Mus., VII, 1911, 181, San
Juan Islands, Puget Sound.
2842. *Acutomentum entomelas* (Jordan and Gilbert).
Pacific coast of America, from Port Harford to Monterey Bay.
Sebastichthys entomelas Jordan and Gilbert, Proc. U. S. Nat. Mus.,
III, 1880, 142, Monterey.
2843. *Acutomentum wilsoni* (Gilbert).
Monterey Bay.
Sebastes wilsoni Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 333,
pl. 16, fig. 8, Monterey Bay.
2844. *Acutomentum rufum* (Eigenmann and Eigenmann).
Pacific coast of America, near San Diego; Cortez Banks.
Sebastes rufus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci.,
2d ser. III, 1890, 13, Point Loma; Cortez Banks.
2845. *Acutomentum macdonaldi* Eigenmann and Beeson.
San Diego, Calif.
Acutomentum macdonaldi Eigenmann and Beeson, Am. Nat., XXVII,
1893, 669, San Diego, Calif.
2846. *Acutomentum eigenmanni* (Cramer).
Coast of California.
Sebastes eigenmanni Cramer, Proc. Calif. Acad. Sci., VI, 1896, 239,
Pl. XLII, Monterey, Calif.
2847. *Acutomentum hopkinsi* (Cramer).
Monterey Bay, Calif.
Sebastes hopkinsi Cramer, Proc. Calif. Acad. Sci., V, 1895, 594, with
plate, Monterey Bay, Calif.
2848. *Acutomentum alutum* (Gilbert).
Pacific coast of America, from Bering Sea to Santa Barbara.
Sebastichthys alutus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 76,
Santa Barbara Islands, Calif.
- Genus 984. **ROSEBASTES** Jordan and Evermann.
Eosebastes Jordan and Evermann, Check List, 1895 430 (*Sebastichthys
aurora* Gilbert).
2849. *Eosebastes aurora* (Gilbert).
Santa Barbara Islands, in deep water.
Sebastichthys aurora Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 80,
Santa Barbara Islands, in 266 fathoms.
2850. *Eosebastes melanostomus* (Eigenmann and Eigenmann).
Coast of California to Alaska.
Sebastes melanostomus Eigenmann and Eigenmann, Proc. Calif.
Acad. Sci., 2d ser., III, 1890, 17, Point Loma, near San Diego, Calif.
2851. *Eosebastes introniger* (Gilbert).
Santa Barbara to Bering Sea.
Sebastichthys introniger Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 81,
Santa Barbara Islands.

2852. *Eosebastes saxicola* (Gilbert).
Southern California to southeast Alaska in moderate depth.
Sebastichthys saxicola Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 78,
Santa Barbara Islands at Albatross stations 2893, 2907, and others,
in 44 to 155 fathoms.
2853. *Eosebastes deani* (Starks).
Puget Sound.
Sebastes deani Starks, Ann. Carnegie Mus., VII, 1911, 178, Puget
Sound.
2854. *Eosebastes crameri* (Jordan).
Coast of Oregon.
Sebastes crameri Jordan, Proc. U. S. Nat. Mus., XIX, 1896, 451,
coast of Oregon, at Albatross station 3091.
2855. *Eosebastes swifti* (Evermann and Goldsborough).
Southern Alaska.
Sebastes swifti Evermann and Goldsborough, Bull. U. S. Bur. Fish.,
XXVI, 1906, 285, Yes Bay and Kasaan Bay, Alaska.
2856. *Eosebastes semicinctus* (Gilbert).
Coast of California.
Sebastes semicinctus Gilbert, Proc. U. S. Nat. Mus., XIX, 1896, 449,
Santa Barbara Channel, Calif.
2857. *Eosebastes diploproa* (Gilbert).
Pacific coast from the Coronado Islands northward to Nanaimo,
British Columbia.
Sebastichthys diploproa Gilbert, Proc. U. S. Nat. Mus., XIII, 1890,
79, Coronado Islands at Albatross station 2935, in 124 fathoms.
2858. *Eosebastes emphæus* (Starks).
Puget Sound, in deep water.
Sebastes emphæus Starks, Ann. Carnegie Mus., VII, 1911, 183, about
San Juan Islands, Puget Sound.
- Genus 985. **SEBASTOMUS** Gill.
Sebastomus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 147 (*Sebastes*
rosaceus Girard).
2859. *Sebastomus constellatus* (Jordan and Gilbert). *Spotted rockfish*.
Coast of California, from San Diego to San Francisco.
Sebastichthys constellatus Jordan and Gilbert, Proc. U. S. Nat. Mus.,
III, 1880, 295, Santa Barbara Channel; San Francisco.
2860. *Sebastomus rosaceus* (Girard). *Corsair*.
Pacific coast from San Diego to Puget Sound.
Sebastes rosaceus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1854, 146,
San Diego, San Francisco.
Sebastes helvomaculatus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859,
26, San Francisco.
2861. *Sebastomus rhodochloris* (Jordan and Gilbert). *Flyfish*.
Off Monterey and San Francisco.
Sebastichthys rhodochloris Jordan and Gilbert, Proc. U. S. Nat. Mus.,
III, 1880, 144, Monterey, Calif.
2862. *Sebastomus chlorostictus* (Jordan and Gilbert). *Pesca vermiglia*.
San Diego; Monterey; San Francisco.
Sebastichthys chlorostictus Jordan and Gilbert, Proc. U. S. Nat. Mus.,
III, 1880, 294, Monterey, Calif.
2863. *Sebastomus umbrosus* (Jordan and Gilbert).
Coast of California, Point Conception to Coronado Islands.
Sebastichthys umbrosus Jordan and Gilbert, Proc. U. S. Nat. Mus., V,
1882, 410, Santa Barbara.
Sebastes æreus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci.,
2d ser., III, 1890, 20, Coronado Islands.
2864. *Sebastomus ayresii* (Gilbert and Cramer).
Coast of southern California at Cortez Banks, near San Diego.
Sebastes ayresii Gilbert and Cramer, Proc. U. S. Nat. Mus., XIX,
1896, 450, Cortez Banks, near San Diego, Calif.

2865. *Sebastomus eos* (Eigenmann and Eigenmann).
Coast of California at Point Loma, near San Diego.
Sebastes eos Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., III, 1890, 18, Point Loma, near San Diego, Calif.
2866. *Sebastomus gillii* (Eigenmann and Eigenmann).
Coast of California at Point Loma, near San Diego.
Sebastes gillii Eigenmann and Eigenmann, Amer. Nat., XXV, February, 1891, 154, Point Loma, near San Diego, Calif.
2867. *Sebastomus rupestris* (Gilbert).
Santa Barbara Islands northward to southeast Alaska.
Sebastichthys rupestris Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 76, Santa Barbara Islands, at Albatross station 2946, in 150 fathoms.
- Genus 986. SEBASTOPYR⁷⁰ Jordan and Evermann.
Sebastopyr Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 506 (*Sebastes ruberrimus* Cramer).
2868. *Sebastopyr ruberrimus* (Cramer). *Red rockfish; Tambor.*
Pacific coast from San Diego to Puget Sound and southeastern Alaska.
Sebastes ruber of early writers, not of Ayres.
Sebastes ruberrimus Cramer, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 597, pls. 62, 63, and 69, Monterey, Calif.
- Genus 987. HISPANISCUS Jordan and Evermann.
Hispaniscus Jordan and Evermann, Check List, 1895, 431 (*Sebastichthys rubrivinctus* Jordan and Gilbert).
2869. *Hispaniscus rubrivinctus* (Jordan and Gilbert). *Spanish flag.*
California coast from San Diego to Monterey; not rare south. A most brilliantly colored fish.
Sebastichthys rubrivinctus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 291, Santa Barbara Channel.
2870. *Hispaniscus levis* (Eigenmann and Eigenmann).
Coast of California, from San Diego to Monterey.
Sebastichthys levis Eigenmann and Eigenmann, West American Scientist, Nov. 9, 1889, 128, San Diego, Calif.
2871. *Hispaniscus zacentrus* (Gilbert).
Coast of California.
Sebastichthys zacentrus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 77, Santa Barbara Islands, at Albatross station 2893 and 2946, in 145 and 150 fathoms.
2872. *Hispaniscus elongatus* (Ayres). *Reina.*
California coast, from San Diego to San Francisco.
Sebastes elongatus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1858-1862 (Oct. 17, 1859), 26, San Francisco.
2873. *Hispaniscus sinensis* (Gilbert).
Gulf of California.
Sebastichthys sinensis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 81, Gulf of California, at Albatross station 3015 in 145 fathoms.
- Genus 988. AUCTOSPINA Eigenmann and Beeson.
Auctospina Eigenmann and Beeson, Amer. Nat., XXVII, 1893, 670 (*Sebastes auriculatus* Girard).
2874. *Auctospina auriculata* (Girard). *Brown rockfish.*
Pacific coast of America, from Cape Mendocino to Cedros Island; common.
Sebastes auriculatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 131, 146, San Francisco.
Sebastes ruber Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854 (Sept. 11), 7, San Francisco.
Sebastes ruber parvus Ayres, loc. cit., I, 1854 (Sept. 11), 7, San Francisco.

⁷⁰ This group is allied to *Sebastomus*, but of much coarser build, the cranial spines in the adult rough with blunt spinules.

2875. *Auctospina dallii* (Eigenmann and Beeson).
Pacific coast of America, from San Francisco to Puget Sound; perhaps a variation of *A. auriculata*.
Pteropodus dallii Eigenmann and Beeson, Amer. Nat., XXVIII, January, 1894, 66, San Francisco.
- Genus 989. **PTEROPODUS** Eigenmann and Beeson.
Pteropodus Eigenmann and Beeson, Amer. Nat., XXVII, 1893, 670 (*Sebastichthys maliger* Jordan and Gilbert); not *Pteropus*, a genus of birds.
2876. *Pteropodus rastrelliger* (Jordan and Gilbert). *Grass rockfish*.
California coast, from San Diego to San Francisco.
Sebastichthys rastrelliger Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 296, Monterey, Calif.
2877. *Pteropodus caurinus* (Richardson).
Pacific coast of America from Puget Sound to Sitka.
Sebastes caurinus Richardson, Voy. *Sulphur*, Ichthyology, 1845, 77, pl. 41, fig. 1, Sitka.
2878. *Pteropodus vexillaris* (Jordan and Gilbert).
California coast, from San Diego to Cape Mendocino, replacing *caurinus* southward.
Sebastichthys vexillaris Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 292, Santa Barbara Channel; San Francisco.
2879. *Pteropodus maliger* (Jordan and Gilbert). *Yellow-backed rock cod*;
Brown rock cod.
Pacific coast of America, from Monterey to Sitka; not rare.
Sebastichthys maliger Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 322, San Francisco.
2880. *Pteropodus gilberti* (Cramer).
Coast of California.
Sebastodes gilberti Cramer, Proc. Calif. Acad. Sci., VI, 1896, 241, Pl. XLIII, San Francisco.
2881. *Pteropodus carnatus* (Jordan and Gilbert). *Flesh-colored rock cod*.
California coast, from San Diego to Point Reyes, replacing *P. chrysomelas* in shallow water.
Sebastichthys carnatus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 73, Monterey, Calif.
2882. *Pteropodus chrysomelas* (Jordan and Gilbert). *Black and yellow rock cod*.
Pacific coast of America, from Puget Sound to San Diego; common in moderate depths.
Sebastichthys chrysomelas Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 455, 465, Monterey, Calif.
2883. *Pteropodus nebulosus* (Ayres). *Yellow-spotted rock cod*.
Pacific coast; southeastern Alaska; Puget Sound region, Vancouver to Point Conception; common.
Sebastes nebulosus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854 (Sept 11), 5, San Francisco.
Sebastes fasciatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 146, Presidio, California; not of Storer.
Sebastichthys fasciolaris Lockington, Proc. U. S. Nat. Mus., III, 1880, 297, San Francisco.
- Genus 990. **SEBASTOCARUS** ⁷¹ Jordan and Evermann.
Sebastocarus Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 507 (*Sebastichthys serriceps* Jordan and Gilbert).
2884. *Sebastocarus serriceps* (Jordan and Gilbert). *Treefish*.
Coast of California, from Point Reyes to Cedros Island; common south.
Sebastichthys serriceps Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 38, Santa Barbara and Santa Catalina Islands, Calif.

⁷¹ This genus is nearly allied to *Sebastichthys* in the more compressed body and especially in the short and high ridges on the head, all of which are free from serrations or accessory tubercles.

Genus 991. SEBASTICHTHYS Gill.

Sebastichthys Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 278, 329
(*Sebastes nigrocinctus* Ayres).

2885. *Sebastichthys nigrocinctus* (Ayres). Black-banded rock cod.

Monterey Bay to Naha Bay, Alaska.

Sebastes nigrocinctus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 25,
fig 6, San Francisco.

Genus 992. METZELAARIA Jordan.

Metzelaaria Jordan, Class. Fishes, 1923, 209 (*Scorpæna tridecimspinosa*
Metzelaar).

2886. *Metzelaaria tridecimspinosa* (Metzelaar).

Caribbean Sea.

Scorpæna tridecimspinosa Metzelaar, Tropisch Atlantische Visschen,
1919, 146, fig. 44, Bonaire, Aruba.

Genus 993. SCORPÆNODES Bleeker.

Scorpænodes Bleeker, Nat. Hist. Ned. Ind., XIII, 1857, 371 (*Scorpæna*
polylepis Bleeker).

Sebastopsis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 278 (*Scorpæna*
polylepis Bleeker).

Sebastes Sauvage, Ann. Sci. Nat. Zool., V, 1873, 1 (*Sebastes minutus*
Cuvier); not *Sebastes* Gill, 1861.

2887. *Scorpænodes xyris* (Jordan and Gilbert).

West coast of Mexico.

Sebastopsis xyris Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
369, Cape San Lucas, Lower California.

Genus 994. HELICOLENUS Goode and Bean.

Helicolenus Goode and Bean, Oceanic Ichth., 1896, 248 (*Scorpæna*
dactylopterus De la Roche).

2888. *Helicolenus dactylopterus* (De la Roche). Seran imperial; Fanegal;

Cardonmera; *Scorfanudu*; *Funal*; *Crabra*; *Redfish*.

Deep waters of Atlantic off both coasts from Narragansett and Chesapeake Bays, and to coasts of Italy and Spain.

Scorpæna dactyloptera De la Roche, Ann. Mus., XIII, 1809, pl. 22,
fig. 2, Ivica, Barcelona.

Sebastes imperialis Cuvier and Valenciennes, Hist. Nat. Poiss., IV,
1829, 246 (336), Nice.

Helicolenus maderensis Goode and Bean, Oceanic Ichth., 1895, 250,
Madeira, and in the Gulf Stream.

Genus 995. SCORPÆNA (Artedi) Linnæus.

Scorpæna (Artedi) Linnæus, Syst. Nat., ed. X, 1758 266 (*S. porcus*
Linnæus).

Parascorpæna Bleeker, Versl. Ak. Amsterdam., IX, pt. 3, 1876, 295
(*Scorpæna picta* Kuhl and van Hasselt in Cuvier and Valenciennes).

Pseudosebastes Sauvage, Nouv. Arch. Mus. Hist. Nat. Paris, II, 1878
(*Sebastes bougainvillei* Cuvier and Valenciennes).

2889. *Scorpæna atlantica* Nichols and Breder.

Coast of Texas

Scorpæna russula atlantica Nichols and Breder, Proc. Biol. Soc. Wash.,
XXXVII, 1924 (Feb. 21), 21, Pl. VII, off Galveston, Tex

2890. *Scorpæna colesi* Nichols.

Coast of North Carolina.

Scorpæna colesi Nichols, Bull. Amer. Mus. Nat. Hist., XXXIII, 1914,
537, fig. 1, Cape Lookout, N. C.

2891. *Scorpæna plumieri* Bloch. Rescacio; Poison grouper; Plumier's pigfoot.

West Indies and Brazil, north to Florida; rarely northward to Woods Hole.

Scorpæna plumieri Bloch, Nya. Handl. Stockh., X, 1789, 234, Martinique.

Scorpæna bufo Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829,
224 (306), Martinique.

Scorpæna rescacio Poey, Memorias, II, 1861, 169, Cuba.

Apistes exul Gosse, Nat. Sojourn Jamaica, 1851, 207, Jamaica.

2892. *Scorpæna mystes* Jordan and Starks. *Lapon*.
Coast of Mexico
Scorpæna mystes Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser. V,
1895, 491, pl. 52, Mazatlan, Mexico.
2893. *Scorpæna grandicornis* Cuvier and Valenciennes *Lionfish, Poison
grouper; Sculpin*.
Florida Keys to Brazil, in the Gulf Stream to Woods Hole, Mass.
Scorpæna grandicornis Cuvier and Valenciennes, Hist. Nat. Poiss., IV,
1829, 227 (309), pl. 86, Martinique, Porto Rico, Habana, San Do-
mingo.
2894. *Scorpæna brasiliensis* Cuvier and Valenciennes.
Atlantic coast, from Charleston to Rio Janeiro.
Scorpæna brasiliensis Cuvier and Valenciennes, Hist. Nat. Poiss., IV,
1829, 224 (305), Brazil.
Scorpæna stearnsi Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 421,
Pensacola, Fla.
2895. *Scorpæna inermis* Cuvier and Valenciennes.
West Indies, north to Florida.
Scorpæna inermis Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829,
228 (311), Martinique
Scorpæna occipitalis Poey, Memorias, II, 1861, 171, Cuba.
Scorpæna calcarata Goode and Bean, Proc. U. S. Nat. Mus., V, 1882,
422, Clearwater Harbor, Fla.
2896. *Scorpæna sonoræ* Jenkins and Evermann.
Guaymas, Gulf of California, and Santa Margarita Island.
Scorpæna sonoræ Jenkins and Evermann, Proc. U. S. Nat. Mus., XI,
1888, 150, Guaymas, Sonora.
2897. *Scorpæna guttata* Girard. *Scorpene; Scorpion; Sculpin*.
Coast of California, from Monterey southward to Cedros Island.
Scorpæna guttata Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 145,
Monterey.
2898. *Scorpæna russula* Jordan and Bollman.
Pacific coast of Colombia.
Scorpæna russula Jordan and Bollman, Proc. U. S. Nat. Mus., XII,
1889, 165, Pacific coast of Colombia.
2899. *Scorpæna histrio* Jenyns.
Pacific coast of America, Panama to Juan Fernandez; Galapagos Islands.
Scorpæna histrio Jenyns, Voy. *Beagle*, Fishes, 1842, 35, pl. 8, Chatham
Island, Galapagos Archipelago.
2900. *Scorpæna pannosa* Cramer.
Panama.
Scorpæna pannosa Cramer, Proc. U. S. Nat. Mus., XIX, 1896, 446,
with plate, Panama.
2901. *Scorpæna albofasciata* Metzelaar.
Caribbean Sea.
Scorpæna albofasciata Metzelaar, Tropisch Atlantische Vissch., 1919,
145, fig. 43; Bonaire, Arubi.
2902. *Scorpæna albifimbria* Evermann and Marsh.
Porto Rico.
Scorpæna albifimbria Evermann and Marsh, Bull. U. S. Fish Com., XX,
1900, 275, fig. 82, Culebra Island; Porto Rico.
2903. *Scorpæna nuttingi* Evermann and Seale.
West Indies (Barbados).
Scorpæna nuttingi Evermann and Seale, University of Iowa Studies
Nat. Hist., X, 1924, 14, 39, Carlisle Bay, Barbados.
2904. *Scorpæna agassizii* Goode and Bean.
Mid-ocean east of Cuba.
Scorpæna agassizii Goode and Bean, Oceanic Ichth., 1896, 247, fig. 243,
latitude 32° 13' N., longitude 39° 10' W.

2905. *Scorpaena cristulata* Goode and Bean.
Atlantic Ocean, off coast of Georgia; perhaps also from northern Spain.
Scorpaena cristulata Goode and Bean, Oceanic Ichth., 1896, 247, fig. 242, off Georgia.
Scorpaena echinita Koehler, Ann. Univ. Lyons, XXVI, 1896, 474, pl. 27, figs 4 and 6, Bay of Biscay, in 700 fathoms.
- Genus 996. **MERINTHE** Snyder.
Merinthe Snyder, Bull. U. S. Fish Com., XXII, 1902, 536 (*Sebastes macrocephalus* Sauvage).
2906. *Merinthe nematophthalma* (Günther).
West Indies, exact locality uncertain.
Sebastes nematophthalmus Gunther, Cat., II, 1861, 99, West Indies.
- Genus 997. **PONTINUS** Poey.
Pontinus Poey, Memorias, II, 1861, 172 (*P. castor* Poey).
2907. *Pontinus castor* Poey.
Cuba.
Pontinus castor Poey, Memorias, II, 1861, 173, Habana, Cuba.
2908. *Pontinus furcirhinus* Garman.
Eastern Pacific in deep water.
Pontinus furcirhinus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 51, Pl. VII, numerous Albatross stations off Panama.
2909. *Pontinus pollux* Poey.
Cuba.
Pontinus pollux Poey, Memorias, II, 1861, 174, Habana, Cuba.
2910. *Pontinus rathbuni* Goode and Bean.
Coast of North Carolina, probably in the Gulf Stream.
Pontinus rathbuni Goode and Bean, Oceanic Ichth., 1895, 255, fig. 245, off Cape Hatteras, N. C.
2911. *Pontinus macrolepis* Goode and Bean.
Yucatan.
Pontinus macrolepis Goode and Bean, Oceanic Ichth., 1895, 257, fig. 247, off Yucatan.
2912. *Pontinus longispinis* Goode and Bean.
Coast of western Florida.
Pontinus longispinis Goode and Bean, Oceanic Ichth., 1895, 258, fig. 246, west coast of Florida, lat. 28° 36' N., long. 85° 33' 30" W., in 111 fathoms.
2913. *Pontinus beanorum* Evermann and Marsh.
Porto Rico.
Pontinus beanorum Evermann and Marsh, Bull. U. S. Fish Com., XX, pt. 1, 1900, 279, fig. 85, San Juan Harbor, Porto Rico.
2914. *Pontinus sierra* (Gilbert).
Gulf of California.
Scorpaena (Sebastoplus) sierra Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 82, Gulf of California, at Albatross stations 2996 and 3011 in 112 and 71 fathoms.
2915. *Pontinus bergii* (Evermann and Marsh).
Porto Rico.
Scorpaena bergii Evermann and Marsh, Bull. U. S. Fish Com., XX, pt. 1, 1900, 276, fig. 83, Mayaguez, Culebra; perhaps not a *Pontinus*.
- Genus 998. **SETARCHES** Johnson.
Setarches Johnson, Proc. Zool. Soc. Lond., 1862, 177 (*S. guntheri* Johnson).
2916. *Setarches parmatatus* Goode.
Coasts of Rhode Island, North Carolina, and western Florida, in deep water.
Setarches parmatatus Goode, Proc. U. S. Nat. Mus., III, 1880, 480, off coast of Rhode Island.

Family 226. ANOPILOPOMIDÆ. Skilfishes; Sablefishes

Genus 999. ANOPILOPOMA Ayres.

Anoplopoma Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 27 (*A. merlangus* Ayres=*Gadus fimbria* Pallas).

Scombrocottus Peters, Monatsber. Akad. Wiss. Berlin, 1872, 569 (*S. salmonicus* Peters=*Gadus fimbria* Pallas).

2917. *Anoplopoma fimbria* (Pallas). Beshow, Coalfish; Skil; Black cod; Sablefish.

Unalaska to San Diego, common northward; a valued food fish in Alaska when it becomes fat.

Gadus fimbria Pallas, Zoogr. Rosso-Asiat., III, 1811, 200, probably Aleutian Islands.

Anoplopoma merlangus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 27, San Francisco market.

Scombrocottus salmonicus Peters, Monatsber. Akad. Wiss. Berlin, 1872, 569, Vancouver Island.

Family 227. ERILEPIDÆ. Fat priests

Genus 1000. ERILEPIS Gill.

Myriolepis Lockington, Proc. U. S. Nat. Mus., III, 1880, 248 (*Myriolepis zonifer* Lockington); name preoccupied in fossil fishes.

Erilepis Gill, Science, Jan. 26, 1894, 54 (*Myriolepis zonifer* Lockington).

Ebisus Jordan and Snyder, Jour. Coll. Sci. Tokyo, XV, 1901, 308 (*E. sagamius* Jordan and Snyder).

2918. *Erilepis zonifer* (Lockington). Aburabozu; Fat priest.

Coast of Japan to Alaska; once taken at Monterey Bay, Calif.

Myriolepis zonifer Lockington, Proc. U. S. Nat. Mus., III, 1880, 248, Monterey.

Ebisus sagamius Jordan and Snyder, Jour. Coll. Sci., Tokyo, XV, 1901, 308, Misaki, Japan.

Family 228. HEXAGRAMMIDÆ. Greenlings

Genus 1001. HEXAGRAMMOS Steller.

Hexagrammos Steller, in Tilesius, Act. Ac. Petrop., II, 1810, 335 (*H. stellera* Tilesius=*H. asper* Tilesius).

Labrax Pallas, Mém. Acad. Sci. Petersb., II, 1810, 382 (*L. lagocephalus* Pallas), not of Klein nor of Cuvier.

Lebius (Steller MS) Pallas, Zoogr. Rosso-Asiat., III, 1811, 279 (*Labrax superciliosus* Pallas).

Chirus Steller, in Pallas, loc. cit., 279 (*Labrax superciliosus* Pallas).

Octogrammus Bleeker, Versl. Akad. Amsterdam (2), VIII, 1874, 370 ("Labrax octogrammus Schlegel"=*Octogrammus pallasi* Bleeker).

Acantholebius Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 166 (*Chiropsis nebulosus* Girard=*Hexagrammos stellera* Tilesius); specimen with broken dorsal spines.

Grammatopleurus Gill, loc. cit., XIII, 1861, 166 (*Labrax lagocephalus* Pallas).

2919. *Hexagrammos decagrammus* (Pallas). Rock trout; Boregat; Bodieron North Pacific, from Point Conception to southern Alaska; common south.

Labrax decagrammus Pallas, Mém. Acad. Petersb., II, 1810, 386, pl. 22, fig. 2, Elias Bay.

Chirus guttatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 132, San Francisco.

Grystes lineatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 9, Bay of San Francisco.

Chiropsis constellatus Girard, U. S. Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 42, San Francisco.

Chirus maculo-seriatus Lockington, Proc. U. S. Nat. Mus., III, 1880, 55, San Francisco.

2920. *Hexagrammos octogrammus* (Pallas). *Alaska greenfish*.
Aleutian Islands, westward to Petropaulski and Robben Island.
Labrax octogrammus Pallas, Zoogr. Rosso-Asiat., III, 1811, 283, Kamchatka and Aleutian Islands; doubtful species.
Chirus ordinatus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 28, Unalaska.
Octogrammus pallas Bleeker, Versl. Akad. Amsterdam, VI, 1870; after Pallas
2921. *Hexagrammos stelleri* Tilesius. *Greenling*.
Kamchatka and Unalaska to San Francisco; abundant in Puget Sound and northward
Hexagrammos stelleri Tilesius, Mém. Acad. St. Petersb., II, 1809, 335, Kamchatka.
Hexagrammos asper Steller in Tilesius, loc. cit., 340, pl. 15, 1810, Kamchatka.
Labrus hexagrammus Pallas, loc. cit., 395, pl. 23, fig. 3, 1810, Kamchatka.
Chiropsus nebulosus Girard, U. S. Pac. R. R. Surv., X, Fishes, 1858, 45, Puget Sound and at Fort Steilacoom.
Chirus trigrammus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 29, Unalaska.
2922. *Hexagrammos superciliosus* (Pallas). *Red rock trout*
Bering Sea to Monterey Bay.
Labrax superciliosus Pallas, Mém. Acad. St. Petersb., II, 1810, 388, Unalaska.
Chirus pictus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 132, San Francisco; Humboldt Bay.
Chirus baltas Cope, Proc. Amer. Phil. Soc., XIII, 1873, 28, Captains Harbor, Unalaska.
Hexagrammus scaber Bean, Proc. U. S. Nat. Mus., IV, 1881, 154, Amchitka, Unalaska.
2923. *Hexagrammos lagocephalus* (Pallas).
Western Bering Sea, Kuril Islands; Alaska; Attu.
Labrax lagocephalus Pallas, Mém. Acad. St. Petersb., II, 1810, 384, Kuril Islands.
- Genus 1002. *PLEUROGRAMMUS* Gill. *Atka mackerels*.
Pleurogrammus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 166 (*Labrax monopterygius* Pallas).
2924. *Pleurogrammus monopterygius* (Pallas). *Atka mackerel*; *Atkafish*; *Yellowfish*.
Northern Pacific, common about Atka and other Aleutian Islands east to Unalaska.
Labrax monopterygius Pallas, Mém. Acad. Sci. St. Petersburg, II, 1810, 391, Unalaska.

Family 229. OPHIODONTIDÆ. Cultus cods

- Genus 1003. *OPHIODON* Girard.
Ophiodon Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 133 (*Ophiodon elongatus* Girard).
Oplopoma Girard, loc. cit., VIII, 1856, 135 (*Oplopoma pantherina* Girard=*Ophiodon elongatus* Girard; young).
2925. *Ophiodon elongatus* Girard. *Cultus cod*; *Blue cod*; *Buffalo cod*; *Lang*.
Pacific coast from Alaska to Santa Barbara.
Ophiodon elongatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 133, San Francisco.
Oplopoma pantherina Girard, loc. cit., VIII, 1856, 135, Cape Flattery, Wash.

Family 230. OXYLEBIIDÆ

- Genus 1004. *OXYLEBIUS* Gill.
Oxylebius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 277 (*O. pictus* Gill).
2926. *Oxylebius pictus* Gill.
From Monterey northward to Puget Sound; not rare.
Oxylebius pictus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 278, California; no definite locality given.

Family 231. ZANIOLEPIDÆ

Genus 1005. ZANIOLEPIS Girard.

Zaniolepis Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 202 (*Z. latipinnus* Girard).

2927. *Zaniolepis latipinnis* Girard.

Coast of California, from San Francisco northward.

Zaniolepis latipinnis Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 202, Fort Steilacoom, Puget Sound

Genus 1006. XANTOCLES Jordan.

Xantocles Jordan, Copeia, Oct. 4, 1917, 88 (*Zaniolepis frenatus* Eigenmann).

2928. *Xantocles frenatus* (Eigenmann).

Shore banks of southern California; rare.

Zaniolepis frenatus Eigenmann, West American Scientist, VI, Nov. 9, 1889, 147, Cortez Banks off San Diego.

Family 232. JORDANIIDÆ

Genus 1007. JORDANIA Starks.

Jordania Starks, Proc. Acad. Nat. Sci. Phila., XLVII, 1895, 410 (*J. zonope* Starks).

2929. *Jordania zonope* Starks.

Puget Sound.

Jordania zonope Starks, Proc. Acad. Nat. Sci. Phila., XLVII, 1895, 410, Port Orchard, Puget Sound.

Genus 1008. PARICELINUS Eigenmann and Eigenmann.

Paricelinus Eigenmann and Eigenmann, West Amer. Sci., VI, 1889, 131 (*P. hopliticus* Eigenmann and Eigenmann).

2930. *Paricelinus hopliticus* Eigenmann and Eigenmann.

Cortez Banks, off San Diego.

Paricelinus hopliticus Eigenmann and Eigenmann, West Amer. Sci., VI, Nov. 9, 1889, 131, Cortez Banks.

Genus 1009. ALCIDEA Jordan and Evermann.

Alcidea Jordan and Evermann, Fishes North and Mid. Amer., 1886, 1893 (*Paricelinus thoburni* Gilbert).

2931. *Alcidea thoburni* (Gilbert).

Coast of Oregon.

Paricelinus thoburni Gilbert, Rept. U. S. Fish Com., XIX, 1893, 432, pl. 30, coast of Oregon, in 75 fathoms.

Family 233. ICELIDÆ. Scaled sculpins

Genus 1010. CHITONOTUS Lockington.

Chitonotus Lockington, Proc. U. S. Nat. Mus., IV, 1881, 141 (*C. megacephalus* Lockington = *Artedius pugetensis* Steindachner).

2932. *Chitonotus pugetensis* (Steindachner).

San Francisco to Puget Sound.

Artedius pugetensis Steindachner, Ichth. Beitr., V, 1876, 133, pl. 14, fig. 2, Puget Sound.

Chitonotus megacephalus Lockington, Proc. U. S. Nat. Mus., IV, 1881, 142, off San Francisco.

Genus 1011. TARANDICHTHYS Jordan and Evermann.

Tarandichthys Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 225 (*Icelinus filamentosus* Gilbert).

2933. *Tarandichthys cavifrons* (Gilbert).

Coast of southern California.

Icelinus cavifrons Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 83, off Santa Barbara Islands, at Albatross stations 2907 and 2945, in 44 and 30 fathoms.

2934. *Tarandichthys filamentosus* (Gilbert).
Coast of southern California; Alaska, off St. Mary's Mission; Hood Canal, Puget Sound.
Icelinus filamentosus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 85, off Santa Barbara Islands, in 145 and 55 fathoms.
2935. *Tarandichthys tenuis* (Gilbert).
Coast of southern California, Alaska, Fort Rupert and Gulf of Georgia.
Icelinus tenuis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 86, off Santa Barbara Islands, at *Albatross* stations 2893, 2946, and others, in 45 to 150 fathoms.
- Genus 1012. *ICELINUS* Jordan.
Icelinus Jordan, Cat. Fish. N. Amer., 1885, 110 (*Artedius quadriseriatus* Lockington).
2936. *Icelinus fimbriatus* Gilbert.
Southern California.
Icelinus fimbriatus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 87, off Santa Barbara Islands, at *Albatross* stations 2893 and 2975, in 145 and 36 fathoms.
2937. *Icelinus oculatus* Gilbert.
Coast of southern California.
Icelinus oculatus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 87, off Santa Barbara Islands, at *Albatross* station 2935, in 124 fathoms.
2938. *Icelinus burchami* Evermann and Goldsborough
Loring and Behm Canal, Alaska.
Icelinus burchami Evermann and Goldsborough, Bull. U. S. Bur. Fish., XXVI, 1906, 297, fig. 48, off Loring, Alaska.
2939. *Icelinus borealis* Gilbert.
Coast of Alaska, north and south of Aleutian Islands and in Bristol Bay.
Icelinus borealis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 415, pl. 25, Aleutian Islands and Bristol Bay at *Albatross* stations 3213, 3214, and others, in 11 to 121 fathoms.
2940. *Icelinus quadriseriatus* (Lockington).
Off San Francisco, between Point Reyes and Golden Gate.
Artedius quadriseriatus Lockington, Proc. U. S. Nat. Mus., II, 1879, 330, off San Francisco.
2941. *Icelinus strabo* Starks.
Puget Sound at Port Ludlow, Wash.
Icelinus strabo Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 551, Port Ludlow, Wash.
2942. *Icelinus fuscescens* Gilbert.
Southern California, Santa Barbara Islands to Monterey.
Icelinus fuscescens Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 340, pl. 17, fig. 9, Santa Barbara Islands, depth 260 to 310 fathoms.
- Genus 1013. *ASTROLYTES* Jordan and Starks.
Astrolytes Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 807 (*Artedius fenestralis* Jordan and Gilbert).
2943. *Astrolytes fenestralis* (Jordan and Gilbert).
Puget Sound to Unalaska.
Artedius fenestralis Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 577, Puget Sound.
Artedius asperulus Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 553, Port Ludlow, Wash.
- Genus 1014. *PARASTROLYTES* Hubbs.
Parastrolytes Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 2 (*Artedius notospilotus* Girard).
2944. *Parastrolytes notospilotus* (Girard).
Coast of California from Straits of Fuca southward; abundant at Santa Barbara and Puget Sound.
Artedius notospilotus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 134, Tomales Bay, Calif.

- Genus 1015. **AXYRIAS** Starks.
Axyrias Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 554 (*Axyrias harringtoni* Starks).
2945. **Axyrias harringtoni** Starks.
 Puget Sound in the vicinity of Port Ludlow, Wash.
Axyrias harringtoni Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 554, pl. 74, Port Ludlow, Wash.
- Genus 1016. **ARTEDIUS** Girard.
Artedius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 134 (*Scorpenichthys lateralis* Girard).
2946. **Artedius lateralis** (Girard).
 Pacific coast of North America, from Puget Sound to Port Harford, Calif.
Scorpenichthys lateralis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 145, San Luis Obispo (Port Harford), Calif.
- Genus 1017. **PARARTEDIUS** Hubbs.
Parartedius Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 3 (*P. hankinsoni* Hubbs).
2947. **Parartedius hankinsoni** Hubbs.
 Tidepool, near end of Point Loma, California.
Parartedius hankinsoni Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 171, Feb. 20, 1926, 4, Tidepool near end of Point Loma, Calif.
- Genus 1018. **ALLARTEDIUS** Hubbs.
Allartedius Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 8 (*A. corallinus* Hubbs).
2948. **Allartedius corallinus** Hubbs.
 Tidepool on Point Lobos, Monterey Co., Calif. Only one specimen known.
Allartedius corallinus Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 8, deep low pool on Point Lobos, Monterey Co., Calif.
- Genus 1019. **ARCHAULUS** Gilbert and Burke.
Archaulus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 36 (*A. biseriatus* Gilbert and Burke).
2949. **Archaulus biseriatus** Gilbert and Burke.
 West Bering Sea.
Archaulus biseriatus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 36, Petrel Bank, Bering Sea.
- Genus 1020. **ARTEDIELLUS** Jordan.
Artediellus Jordan, Cat. Fish. N. Amer., 1885, 110 (*Cottus uncinatus* Reinhardt).
2950. **Artediellus uncinatus** (Reinhardt).
 Greenland to Labrador.
Cottus uncinatus Reinhardt, K. Dansk. Vid. Selsk. Natur. Math. Afh., VI, 1833; XLIV, 1837, Greenland.
Artediellus atlanticus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1906, Massachusetts Bay.
2951. **Artediellus pacificus** Gilbert.
 Coast of Alaska; in Bristol Bay, south of Sannak Island, and north of Unalaska.
Artediellus pacificus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 416, Unalaska, Sannak, Bristol Bay, at Albatross station 3216, south of Sannak Island, many other stations in Bristol Bay, and station 3323 north of Unalaska, in 8 to 61 fathoms.
2952. **Artediellus clarki** (Evermann and Goldsborough).
 Bering Sea.
Blennicottus clarki Evermann and Goldsborough, Bull. U. S. Fish Com., XXVI, 1906, 323, fig. 85; Bering Sea. Young, perhaps of *Artediellus pacificus*.

2953. *Artediellus miacanthus* Gilbert and Burke.
Western Bering Sea.
Artediellus miacanthus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 47, fig. 7, east coast of Kamchatka.
2954. *Artediellus camchaticus* Gilbert and Burke.
Western Bering Sea.
Artediellus camchaticus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 46, fig. 6, Avatcha Bay, east coast of Kamchatka.
- Genus 1021. *RUSCARIUS* Jordan and Starks.
Ruscarius Jordan and Starks, Proc. Calif. Acad. Sci., 2d series, V, 1895, 805 (*Ruscarius meanyi* Jordan and Starks).
2955. *Ruscarius meanyi* Jordan and Starks.
Puget Sound, at Port Orchard near Seattle.
Ruscarius meanyi Jordan and Starks, Proc. Calif. Acad. Sci., 2d series, V, 1895, 805, pl. 80, Port Orchard, Puget Sound.
- Genus 1022. *RUSCARIOPS* Hubbs.
Ruscariops Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 12 (*R. creaseri* Hubbs).
2956. *Ruscariops creaseri* Hubbs.
Low rock pool at Bird Rock, San Diego County, Calif.
Ruscariops creaseri Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 12, very low rock pool at Bird Rock, San Diego Co., Calif.
- Genus 1023. *RASTRINUS* Jordan and Evermann.
Rastrinus Jordan and Evermann, Check List, 1895, 437 (*Icelus scutiger* Bean).
2957. *Rastrinus scutiger* (Bean).
South of the Alaskan Peninsula.
Icelus scutiger Bean, Proc. U. S. Nat. Mus., XIII, 1890, 41, Trinity Islands, Alaska, at *Albatross* station 2853, in 159 fathoms.
- Genus 1024. *ICELUS* Krøyer.
Icelus Krøyer, Natur. Hist. Tidsskr., I, 1845, 253 (*Icelus hamatus* Krøyer=*Cottus bicornis* Reinhardt).
2958. *Icelus bicornis* (Reinhardt).
Arctic seas, circumpolar; Spitzbergen to northern Russia, Finland, Alaska, Greenland, Labrador, and Cape Cod.
Cottus bicornis Reinhardt, Vid. Selsk. Natur. og Math. Afh., VIII, 1841, LXXV, Greenland.
Icelus hamatus Krøyer, Nat. Hist. Tidsskr., I, 1845, 253, Belsund in Spitzbergen.
Icelus furciger Malm, Forhandl. Skand. Naturf., 1865, 9 Mote, 410, Bohuslan, Sweden.
2959. *Icelus spiniger* Gilbert.
Coast of Alaska in the vicinity of Unalaska Island and in Bristol Bay.
Icelus spiniger Gilbert, Rept. U. S. Fish Com., XIX, 1893, 412, pl. 24, fig. 1, Bristol Bay and Unalaska, at *Albatross* stations 3216, 3223, and many others, in 17 to 121 fathoms.
2960. *Icelus euryops* Bean.
Bering Sea, off Trinity Islands, Alaska.
Icelus euryops Bean, Proc. U. S. Nat. Mus., XIII, 1890, 41, off Trinity Islands, at *Albatross* station 2853, in 159 fathoms.
2961. *Icelus vicinalis* Gilbert.
Bristol Bay, Alaska.
Icelus vicinalis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 413, Bristol Bay at *Albatross* stations 3324, 3330, 3331, and 3332, in 109 to 406 fathoms.
2962. *Icelus canaliculatus* Gilbert.
Bering Sea north of Unalaska.
Icelus canaliculatus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 412, pl. 24, fig. 2, off Unalaska, at *Albatross* station 3329, in 399 fathoms.

2963. *Icelus uncinialis* Gilbert and Burke.
Coast Bering Sea, off Commander Islands.
Icelus uncinialis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 39, fig. 2, Petrel Bank, near Commander Islands.
2964. *Icelus spatula* Gilbert and Burke.
East coast of Kamchatka.
Icelus spatula Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 41, fig. 3, Avatcha Bay.
2965. *Icelus australis* (Eigenmann and Eigenmann)
Cortez Banks off San Diego.
Icelinus australis Eigenmann and Eigenmann, West Amer. Sci., 1889, 131, Cortez Banks, Calif.
- Genus 1025. **STELGIDONOTUS** Gilbert and Thompson.
Stelgidonotus Gilbert and Thompson, Proc. U. S. Nat. Mus., XXVIII, 1905, 977 (*S. latifrons* Gilbert and Thompson).
2966. *Stelgidonotus latifrons* Gilbert and Thompson.
Puget Sound region, Admiralty Head, Friday Harbor.
Stelgidonotus latifrons Gilbert and Thompson, Proc. U. S. Nat. Mus., XXVIII, 1905, 977, Friday Harbor, Wash.
- Genus 1026. **RADULINUS** Gilbert.
Radulinus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 88 (*R. asprellus* Gilbert).
2967. *Radulinus boleoides* Gilbert.
Off Santa Catalina Island in shallow water.
Radulinus boleoides Gilbert, Report U. S. Fish Com., XXIV, 1898, 26, pl. 1, and in Jordan and Evermann, Fishes North and Mid. Amer., II, 1898, 1919, Santa Catalina Island, Calif.
2968. *Radulinus asprellus* Gilbert.
Coast of Oregon and Washington, and in Puget Sound.
Radulinus asprellus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 88, off Oregon and Washington, at *Albatross* stations 3046, 3057, and others, in 43 to 77 fathoms.
- Genus 1027. **STELGISTROPS** Hubbs.
Stelgistrops Hubbs, Occ. Papers, Mus. Zool. Univ. Mich. No. 170, Feb. 20, 1926, 8 (*Stelgistrum beringianum* Gilbert and Burke).
2969. *Stelgistrops beringiana* (Gilbert and Burke).
South Bering Sea.
Stelgistrum beringianum Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 52, fig. 9, Petrel Bank, Aleutian Islands.
- Genus 1028. **THYRISCUS** Gilbert and Burke.
Thyriscus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 43 (*T. anoplus* Gilbert and Burke).
2970. *Thyriscus anoplus* Gilbert and Burke.
Bering Sea.
Thyriscus anoplus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 43, fig. 4, Attu Island.
- Genus 1029. **TRIGLOPS** Reinhardt.
Triglops Reinhardt, Vid. Selsk. Natur. Math. Afh., V, 1832, LII (*T. pingeli* Reinhardt).
2971. *Triglops pingeli* Reinhardt.
North Atlantic, from Spitzbergen and western Norway to Greenland.
Triglops pingeli Reinhardt, Vid. Selsk. Natur., V, 1832, LII, Greenland.
Triglops pleurostictus Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 81, Godhaven, Greenland.
2972. *Triglops ommatistius* Gilbert.
North Atlantic coast of North America, Newfoundland to Cape Ann.
Triglops ommatistius Gilbert, Proc. U. S. Nat. Mus., XLIV, 1913, 465, Gloucester, Mass.; heretofore wrongly called *Triglops pingeli*, which is mainly a European species.

2973. *Triglops terræ-novæ* Gilbert.
Coast of Newfoundland.
Triglops ommatistius terræ-novæ Gilbert, Proc. U. S. Nat. Mus., XLIV, 1913, 467, off Newfoundland, in 39 fathoms.
2974. *Triglops metopias* Gilbert and Burke.
West Bering Sea.
Triglops metopias Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 50, fig. 8, Petrel Bank, Bering Sea.
2975. *Triglops beani* Gilbert.
Alaska, both north and south of the Aleutian Islands and in Bristol Bay.
Triglops beani Gilbert, Rept. U. S. Fish Com., XIX, 1893, 426, pl. 28, upper fig., Aleutian Islands and Bristol Bay, at *Albatross* stations 3214, 3217, and many others, in $7\frac{1}{4}$ to 42 fathoms.
2976. *Triglops scepticus* Gilbert
Aleutian Islands, south of Sannak and north of Unalaska Island.
Triglops scepticus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 428, pl. 28, lower fig., Aleutian Islands, at *Albatross* stations 3215, 3222, and others, in 43 to 138 fathoms.
- Genus 1030. **PRIONISTIUS** Bean.
Prionistius Bean, Proc. U. S. Nat. Mus., VI, 1883, 355 (*P. macellus* Bean).
2977. *Prionistius macellus* Bean.
British Columbia, in Carter Bay.
Prionistius macellus Bean, Proc. U. S. Nat. Mus., VI, 1883, 355, Carter Bay, British Columbia.
- Genus 1031. **ELANURA** Gilbert.
Elanura Gilbert, Rept. U. S. Fish Com., XIX, 1893, 429 (*E. forficata* Gilbert).
2978. *Elanura forficata* Gilbert
Aleutian Islands, south of Sannak and north of Unimak Island.
Elanura forficata Gilbert, Rept. U. S. Fish Com., XIX, 1893, 430, pl. 30, fig. 1, Aleutian Islands, at *Albatross* stations 3213, 3214, and 3222, in 33 to 50 fathoms.
- Genus 1032. **STERNIAS** Jordan and Evermann.
Sternias Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1926 (*Triglops xenostethus* Gilbert).
2979. *Sternias xenostethus* (Gilbert).
Bering Sea, north to Unalaska.
Triglops xenostethus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 429, pl. 29, two figures, north of Unalaska, at *Albatross* station 3230, in 34 fathoms.
- Genus 1033. **PTERYGIOCOTTUS** Bean and Weed.
Pterygiocottus Bean and Weed, Trans. Roy. Soc. Canada, XII, 1920, 73 (*Pterygiocottus macouni* Bean and Weed).
2980. *Pterygiocottus macouni* Bean and Weed.
Vancouver Island.
Pterygiocottus macouni Bean and Weed, Trans. Roy. Soc. Canada, XIII, 1920, 73, Vancouver Island.
- Genus 1034. **MELLETES** Bean.
Melletes Bean, Proc. U. S. Nat. Mus., II, 1879, 354 (*M. papilio* Bean).
2981. *Melletes papilio* Bean.
Aleutian Islands.
Melletes papilio Bean, Proc. U. S. Nat. Mus., II, 1879, 354, St. Paul Island, Alaska.
- Genus 1035. **HEMILEPIDOTUS** Cuvier.
Hemilepidotus Cuvier, Règne Animal, ed. 2, II, 1829, 165 (*Cottus hemilepidotus* Tilesius).
Temnistia Richardson, Fauna Bor.-Amer., III, 1831, 59 (*Blepsias ventriosus* Eschscholtz = *Cottus hemilepidotus* Tilesius).

2982. *Hemilepidotus hemilepidotus* (Tilesius). *Red sculpin*.
Kamchatka to San Francisco; not common in Bering Sea; abundant from Sitka to Puget Sound.
? *Cottus hemilepidotus* Tilesius, Mém. Acad. Petersb., III, 1810, 262, Sea of Okhotsk.
Cottus trachurus Pallas, Zoogr. Ross.-Asiat., III, 1811, 138, Kuril Islands
Hemilepidotus tilesii Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 203 (276), pl. 85, after Tilesius and Pallas.
? *Blepsias ventricosus* Eschscholtz, Zool. Atlas, 3d Heft, 1829, 14, t. 13, Norfolk Sound and Sitka
Hemilepidotus gibbsii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 13, no definite locality; Northwest boundary survey.
2983. *Hemilepidotus jordani* Bean. *Irish Lord; Yellow sculpin*.
Alaska; Unalaska.
Hemilepidotus jordani Bean, Proc. U. S. Nat. Mus., IV, 1881, 153, Unalaska
2984. *Hemilepidotus zapus* Gilbert and Burke.
Bering Sea.
Hemilepidotus zapus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 54, fig. 10, Attu Island.
- Genus 1036. *CALYCILEPIDOTUS* Ayres.
Calycilepidotus Ayres, Proc. Calif. Acad. Nat. Sci., ed. 2, I, 1855, 81 (76) (*C. spinosus* Ayres).
2985. *Calycilepidotus spinosus* Ayres.
Coast of California; known only from about Monterey and San Francisco.
Calycilepidotus spinosus Ayres, Proc. Calif. Acad. Nat. Sci., ed. 2, I, 1855, 81 (76), San Francisco.
- Genus 1037. *ENOPHRYS* Swainson.
Enophrys Swainson, Class. Fishes, II, 1839, 271 (*Cottus claviger* Cuvier and Valenciennes).
2986. *Enophrys claviger* (Cuvier and Valenciennes).
Bering Sea.
Cottus claviger Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 143 (195), pl. 79, Kamchatka.
Cottus elegans Gray, in Cuvier and Valenciennes, loc. cit., 143 (195), same type.
- Genus 1038. *ASPICOTTUS* Girard.
Aspicottus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 130 (*A. bison* Girard)
Clypeocottus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 12 (*C. robustus* Ayres = *Aspicottus bison* Girard).
2987. *Aspicottus bison* Girard. *Buffalo sculpin*.
San Francisco to Alaska; abundant about Puget Sound.
Aspicottus bison Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 130, San Francisco.
Clypeocottus robustus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 11, San Francisco.
2988. *Aspicottus taurinus* (Gilbert).
Monterey Bay.
Enophrys taurinus Gilbert, Proc. U. S. Nat. Mus., XLVII, 1915, 135, Monterey Bay.
- Genus 1039. *CERATOCOTTUS* Gill.
Ceratocottus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 165 (*Cottus diceraus* Pallas).
2989. *Ceratocottus diceraus* (Pallas).
Alaska and Kamchatka.
Cottus diceraus Pallas, Nov. Act. Petropol., 1783, 354, pl. 10, fig. 7, Petropaulski, Kamchatka.
Synanceia cervus Tilesius, Mém. Acad. Petersb., III, 1811, 278, pl. 13, Petropaulski.
Cottus stelleri Bloch and Schneider, Syst. Ichth., 1801, 63, after Steller.

2990. *Ceratocottus lucasi* Jordan and Gilbert.

Bering Sea.

Ceratocottus lucasi Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1940, St. Paul Island, Bering Sea.**Family 234. BLEPSIIDÆ****Genus 1040. HISTIOCOTTUS Gill.***Peropus* Lay and Bennett, Beechey's Voy., Zool., Fish., 1839, 50 (*Blepsias bilobus* Cuvier and Valenciennes); name preoccupied.*Histiocottus* Gill, Proc. U. S. Nat. Mus., XI, 1888, 573 (*Blepsias bilobus* Cuvier and Valenciennes).**2991. *Histiocottus bilobus* (Cuvier and Valenciennes).**

Coast of Alaska and Kamchatka.

Blepsias bilobus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 279 (379), no locality, somewhere off Kamchatka.**Genus 1041. BLEPSIAS Cuvier.***Blepsias* Cuvier, Règne Anim., ed. 2, II, 1829, 167 (*Trachinus cirrhosus* Pallas).**2992. *Blepsias cirrhosus* (Pallas). *Nee-kao-kaiung*.**

Alaska to San Francisco; common at Unalaska, but not often south of Puget Sound.

Trachinus cirrhosus Pallas, Zoogr. Rosso-Asiat., III, 1811, 237, Bering Sea.*Blepsia trilobus* Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 276 (375); same types.**Genus 1042. NAUTICHTHYS Girard.***Nautichthys* Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 74 (*Blepsias oculoasciatus* Girard).*Nautiscus*⁷² Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2019 (*Nautichthys pribilovius* Jordan and Gilbert).**Subgenus NAUTICHTHYS Girard.****2993 *Nautichthys oculoasciatus* (Girard).**

Pacific coast, Alaska to Monterey, chiefly northward; common in Bristol Bay and about Unalaska.

Blepsias oculoasciatus Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 202, Fort Steilacoom, Wash.**Subgenus NAUTISCUS Jordan and Gilbert.****2994. *Nautichthys pribilovius* (Jordan and Gilbert).**

Bering Sea, Alaska.

Nautiscus pribilovius Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2020, off St. George Island, Bering Sea.**Family 235. SCORPÆNICHTHYIDÆ. Cabezones****Genus 1043. SCORPÆNICHTHYS Girard.***Scorpænichthys* Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 131 (*Hemitripterus marmoratus* Ayres).**2995 *Scorpænichthys marmoratus* (Ayres). *Cabezon*.**

Pacific coast of America from Puget Sound to San Diego.

Hemitripterus marmoratus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1851, 4, San Francisco.⁷² The genus *Nautiscus* is now regarded as scarcely distinct from *Nautichthys*.

Family 236. COTTIDÆ. Scaleless sculpins

Genus 1044. COTTUS⁷³ Linnæus. *Miller's thumbs*.

Cottus Linnæus, Syst. Nat., Ed. X, 1758, 264 (*C. gobio* Linnæus).

Pegedictis Rafinesque, Ichth. Ohiensis, 1820, 85 (*P. ictalops* Rafinesque = ?*Cottus bairdi* Girard, apparently also confused with the darter *Catostomus flabellaris*).

Uranidea DeKay, New York Fauna: Fishes, 1842, 61 (*U. quiescens* DeKay = *Cottus gracilis* Heckel).

Cottopsis Girard, Proc. Bost. Soc. Nat. Hist., III, 1850, 303 (*Cottus asper* Richardson).

Potamocottus Gill, loc. cit., VIII, 1861, 40 (*P. carolinæ* Gill).

Tauridea Jordan and Rice, in Jordan, Man. Vert., Ed. II, 1878, 255 (*Cottopsis ricei* Nelson).

2996. *Cottus gulosus* (Girard).

Streams of the Coast Ranges in California; Columbia, Klamath, and Sacramento River regions; Alaska; Loring and Boca de Quadra.

Cottopsis gulosus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 120, San Mateo Creek and San Joaquin River, Calif.

Cottopsis parvus Girard, loc. cit., 1854, 144, Monterey, Presidio, Fort Reading, Petaluma.

Uranidea semiscabra centropoleura Eigenmann and Eigenmann, West Amer. Scientist, VI, No. 49, 1889, 149, Allen Springs, Lake Co., Calif.

Subgenus COTTOPSIS Girard.

2997. *Cottus asper* Richardson. *Prickly bullhead*.

Streams of the Cascade Range, southeastern Alaska to Oregon; south to Sacramento River.

Cottus asper Richardson, Fauna Bor.-Amer., Fish., 1836, 295, Columbia River at Fort Vancouver.

Trachidermis richardsoni Heckel, Ann. Wiener Mus., II, 1840, 162, Columbia River.

2998. *Cottus evermanni* Gilbert.

Lost River, Oreg.

Cottus evermanni Gilbert, Bull. U. S. Fish Com., XVII, 1897, 11, with figure, Lost River, Lostine, Oreg.

2999. *Cottus rhothea* (Rosa Smith).

Hangman Creek, Tekoa, Washington, and elsewhere in Columbia River Basin

Uranidea rhothea Rosa Smith, Proc. U. S. Nat. Mus., V, 1882, 347, Spokane Falls, Wash.

3000. *Cottus shasta* Jordan and Starks.

Upper Sacramento Basin, about Mount Shasta.

Cottus shasta Jordan and Starks, Proc. Calif. Acad. Sci. 2d ser., VI, 1896, 224, McCloud River, Baird, Shasta County, Calif.

3001. *Cottus punctulatus* (Gill).

Headwaters of Green River, Wyo.; Silver Creek, Harney County, Oreg.

Potamocottus punctulatus Gill, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 40, Bridger Pass, Wyo.

⁷³ In every cold stream of northern Europe, northern Asia, and northern America species of this genus abound, lurking on the bottom and devouring the eggs of trout which inhabit the same waters. The species are all differentiated and like those of trout subject to much individual variation. Those of the United States and Canada are not yet well known and the number of species has probably been unduly expanded. Efforts to subdivide the group have not been very successful and we place all the American forms in a single genus, *Cottus*. Western species, allies of *Cottus asper*, termed *Cottopsis*, are in general larger in size, more prickly, with well-developed palatine teeth and the ventral rays 1, 4. Those of the middle west allied to *Cottus bairdi*, called *Potamocottus*, show similar characters but are for the most part smaller and smoother, the colors more mottled. Rafinesque's *Pegedictis ictalops*, the "Cat's-eye Spring fish," seems to us a *Cottus* described carelessly or more likely from memory. In New England and south to Pennsylvania the allies of *Cottus gracilis* called *Uranidea* are small, smooth, without palatine teeth and the ventral rays 1, 3. But all these traits are subject to intergrading. A few species in deeper waters in Lake Michigan and Canada, allies of *Cottus ricei* called *Tauridea*, are more or less rough and have the preopercular spine larger, hooked and somewhat spiral, otherwise much as in *Cottopsis*.

3002. *Cottus semiscaber* (Cope). *Rocky Mountain bullhead*.
Rocky Mountain region in clear streams on both slopes; abundant in Colorado, northern New Mexico, Wyoming, Montana, Idaho, Utah, eastern Washington, and Oregon.
Collopsis semiscaber Cope, Hayden Surv. Mont., 1871, 476, Fort Hall, Idaho.
Uranidea wheeleri Cope, Proc. Amer. Phil. Soc. Phila., XIV, 1874, 138, Beaver River, Utah.
3003. *Cottus asperimus* Rutter.
California; Sacramento-San Joaquin Basin; Fall River
Cottus asperimus Rutter, Bull. U. S. Bur. Fish., XXVII, 1907, 144, fig. 3, Fall River, Calif.
3004. *Cottus beldingii* Eigenmann and Eigenmann.
Streams of Columbia River Basin, south to Lake Lahontan; abundant east of the Cascades and in Lake Tahoe; fossil in Lahontan Basin, in Pleistocene deposits.
Cottus beldingii Eigenmann and Eigenmann, Amer. Nat., XXV, 1891, 1132, Lake Tahoe
3005. *Cottus annæ* Jordan and Starks.
Eagle River, a tributary of Grand River, Colo., Colorado River Basin.
Cottus annæ Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 223, Pl. XXVII, Eagle River at Gypsum, Colo.
3006. *Cottus philonips* Eigenmann and Eigenmann.
Columbia River Basin at Field, British Columbia, and in Idaho.
Cottus philonips Eigenmann and Eigenmann, Amer. Nat., XXVI, 1892, 963, Kicking Horse River, Field, British Columbia.
3007. *Cottus klamathensis* Gilbert.
Klamath Lakes, Oreg.
Cottus klamathensis Gilbert, Bull. U. S. Fish. Com., XVII, 1897, 10, figure, Upper Klamath Lake, Oreg.
3008. *Cottus macrops* Rutter.
Fall River, Calif.
Cottus macrops Rutter, Bull. U. S. Bur. Fish., XXVII, 1907, 146, fig. 4, Fall River, Calif.
3009. *Cottus perplexus* Gilbert and Evermann.
Skookumchuck and Newaukum Rivers, near Chehalis, Wash.
Cottus perplexus Gilbert and Evermann, Bull. U. S. Fish. Com., XIV, 1894, 202, pl. 20, fig. 1, Skookumchuck River, near Chehalis, Wash.
3010. *Cottus princeps* Gilbert.
Klamath Lakes, Oreg.
Cottus princeps Gilbert, Bull. U. S. Fish. Com., XVII, 1897, 12, figure, Upper Klamath Lake, Oreg.
3011. *Cottus bendirei* (Bean).
Walla Walla, Wash.; Rattlesnake Creek near Camp Harney, Oreg., and Goose Creek near Meadows, Idaho.
Potamocottus bendirei Bean, Proc. U. S. Nat. Mus., IV, 1881, 27, Rattlesnake Creek near Camp Harney, Oreg.
3012. *Cottus greeniei* (Gilbert and Culver).
Snake River Basin, Idaho
Uranidea greeniei Gilbert and Culver, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1965, Thousand Springs, Snake River, Idaho.
3013. *Cottus marginatus* (Bean).
Tributaries of Walla Walla River, Wash.
Uranidea marginata Bean, Proc. U. S. Nat. Mus., IV, 1881, 26, Walla Walla, Wash.
3014. *Cottus tenuis* (Evermann and Meek).
Klamath Lakes, Oreg.
Uranidea tenuis Evermann and Meek, Bull. U. S. Fish. Com., XVII, 1897, 83, fig. 6, Upper Klamath Lake, Klamath Falls, Oreg.

3015. *Cottus leiopomus* Gilbert and Evermann.
Wood River, Shoshone, Idaho.
Cottus leiopomus Gilbert and Evermann, Bull. U. S. Fish Com., XIV, 1894, 203, pl. 20, fig. 2, Wood River, Shoshone, Idaho.
3016. *Cottus aleuticus* Gilbert.
Unalaska; Departure Bay, Vancouver Island; southward to Monterey in the Coast Range.
Uranidea microstoma Lockington, Proc. U. S. Nat. Mus., III, 1880, 58, St. Paul, Kadiak Island; name preoccupied in *Cottus*.
Cottus aleuticus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 418, streams at Unalaska; also in Departure Bay, Vancouver Island.
3017. *Cottus chamberlaini* Evermann and Goldsborough.
Southeastern Alaska.
Cottus chamberlaini Evermann and Goldsborough, Bull. U. S. Bur. Fish., XXVI, 1906, 308, fig. 66, Loring, Alaska.

Subgenus PEGEDICTIS Rafinesque.

3018. *Cottus ictalops* (Rafinesque.) *Müller's thumb; Mufflejaws; Springfish; Blob; Bullhead.*
Middle and northern States, east of the Dakotas and Nebraska; Lake Superior to the Ohio and Potomac Rivers.
Pegedictis ictalops Rafinesque, Ichth. Ohnensis, 85, supplement, 1820. Falls of the Ohio; perhaps confused with *Catostomus flabellaris*.
Cottus richardsoni Agassiz, Lake Superior, 1850, 300, northern shore of Lake Superior; not *Trachidermis richardsoni* Heckel, also a species of *Cottus*.
Cottus bairdii Girard, Proc. Amer. Assoc. Adv. Sci., II, 1850, 410, Mahoning River, Poland, Ohio.
Cottus wilsoni Girard, Monogr. Cottoids, in Smithsonian. Contrib. to Knowledge, 1850, 42, Pittsburgh, Pa.
Cottus alboradus Girard, loc. cit., 46, Fort Gratiot, Lake Huron.
Uranidea spilota Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 82, Grand River at Grand Rapids, Mich.
3019. *Cottus meridionalis* Girard. *Southern miller's thumb; Blob; Mufflejaw*
Cottus meridionalis Girard, Proc. Am. Assn. Adv. Sci., II, 1850, 410. James River, Va.
Potamocottus carolinæ Gill, Proc. Bost. Soc. Nat. Hist., 1861, 40. Carolinas.
Potamocottus zopherus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 320, small tributaries of the Etowah and Oostanaula Rivers near Rome, Ga.

Subgenus URANIDEA DeKay

3020. *Cottus franklini* Agassiz.
Lake Superior and Lake Michigan in deep water.
Cottus franklini Agassiz, Lake Superior, 1850, 303, north and east shores of Lake Superior.
Uranidea hoyi Putnam, in Nelson, Bull. Ill. Lab. Nat. Hist., I, No. 1, 1876, 41, Lake Michigan, in deep water.
Uranidea kumtzeni Hoy, in Nelson, loc. cit., 41, Lake Michigan, in deep water.
3021. *Cottus gracilis*⁷⁴ Heckel. *Müller's thumb; Rock cusk; Little stargazer.*
Streams of New England, Maine to New York; recorded from tributaries of the Connecticut, Lake Champlain, Hudson, Delaware, and Susquehanna.
Cottus gracilis Heckel, Ann. Wien Mus., II, 1837, 148, New York.
Cottus viscosus Haldeman, Suppl. Mongr. Linnæa, 1840, 3, Susquehanna River.
Uranidea quiescens DeKay, New York Fauna: Fishes, 1842, 61, Lake Pleasant, Adirondack region.
Cottus gobioides Girard, Proc. Amer. Assoc. Adv. Sci., II, 1850, 411, Lamville River, Johnson, Vt.
Cottus boeoides Girard, loc. cit., II, 411, Windsor, Vt.

⁷⁴ May not be distinct from the northern *C. cognatus*.

- Cottus copei* Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 326, probably New Jersey.
Cottus gracilis cayuga Meek, Ann. New York Acad. Sci., IV, 1888, 315, Cayuga Lake, N Y
3022. *Cottus formosus* Girard.
 Lake Ontario, in deep water.
Cottus formosus Girard, Monogr. Cott., in Smithson. Contrib., III, 1850, 58, Lake Ontario, off Oswego, in stomach of *Lota maculosa*.
 Subgenus TAURIDEA Jordan and Rice.
3023. *Cottus ricei* (Nelson).
 Lake Michigan; Lake Ontario.
Cottopsis ricei Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 40, Lake Michigan off Evanston.
Urandeia pollicaris Jordan and Gilbert, Proc. U.S. Nat. Mus., V, 1882, 222, Lake Michigan, off Racine, Wis.
3024. *Cottus onychus* Eigenmann and Eigenmann.
 Saskatchewan Basin.
Cottus onychus Eigenmann and Eigenmann, Am. Nat., XXVI, November, 1892, 963, Bow River at Calgary, a tributary of the South Saskatchewan, Alberta Territory.
3025. *Cottus cognatus* Richardson. *Northern miller's thumb*.
 Northern United States and northward, through Canada.
Cottus cognatus Richardson, Fauna Bor.-Amer., III, 1836, 40, Great Bear Lake.
- Genus 1045. ACANTHOCOTTUS Girard. *Daddy sculpin*
Acanthocottus Girard, Proc. Bost. Soc. Nat. Hist., III, 1849, 185 (*Cottus granlandicus* Cuvier and Valenciennes).
3026. *Acanthocottus mitchilli*⁷⁵ (DeKay).
 Coast of southern New England and New York.
Cottus mitchilli DeKay, in Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 188, New York; after *Cottus scorpio* Mitchill.
3027. *Acanthocottus bubalis* (Euphrasen). *Fatherlasher, Lucky proach*.
 Barbados.
Cottus bubalis Euphrasen, Kong. Vet. Nya Abhandl., VII, 1786, 65, taf. 3, fig 8, 1885, Barbados
Cottus maculatus Fischer, Jahrb. Wissensch. Aust. Hamburg, II, 1885, 78, taf. 2, fig. 8, Barbados.
3028. *Acanthocottus æneus* (Mitchill). *Grubby; Brassy sculpin; Pigmy sculpin*.
 Coast of southern New England and New York.
Cottus æneus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 380, New York.
Cottus anceps Sauvage, Nouv. Archiv Mus. Paris, I, 1878, 145, pl. 1, fig. 13, New York.
 ?*Cottus nigricans* LeSueur, in Vaillant, Bull. Soc. Philom., serie 8, tome VIII, No. 1, 1895, 12, no locality.
3029. *Acanthocottus octodecemspinosus* (Mitchill). *Long-spined sculpin; Common sculpin; Bullhead; Eighteen-spined sculpin; Hacklehead*.
 Atlantic coast south to Virginia; common northward.
Cottus octodecemspinosus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 380, New York.
3030. *Acanthocottus scorpius* (Linnæus). *European sculpin; Ulke*.
 Northern Europe and Arctic regions, not common on our coasts; recorded by Doctor Lütken from the Baltic, Finland, Spitzbergen, Nova Zembla, coasts of England and northern Asia; also recorded from Eastport, Me.
Cottus scorpius Linnæus, Syst. Nat., Ed. X, 1758, 265, in Atlantic Ocean off Europe.
3031. *Acanthocottus scorpioides* (Fabricius). *Arctic sculpin*.
 Arctic regions of America, Greenland to Siberia.
Cottus scorpioides Fabricius, Fauna Grœnl., 1780, 157, Greenland.
Cottus pachypus Günther, Cat., II, 1860, 161, Port Leopold

⁷⁵ Differentiated by Nichols, Copeia, No 31, 40, 1916

- 3032. *Acanthocottus grœnlandicus* (Cuvier and Valenciennes). *Daddy sculpin*.**
 New York to Greenland; according to Fabricius it is abundant in all the bays and inlets of Greenland.
Cottus grœnlandicus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 135 (156), no locality given, probably Greenland; after Fabricius.
Cottus porosus Cuvier and Valenciennes, loc. cit., VIII, 1831, 367 (498), Baffins Bay.
Acanthocottus mucosus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854 (Oct. 9), 12, coast of New England.
Acanthocottus ocellatus Storer, Boston Jour. Nat. Hist., 1850, 253, Labrador.
Cottus glacialis Richardson, in Belcher, Last Arctic Voy. H. M. S. Assistant, App. 3, 1855, 349, pl. 24, Northumberland Sound.
- Genus 1046. MYOXOCEPHALUS (Steller) Tilesius.**
Myoxocephalus Tilesius, Mém. Acad. Sci. Petersb., IV, 1811, 273 (*Myoxocephalus stelleri* Tilesius)
Boreocottus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 166 (*B. axillaris* Gill).
- 3033. *Myoxocephalus polyacanthocephalus* (Pallas). *Great sculpin; Kalog.***
 Alaska to Kamchatka; abundant, Bering Sea to Puget Sound, tide pools of Karluk.
Cottus polyacanthocephalus Pallas, Zoogr. Rosso.-Asiat., III, 1811, 133, Aleutian Islands.
- 3034. *Myoxocephalus jaok* (Cuvier and Valenciennes).**
 Kamchatka.
Cottus jaok Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 126 (172), Kamchatka.
Cottus humilis Bean, Proc. U. S. Nat. Mus., IV, 1881, 149, Chamisso Island, Bering Straits.
- 3035. *Myoxocephalus verrucosus* (Bean).**
 Plover Bay, Siberia, near Bering Straits; Unalaska; Sitka; Bristol Bay; Kings Island; Grantley Harbor, Port Clarence; Litnik.
Cottus verrucosus Bean, Proc. U. S. Nat. Mus., IV, 1881, 152, Plover Bay, Siberia.
- 3036. *Myoxocephalus axillaris* (Gill).**
 Bering Straits.
Boreocottus axillaris Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 166, Bering Straits.
- 3037. *Myoxocephalus stelleri* Tilesius.**
 Western shore of Bering Sea and Okhotsk Sea.
Myoxocephalus stelleri Tilesius, Mém. Acad. Petersb., IV, 1811, 273, with plate, Petropaulski.
Cottus decastris Kner, Denk. Kais. Acad. Wissen., XXIV, 1865, 2, taf. 2, figs. 1, 1a, Decastris Bay, near mouth of the Amur.
 ?*Cottus marmoratus* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 366 (497), Petropaulski.
- 3038. *Myoxocephalus mednius* Bean.**
 Medni Island, Bering Sea.
Myoxocephalus mednius Bean, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1983, Medni Island, Bering Sea.
- 3039. *Myoxocephalus parvulus* Gilbert and Burke.**
 Medni Island, Commander Group.
Myoxocephalus parvulus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 59, Preobazenskoi Bar, Medni (Copper) Island.
- 3040. *Myoxocephalus niger* (Bean). *Black sculpin.***
 Pribilof Islands
Cottus niger Bean, Proc. U. S. Nat. Mus., IV, 1881, 151, St. Paul Island, Bering Sea.
 ?*Cottus mertensii* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 366 (496), Petropaulski.
- Genus 1047. MEGALOCOTTUS Gill.**
Megalocottus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 166 (*Cottus platycephalus* Pallas).

3041. *Megalocottus platycephalus* (Pallas).
Kamchatka.
Cottus platycephalus Pallas, Zoogr. Rosso.-Asiat., III, 1811, 135, Kamchatka; after Steller
Cottus tæniopterus Kner, Sitzungsber. Ak. Wiss. Wien, LXIII, 1868, 28, Taf. 14, fig. 10, Decastria Bay, near mouth of Amur River.
3042. *Megalocottus laticeps* (Gilbert).
Eastern shore of Bering Sea
Acanthocottus laticeps Gilbert, Rept. U. S. Fish. Com., XIX, 1893, 422, pls. 26 and 27, Nushagak River and Herendeen Bay, Alaska.
- Genus 1048. **POROCOTTUS** Gill.
Porocottus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 166 (*P. quadrifilis* Gill).
3043. *Porocottus sellaris* (Gilbert).
Bristol Bay, Alaska.
Acanthocottus sellaris Gilbert, Rept. U. S. Fish. Com. XIX, 1893, 419, Bristol Bay, at Albatross stations 3229, 3231, and others, in 5 to 17 fathoms.
3044. *Porocottus quadratus* Bean.
Bering Island.
Porocottus quadratus Bean, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1998, Bering Island.
3045. *Porocottus polaris* (Sabine).
East side of peninsula of Boothia.
Cottus polaris Sabine, App. Parry's First Voyage. CCXIII, 1824, north Georgia.
3046. *Porocottus quadrifilis* Gill.
Bering Straits.
Porocottus quadrifilis Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 166, Bering Straits.
3047. *Porocottus bradfordi* Rutter.
Alaska; rock pools at Karluk.
Porocottus bradfordi Rutter, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2862, Karluk.
- Genus 1049. **MALACOCOTTUS** Bean.
Malacocottus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 42 (*M. zonurus* Bean).
3048. *Malacocottus zonurus* Bean.
Coast of Alaska off Trinity Islands.
Malacocottus zonurus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 43, off Trinity Islands, Alaska, 56° N., 154° W., at Albatross station 2853, in 159 fathoms.
3049. *Malacocottus kincaidii* Gilbert and Thompson.
Puget Sound.
Malacocottus kincaidii Gilbert and Thompson, Proc. U. S. Nat. Mus., XXVIII, 1905, 979, fig. 2, Brannon, Hood's Canal, Wash.
- Genus 1050. **ONCOCOTTUS** Gill.
Oncocottus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 13 (*Cottus quadricornis* Linnæus).
3050. *Oncocottus quadricornis* (Linnæus).
Arctic America to Baltic Sea, chiefly northward; abundant in eastern Baltic and in Lakes Ladoga and Onega; north to White Sea and Nova Zembla; rare in England and eastern Greenland; unknown in western Greenland.
Cottus quadricornis Linnæus, Syst. Nat., ed. X, 1758, 264, Baltic Sea.
3051. *Oncocottus hexacornis* (Richardson).
Arctic Sea about Hershel Island.
Cottus hexacornis Richardson, Franklin's Journal, 1823, 726, mouth of Tree River near Coppermine River.
Acanthocottus labradoricus Girard, Bost. Jour. Nat. Hist., VI, 1850, 247, pl. 7, fig. 3, coast of Labrador.

3052. *Oncocottus gilberti* Fowler.
Point Barrow, Alaska; Arctic Sea.
Oncocottus hexacornis gilberti Fowler, Proc. Acad. Nat. Sci. Phila.,
LVII, 1905, 366, figs. 2, 3, 4, and 5, Point Barrow, Alaska.
- Genus 1051. **TRIGLOPSIS** Girard.
Triglopsis Girard, Proc. Bost. Soc. Nat. Hist., IV, 1851, 16 (*T. thompsoni*
Girard).
3053. *Triglopsis thompsoni* Girard. *Lake sculpin*.
Deep waters of Lake Ontario
Triglopsis thompsoni Girard, Proc. Bost. Soc. Nat. Hist., IV, 1851, 19,
Lake Ontario off Oswego, N. Y.
3054. *Triglopsis stimpsoni* Hoy.
Lake Michigan in deep water.
Triglopsis stimpsoni Hoy, Trans. Wis. Acad. Sci., 1872, 99, Lake Mich-
igan, off Chicago.
3055. *Triglopsis ontariensis* Jordan and Thompson.
Deep waters of Lake Ontario
Triglopsis ontariensis Jordan and Thompson, Proc. U. S. Nat. Mus.,
XXXVIII, 1911, 75, Lake Ontario, off Toronto
- Genus 1052. **GYMNOCANTHUS** Swainson.
Gymnocanthus Swainson, Class. Fishes, Amphib., etc., II, 1839, 271
(*Cottus ventralis* Cuvier and Valenciennes = *Cottus pistilliger* Pallas).
Phobetor Krøyer, Naturh. Tidskr., I, 1844, 263 (*Cottus tricuspis* Rein-
hardt).
Elaphocottus Sauvage, Nouv. Arch. Mus. Paris (2), 1, 1878, 142
(*Cottus pistilliger* Pallas).
Gymnacanthus of many authors. Corrected spelling.
3056. *Gymnocanthus pistilliger* (Pallas).
Coasts of Alaska; taken abundantly in Bristol Bay.
Cottus pistilliger Pallas, Zoogr. Rosso-Asiat., III, 1811, 143, Unalaska
Island.
Cottus ventralis Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829,
142 (194), Kamchatka.
Cottus cephaloides Gray, in Cuvier and Valenciennes, Hist. Nat. Poiss.,
IV, 1829, 142 (194), Kamchatka.
3057. *Gymnocanthus tricuspis* (Reinhardt)
Arctic seas, south to Norway and Labrador; not very common on our
coasts.
Cottus tricuspis Reinhardt, Vidensk. Selsk. Nat. Math., V, 1832, LII,
Greenland.
Acanthocottus patris Storer, Bost. Jour. Nat. Hist., VI, 1857, 250,
Labrador.
3058. *Gymnocanthus galeatus* (Bean).
Aleutian Islands.
Gymnacanthus galeatus Bean, Proc. U. S. Nat. Mus., IV, 1881, 153,
Unalaska.
- Genus 1053. **CROSSIAS** Jordan and Starks.
Crossias Jordan and Starks, Proc. U. S. Nat. Mus., XXVII, 1904, 296
(*C. allisi* Jordan and Starks).
3059. *Crossias albomaculatus* Schmidt.
Bering Sea.
Crossias albomaculatus Schmidt, Ann. Mus. Zool. Russ., XX, 1923,
614, fig. 2, Bering Sea.
3060. *Crossias beringi* Soldatov.
Bering Sea.
Crossias beringi Soldatov, Ann. Mus. Zool. Russ., XXI, 1924, 218,
Bering Sea.
3061. *Crossias camtschaticus* Schmidt.
Kamchatka.
Crossias camtschaticus Schmidt, Ann. Mus. Zool. Russ., XX, 1923, 17,
Kamchatka.

Genus 1054. **LEIOCOTTUS** Girard.

Leiocottus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 133 (*L. hirundo* Girard).

3062. ***Leiocottus hirundo*** Girard.

Coast of southern California, chiefly about the Santa Barbara Islands; extremely local.

Leiocottus hirundo Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 133, San Miguel Island near Santa Barbara, Calif

Genus 1055. **LEPTOCOTTUS** Girard.

Leptocottus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 130 (*L. armatus* Girard).

3063. ***Leptocottus armatus*** Girard. *Smooth cabezon*.

Pacific coast, from Kodiak to San Diego, abundant northward.

Leptocottus armatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 131, Cape Flattery, Fort Steilacoom, Shoalwater Bay, Humboldt Bay, San Francisco, Monterey, San Pedro, Fort Point, San Diego, and Tomales Bay.

Genus 1056. **ASEMATICHTHYS** Gilbert.

Asematichthys Gilbert (*Asematichthys taylori* Gilbert).

3064. ***Asematichthys taylori*** Gilbert.

Vancouver Island.

Asematichthys taylori Gilbert, Departure Bay, Nanaimo, Vancouver Island.

Genus 1057. **CLINOCOTTUS** Gill.

Clinocottus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 166 (*Oligocottus analis* Girard).

Rusulus Starks and Mann, Univ. Calif. Pub. Zool., VIII, 1911, 14 (*Clinocottus sabarræ* Starks and Mann; young).

3065. ***Clinocottus analis*** (Girard).

Coast of California from Point Conception northward.

Oligocottus analis Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 201, Monterey.

Cottus criniger Gunther, Cat., II, 1860, 522, Monterey, Calif.

Rusulus sabarræ Starks and Mann, Univ. Calif. Pub. Zool., VIII, 1911, 14, Ballast Point, off mouth of San Diego Bay; young.

3066. ***Clinocottus australis*** Hubbs.

Southern California from Los Angeles County and Santa Catalina Island southward to Todos Santos, San Martin, San Benito, and Guadalupe Islands. Probably varies into *C. analis*, of which it is the southern form.

Clinocottus analis australis Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 171, Feb. 20, 1926, 11, Point Loma, San Diego, Calif.

Genus 1058. **OLIGOCOTTUS** Girard.

Oligocottus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 132 (*O. maculosus* Girard).

3067. ***Oligocottus maculosus*** Girard. *Johnny*.

San Francisco to central Alaska.

Oligocottus maculosus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 133, Tomales Bay, San Francisco.

Oligocottus borealis Jordan and Snyder, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 225, Neah Bay, Wash.

Genus 1059. **GREELEYA** ⁷⁸ Jordan.

Eximia Greeley, Bull. U. S. Fish Com., XIX, 1899, 18 (*Eximia rubellio* Greeley); name preoccupied.

Greeleya Jordan, Genera of Fishes, IV, 1920, 493 (*Eximia rubellio* Greeley).

⁷⁸ The separation of this genus from *Oligocottus* has been questioned by Starks and approved by Hubbs.

3068. *Greeleya rubellio* (Greeley).
Monterey Bay, in deep tide pools.
Eximia rubellio Greeley, Bull. U. S. Fish Com., XIX, 1899, 18, fig. 5,
Pacific Grove, Calif.
- Genus 1060. *DIALARCHUS* Greeley.
Dialarchus Greeley, Bull. U. S. Bur. Fish., XIX, 1899, 14 (*Oligocottus*
snyderi Greeley).
3069. *Dialarchus snyderi* (Greeley).
Coast of California, Crescent City to San Luis Obispo, in tide pools.
Oligocottus snyderi Greeley, in Jordan and Evermann, Fishes North and
Mid. Amer., 1898, 2871, Pacific Grove, Calif.
- Genus 1061. *SIGMISTES* Rutter.
Sigmistes Rutter, in Jordan and Evermann, Fishes North and Mid.
Amer., 1898, 2863 (*S. caulias* Rutter).
3070. *Sigmistes caulias* Rutter.
Rock pools at Kodiak Island, Alaska.
Sigmistes caulias Rutter, in Jordan and Evermann, Fishes North and
Mid. Amer., 1898, 2863, rock pools at Karluk, Kodiak Island, Alaska.
- Genus 1062. *OXYCOTTUS* Jordan and Evermann.
Oxycottus Jordan and Evermann, Fishes North and Mid. Amer., 1898,
2015 (*Oligocottus acuticeps* Gilbert).
3071. *Oxycottus acuticeps* (Gilbert).
Alaska south to Cape Flattery, Departure Bay, Vancouver Island,
Kodiak.
Oligocottus acuticeps Gilbert, Rept. U. S. Fish Com., XIX, 1893, 432,
Unalaska.
- Genus 1063. *ALLOCOTTUS* Hubbs.
Allocottus Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 171, 1926,
14 (*Oligocottus embryum* Jordan and Starks).
3072. *Allocottus embryum* (Jordan and Starks).
Washington (Neah Bay), Point Lobos, Calif., Karluk and Sitka,
Alaska.
Oligocottus embryum Jordan and Starks, Proc. Calif. Acad. Sci., 2d
ser., V, 1895, 808, pl. 82, Neah Bay, Wash.
- Genus 1064. *RUSCICULUS* Greeley.
Rusciculus Greeley, Bull. U. S. Fish Com., XIX, 1899, 13 (*R. rimensis*
Greeley).
3073. *Rusciculus rimensis* Greeley.
Monterey Bay in tide pools.
Rusciculus rimensis Greeley, Bull. U. S. Fish Com., XIX, 1899, 13,
fig. 3, Point Lobos, near Carmel, Calif.
- Genus 1065. *BLENNICOTTUS* Gill.
Blennicottus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 166 (*Oligo-*
cottus globiceps Girard).
3074. *Blennicottus globiceps* (Girard).
Pacific coast from Coronado Islands northward to Cape Mendocino.
Oligocottus globiceps Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1853,
58, South Farallons.
Blennicottus globiceps bryosus Jordan and Starks, Proc. Calif. Acad.
Sci., 2d ser., V, 1895, 808, Neah Bay, Wash.
- Genus 1066. *MONTEREYA* Hubbs.
Montereya Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 171,
1926, 16 (*Blennicottus recalvus* Greeley).
3075. *Montereya recalva* (Greeley).
Monterey Bay to San Diego, in tide pools.
Blennicottus recalvus Greeley, Bull. U. S. Fish Com., XIX, 1899
(Dec. 13), 9, fig. 1, Pacific Grove, Calif.
- Genus 1067. *ORTHONOPIAS* Starks and Mann.
Orthonopias Starks and Mann, Univ. Cal. Publ. Zool., VIII, 1911, 11,
(*O. triacis* Starks and Mann).

3076. *Orthonopias triacis* Starks and Mann.
Coast of California, Monterey to San Diego.
Orthonopias triacis Starks and Mann, Univ. Cal. Publ. Zool., VIII, 1911, 11, fig. 1, Cortez Banks off San Diego.
- Genus 1068. *DASYCOTTUS* Bean.
Dasycottus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 42 (*D. setiger* Bean).
3077. *Dasycottus setiger* Bean.
Puget Sound to Bering Sea.
Dasycottus setiger Bean, Proc. U. S. Nat. Mus., XIII, 1890, 42, off Sitkalidak Island, Alaska.
- Genus 1069. *ZESTICELUS* Jordan and Evermann.
Zesticelus Jordan and Evermann, Check List, 1895, 443 (*Acanthocottus profundorum* Gilbert).
3078. *Zesticelus profundorum* (Gilbert).
Bering Sea, north of Unalaska.
Acanthocottus profundorum Gilbert, Rept. U. S. Fish Com., XIX, 1893, 423, pl. 27, upper fig., Bering Sea, north of Unalaska, at Albatross station 3328, in 399 fathoms.
- Genus 1070. *THECOPTERUS* Smith.
Thecopterus Smith (H. M.), Proc. Biol. Soc. Wash., XVII, 1904, 163 (*Thecopterus aleuticus* Smith).
3079. *Thecopterus aleuticus* Smith.
Bering Sea.
Thecopterus aleuticus Smith, Proc. Biol. Soc. Wash., XVII, 1904, 163, near Rat Islands, in 270 fathoms.
- Genus 1071. *COTTUNCULUS* Collett.
Cottunculus Collett, Norges Fiske, 1875, 20 (*C. microps* Collett).
3080. *Cottunculus microps* Collett.
Deep waters off coasts of Norway and Rhode Island.
Cottunculus microps Collett, Norges Fiske, 1875, 20, pl. I, figs. 1-3, Hasvig near Hammerfest, Norway, in 200 fathoms.
3081. *Cottunculus thomsonii* (Günther).
North Atlantic, in deep water.
Cottus thomsoni Günther, Proc. Royal Soc. Edinburgh, XI, 1882, 679, Faroe Channel, in 535 fathoms.
Cottunculus torvus Goode, Proc. U. S. Nat. Mus., III, 1880, 479, Gulf Stream off Rhode Island. *A nomen nudum*.
- Genus 1072. *ASCELICHTHYS* Jordan and Gilbert.
Ascelichthys Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 264 (*A. rhodorus* Jordan and Gilbert).
3082. *Ascelichthys rhodorus* Jordan and Gilbert.
Pacific coast from Sitka to Cape Mendocino; abundant in Neah Bay, Wash.
Ascelichthys rhodorus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 264, Waadda Island, Neah Bay, near Cape Flattery.
- Genus 1073. *PSYCHROLUTES* Günther.
Psychrolutes Günther, Cat., III, 1861, 516 (*P. paradoxus* Günther).
3083. *Psychrolutes paradoxus* Günther.
Puget Sound to Kodiak Island.
Psychrolutes paradoxus Günther, Cat., III, 1861, 516, Gulf of Georgia.
Psychrolutes zebra Bean, Proc. U. S. Nat. Mus., XIII, 1890, 43, Aleutian Islands, between Unga and Nagai Islands, in 110 fathoms.
- Genus 1074. *GILBERTIDIA* Berg.
Gilbertina Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 811 (*Gilbertina sigalutes* Jordan and Starks); name preoccupied.
Gilbertidia Berg, Comun del Museo Nac. de Buenos Aires, I, 1898, 42 (*Gilbertina sigalutes* Jordan and Starks).

3084. *Gilbertidia sigalutes* (Jordan and Starks).
 Puget Sound.
Gilbertidia sigalutes Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser.,
 V, 1895, 811, pl. 86, Port Orchard, near Seattle.

Family 237. SYNCHIRIDÆ

- Genus 1075. *SYNCHIRUS* Bean.
Synchirus Bean, Proc. U. S. Nat. Mus., XII, 1889, 641 (*S. gilli* Bean).
 3085. *Synchirus gilli* Bean.
 Puget Sound; Barclay Sound, Port Townsend; rare.
Synchirus gilli Bean, Proc. U. S. Nat. Mus., XII, 1889, 642, Barclay
 Sound, British Columbia.

Family 238. RHAMPHOCOTTIDÆ

- Genus 1076. *RHAMPHOCOTTUS* Günther.
Rhamphocottus Günther, Ann. Mag. Nat. Hist., Series 4, XIV, 1874,
 369 (*R. richardsoni* Günther).
 3086. *Rhamphocottus richardsoni* Günther.
 North Pacific, Alaska to Monterey; not infrequent in Puget Sound.
Rhamphocottus richardsoni Günther, Ann. Mag. Nat. Hist., XIV,
 1874, 369, Fort Rupert, British America.
 Genus 1077. *ULCA* Jordan and Evermann.
Ulca Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896,
 227 (*Hemitripteris marmoratus* Bean).
 3087. *Ulca marmorata* (Bean).
 Alaska, Bering Sea and off Sitkalidak Island.
Hemitripteris marmoratus Bean, Proc. U. S. Nat. Mus., XIII, 1890,
 43, off Sitkalidak Island.

Family 239. HEMITRIPTERIDÆ. Sea ravens

- Genus 1078. *HEMITRIPTERUS* Cuvier.
Hemitripteris Cuvier, Règne Anim., ed. 2, II, 1829, 164 (*Scorpena
 americanus* Gmelin).
 3088. *Hemitripteris americanus* (Gmelin). *Sea raven; Yellow sculpin.*
 Atlantic coast of America from Tortugas to Labrador, chiefly northward.
Scorpena americana Gmelin, Syst. Nat., ed. XIV, 1788, 1220, after
 "Diable en Crapand de l'Amérique" of Duhamel de Moreau.
Cottus acadian Walbaum, Arted. Pisc., 1792, 392, 1784, Nova Scotia;
 after Pennant
Cottus hispidus Bloch and Schneider, Syst. Ichth., 1801, 63, New York.
Cottus tripterygius Bloch and Schneider, Syst. Ichth., 1801, 63, Nova
 Scotia; after Pennant.
 3089. *Hemitripteris villosus* (Pallas).
 Coast of Alaska.
Cottus villosus Pallas, Zoogr. Rosso-Asiat., III, 1811, 129, Cape Kronok,
 Kamchatka, Itscha River.
Hemitripteris cavifrons Lockington, Proc. Acad. Nat. Sci. Phila.,
 XXXII, 1880, 233, Kodiak Island, Alaska.

Family 240. AGONIDÆ. Sea poachers; Alligatorfishes

- Genus 1079. *HYPAGONUS* Gill.
Hypsagonus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 167 (*Aspidophorus
 quadricornis* Cuvier and Valenciennes).
Cheiragonus Herzenstein, Bull. Acad. Sci. St. Petersburg., XIII, 1890, 116
 (*Hypsagonus gradiens* Herzenstein = *Aspidophorus quadricornis* Cuvier
 and Valenciennes).
 3090. *Hypsagonus quadricornis* (Cuvier and Valenciennes).
 North Pacific; Kamchatka; Bering Sea and coasts of Alaska south to
 Bristol Bay and Puget Sound.
Aspidophorus quadricornis Cuvier and Valenciennes, Hist. Nat. Poiss.,
 IV, 1829, 162 (221), Kamchatka.
Hypsagonus gradiens Herzenstein, Bull. Acad. Int. Sci. St. Petersburg.,
 XIII, 1890, 116, Kamchatka, Sinus Awatcha Bay, Petropaulski.

Genus 1080. *STELLERINA* Cramer.

Stellerna Cramer, in Jordan and Evermann, Check List, 1895, 447
(*Brachyopsis xyosternus* Jordan and Gilbert).

3091. *Stellerna xyosterna* (Jordan and Gilbert).

Coast of California and Oregon.

Brachyopsis xyosternus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, July 2, 1880, 152, Santa Cruz, Calif.

Agonus annæ Steindachner, Ichth. Beitr., IX, 1880, 254, pl. 6, figs. 1, 1a, Sitzb. K. Acad. Wiss., LXXXII, San Francisco, Calif.

Genus 1081. *OCCA* Jordan and Evermann.

Occa Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2043
(*Brachyopsis verrucosus* Lockington).

3092. *Occa verrucosa* (Lockington).

Coast of California, south to San Francisco.

Brachyopsis verrucosus Lockington, Proc. U. S. Nat. Mus., III, May 6, 1880, 60, Drake Bay, near San Francisco.

Agonus barkani Steindachner, Ichth. Beitr., IX, 1880, 253, pl. 5, Sitzb. K. Acad. Wiss., LXXXII, San Francisco, Calif.

Genus 1082. *OCCELLA* Jordan and Hubbs.

Ocella Jordan and Hubbs, Mem. Carnegie Mus., X, No. 2, 1925, 291
(*Agonus dodecaedron* Tilesius).

3093. *Ocella dodecaëdron* (Tilesius).

North Pacific, Kamchatka, Bristol Bay, coast of Alaska.

Agonus dodecaëdrus Tilesius, Mém. Acad. Petersb., IV, 1810, pl. 13, Kamchatka.

Phalangistes lorincatus Pallas, Zoog. Rosso-Asiat., III, 1811, 114, tab. 19, Kamchatka.

Genus 1083. *PALLASINA* Cramer.

Pallasina Cramer, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 815 (*Siphonogonus barbata* Steindachner).

3094. *Pallasina barbata* (Steindachner).

Bering Sea on both coasts to Hokkaido, northern Japan.

Siphonogonus barbatus Steindachner, Ichth. Beitr., V, July, 1876, 140, pl. 5, Sitzb. der K. Acad. Wiss., LXXIV, Hakodate, Japan; the locality of Nagasaki also mentioned is impossible.

3095. *Pallasina aix* Starks.

Puget Sound to the Aleutian Islands.

Pallasina aix Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 558, pl. 75, Puget Sound near Port Ludlow, Wash.

Genus 1084. *LEPTAGONUS* Gill.

Leptagonus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 167, 259
(*Aspidophorus spinosissimus* Krøyer = *Aspidophorus decagonus* Cuvier and Valenciennes).

3096. *Leptagonus decagonus* (Bloch and Schneider).

Arctic Ocean, south to Newfoundland and Norway.

Agonus decagonus Bloch and Schneider, Syst. Ichth., I, 1801, 105, pl. 27, erroneously recorded from the East Indies; the type from Greenland.
Aspidophorus spinosissimus Krøyer, Naturhist. Tidsskr., I, 1844, 250, Greenland.

Genus 1085. *PODOTHECUS* Gill.

Podothecus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 77, 259 (*P. peristethus* Gill = *Agonus acipenserinus* Tilesius).

Paragonus Gill, loc. cit., 167, 259 (*Agonus acipenserinus* Tilesius).

3097. *Podothecus acipenserinus* (Tilesius). Common alligatorfish.

Kamchatka to Puget Sound.

Agonus acipenserinus Tilesius, Mém. Acad. Petersb., IV, 1811, 422, pl. 11, fig. 163, Unalaska.

Podothecus peristethus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 259, Simeahmoo, Washington Territory.

3098. *Podothecus gilberti* Collett.
Kamchatka.
Podothecus gilberti Collett, Proc. Zool. Soc. London, 1894, 67, pl. 45, Kamchatka.
- Genus 1086. AGONUS Bloch and Schneider.
Agonus Bloch and Schneider, Syst. Ichth., 1801, 104 (*Cottus cataphractus* Linnæus).
Aspidophorus Lacépède, Hist. Nat. Poiss., III, 1802, 221 (*A. armatus* Lacépède = *Cottus cataphractus* Linnæus).
Phalangistes Pallas, Zoog. Rosso-Asiat., III, 1811, 113 (*Cottus cataphractus* L.).
3099. *Agonus cataphractus* (Linnæus). *Sea poacher*; *Pogge*.
Northern Europe to western Greenland.
Cottus cataphractus Linnæus, Syst. Nat., Ed. X, 1758, 264; after Artedi.
Cottus broadus Bonnaterre, Encycl. Meth., 1788, 67, northern seas.
Aspidophorus armatus Lacépède, Hist. Nat. Poiss., III, 1802, 222 Atlantic Ocean.
Cataphractus schoneveldi Fleming, British Animals, 1842, 216, Great Britain.
Aspidophorus europæus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 147 (201), sandy shores of northern Europe.
- Genus 1087. STELGIS Cramer.
Stelgis Cramer, in Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 821 (*Agonus vulsus* Jordan and Gilbert).
3100. *Stelgis vulsus* (Jordan and Gilbert).
Off San Francisco.
Agonus vulsus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 330, Point Reyes, near San Francisco.
- Genus 1088. AVERRUNCUS Jordan and Starks.
Averruncus Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 821, 824 (*A. emmelane* Jordan and Starks).
3101. *Averruncus emmelane* Jordan and Starks.
Puget Sound, near Port Orchard.
Averruncus emmelane Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 821, pl. 91, Port Orchard, Puget Sound.
3102. *Averruncus sterletus* Gilbert.
Coast of southern California.
Averruncus sterletus Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2072, off Coronado Island.
- Genus 1089. XYSTES Jordan and Starks.
Xystes Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 824 (*X. axinophrys* Jordan and Starks).
3103. *Xystes axinophrys* Jordan and Starks.
Puget Sound to Monterey.
Xystes axinophrys Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 824, pl. 92, Port Orchard, Admiralty Inlet.
- Genus 1090. SARRITOR Cramer.
Sarritor Cramer, in Jordan and Evermann, Check List, 1895, 448, and Fishes of North and Mid. Amer., 1898, 2072 (*Odontopyxis frenatus* Gilbert).
3104. *Sarritor frenatus* (Gilbert).
Coast of Alaska.
Odontopyxis frenatus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 435; pl. 30, fig. 3. Alaska and Aleutian Islands, at Albatross stations 3219, 3225 and many others, in 16 to 351 fathoms.
3105. *Sarritor leptorhynchus* (Gilbert).
Coast of Alaska.
Odontopyxis leptorhynchus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 437, north and south of the Alaska peninsula, at Albatross stations 3215, 3219, and others in 32 to 59 fathoms.
- Genus 1091. BATHYAGONUS Gilbert.
Bathyagonus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 89 (*B. nigr-pinus* Gilbert).

- 3106. *Bathyagonus nigripinnis* Gilbert.**
North Pacific, known from the Aleutian Islands, Bering Sea, south to the coast of Washington
Bathyagonus nigripinnis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 89, coast of Washington, at *Albatross* station 3073, in 477 fathoms.
- Genus 1092. XENERETMUS Gilbert.**
Xenochirus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 90 (*X. triacanthus* Gilbert); name preoccupied.
Xeneretmus Gilbert in Jordan, Amer. Nat., 1903, 360 (*Xenochirus triacanthus* Gilbert).
- 3107. *Xeneretmus triacanthus* (Gilbert).**
Pacific coast, from Santa Barbara to Washington.
Xenochirus triacanthus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 91, off coast of California, Oregon, and Washington, at *Albatross* stations 2893, 2973, and 3059, in 145, 68, and 77 fathoms.
- Genus 1093. ASTEROTHECA Gilbert.**
Asterothea Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 343 (*Xenochirus pentacanthus* Gilbert).
- 3108. *Asterothea pentacantha* (Gilbert).**
Coast of Washington.
Xenochirus pentacanthus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 91, coast of Washington, at *Albatross* station 3076, in 178 fathoms.
- 3109. *Asterothea alascana* (Gilbert).**
Puget Sound to Bering Sea; coast of Washington.
Xenochirus alascanus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 438, Unimak Pass, Aleutian Islands, at *Albatross* stations 3216, 3219, and others, in 35 to 138 fathoms.
- 3110. *Asterothea infraspinata* (Gilbert).**
Off coast of Washington.
Xeneretmus infraspinatus Gilbert, Proc. Calif. Acad. Sci., 3d ser., III, 1904, 262, pl. 27, off Cape Flattery, Wash., in 77 fathoms.
- Genus 1094. XENOPYXIS Gilbert.**
Xenopyxis Gilbert, Proc. U. S. Nat. Mus., XLIII, 1915, 345 (*Xeneretmus latifrons* Gilbert).
- 3111. *Xenopyxis latifrons* (Gilbert).**
North Pacific coast from Oregon to San Diego.
Xenochirus latifrons Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 92, off coast of Oregon and San Diego, at *Albatross* stations 2898, 2935, etc., in 61 and 158 fathoms.
- 3112. *Xenopyxis leiops* (Gilbert).**
Off southern California.
Xeneretmus leiops Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 348, pl. 17, fig. 10, off Santa Catalina Island.
- 3113. *Xenopyxis ritteri* (Gilbert).**
Off San Diego, Calif.
Xeneretmus ritteri Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 350, pl. 17, fig. 11, off San Diego, in 176 to 181 fathoms.
- Genus 1095. ODONTOPYXIS Lockington.**
Odontopyxis Lockington, Proc. U. S. Nat. Mus., II, 1879, 328 (*O. trispinosus* Lockington).
- 3114. *Odontopyxis trispinosus* Lockington.**
Pacific coast, Alaska to Santa Barbara.
Odontopyxis trispinosus Lockington, Proc. U. S. Nat. Mus., II, 1879, 328, San Francisco.
- Genus 1096. BOTHRAGONUS Gill.**
Bothragonus Gill, in Jordan and Gilbert, Synopsis, 1883, 728 (*Hypsagonus swanii* Steindachner).
- 3115. *Bothragonus swanii* (Steindachner).**
Puget Sound.
Hypsagonus swanii Steindachner, Ichth. Beitr., V, July, 1876, 144, pl. 4, Sitzb. Akad. Wiss. Wien, LXXIV, Port Townsend, Wash.

Family 241. ASPIDOPHOROIDIDÆ

Genus 1097. *ULCINA* Cramer.

Ulcina Cramer, in Jordan and Evermann, Check List, 1895, 449, 1896
(*Aspidophoroides olriki* Lutken)

3116. *Ulcina olriki* (Lutken).

Arctic Ocean; known from west coast of Greenland, Davis Strait, Kara Sea, Barents Bay, and Nova Zembla
Aspidophoroides olriki Lutken, Forelob. Meddel. om Nord Ulkefiske Vidsensk. Meddel. Naturhist. Foren. Kjøb., 1876, 386, 3 figs, Greenland.

3117. *Ulcina güntheri* (Bean).

Coast of Alaska.

Aspidophoroides guntheri Bean, Proc. U. S. Nat. Mus., VIII, 1885, 74, Alaska.

Genus 1098. ASPIDOPHOROIDES Lacépède.

Aspidophoroides Lacépède, Hist. Nat. Poiss., III, 1802, 228 (*A. tranquebar* Lacépède = *Cottus monopterygius* Bloch).

Canthirhynchus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 272 (*A. monopterygius* Bloch).

3118. *Aspidophoroides monopterygius* (Bloch). *Sea poacher*.

Greenland to Cape Cod; accidental to New York

Cottus monopterygius Bloch, Ichth., II, 1786, 156, pl. 178, figs 1, 2, Tranquebar; an error.

Cottus indicus Bonnaterre, Tableau Encycl. Meth., 1788, 68, pl. 87, fig. 367, India; after Bloch

Aspidophoroides tranquebar Lacépède, Hist. Nat. Poiss., III, 1802, 228, Tranquebar

Aspidophoroides grænländicus Valenciennes, in Cuvier, Règne Anim., ed. 2, II, 1829, pl. 21, fig. 3, Greenland.

3119. *Aspidophoroides bartoni* Gilbert.

Aleutian Islands.

Aspidophoroides bartoni Gilbert, Rept. U. S. Fish Com., XIX, 1893, 434, Aleutian Islands and Bristol Bay, Alaska, at Albatross stations 3213, 3223, and many others, in 11½ to 121 fathoms.

Genus 1099. ANOPLAGONUS Gill.

Anoplagonus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 259 (*Aspidophoroides inermis* Gunther).

3120. *Anoplagonus inermis* (Günther).

Coasts of Alaska, south to Vancouver Island.

Aspidophoroides inermis Gunther, Cat., II, 1860, 524, Vancouver Island.

Family 242. CYCLOPTERIDÆ

Genus 1100. CYCLOPTERUS Linnæus.

Cyclopterus Linnæus, Syst. Nat., Ed. X, 1758, 260 (*C. lumpus* Linnæus).

Lumpus Oken, Isis, 1817, 1782; based on "les Lumps" Cuvier (*Cyclopterus lumpus* Linnæus).

3121. *Cyclopterus lumpus* Linnæus. *Lumpfish; Cock and hen paddle; Sea owl; Paddle; Licorne de mer; Lump sucker.*

North Atlantic; rocky shores of both coasts, south to Cape Cod and France.

Cyclopterus lumpus Linnæus, Syst. Nat., Ed. X, 1758, 260, Baltic and North Sea.

Cyclopterus minutus Pallas, Spicilegium, Zool., VII, 1769, 12, Atlantic Ocean.

Cyclopterus pavoninus Shaw, Nat. Hist., IX, 1797, pl. 310.

Cyclopterus pyramidatus Shaw, Gen. Zool., V, 1804, 390.

Cyclopterus canuleus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 480, New York Harbor

Cyclopterus coronatus Couch, Nat. Hist. Cornwall, in Trans. Linn. Soc. London, XIV, 1823, 47, Cornwall.

Lumpus vulgaris McMurtrie, Eng. ed. Cuvier, Anim. King, 1831; after
C. lumpus Linnæus.
Lumpus anglorum DeKay, N. Y. Fauna: Fishes, IV, 1842, 305, after
 Linnæus.

Genus 1101. **EUMICROTREMUS** Gill.

Eumicrotremus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 190
 (*Cyclopterus spinosus* Müller).

3122. *Eumicrotremus spinosus* (Müller). *Spinous lumpfish*.
 North Atlantic and Arctic Oceans, south to Maine and Denmark; and
 Eastport, Me.
Cyclopterus spinosus Müller, Prodr. Zool. Dan., IX, 1777, Denmark.

3123. *Eumicrotremus orbis* (Günther).
 Aleutian Islands, south to Vancouver Island.
Cyclopterus orbis Günther, Cat., III, 1861, 158, Vancouver Island.

3124. *Eumicrotremus phrynoides* Burke.
 Bering Sea.
Eumicrotremus phrynoides Burke, Bull. U. S. Bur. Fish., XXX, 1910,
 69, Petrel Bank, Alaska.

Genus 1102. **LETHOTREMUS** Gilbert.

Lethotremus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 449 (*L. muticus*
 Gilbert).

3125. *Lethotremus muticus* Gilbert.
 Aleutian Islands.
Lethotremus muticus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 449,
 pl. 31, Unimak Pass, Alaska, at Albatross stations 3223 and 3258,
 in 56 to 70 fathoms.

3126. *Lethotremus vinolentus* Jordan and Starks.
 Puget Sound, Wash.
Lethotremus vinolentus Jordan and Starks, Proc. Calif. Acad. Sci., 2d
 ser., V, 1895, 827, pl. 94, Puget Sound, near Seattle, Wash.

Genus 1103. **CYCLOPTEROIDES** Garman.

Cyclopteroides Garman, Mem. Mus. Comp. Zool., XIV, 1892, 37 (*C.*
gyrinops Garman).

3127. *Cyclopteroides gyrinops* Garman.
 Pribilof Islands, Alaska.
Cyclopteroides gyrinops Garman, Mem. Mus. Comp. Zool., XIV, 1892,
 37, St. Paul Island.

Family 243. **LIPAROPIDÆ**

Genus 1104. **APTOCYCLUS** De la Pylaie.

Aptocyclus De la Pylaie, Congrès Scientifique Potiers, 1834, 529 (*Cy-*
clopterus ventricosus Pallas; misprinted *Cyclogasterus*).
Cycloptericthys Steindachner, Ichth. Beitr., X, 1881, 14, 192 (*C.*
glaber Steindachner=*Cyclopterus ventricosus* Pallas).

3128. *Aptocyclus ventricosus* (Pallas).
 Kamchatka; Sea of Okhotsk.
Cyclopterus ventricosus Pallas, Spicilegium Zool., VII, 1770, 15, t. 2, no
 locality given, somewhere about Kamchatka.
Cycloptericthys glaber Steindachner, Ichth. Beitr., X, 1881, 14, pl. 8,
 Sea of Okhotsk.

Genus 1105. **LIPAROPS** Garman.

Liparops Garman, Mem. Mus. Comp. Zool., XV, 1892, 42 (*Cyclopterus*
stelleri Pallas).

3129. *Liparops stelleri* (Pallas).
 Peter and Paul Harbor, Kamchatka.
Cyclopterus stelleri Pallas, Zoogr. Rosso-Asiatica, III, 1811, 73, Peter
 and Paul Harbor, Kamchatka.

Family 244. LIPARIDÆ. Sea snails

Genus 1106. POLYPERA Burke.

Polypera Burke, Proc. U. S. Nat. Mus., XLIII, 1913, 567 (*Liparis greeni* Jordan and Starks).

3130. *Polypera greeni* (Jordan and Starks).

Puget Sound at Victoria, British Columbia.

Neoliparis greeni Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 829, pl. 96, figs. 2 and 3, Victoria, British Columbia.

Genus 1107. NEOLIPARIS Steindachner.

Neoliparis Steindachner, Ichth. Beitr., III, 1875, 54 (*Liparis mucosus* Ayres).

3131. *Neoliparis atlanticus* Jordan and Evermann. Sea snail; Lump sucker.

Common along rocky shores from Newfoundland to Cape Cod.

Neoliparis atlanticus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2107, Godbout, Quebec.

Liparis donlagui of authors, not of Donovan, the European cognate.

3132. *Neoliparis mucosus* (Ayres).

Pacific coast from Puget Sound to San Francisco; rare.

Liparis mucosus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 22, San Francisco.

Neoliparis floræ Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 830, pl. 96, fig. 1, Waadda Island, Straits of Juan de Fuca.

3133. *Neoliparis fissuratus* Starks.

Puget Sound.

Neoliparis fissuratus Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 560, Port Ludlow, Puget Sound.

3134. *Neoliparis rutteri* Gilbert and Snyder.

Kodiak Island, Alaska.

Neoliparis rutteri Gilbert and Snyder, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2108, Ugah Bay, Kodiak Island.

3135. *Neoliparis bristolensis* (Burke).

Bristol Bay, Bering Sea.

Cyclogaster bristolensis Burke, Proc. U. S. Nat. Mus., XLIII, 1913, 567, Bristol Bay in about 20 fathoms.

3136. *Neoliparis calliodon* (Pallas).

Coasts of Alaska, north to Kamchatka and Bering Sea; recorded from St. Paul, Kamchatka, Plover Bay, Siberia, Unalaska, and Kodiak.

Cyclopterus calliodon Pallas, Zoogr. Rosso-Asiat., III, 1811, 75, Kamchatka and Aleutian Islands.

3137. *Neoliparis grebnitzkii* Schmidt.

Commander Islands.

Neoliparis grebnitzkii Schmidt, Fauna Mare Orientale, 1875, 61, Bering Island.

3138. *Neoliparis beringianus* (Gilbert and Burke).

Bering Island.

Cyclogaster (*Neoliparis*) *beringianus* Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 72; Nikolski, Bering Island.

3139. *Neoliparis micraspidophorus* (Burke).

Bering Island.

Cyclogaster (*Neoliparis*) *micraspidophorus* Burke, Bull. U. S. Bur. Fish., XXX, 1910, 71, Nikolski, Bering Island.

Genus 1108. LIPARIS Scopoli.

Cyclogaster ⁷⁷ Gronow, Act. Helvet., IV, 1760, 265 (*Cyclopterus liparis* Linnæus); nonbinomial.

Cyclogaster Gronow, Zoophyl., No. 198, 1763; nonbinomial.

Liparis (Artedi) Scopoli, Introd. Hist. Nat., 1777, 453 (*Cyclopterus liparis* Linnæus).

Cyclogaster Gronow, Cat. Fishes, ed. Gray, 1854, 40 (*Cyclopterus liparis* L.).

⁷⁷ We do not find that Gronow used the name *Cyclogaster* as binomial. Although later than Linnæus (1758), he had not then adopted, most likely had not heard of, the Linnæan system.

3140. *Liparis liparis* (Linnaeus). *Sea snail; Striped sea snail.*
 North Atlantic, on both shores, south to Connecticut and France,
 common north; very variable, a dozen color varieties having received
 names.
Cyclopterus liparis Linnaeus, Syst. Nat., Ed. XII, I, 1766, 414, northern
 ocean; after Artedi and Gronow.
Cyclopterus lineatus Lepechin, Nov. Comm. Petropol., XVIII, 1774,
 520, White Sea.
Cyclopterus liparis minor Fabricius, Fauna Greenlandica, 1780, 135,
 Greenland
Cyclopterus musculus Lacépède, Hist. Nat. Poiss., IV, 1802, 683, near
 Havre.
3141. *Liparis osborni* Townsend and Nichols
 Off southern California in 440 fathoms
Liparis osborni Townsend and Nichols, Bull. Amer. Mus. Nat. Hist.,
 LII, 1925, 14, fig. 4, off southern California.
3142. *Liparis cyclopus* Gunther
 Aleutian Islands to Puget Sound.
Liparis cyclopus Gunther, Cat., III, 1861, 162, Esquimault Harbor,
 Vancouver Island.
3143. *Liparis megacephalus* (Burke).
 Southeastern Bering Sea.
Cyclogaster megacephalus Burke, Proc. U. S. Nat. Mus., XLIII, 1913,
 569, southeastern Bering Sea at 37 fathoms.
3144. *Liparis fucensis* Gilbert
 Straits of Fuca
Liparis fucensis Gilbert, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 837,
 and Rept U S Fish Com., XIX, 1893, 447, Port Angeles, Straits of
 Juan de Fuca.
- Genus 1109. *CARELIPARIS* Garman.
Careliparis Garman, Mem. Mus. Comp. Zool., XIV, 1892, 62 (*Liparis*
agassizii Putnam).
3145. *Careliparis tunicatus* (Reinhardt).
 Greenland.
Liparis tunicata Reinhardt, Overs. Kong. Danske Vidensk. Selsk., VI,
 CXI, 1836, Greenland; after Fabricius
Liparis fabricii Krøyer, Naturh. Tidsskr., II, 1847, 274, Greenland;
 D. XIII, 27; A. II, 30; P. 34; C. 9; Vert. 46.
Liparis arctica Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 191, Port
 Foulke, Greenland.
3146. *Careliparis agassizii* (Putnam)
 Bering Sea; Sakhalin Island.
Liparis agassizii Putnam, Proc. Amer. Assoc. Adv. Sci., 1874, 339,
 Sakhalin Channel of Tartary.
Liparis gibbus Bean, Proc. U. S. Nat. Mus., IV, 1881, 148, Unalaska,
 St. Paul Island, Indian Point, Cape Chaplin, and Plover Bay, Siberia.
3147. *Careliparis herschelinus* (Scofield)
 Arctic Ocean.
Liparis herschelinus Scofield, in Jordan and Evermann, Fishes North
 and Mid Amer., 1898, 2123, Herschel Island, Arctic Ocean, near
 mouth of the Mackenzie.
3148. *Careliparis dennyi* (Jordan and Starks).
 North Pacific, from Aleutian Islands to Puget Sound.
Liparis dennyi Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V,
 1895, 835, pl. 98, Admiralty Inlet near Seattle.
- Genus 1110. *LYOLIPARIS* Jordan and Evermann.
Lyoliparis Jordan and Evermann, Check List, 1895, 451 (*Liparis*
pulchellus Ayres).
3149. *Lyoliparis pulchellus* (Ayres).
 Point Reyes, Calif.; northern Pacific, Alaska to Monterey.
Liparis pulchellus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 22, San
 Francisco.

- Genus 1111. ACTINOCHIR** Gill.
Actinochir Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 190, 193 (*A. major* Gill)
- 3150. Actinochir major** Gill.
 Coasts of Greenland.
Actinochir major Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 193, Greenland
Cyclopterus liparis major Fabricius, Fauna Groenland, 1780, 136, Greenland; *major* is here simply an adjective.
- Genus 1112. CRYSTALLICHTHYS** Jordan and Gilbert.
Crystallichthys Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2364 (*C. mirabilis* Jordan and Gilbert).
- 3151. Crystallichthys mirabilis** Jordan and Gilbert.
 Bering Sea.
Crystallichthys mirabilis Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2365, Provostmaya, Kamchatka.
- 3152. Crystallichthys cyclospilus** Gilbert and Burke.
 Bering Sea, Petrel Bank, St. Paul Island.
Crystallichthys cyclospilus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 74, Petrel Bank.
- 3153. Crystallichthys cyclostigma** (Gilbert).
 Bristol Bay, Alaska
Liparis cyclostigma Gilbert, Rept. U. S. Fish Com., XIX, 1893, 446, Bristol Bay, Alaska, at *Albatross* station 3252, in 29½ fathoms.
- Genus 1113. BATHYPHASMA** Gilbert.
Bathyphasma Gilbert, Rept. U. S. Fish Com., XIX, 1893, 447 (*B. ovigerum* Gilbert).
- 3154. Bathyphasma ovigerum** Gilbert.
 Off Queen Charlotte Island, British Columbia.
Bathyphasma ovigerum Gilbert, Rept. U. S. Fish Com., XIX, 1893, 448, Queen Charlotte Islands, at *Albatross* station 3342 in 1,588 fathoms.
- Genus 1114. CAREMITRA** Jordan and Evermann.
Caremitra Jordan and Evermann, Check List, 1895, 452 (*Careproctus simus* Gilbert).
- 3155. Caremitra sima** (Gilbert).
 Off Unalaska Island
Careproctus simus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 444, off Unalaska, at *Albatross* station 3331 in 350 fathoms.
- Genus 1115. LIPARISCUS** Gilbert.
Lipariscus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 358 (*L. nanus* Gilbert).
- 3156. Lipariscus nanus** Gilbert
 Monterey Bay, Calif.
Lipariscus nanus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 358, pl. 19, Monterey Bay in 32 to 309 fathoms.
- Genus 1116. EURYMEN** Gilbert and Burke.
Eurymen Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 64 (*E. gyrinus* Gilbert and Burke).
- 3157. Eurymen gyrinus** Gilbert and Burke
 Bering Sea.
Eurymen gyrinus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 64, Avatcha Bay, east coast of Kamchatka.
- Genus 1117. ACANTHOLIPARIS** Gilbert and Burke.
Acantholiparis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 83 (*A. opercularis* Gilbert and Burke).
- 3158. Acantholiparis opercularis** Gilbert and Burke.
 East coast of Kamchatka
Acantholiparis opercularis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 83, Starichkof Island.

- Genus 1118. *NECTOLIPARIS* Gilbert and Burke.
Nectoliparis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 82 (*N. pelagicus* Gilbert and Burke).
3159. *Nectoliparis pelagicus* Gilbert and Burke.
 Bering Sea to Hokkaido, Japan; pelagic.
Nectoliparis pelagicus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 82, fig. 27, Bering Sea.
- Genus 1119. *ELASSODISCUS* Gilbert and Burke.
Elassodiscus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 81 (*E. tremebundus* Gilbert and Burke).
3160. *Elassodiscus tremebundus* Gilbert and Burke.
 Coast of Kamchatka.
Elassodiscus tremebundus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 81, Avatcha Bay.
- Genus 1120. *CAREPROCTUS* Krøyer.
Careproctus Krøyer, Naturh. Tidskr., I, 1862, 253-257 (*Liparis reinhardi* Krøyer).
Allochir Jordan and Evermann, Check List, 1895, 452 (*Careproctus melanurus* Gilbert).
Allurus Jordan and Evermann, loc. cit., 452 (*Careproctus ectenes* Gilbert).
Alnectes Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2866 (*Careproctes ectenes* Gilbert, substitute for *Allurus*, preoccupied).
- Subgenus *CAREPROCTUS* Krøyer.
3161. *Careproctus reinhardi* (Krøyer).
 Greenland, Jan Mayen, and Beeren Island.
Liparis reinhardi Krøyer, Naturh. Tidskr., I, 1862, 252, Greenland.
3162. *Careproctus colletti* Gilbert.
 South of Alaska Peninsula to Sea of Okhotsk.
Careproctus colletti Gilbert, Rept. U. S. Fish Com., XIX, 1893, 442, south of Alaska Peninsula, at Albatross station 3338 in 625 fathoms.
3163. *Careproctus phasma* Gilbert.
 Bristol Bay, Alaska.
Careproctus phasma Gilbert, Rept. U. S. Fish Com., XIX, 1893, 443, Bristol Bay, Alaska, at Albatross stations 3254 and 3256 in 46 and 49 fathoms.
3164. *Careproctus spectrum* Bean.
 Between Unga and Nagai Islands, Alaska.
Careproctus spectrum Bean, Proc. U. S. Nat. Mus., XIII, 1890, 40, between Unga and Nagai Islands, 160° 18' W., 55° 10' N., at Albatross station 2848 in 110 fathoms.
3165. *Careproctus ranulus* (Goode and Bean).
 Off Halifax Harbor.
Liparis ranula Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 46, Halifax.
3166. *Careproctus gelatinosus* (Pallas).
 Peter and Paul Harbor, Kamchatka.
Cyclopterus gelatinosus Pallas, Spicilegia, VII, 1769, 19, Peter and Paul Harbor, Kamchatka.
3167. *Careproctus ostentum* Gilbert.
 Unalaska Island.
Careproctus ostentum Gilbert, Rept. U. S. Fish Com., XIX, 1893, 444, north of Unalaska Island, at Albatross stations 3324 and 3331 in 109 and 350 fathoms.
3168. *Careproctus furcellus* Gilbert and Burke.
 Bering Sea.
Careproctus furcellus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 80, between Petrel Bank and Agattu Island, Bering Sea.
3169. *Careproctus gilberti* Burke.
 Southeastern Alaska and British Columbia.
Careproctus gilberti Burke, Proc. U. S. Nat. Mus., XLIII, 1913, 570, Kodiak Island and neighborhood in 102 to 482 fathoms.

3170. *Careproctus candidus* Gilbert and Burke.
Attu Island, Aleutian Chain.
Careproctus candidus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX,
1910, 77, fig. 22, East Cape, Attu Island, Alaska.
3171. *Careproctus mollis* Gilbert and Burke.
Attu Island, Aleutian Chain
Careproctus mollis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX,
1910, 77, fig. 21, East Cape, Attu Island, Alaska.
3172. *Careproctus bowersianus* Gilbert and Burke.
Bering Sea.
Careproctus bowersianus Gilbert and Burke, Bull. U. S. Bur. Fish.,
XXX, 1910, 76, fig. 20, Bowers Bank, Bering Sea.
3173. *Careproctus opisthotremus* Gilbert and Burke.
Bering Sea.
Careproctus opisthotremus Gilbert and Burke, Bull. U. S. Bur. Fish.,
XXX, 1910, 78, fig. 23, Petrel Bank, between Attu and Agattu
Islands, Alaska.
3174. *Careproctus longifilis* Garman.
Off Galapagos Islands and Panama in deep water.
Careproctus longifilis Garman, Mem. Mus. Comp. Zool., XIV, 1892, 9,
southwest Panama.
3175. *Careproctus longipinnis* Burke.
Arctic Ocean; Faroe Islands to Beeren Island.
Careproctus longipinnis Burke, Ann. Mag. Nat. Hist., 8th ser., IX,
1912, 510, Ingolf station 139, north of the Faroe Islands, depth 702
Danish fathoms.
3176. *Careproctus melanurus* Gilbert.
Off coast of California and Oregon.
Careproctus melanurus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 560,
off coasts of California and Oregon, at Albatross stations 2840, 2891,
and others in 178 to 339 fathoms.
3177. *Careproctus attenuatus* Gilbert and Burke.
Bering Sea.
Careproctus attenuatus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX,
1910, 79, Petrel Bank and Agattu Island, Alaska in 482 fathoms.
- Subgenus ALLINECTES Jordan and Evermann.
3178. *Careproctus ectenes* Gilbert.
Unalaska Island.
Careproctus ectenes Gilbert, Rept. U. S. Fish Com., XIX, 1893, 442,
north of Unalaska, at Albatross station 3331 in 350 fathoms.
- Genus 1121. PROGNURUS Jordan and Evermann.
Prognurus Jordan and Evermann, Fishes North and Mid. Amer., 1896,
2866 (*P. cypselurus* Jordan and Gilbert).
3179. *Prognurus cypselurus* Jordan and Gilbert.
Bering Sea and north Pacific.
Prognurus cypselurus Jordan and Gilbert, in Jordan and Evermann,
Fishes North and Mid. Amer., 1898, 2866, Bogoslof Island, Bering
Sea.
- Genus 1122. GYRINICHTHYS Gilbert.
Gyrinichthys Gilbert, Rept. U. S. Fish Com., XIX, 1893, 444 (*G. miny-
tremus* Gilbert).
3180. *Gyrinichthys minytremus* Gilbert.
Off Unalaska Island, Alaska.
Gyrinichthys minytremus Gilbert, Rept. U. S. Fish Com., XIX, 1893,
444, north of Unalaska at Albatross station 3331 in 350 fathoms.

Genus 1123. PARALIPARIS⁷⁸ Collett.

- Paraliparis* Collett, Vid. Selsk. Forsk. Christiania, No. 14, 1878, 32 (*Paraliparis bathybi* Collett); species allied to *P. cephalus* with narrow gill openings.
- Amitra* Goode, Proc. U. S. Nat. Mus., III, 1880, 478 (*A. liparina* Goode).
- Monomitra* Goode, loc. cit., VI, 1883, 109 (*Amitra liparina* Goode); substitute for *Amitra* regarded as preoccupied by *Amitrus* Schonherr, a beetle.
- Hilgendorfia* Goode and Bean, Oceanic Ichthyol., 1895, 280 (*Paraliparis membranaceus* Gunther).
- Amitrichthys* Jordan and Evermann, Check List, 1895, 453 (*Paraliparis cephalus* Gilbert).
3181. *Paraliparis holomelas* Gilbert.
North of Unalaska Island, Alaska.
Paraliparis holomelas Gilbert, Rept. U. S. Fish Com., XIX, 1893, 441, north of Unalaska, at Albatross stations 3308 and 3332 in 406 and 1,625 fathoms.
3182. *Paraliparis cephalus* Gilbert.
Alaska to California.
Paraliparis cephalus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 561, off California and Oregon, at Albatross station 2919 in 984 fathoms.
3183. *Paraliparis mento* Gilbert.
Coast of Oregon.
Paraliparis mento Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 562, off coast of Oregon, at Albatross station 3071 in 685 fathoms.
3184. *Paraliparis albescens* Gilbert.
Off Monterey Bay, Calif.
Paraliparis albescens Gilbert, Proc. U. S. Nat. Mus., XLVIII, 355, pl. 18, fig. 13, off Monterey Bay.
3185. *Paraliparis caudatus* Gilbert.
Monterey Bay, Calif.
Paraliparis caudatus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 356, pl. 18, fig. 14, Monterey Bay.
3186. *Paraliparis deani* Burke.
Bering Sea; once taken in Monterey Bay.
Paraliparis deani Burke, Proc. U. S. Nat. Mus., XLIII, 1912, 571, Stevens Passage, Alaska.
3187. *Paraliparis dactylosus* Gilbert.
Off Santa Cruz, Calif.
Paraliparis dactylosus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 469, pl. 34, fig. 2, off Santa Cruz, Calif., at Albatross station 3112, in 296 fathoms.
3188. *Paraliparis grandiceps* Garman.
Off Lower California in deep water.
Paraliparis grandiceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, off Lower California.
3189. *Paraliparis rosaceus* Gilbert.
Pacific coast of Mexico.
Paraliparis rosaceus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 93, Pacific coast of Mexico, at Albatross stations 2898, 2935, and others in 61 to 158 fathoms.
3190. *Paraliparis fimbriatus* Garman.
Off Colombia and Galapagos Islands in deep water.
Paraliparis fimbriatus Garman, in Mem. Mus. Comp. Zool., XIV, 1892, 9, southwest of Panama.
3191. *Paraliparis attenuatus* Garman.
Off Bay of Panama in deep water.
Paraliparis attenuatus Garman, Deep Sea Fishes, Mem. Mus. Comp. Zool., XXIV, 1899, 118, pl. XXVII, off Bay of Panama.

⁷⁸ This group of degenerate liparids is excessively abundant in all northern sea bottoms at considerable depth. It should, no doubt, be subdivided, but the present knowledge of the group hardly permits this.

3192. *Paraliparis angustifrons* Garman.
Off Bay of Panama in deep water.
Paraliparis angustifrons Garman, Mem. Mus. Comp. Zool. XXIV, 1899, 119, Pl. XXVII, Albatross station 3394, off Bay of Panama.
3193. *Paraliparis latifrons* Garman.
Off Bay of Panama in deep water.
Paraliparis latifrons Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 120, Pl. XXVII, off Bay of Panama.
3194. *Paraliparis garmani* Burke
Off coast of New England.
Paraliparis garmani Burke, Proc. U. S. Nat. Mus., XLIII, 1912, 572, off coast of New England in 328 to 542 fathoms.
3195. *Paraliparis copei* Goode and Bean
Gulf Stream
Paraliparis copei Goode and Bean, Oceanic Ichth., 1896, 279, fig. 353, Gulf Stream, at Albatross station 2232, 39° 12' 17" N., 72° 9' 30" W., in 520 fathoms.
- Subgenus *AMITRA* Goode and Bean.
3196. *Paraliparis liparinus* (Goode)
Atlantic Ocean, off Rhode Island.
Amitra liparina Goode, Proc. U. S. Nat. Mus., III, 1880, 478, off Rhode Island.
- Subgenus *HILGENDORFIA* Goode and Bean.
3197. *Paraliparis ulochir* Gilbert.
Gulf of California to Unalaska
Paraliparis ulochir Gilbert, Rept. U. S. Fish Com., XIX, 1893, 441, Gulf of California, at Albatross station 3010 in 1005 fathoms.
- Genus 1194. *RHINOLIPARIS* Gilbert.
Rhinoliparis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 445 (*R. barbulifer* Gilbert).
3198. *Rhinoliparis barbulifer* Gilbert.
Bering Sea.
Rhinoliparis barbulifer Gilbert, Rept. U. S. Fish Com., XIX, 1893, 445, north of Unalaska, Alaska, at Albatross stations 3237, 3325, and others, in 225 to 576 fathoms.
3199. *Rhinoliparis attenuatus* Burke.
Bering Sea.
Rhinoliparis attenuatus Burke, Proc. U. S. Nat. Mus., XLIII, 1912, 573, Bering Sea in 576 fathoms.

Family 245. *PERISTEDIIDÆ*

- Genus 1125. *PERISTEDION* Lacépède.
Peristedion Lacépède, Hist. Nat. Poiss., III, 1802, 368 (*P. malarma* Lacépède = *Trigla cataphracta* L.).
Peristethus Kaup, amended spelling.
3200. *Peristedion brevirostre* (Günther).
Caribbean Sea.
Peristethus brevirostre Günther, Cat., II, 1860, 217, West Indies.
3201. *Peristedion miniatum* Goode.
Gulf Stream.
Peristedion miniatum Goode, Proc. U. S. Nat. Mus., III, 1880, 349, Gulf Stream, off Rhode Island.
3202. *Peristedion longispathum* (Goode and Bean).
West Indies, off Cuba and Barbados.
Peristedum longispatha Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, No. 5, 166, off Santa Cruz Island, West Indies, and off Barbados.
3203. *Peristedion gracile* Goode and Bean.
Gulf of Mexico and Porto Rico
Peristedion gracile Goode and Bean, Oceanic Ichth., 1896, 473, Gulf of Mexico.

3204. *Peristedion platycephalum* (Goode and Bean).
West Indies; Barbados
Peristedium platycephalum Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, No 5, 167, off Barbados.
3205. *Peristedion barbiger* (Garman).
Off Bay of Panama in deep water.
Peristedium barbiger Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 110, off Bay of Panama.
3206. *Peristedion crustosum* (Garman).
Off Bay of Panama in deep water
Peristedium crustosum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 112, Pl. A, off Bay of Panama.
- Genus 1126. *VULSICULUS* Jordan and Evermann.
Vulsiculus Jordan and Evermann, Check List, 1895, 489 (*Peristedion imberbe* Poey).
3207. *Vulsiculus imberbis* (Poey).
Gulf of Mexico
Pterystedion imberbe Poey, Memorias, II, 1861, 367, Cuba; apparently never described.
Peristedion micronemus Poey, Ann. Lyc. Nat. Hist. N. Y., IX, 1870, 321, Cuba; substitute name on discovery of the minute barbels.

Family 246. TRIGLIDÆ

- Genus 1127. *COLOTRIGLA* Gill.
Colotrigla Gill, Proc. U. S. Nat. Mus., XXVII, 1905, 240 (*Prionotus stearnsi* Jordan and Swain).
3208. *Colotrigla stearnsi* (Jordan and Swain).
Gulf of Mexico.
Prionotus stearnsi Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 541, Pensacola, Fla.
- Genus 1128. *GURNARDUS* Jordan and Evermann.
Gurnardus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2152 (*Prionotus gymnotethus* Gilbert).
3209. *Gurnardus birostratus* (Richardson).
Gulf of Fonseca, Pacific coast of Central America.
Prionotus birostratus Richardson, Voy. Sulphur, Ichth., pt. II, 1845, 81, April, Gulf of Fonseca, Pacific coast of Central America.
3210. *Gurnardus gymnotethus* (Gilbert).
Gulf of California.
Prionotus gymnotethus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 559, Gulf of California, in shallow water.
3211. *Gurnardus xenisma* (Jordan and Bollman).
Pacific Ocean, off coast of Colombia.
Prionotus xenisma Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 169, Pacific coast of Colombia.
3212. *Gurnardus loxias* Jordan.
Pacific coast of Central America.
Prionotus loxias Jordan, Proc. U. S. Nat. Mus., XIX, 1896, 452, Bay of Panama.
3213. *Gurnardus ruscarius* (Gilbert and Starks).
Panama Bay; Mazatlan; Magdalena Bay.
Prionotus ruscarius Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 165, Pl. XXVII, Panama Bay.
- Genus 1129. *MERULINUS* Jordan and Evermann.
Merulinus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2148 (*Trigla carolina* Linnaeus).

- 3214. *Merulinus carolinus* (Linnæus).** *Common gurnard; Red-winged sea robin; Pigfish; Rock pigfish; Flyingfish; Spinous gurnard; Small-scaled sea robin; Web-fingered sea robin; Carolina robin; Carolina sea robin.*
Cape Ann to South Carolina, chiefly northward.
Trigla carolina Linnæus, Mantissa, II, 1771, 528, Carolina.
Trigla palmipes Mitchell, Trans. Lit. Phil. Soc. New York, 1815, 431, pl. 4, fig. 5, New York Harbor.
Prionotus pileatus Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 77, Massachusetts Bay.
- 3215. *Merulinus scitulus* (Jordan and Gilbert)**
South Atlantic coast of United States, Beaufort to St. Augustine.
Prionotus scitulus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 288, Beaufort, N. C.
- 3216. *Merulinus roseus* (Jordan and Evermann)**
Gulf of Mexico; Tampa Bay, Pensacola; Tortugas.
Prionotus roseus Jordan and Evermann, Proc. U. S. Nat. Mus., IX, 1886, 470, off Tampa Bay, Fla.
- 3217. *Merulinus salmonicolor* Fowler.**
New River, Dade County, Fla.
Merulinus salmonicolor Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 333, New River, Fla.
- Genus 1130. *PRIONOTUS* Lacépède.** *Gurnards; Sea robins.*
Prionotus Lacépède, Hist. Nat. Poiss., III, 1802, 337 (*Trigla evolans* Linnæus).
Ornichthys Swainson, Classn. Fishes, Amphib., etc., 1839, 262 (*Trigla punctata* Bloch).
Chriolax Jordan and Gilbert, Proc. U. S. Nat. Mus., I, 1878, 374 (*Trigla evolans* Linnæus).
- 3218. *Prionotus evolans* (Linnæus).** *Southern striped gurnard; Lined sea robin; Flyingfish; Striped sea robin.*
South Atlantic coast of United States.
Trigla evolans Linnæus, Syst. Nat., Ed. XII, 1766, 498, Carolinas.
Prionotus sarritor Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 615, Charleston; Beaufort, N. C.
- 3219. *Prionotus strigatus* (Cuvier).** *Northern striped gurnard; Brown-winged sea robin; Sea robin; Sand pigfish; Pigfish; Flyingfish; Red-winged gurnard.*
Atlantic coast of northern States, Cape Cod to Virginia; replacing *P. evolans* northward.
Trigla strigata Cuvier, Règne Anim., ed. 2, II, 1829, 161, New York; after *lineata* of Mitchill.
- 3220. *Prionotus rubio* Jordan.** *Rubio volador.*
West Indies.
Prionotus rubio Jordan, Proc. U. S. Nat. Mus., IX, 1886, 50, Habana.
Rubio volador Parra, Descr. Dif. Piezas de Hist. Nat., 1787, 98, lám. 38, Habana.
- 3221. *Prionotus punctatus* Bloch.** *Spotted sea robin; Rubio volador.*
West Indies and Caribbean Sea; Cameron, La.
Prionotus punctatus Bloch, Ichth., 1793, pl. 353, Martinique; on a drawing by Plumier.
- 3222. *Prionotus ophryas* Jordan and Swain.**
Gulf of Mexico; Snapper Banks off Pensacola, Fla.
Prionotus ophryas Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 542, Snapper Banks off Pensacola, Fla.
- 3223. *Prionotus pectoralis* (Nichols and Breder).**
Coast of Texas.
Prionotus miles pectoralis Nichols and Breder, Proc. Biol. Soc. Wash., XXXVII, 1924, 22, Gulf of Mexico, off Aransas Pass, Tex.
- 3224. *Prionotus stephanophrys* Lockington.**
Deep water off San Francisco, Point Reyes, and Monterey; very rare.
Prionotus stephanophrys Lockington, Proc. U. S. Nat. Mus., III, 1880, 529, Point Reyes, near San Francisco.

3225. *Prionotus quiescens* Jordan and Bollman
Pacific Ocean off the coast of Colombia; Gulf of California.
Prionotus quiescens Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 166, off Pacific coast of Colombia.
3226. *Prionotus frontalis* Garman.
Off Bay of Panama in deep water.
Prionotus frontalis Garman, Deep Sea Fishes, in Mem. Mus. Comp. Zool., XXIV, 1899, 108, off Bay of Panama.
3227. *Prionotus albirostris* Jordan and Bollman.
Pacific Ocean off the coast of tropical America.
Prionotus albirostris Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 168, Pacific Ocean off the coast of Colombia.
3228. *Prionotus beanii* Goode.
Off Trinidad.
Prionotus beanii Goode, in Goode and Bean, Oceanic Ichth., 1896, 468, off Trinidad.
3229. *Prionotus tribulus* Cuvier and Valenciennes. *Big-headed gurnard*; *Sea robin*.
South Atlantic coast from Long Island to Brazos, Tex.
Prionotus tribulus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 72 (98), pl. 74, New York; Carolinas.
? *Trigla carolina* Bloch, Ichth., 1793, 352, Carolinas; not of Linnæus.
3230. *Prionotus horrens* Richardson.
Pacific coast of tropical America.
Prionotus horrens Richardson, Voy. Sulphur, Ichth., 1843, 79, pl. 42, figs 1-3, Gulf of Fonseca.
- Genus 1131. *FISSALA* Gill.
Fissala Gill, Proc. U. S. Nat. Mus., XXVIII, 1905, 342 (*Prionotus alatus* Goode and Bean).
3231. *Fissala alata* (Goode and Bean).
Off Charleston, S. C.
Prionotus alatus Goode and Bean, Bull. Mus. Comp. Zool., X, 1883, 210, deep sea off Charleston, S. C.
- Genus 1132. *BELLATOR* Jordan and Evermann.
Bellator Jordan and Evermann, Check List, 1895, 488 (*Prionotus militaris* Goode and Bean).
3232. *Bellator militaris* (Goode and Bean).
Off Cape Catoche, Yucatan; Gulf of Mexico.
Prionotus militaris Goode and Bean, Oceanic Ichth., 1896, 464, figs. 380 and 384, off Cape Catoche, Yucatan.
3233. *Bellator egretta* (Goode and Bean).
Barbados.
Prionotus egretta Goode and Bean, Oceanic Ichth., 1896, 465, fig. 381, off Barbados, at Blake station 64.

Family 247. DACTYLOPTERIDÆ

- Genus 1133. *DACTYLOPTERUS* Lacépède. *Flying gurnards*.
Cephalacanthus Lacépède, Hist. Nat. Poiss., III, 1802, 323 (*C. spinarella* Lacépède); young examples.
Dactylopterus Lacépède, loc. cit., 325 (*D. pirapeda* Lacépède = *Trigla volitans* Linnæus); adult form.
Cephacandia Rafinesque, Analyse Nature, 1815, 85; substitute for *Cephalacanthus*; too long a name.
Gonocephalus Gronow, Cat. Fish., ed. Gray, 1854, 106 (*G. macrocephalus* Gray = *Trigla volitans* Linnæus).

3234. *Dactylopterus volitans* (Linnaeus). *Flying robin*, *Batfish*; *Volador*, *Flying gurnard*; *Murcielago*; *Civetta de mer*.
Atlantic Ocean on both coasts; also Gulf of Mexico.
Trigla volitans Linnaeus, Syst. Nat., Ed. X, I, 1758, 302; after Artedi.
Hirundo Catesby, Nat. Hist. Carol., II, 1771, Tab. 8, p. 8, Bahamas.
Trigla tentabunda Walbaum, Artedi Piscium, III, 1792, 362; after Klein.
Trigla fasciata Bloch and Schneider, Syst. Ichth., 1801, 16, tab. 3, fig. 1; after Klein.
Dactylopterus pirapeda Lacépède, Hist. Nat. Poiss., III, 1802, 326, Mediterranean and almost all warm seas.
Polynemus sexradiatus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, pl. 4, fig. 10, New York.
Callionymus pelagicus Rafinesque, Amer. Month. Mag., 1818, 205, Atlantic Ocean.
Gonocephalus macrocephalus Gronow, Cat. Fish., ed. Gray, 1854, 106; pelagic.

Order SS. HOLCONOTI

Family 248. EMBIOTOCIDÆ. Surffishes; Viviparous perches

Genus 1134. MICROMETRUS Gibbons.

- Micrometrus* Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer Times and Transcript, May 30, 1854 (*M. minimus* Gibbons).
Abeona Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 322 (*A. trowbridgei* Girard).

3235. *Micrometrus minimus* (Gibbons).

- San Francisco to San Diego.
Cymatogaster minimus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854, 106, San Francisco Bay.
Holoconotus trowbridgei Girard, loc. cit., 152, no locality given.

Genus 1135. AMPHIGONOPTERUS Hubbs.

- Amphigonopterus* Hubbs, Proc. Biol. Soc. Wash., XXXI, May 16, 1918, 13 (*Abeona aurora* Jordan and Gilbert).

3236. *Amphigonopterus aurora* (Jordan and Gilbert).

- Monterey Bay, Calif.
Abeona aurora Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 299, Monterey Bay, Calif.

Genus 1136. CYMATOGASTER Gibbons.

- Cymatogaster* Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer Times and Transcript, May 18, 1854 (*C. aggregatus* Gibbons).
Metrogaster Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 128 (*Cymatogaster aggregatus* Gibbons).
Sema Jordan, Bull. U. S. Geol. Surv., IV, 1878, 399 (*S. signifer* Jordan; larvæ with very high vertical fins and closely compressed body).

3237. *Cymatogaster aggregatus* Gibbons. *Sparada*; *Viviparous perch*; *Perch*

- Pacific coast from Port Wrangel, Alaska, to Todos Santos Bay.
Cymatogaster aggregatus Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer Times and Transcript, May 18, 1854, San Francisco.
Sema signifer Jordan, Bull. Hayden's Geol. Surv., IV, 1878, 399, Rio Grande, Tex.; error for San Francisco.
Metrogaster lineolatus Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 129, no locality.

Genus 1137. BRACHYISTIUS Gill.

- Brachyistius* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 275 (*B. frenatus* Gill)

3238. *Brachyistius frenatus* Gill.

- Vancouver Island to San Diego.
Brachyistius frenatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 275, California coast.
Ditrema brevipinne Günther, Cat., IV, 1862, 248, Esquimaux Harbor, Vancouver Island.

- Genus 1138. **ZALEMBIUS** Jordan and Evermann.
Zalemibus Jordan and Evermann, Fishes North and Mid. Amer., 1896,
 1499 (*Cymatogaster rosaceus* Jordan and Gilbert)
3239. **Zalemibus rosaceus** (Jordan and Gilbert).
 San Francisco, Calif.
Cymatogaster rosaceus Jordan and Gilbert, Proc. U. S. Nat. Mus., III,
 1880, 303, off San Francisco in rather deep water; the other species of
 this family live on sandy shores.
- Genus 1139. **HYPOCRITICHTHYS** Gill.
Hypocritichthys Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 275
 (*Hyperprosopon analis* Alexander Agassiz).
3240. **Hypocritichthys analis** (Agassiz).
 San Francisco to Point Conception
Hyperprosopon analis Alexander Agassiz, Proc. Bost. Soc. Nat. Hist.,
 VIII, 1861, 133, San Francisco.
- Genus 1140. **TOCICHTHYS** Hubbs.
Tocichthys Hubbs, Proc. Biol. Soc. Wash., XXXI, May 16, 1918, 12
 (*Hyperprosopon agassizii* Gill).
3241. **Tocichthys agassizii** (Gill).
 Coast of California, San Francisco to Santa Barbara; most common
 along San Luis Obispo County.
Hyperprosopon agassizii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
 276, California.
- Genus 1141. **HYPERPROSOPON** Gibbons.
Hyperprosopon Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer
 Times and Transcript, May 18, 1854 (*H. argenteum* Gibbons).
Eunichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1855, 322 (*E.*
megalops Girard).
Bramopsis Alexander Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861,
 132 (*B. mento* Agassiz = *Hyperprosopon argenteus* Girard).
3242. **Hyperprosopon argenteus** Gibbons. *Wall-eyed surf-fish, White perch.*
 Coast of California, Cape Disappointment to Todos Santos Bay.
Hyperprosopon argenteum Gibbons, Proc. Acad. Nat. Sci. Phila., VII,
 1854, 105, San Francisco.
Hyperprosopon argenteum punctatum Gibbons, loc. cit., 106, San Fran-
 cisco.
Hyperprosopon arcuatum Gibbons, loc. cit., 125, San Francisco.
Holconotus megalops Girard, loc. cit., Presidio, Humboldt Bay; Astoria.
Bramopsis mento Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 133,
 no locality.
- Genus 1142. **HOLCONOTUS** Agassiz.
Holconotus Agassiz, Amer. Jour. Sci. Arts., XVII, May, 1854, 367 (*H.*
rhodotus Agassiz).
Cymatogaster Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer
 Times and Transcript, June 21, 1854 (*C. pulchellus* Gibbons); not
 of Gibbons, May 18, 1854.
3243. **Holconotus rhodotus** Agassiz.
 Coast of California, San Francisco to San Diego.
Holconotus rhodotus Agassiz, Amer. Jour. Sci. Arts., XVII, May,
 1854, 368, San Francisco.
Cymatogaster pulchellus Gibbons, Proc. Acad. Nat. Sci. Phila., VII,
 July, 1854, 123, San Francisco.
Cymatogaster larkinsi Gibbons, loc. cit., 123, San Francisco.
Cymatogaster ellipticus Gibbons, loc. cit., 124, San Francisco.
- Genus 1143. **AMPHISTICHUS** Agassiz.
Amphistichus Agassiz, Amer. Jour. Sci. Arts., May, 1854, 367 (*A. argen-
 teus* Agassiz).
Mytilophagus Gibbons, Proc. Acad. Nat. Sci. Phila., VI, July, 1854,
 125 (*M. fasciatus* Gibbons = *A. argenteus* Agassiz).

- 3244. *Amphistichus argenteus* Agassiz.**
Pacific coast, from Cape Flattery to San Diego; very abundant on sandy shores
Amphistichus argenteus Agassiz, Amer. Jour. Sci. Arts, XVII, May, 1854, 367, San Francisco
Mytilophagus fasciatus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, July, 1854, 125, San Francisco.
Amphistichus heermanni Girard, loc. cit., 135, Cape Flattery, San Francisco.
Amphistichus similis Girard, loc. cit., 153, San Francisco.
- Genus 1144. EMBIOTOCA Agassiz.**
Embiotoca Agassiz, Amer. Jour. Sci. Arts, XVI, 1853 (November), 386 (*E. jacksoni* Agassiz).
- 3245. *Embiotoca jacksoni* Agassiz. Black surfish.**
Vancouver Island to Todos Santos Bay; abundant and variable.
Embiotoca jacksoni Agassiz, Amer. Jour. Sci. Arts, XVII, 1853, 387, and XVIII, 1854, 366, San Francisco.
Holconotus fuliginosus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854, 123, San Francisco.
Embiotoca cassidy Girard, loc. cit., 151, San Diego.
?Embiotoca webber Girard, loc. cit., VIII, 1855, 320, San Diego.
- Genus 1145. TÆNIOTOCA Alexander Agassiz.**
Holconotus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854 (July), 122 (*H. agassizii* Gibbons = *Embiotoca lateralis* Agassiz); not *Holconotus* Agassiz.
Tæniotoca Alexander Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 133 (*Embiotoca lateralis* Agassiz).
- 3246. *Tæniotoca lateralis* (Agassiz). Striped surfish.**
Vancouver Island to Todos Santos Bay; abundant southward.
Embiotoca lateralis Agassiz, Amer. Jour. Sci. Arts., XVIII, 1854 (May), 356, San Francisco.
Holconotus agassizii Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854 (July), 122, San Francisco.
Embiotoca lineata Girard, loc. cit., (Aug.), 134, San Francisco, Tomales Bay, Presidio.
Embiotoca ornata Girard, loc. cit., VII, 1855, 321, San Diego, Calif.
Embiotoca perspicabilis Girard, loc. cit., 321, Puget Sound.
- Genus 1146. PHANERODON Girard.**
Phanerodon Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 153 (*P. furcatus* Girard).
- 3247. *Phanerodon furcatus* Girard. White surfish.**
Pacific coast from Vancouver Island to Todos Santos Bay; abundant southward.
Phanerodon furcatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 153, Presidio, San Francisco, Calif.
- 3248. *Phanerodon atripes* (Jordan and Gilbert).**
Monterey to San Diego; scarce.
Ditrema atripes Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 320, Monterey Bay, Calif.
Ditrema orthonotus Eigenmann and Eigenmann, West American Scientist, October, 1889, 127, Cortez Banks, off San Diego.
- Genus 1147. RHACCOCHILUS Agassiz.**
Rhacochilus Agassiz, Amer. Jour. Sci. Arts, XVIII, 1854 (May), 367 (*R. toxotes* Agassiz).
Pachylabrus Gibbons, Proc. Calif. Acad. Sci., in Daily Placer Times and Transcript, June 21, 1854 (*P. variegatus* Gibbons = *R. toxotes* Agassiz).
- 3249. *Rhacochilus toxotes* Agassiz. Bigmouth surfish; Alfonsa.**
Rhacochilus toxotes Agassiz, Amer. Jour. Sci. Arts, XVIII, 1854 (May), 367, San Francisco.
Pachylabrus variegatus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854 (July), 126, San Francisco.

Genus 1148. **HYP SURUS** Alexander Agassiz.

Hypsurus Alexander Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 133 (*Embiotoca caryi* Agassiz).

3250. **Hypsurus caryi** (Agassiz). *Bugara*.

Cape Mendocino to San Diego; common northward.

Embiotoca caryi Agassiz, Amer. Jour. Sci. Arts, XVI, 1853, 389, and XVII, 1854, 366, San Francisco.

Holconotus gibbonsi (Gibbons MS., Proc. Calif. Acad. Sci., 1854), Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854, 12, San Francisco.

Genus 1149. **DAMALICHTHYS** Girard.

Damalichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1855, 321 (*D. vacca* Girard = *D. argyrosoma* Girard).

3251. **Damalichthys argyrosoma** Girard.

Vancouver Island to Tochos Santos Bay; very abundant northward.

Damalichthys argyrosoma Girard, Proc. Acad. Nat. Sci. Phila., VII, 1855, 136, San Francisco; young individual in bad condition.

Damalichthys vacca Girard, loc. cit., 321, Puget Sound.

Genus 1150. **HYSTEROCARPUS** Gibbons.

Hysteroecarpus Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer Times and Transcript, May 18, 1854, and in Proc. Acad. Nat. Sci. Phila., VII, 1854, 124 (*H. traski* Gibbons).

Sargosomus Agassiz, in Alexander Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 128 (*S. fluviatilis* Agassiz = *Hysteroecarpus traski* Gibbons).

Dacentrus Jordan, Bull. U. S. Geol. Surv., IV, 1878, 667 (*D. lucens* Jordan; very young).

3252. **Hysteroecarpus traski** Gibbons.

Rivers of central California, chiefly in the Sacramento-San Joaquin Valley from Lake County to Santa Clara County.

Hysteroecarpus traski Gibbons, Proc. Acad. Nat. Sci., Phila., VII, 1854, 105, lower Sacramento River.

Sargosomus fluviatilis Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 130, Sacramento.

Dacentrus lucens Jordan, Bull. U. S. Geol. Surv., IV, 1878, 667, "Rio Grande, Tex." locality erroneous, should be San Francisco.

Order TT. **CHROMIDES**Family 249. **POMACENTRIDÆ**Genus 1151. **AZURINA** Jordan and McGregor.

Azurina Jordan and McGregor, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1544 (*A. hirundo* Jordan and McGregor).

3253. **Azurina hirundo** Jordan and McGregor.

Pacific coast of Mexico.

Azurina hirundo Jordan and McGregor, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1544, Guadalupe Island, off Lower California.

Genus 1152. **FURCARIA** Poey.

Furcaria Poey, Memorias, II, 1860, 194 (*F. puncta* Poey = *Heliasas multilineatus* Guichenot in Ramon de la Sagra, Hist. fisica Cuba, IV).

3254. **Furcaria atrilobata** (Gill).

Pacific coast of America, from Cape San Lucas southward.

Chromis (*Furcaria*) *atrilobata* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 149, Cape San Lucas, Lower California.

3255. **Furcaria cyanea** Poey.

Cuba.

Furcaria cyanea Poey, Memorias, II, 1860, 196, Cuba.

3256. **Furcaria multilineata** (Guichenot).

Cuba.

Heliasas multilineatus Guichenot, in Ramon de la Sagra, Poiss. Cuba, 1853, 76, pl. 2, fig. 2, Habana.

Furcaria puncta Poey, Memorias, II, 1860, 195, Cuba.

3257. *Furcaria bermudæ* (Nichols).
Bermuda.
Chromis bermudæ Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 60, Bermuda.
- Genus 1153. **HELIASES** Cuvier and Valenciennes.
Heliases Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 369 (495)
(*H. insolatus* Cuvier and Valenciennes).
Heliastes Gunther, revised spelling.
3258. *Heliases insolatus* Cuvier and Valenciennes. *Chauffe soleil*.
West Indies, north to Pensacola.
Heliases insolatus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 370 (494), pl. 137, Martinique.
3259. *Heliases enchrysurus* (Jordan and Gilbert).
Snapper Banks off Pensacola and Tampa, Fla.
Chromis enchrysurus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 286, Pensacola, Fla.
3260. *Heliases marginatus* Castelnau.
Brazil; St. Croix Island, West Indies.
Heliases marginatus Castelnau, Anim. Amer. Sud, Poiss., 1855, 9, Pl. III, fig. 1, Bahia.
- Genus 1154. **AYRESIA**⁷⁹ Cooper.
Ayresia Cooper, Proc. Calif. Acad. Nat. Sci., 1st ser., III, 1863, 73 (*A. punctipinnis* Cooper).
3261. *Ayresia punctipinnis* Cooper. *Blacksmith*.
Coast of California, from Point Conception to Cedros Island.
Ayresia punctipinnis Cooper, Proc. Calif. Acad. Nat. Sci., 1st ser., III, 1863, 73, San Diego Bay, San Pedro, and Santa Barbara.
- Genus 1155. **EUPOMACENTRUS**⁸⁰ Bleeker.
Eupomacentrus Bleeker, Nat. Verh. Holl. Maats., Wetensk., II, 1877, 73 (*Pomacentrus lividus* Bleeker); teeth in one row, truncate; snout scaly; lower jaw naked; dorsal not notched.
Amblypomacentrus Bleeker, loc. cit., 68 (*Pomacentrus breviceps*); similar, snout naked.
3262. *Eupomacentrus fuscus* (Cuvier and Valenciennes). *Maria molle*.
West Indies and coast of Brazil north to Key West.
Pomacentrus fuscus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 324 (432), Brazil
Pomacentrus variabilis Castelnau, Anim. Nouv. ou Rares, Poiss., 1855, 9, pl. 3, fig. 3, Bahia.
Pomacentrus atro-cyanus Poey, Memorias, II, 1861, 190, Cuba.
3263. *Eupomacentrus diencæus* Jordan and Rutter.
Jamaica.
Eupomacentrus diencæus Jordan and Rutter, Proc. Acad. Nat. Sci. Phila., XLIX, 1897, 116, Jamaica.
3264. *Eupomacentrus leucostictus* (Müller and Troschel). *Cockeye pilot*; *Beau Gregoire*; *Black pilot*.
West Indies to Snapper Banks, west Florida; accidental north to Woods Hole, Mass.
Pomacentrus leucostictus Müller and Troschel, in Schomburgk's Hist. Barbados, 1848, 674, Barbados.
Pomacentrus xanthurus Poey, Memorias, II, 1861, 190, Cuba.
Pomacentrus caudalis Poey, Synopsis, 1867, 328, Cuba.
Pomacentrus dorso-punicans Poey, loc. cit., 328, Cuba.
3265. *Eupomacentrus analis* (Poey).
Cuba, Key West, and Porto Rico.
Pomacentrus analis Poey, Synopsis, 1867, 327, Cuba.

⁷⁹ This genus is near the Mediterranean genus *Chromis* Cuvier, based on *Sparus chromis* Linnaeus, a species with 14 dorsal spines, the head high and narrow, the tail short, the caudal not deeply forked. *Furcaria* and *Heliases* are also closely related and usually the four groups have been united into one as *Chromis* or "*Heliastes*."

⁸⁰ In the closely related old world genus *Pomacentrus* the teeth are in two rows. Several of the species here accepted are of very doubtful validity, and the genera of Bleeker derived from the subdivisions of *Pomacentrus* need further study.

3266. *Eupomacentrus otophorus* (Poey).
Cuba.
Pomacentrus otophorus Poey, Memorias, II, 1861, 188, Cuba.
3267. *Eupomacentrus chrysus* Bean.
Bermudas
Eupomacentrus chrysus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 32,
Whites Flat Channel, Bermudas
3268. *Eupomacentrus flaviventer* (Troschel).
Atlantic coast of Mexico.
Pomacentrus flaviventer Troschel, in J. W. von Müller's Reisen in den
Vereinigten Staaten, Canada, und Mexico, III, 1865, 633, Atlantic
Ocean
3269. *Eupomacentrus gillii* (Gilbert and Starks).
Panama.
Pomacentrus gillii Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904,
141, pl. XXII, fig. 44, Panama.
3270. *Eupomacentrus partitus* (Poey).
Cuba.
Pomacentrus partitus Poey, Synopsis, 1867, 327, Cuba.
3271. *Eupomacentrus planifrons* (Cuvier and Valenciennes). *Petite jaquette*.
Tortugas; West Indies, Jamaica to Martinique.
Pomacentrus planifrons Cuvier and Valenciennes, Hist. Nat. Poiss., V,
1830, 323 (431), Martinique.
3272. *Eupomacentrus leucurus* (Gilbert).
Socorro Island, one of the Revillagigedo group, off Pacific coast of
Mexico.
Pomacentrus leucurus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 554,
Socorro Island.
3273. *Eupomacentrus adustus* (Troschel).
Cuba.
Pomacentrus adustus Troschel, in J. W. von Müller's Reisen in den
Vereinigten, Staaten, Canada und Mexico, III, 1865, 633, Mexico.
Pomacentrus obscuratus Poey, Enumeratio, 1876, 101, Habana.
3274. *Eupomacentrus rectifrenum* (Gill). *Pescado azul de dos colores*.
Pacific coast of America from Cape San Lucas to Panama; common.
Pomacentrus rectifrenum Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
148, Cape San Lucas, Lower California.
Pomacentrus analigutta Gunther, Cat., IV, 1862, 27, Cape San Lucas.
3275. *Eupomacentrus flavilatus* (Gill). *Pescado azul de dos colores*.
Pacific coast of America, from Cape San Lucas to Mazatlan and beyond.
Pomacentrus flavilatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 148,
Cape San Lucas, Lower California.
3276. *Eupomacentrus nepenthe* Nichols.
Bahamas.
Eupomacentrus nepenthe Nichols, Amer. Mus. Novitates, XXVI, 1921,
1, fig. 1, Bahamas.
- Genus 1156. *NEXILARIUS* Gilbert.
Nexilarius Gilbert, in Jordan and Evermann, Check List, 1895, 512
(*Euschistodus concolor* Gill); misprinted *Nexilaris*.
3277. *Nexilarius concolor* (Gill).
Panama.
Euschistodus concolor Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
145, Panama.
- Genus 1157. *ABUDEFDUF*⁵¹ Forskål. *Pintanos*.
Abudefduf Forskål, Descr. Anim., etc., 1775, 59 (*Chætodon sordidus*
Forskål).
Glyphisodon Lacépède, Hist. Nat. Poiss., IV, 1803, 542 (*Chætodon saxatilis*
Linnæus).
Euschistodus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 145 (*E.*
declivifrons Gill).

⁵¹ The small brilliantly colored species of coral reefs in the South Seas constituting the genus *Chrysiptera* Swainson are probably separable from *Abudefduf*.

- 3278. *Abudefduf marginatus* (Bloch).** *Pintano; Cow pilot; Jaqueta; Mojarra*
ravado; Demoiselle
 Atlantic coast of tropical America, West Indies, to Brazil; once reported
 from Newport, R. I.
Jaguacanguare Marcgrave, Hist. Brazil, 1648, 156, Brazil
Chætodon marginatus Bloch, Ichthyol., III, 1787, 98, Martinique;
 after Plumier.
Chætodon sargoides Lacépède, Hist. Nat. Poiss., IV, 1803, 452, Mar-
 tinique.
Glyphisodon moucharra Lacépède, loc. cit., 542, Brazil.
Glyphidodon saxatilis of authors, not of Linnæus, an Asiatic species.
 ?*Chætodon mauritii* Bloch, Naturgesch. Ausl. Fische, III, 1787, 109,
 Brazil, on a drawing by Prince Maurice. A doubtful species,
 apparently the same as *A. marginatus*.
- 3279. *Abudefduf troschelii* (Gill).**
 Pacific coast of Mexico; Cape San Lucas; Socorro Island; Cocos
 Islands; Galapagos.
Glyphidodon troschelii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
 150, Cape San Lucas.
- 3280. *Abudefduf declivifrons* (Gill).**
 Pacific coast of tropical America, from Cape San Lucas southward.
Euschistodus declivifrons Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
 145, 146, Cape San Lucas.
- 3281. *Abudefduf analogus* (Gill).**
 Atlantic coast of Central America.
Euschistodus analogus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 220,
 Aspinwall.
- 3282. *Abudefduf taurus* (Müller and Troschel).** *Dovetail fish.*
Tortugas; Barbados.
Glyphidodon taurus Müller and Troschel, in Schomburgk, Hist. Bar-
 bados, 1848, 674, Barbados.
- Genus 1158. **CENTROCHROMIS** Norman.
Centrochromis Norman, Ann. Mag. Nat. Hist., ser. 9, IX, 1922, 534
 (*Glyphidodon rudis* Poey.)
- 3283. *Centrochromis rudis* (Poey).**
 Cuba.
Glyphidodon rudis Poey, Memorias, II, 1861, 191, Cuba.
- Genus 1159. **HYPSTYPOPS** Gill. *Garibaldi.*
Hypsypops Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 165 (*Gly-
 phisodon rubicundus* Girard).
- 3284. *Hypsypops rubicundus* (Girard).** *Garibaldi; Sea goldfish; a magnificent
 and hardy aquarium fish.*
 Coast of California south of Monterey; abundant about Santa Catalina
 Island.
Glyphisodon rubicundus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854,
 148, Monterey, Calif.
- Genus 1160. **STEGASTES** Jenyns.
Stegastes Jenyns, Voy. Beagle, Fishes, 1842, 63 (*S. imbricatus* Jenyns).
Microspathodon Gunther, Cat., IV, 1862, 35 (*Glyphidodon chrysurus*
 Cuvier and Valenciennes).
Pomataprion Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 214, 216
 (*Hypsypops dorsalis* Gill).
- 3285. *Stegastes chrysurus* (Cuvier and Valenciennes).**
Tortugas; West Indies; Cuba and St. Thomas.
Glyphisodon chrysurus Cuvier and Valenciennes, Hist. Nat. Poiss., V,
 1830, 356 (476), St. Thomas.
Pomacentrus denegatus Poey, Memorias, II, 1861, 190, Cuba.
- 3286. *Stegastes bairdii* (Gill).**
 Pacific coast of tropical America, from Cape San Lucas to Panama.
Pomacentrus bairdii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 149,
 Cape San Lucas.

3287. *Stegastes niveatus* (Poey).
Cuba.
Pomacentrus niveatus Poey, Enumeratio, 1876, 102, Habana.
3288. *Stegastes dorsalis* (Gill).
Pacific coast of America, from Cape San Lucas to Mazatlan.
Hyssypops dorsalis Gill, Proc. Acad. Nat. Sci. Phila. XIV, 1862, 147, Cape San Lucas; young
Pomacentrus quadrigutta Gill, loc. cit., 149, Cape San Lucas
Microspathodon cinereus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 71, Clarion and Socorro Islands
Microspathodon czurissimus Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 478, pl. 44, Venados Islands, near Mazatlan; adult.

Family 250. CICHLIDÆ.⁸² Mojarra de rio

Genus 1161. CICHLAURUS Swainson.

Cichlaurus Swainson, Nat. Hist. Classn. Fishes, etc., II, 1839, 173, definition, but no type named; this is given on p. 230, where, without assigned reason, the name is changed to *Cichlasoma*.

Cichlasoma Swainson, Nat. Hist., etc., loc. cit., 230 (*Labrus punctatus* Bloch).

Heros Heckel, Natterers Reise, II, 1840, 362 (*Heros severus* Heckel).

3289. *Cichlaurus bifasciatus* (Steindachner).
Mexico.
Heros bifasciatus Steindachner, Denks. Ak. Wiss. Wien, XXIII, 1864, 60, pl. II, Mexico.
3290. *Cichlaurus angulifer* (Günther).
East slope of Guatemala.
Heros angulifer Günther, Cat., IV, 1862, 298, River de Santa Isabel, Guatemala.
3291. *Cichlaurus microphthalmus* (Günther).
Rio Motagua, Guatemala
Heros microphthalmus Günther, Cat., IV, 1862, 295, Rio Motagua, Guatemala.
3292. *Cichlaurus guttulatus* (Günther).
Pacific drainage of Guatemala.
Heros guttulatus Günther, Proc. Zool. Soc. Lond., 1864, 152, Lake Amatitlan, Guatemala.
3293. *Cichlaurus meeki* (Hildebrand). *Mojarra; Mojarra negra; Mojarra plateada*.
Lakes and rivers of El Salvador, Pacific slope of Central America.
Cichlasoma meeki Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925, 275, fig. 20, Lake Guiga, El Salvador.
3294. *Cichlaurus melanurus* (Günther).
Lake Peten, Guatemala.
Heros melanurus Günther, Cat., IV, 1862, 288, Lake Peten, Guatemala.
3295. *Cichlaurus fenestratum* (Günther).
Atlantic slope of southern Mexico.
Chromis fenestrata Günther, Proc. Zool. Soc. Lond., 1860, 318, Mexico.
3296. *Cichlaurus godmanni* (Günther).
Atlantic slope of Guatemala.
Heros godmanni Günther, Cat., IV, 1862, 296, Rio Cahabon, Guatemala.
3297. *Cichlaurus intermedius* (Günther).
Lake Peten, Guatemala
Heros intermedius Günther, Cat., IV, 1862, 298, Lake Peten, Guatemala.
3298. *Cichlaurus deppii* (Heckel).
Mexico.
Heros deppii Heckel, Ann. Wiener Mus., II, 1840, 382, Mexico.

⁸² The many species of this genus in the Mexican region are chiefly the overflow from the vast Brazilian fauna. The larger genera as here recognized are very closely related and were recently regarded as subgenera of *Cichlasoma* (*Cichlaurus*) = *Heros* Heckel; several of the species are only provisionally located and a few (*godmanni*, *hedricki*, *lentiginosus*) may represent new groups not yet defined.

3299. *Cichlaurus calobrensis* (Meek and Hildebrand).
Streams of Panama
Cichlasoma calobrense Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 90, Rio Calobra, Panama.
3300. *Cichlaurus umbrifer* (Meek and Hildebrand).
Rivers of Panama
Cichlasoma umbriferum Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 88, Rio Cupe, Cituro, Panama.
- Genus 1162. *THERAPS* Günther.
Theraps Günther, Cat., IV, 1862, 284 (*T. irregularis* Günther).
?*Barodon* Agassiz, Proc. Bost. Soc. Nat. Hist., VI, 1858, 401 (*B. fasciatus* Agassiz); scanty description; said to be slender as in the Sparoid genus *Boops* or *Box*.
3301. *Theraps irregularis* Günther.
Atlantic slope of Guatemala and southern Mexico.
Theraps irregularis Günther, Cat., IV, 1862, 284, Guatemala.
3302. *Theraps mojarra* (Meek).
Pacific slope of Isthmus of Tehuantepec.
Cichlasoma mojarra Meek, Field Mus., Zool., V, 1904, 93, 217, San Geronimo, Oaxaca, Mexico.
3303. *Theraps nebulifer* (Günther).
Atlantic lowland streams of the Isthmus of Tehuantepec.
Heros nebulifer Günther, Proc. Zool. Soc. Lond., 1860, 318, Mexico.
3304. *Theraps gibbiceps* (Steindachner).
Rivers of Tabasco, Mexico.
Heros gibbiceps Steindachner, Denks. Ak. Wiss. Wien, XXIII, 1864, 66, Pl. V, figs. 1 and 2, Rio Teapa, Tabasco, Mexico.
3305. *Theraps zonatus* (Meek).
Pacific slope of Isthmus of Tehuantepec, Mexico.
Cichlasoma zonatum Meek, Proc. Biol. Soc. Wash., XVIII, 1905, 245, Niltpec, Oaxaca, Mexico.
Barodon fasciatus Agassiz, Proc. Bost. Soc. Nat. Hist., VI, 1858, 408, Lake Nicaragua.
3306. *Theraps eigenmanni* (Meek).
Rio Tehuacan, Mexico
Cichlasoma eigenmanni Meek, Field Col. Mus., Zool., III, 1902, 119, Rio Tehuacan, Mexico.
3307. *Theraps teapæ* (Evermann and Goldsborough).
Rio Teapa, Mexico.
Cichlasoma teapæ Evermann and Goldsborough, Bull. U. S. Fish Com., XXI, 1901, 156, fig. 6, Rio Teapa, Mexico.
3308. *Theraps frontalis* (Meek).
Turrubales, Costa Rica.
Cichlasoma frontale Meek, Field Col. Mus., Zool., VIII, 1909, 210, Turrubales, Costa Rica.
3309. *Theraps punctatum* (Meek).
Buenos Aires de Terraba, Costa Rica.
Cichlasoma punctatum Meek, Field Mus. Nat. Hist., Zool., VII, 1909, 210, Buenos Aires de Terraba, Costa Rica; name preoccupied in *Cichlasoma* but not in *Theraps*.
Theraps terrabæ Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 507, substitute for *Cichlasoma punctatum* Meek, preoccupied.
3310. *Theraps gadovii* (Regan).
Atlantic slope of southern Mexico.
Cichlosoma gadovii Regan, Ann. Mag. Nat. Hist., 1905, 232, Motozongongo, Mexico.
3311. *Theraps milleri* (Meek).
El Rancho, Guatemala.
Cichlasoma milleri Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 142, El Rancho, Guatemala.

3312. *Theraps balteatus* (Gill and Bransford).
Great Lakes of Nicaragua.
Heros balteatus Gill and Bransford, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 184, Lake Nicaragua.
3313. *Theraps nicaraguensis* (Gunther).
Great Lakes of Nicaragua.
Heros nicaraguensis Gunther, Proc. Zool. Soc. Lond., 1864, 153, Lake Nicaragua.
3314. *Theraps nigrinus* (Meek).
Lake Nicaragua.
Cichlasoma nigrinum Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 128, Lake Nicaragua.
3315. *Theraps lentiginosus* (Steindachner).
Rivers of Mexico
Heros lentiginosus Steindachner, Denks. Akad. Wiss. Wien, XXIII, 1864, 62, Pl. III, fig. 1, Mexico
- Genus 1163. **PARAPETENIA** Regan.
Hypsophrys Agassiz, Proc., Bost. Soc. Nat. Hist., VI, 1858, 401 (*H. unimaculatus* Agassiz); no description except that it resembles *Chrysophrys* (*Sparus*) and has a large round black spot on the shoulder; lakes of Nicaragua.
? *Parachromis* Agassiz, Proc., Bost. Soc. Nat. Hist., VI, 1858, 401 (*P. gulosus* Agassiz); not described but said to resemble *Dentex*; lakes of Nicaragua.
Parapetenia Regan, Ann. Mag. Nat. Hist., XVI, 7th ser., 1905, 324 (*Acara adpersum* Gunther).
3316. *Parapetenia adpersa* (Gunther).
Barbados.
Acara adpersum Günther, Cat., IV, 1862, 282, Barbados.
3317. *Parapetenia dovii* (Gunther).
Great Lakes of Nicaragua.
Heros dovii Gunther, Proc. Zool. Soc. Lond., 1864, 154, Lake Nicaragua.
3318. *Parapetenia managuense* (Gunther).
Great Lakes of Nicaragua.
Heros managuensis Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 392, 463, pl. 77, fig. 3, Lake Managua, Nicaragua.
3319. *Parapetenia motaguensis* (Günther).
Atlantic slope of Guatemala.
Heros motaguensis Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 391, 462, pl. 77, fig. 2, Lake Motagua, Guatemala.
3320. *Parapetenia friedrichstahli* (Heckel).
Lake Peten, Guatemala.
Heros friedrichstahl Heckel, Ann. Mus. Wien, II, 1840, 381, Lake Peten, Guatemala.
3321. *Parapetenia multifasciata* (Regan).
Guatemala; British Honduras.
Cichlasoma multifasciatum Regan, Ann. Mag. Nat. Hist., 7th ser., XVI, 1905, 335, Lake Peten, Guatemala.
3322. *Parapetenia salvini* (Günther).
Atlantic slope of southern Mexico to Guatemala.
Heros salvini Gunther, Cat., IV, 1862, 294, Rio de Santa Isabel, Guatemala.
Heros triagramma Steindachner, Denkschr. Akad. Wiss. Wien, XXIII, 1864, 70, Taf. 3, fig. 2, central Mexico.
3323. *Parapetenia urophthalma* (Gunther).
Atlantic slope of southern Mexico to Guatemala; British Honduras and Yucatan.
Heros urophthalmus Günther, Cat., IV, 1862, 291, Lake Peten.
3324. *Parapetenia istlana* (Jordan and Snyder).
Rio Balsas, Mexico.
Heros istlanus Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 144, fig. 21, Rio Ixtla at Puente de Ixtla, Mexico.

3325. *Parapetenia tetracantha* (Cuvier and Valenciennes). *Viajaca*.
Rivers of Cuba.
Centrarchus tetracanthus Cuvier and Valenciennes, Hist. Nat. Poiss.,
VII, 1831, 346 (460), Habana.
Chromis fusco-maculatus Guichenot, in Ramon de la Sagra, Hist. Ile
Cuba, Poiss., 1853, 78, pl. 2, fig. 3, Habana.
3326. *Parapetenia tetracantha torralbasi* (Eigenmann).
Cuba; Calabazar, Almendares River; San Cristobal; Los Palacios.
Heros tetracanthus torralbasi Eigenmann, Bull. U. S. Fish Com., XXII,
1902, 230, fig. 11, Calabazar Cuba
3327. *Parapetenia tetracantha grisea* (Eigenmann).
San Antonio, Cuba.
Heros tetracanthus griseus Eigenmann, Bull. U. S. Fish Com., XXII,
1902, 233, San Antonio, Cuba.
3328. *Parapetenia tetracantha lata* (Eigenmann).
San Juan, Cuba.
Heros tetracanthus latus Eigenmann, Bull. U. S. Fish Com., XXII,
1902, 234, San Juan, Cuba.
3329. *Parapetenia tetracantha cincta* (Eigenmann).
Paso Real, Cuba.
Heros tetracanthus cinctus Eigenmann, Bull. U. S. Fish Com., XXII,
1902, 234, fig. 16, Paso Real, Cuba.
3330. *Parapetenia parma* (Gunther).
Mexico and Guatemala.
Heros parma Gunther, Cat., IV, 1862, 285, Mexico and Guatemala
3331. *Parapetenia nigricans* (Eigenmann).
Pinar del Rio, Cuba.
Heros nigricans Eigenmann, Bull. U. S. Fish Com., XXII, 1902, 235,
fig. 17, Pinar del Rio, Cuba.
3332. *Parapetenia steindachneri* (Jordan and Snyder).
Lower portion of the basin of the Rio Panuco and Rio Verde, Mexico.
Cichlasoma steindachneri Jordan and Snyder, Bull. U. S. Fish Com.,
XIX, 1899, 143, fig. 20, Rio Verde, Mexico.
3333. *Parapetenia bartoni* (Bean).
Upland portion of the Rio Verde in San Luis Potosi, Mexico.
Acara bartoni Bean, Proc. U. S. Nat. Mus., XV, 1892, 286, Pl. XLIV,
fig. 3, Huasteca Potosina, San Luis Potosi, Mexico.
3334. *Parapetenia labridens* (Pellegrin).
Upland portion of the Rio Verde in San Luis Potosi, Mexico.
Heros labridens Pellegrin, Bull. Mus. Hist. Nat. Paris, 1903, 122,
Huasteca, Potosi, Mexico.
3335. *Parapetenia beani* (Jordan). *Mojarra verde*.
Lowland streams of Sinaloa and Jalisco.
Heros beani Jordan, Proc. U. S. Nat. Mus., XI, 1888, 332, Rio Presidio,
near Mazatlan, Mexico.
3336. *Parapetenia tenuis* (Meek).
Atlantic slope of the isthmus of Tehuantepec; Achatol; Vera Cruz,
Mexico.
Cichlasoma tenue Meek, Field Mus. Nat. Hist., Zool., VII, 1906, 94,
Achatol, Vera Cruz, Mexico.
3337. *Parapetenia centralis* (Meek).
Pacific slope of Guatemala; Caballo Blanco.
Cichlasoma centrale Meek, Field Mus. Nat. Hist., Zool., VII, 1906, 94,
Caballo Blanco, Guatemala.
3338. *Parapetenia alfari* (Meek).
Atlantic slope of Costa Rica; Turrialba.
Cichlasoma alfari Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 148,
Turrialba, Costa Rica.
3339. *Parapetenia mento* (Vaillant and Pellegrin).
Southern Mexico.
Heros mento Vaillant and Pellegrin, Bull. Mus. Hist. Nat. Paris, 1902,
88, Rio Negro, southern Mexico.

3340. *Parapetenia trimaculata* (Günther).
Pacific coast streams of southern Mexico and Guatemala.
Heros trimaculatum Günther, Trans. Zool. Soc. Lond., VI, 1868, 461,
pl. 76 (Nov. 27), Chiapam and Huamuchal, Mexico.
- Genus 1164. *ARCHOCENTRUS* Gill.
Archocentrus Gill, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 186
(*Heros centrarchus* Gill and Bransford).
3341. *Archocentrus centrarchus* (Gill and Bransford).
Great Lakes of Nicaragua.
Heros centrarchus Gill and Bransford, Proc. Acad. Nat. Sci. Phila.,
XXIX, 1877, 185, Lake Nicaragua.
3342. *Archocentrus nigrofasciatus* (Günther).
Lakes Amatitlan and Atitlan, Guatemala.
Heros nigrofasciatus Günther, Trans. Zool. Soc. Lond., VI, 1868, 391,
452, pl. 74, fig. 3, Lakes Amatitlan and Atitlan.
3343. *Archocentrus spilurus* (Günther).
Rio Motagua and Lake Isabel, Guatemala.
Heros spilurus Günther, Cat., IV, 1862, 289, Lake Yzabal, Guatemala.
3344. *Archocentrus spinosissimus* (Vaillant and Pellegrin).
Rio Palochic, Guatemala.
Heros (Cichlasoma) spinosissimus Vaillant and Pellegrin, Bull. Mus.
Hist. Nat. Paris, 1902, 87, Rio Palochic, Guatemala.
3345. *Archocentrus immaculatus* (Pellegrin).
Rio Palochic, Guatemala.
Cichlasoma spinosissimum immaculatum Pellegrin, Mém. Soc. Zool.
France, XVI, 1903, 225, Rio Palochic, Guatemala.
- Genus 1165. *ASTATHEROS*⁸³ Pellegrin.
Astatheros Pellegrin, Mém. Soc. Zool. France, 1833, 205 (*Heros hetero-*
dontus Vaillant and Pellegrin).
Erythrictithys Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 118, 121
(*Heros citrinellus* Günther); name twice preoccupied as *Erythrictithys*.
3346. *Astatheros heterodontus* (Vaillant and Pellegrin).
Isthmus of Tehuantepec, Mexico.
Heros heterodontus Vaillant and Pellegrin, Bull. Mus. Nat. Hist., Paris,
1902, 86, Isthmus of Tehuantepec.
3347. *Astatheros evermanni* (Meek).
Pacific slope of Isthmus of Tehuantepec.
Cichlasoma evermanni Meek, Field Col. Mus. Zool., V, 1904, 214, fig.
70, Rio Tehuantepec, Tehuantepec, Mexico.
3348. *Astatheros robertsoni* (Regan).
British Honduras.
Cichlasoma robertsoni Regan, Ann. Mag. Nat. Hist., ser. 7, XVI,
1905, 239, Stann Creek, British Honduras.
3349. *Astatheros lethrinus* (Regan).
Rio Iroquois, Costa Rica.
Cichlasoma lethrinus Regan, Ann. Mag. Nat. Hist., ser. 8, II, 1908, 462,
Rio Iroquois, Costa Rica.
3350. *Astatheros dorsatus* (Meek).
Lake Managua, Nicaragua.
Cichlasoma dorsatum Meek, Field Col. Mus. Nat. Hist., Zool., VII,
1907, 123, Lake Managua.
3351. *Astatheros longimanus* (Günther).
Lake Nicaragua.
Heros longimanus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27),
391, 453, pl. 72, fig. 2, Lake Nicaragua.
3352. *Astatheros altifrons* (Kner and Steindachner).
Panama and Colombia.
Heros altifrons Kner and Steindachner, Sitz. Bayer. Akad., 1863, 223,
Colombia.

⁸³ This group is usually united with the two large-lipped species *labiatus* and *lobochilus*, to which the name *Amphilophus* was given earlier. *Citrinellus* and *Erythræus* seem intermediate.

3353. *Astatheros lauræ* (Regan).
Tampico, Mexico.
Cichlasoma lauræ Regan, Ann. Mag. Nat. Hist., 8th ser., 2, 1908, 223,
Tampico, Mexico.
3354. *Astatheros margaritifer* (Günther).
Lake Peten, Guatemala.
Heros margaritifer Günther, Cat., IV, 1862, 287, Lake Peten.
Heros melanopogon Steindachner, Denks. Akad. Wiss. Wien, XXIII,
1864, 72, pl. 1, fig. 3, Central America.
3355. *Astatheros sexfasciatus* (Regan).
Guapote, Mexico.
Cichlasoma sexfasciatum Regan, Ann. Mag. Nat. Hist., (7), XVI, 1905,
230, Guapote, Mexico.
3356. *Astatheros septemfasciatus* (Regan).
Costa Rica.
Cichlasoma septemfasciatum Regan, Ann. Mag. Nat. Hist., II, ser. 8,
1908, 461, Rio Iroquois, Costa Rica.
3357. *Astatheros basilaris* (Gill and Bransford).
Lake Nicaragua.
Heros basilaris Gill and Bransford, Proc. Acad. Nat. Sci. Phila., XXIX,
1877, 182, Lake Nicaragua.
3358. *Astatheros rectangularis* (Steindachner).
Mexico.
Acara rectangulare Steindachner, Chromiden Mejicos, 1864, pl. 1, fig. 1,
1, Mexico.
3359. *Astatheros maculicauda* (Regan).
Atlantic slope of Guatemala and Panama.
Cichlasoma maculicauda Regan, Ann. Mag. Nat. Hist., XVI, ser. 7,
1905, 227, Lake Yzabal, Rio Motagua, and Rio Chagres.
3360. *Astatheros tuyrensis* (Meek and Hildebrand).
Rivers of Panama.
Cichlasoma tuyrensis Meek and Hildebrand, Field Mus. Nat. Hist.,
Zool., X, 1913, 89, Rio Tuyra, Boca de Cupa, Panama.
3361. *Astatheros oblongus* (Günther).
Rio Motagua; Guatemala.
Heros oblongus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27),
392, 464, Rio Motagua, Guatemala.
3362. *Astatheros montezuma* (Heckel).
Mexico.
Heros montezuma Heckel, Ann. Wiener Mus., II, 1840, 382, Mexico.
3363. *Astatheros globosum* (Miller).
Puerto Barrios, Guatemala.
Cichlasoma globosum Miller, Bull. Amer. Mus. Nat. Hist., XXIII,
1907, 114, fig. 4, Puerto Barrios, Guatemala.
3364. *Astatheros rostratum* (Gill and Bransford).
Great Lakes of Nicaragua.
Heros rostratus Gill and Bransford, Proc. Acad. Nat. Sci. Phila., XXIX,
1877, 181, Lake Nicaragua.
3365. *Astatheros hedricki* (Meek).
Large Atlantic slope rivers of Mexico, south of Vera Cruz.
Cichlasoma hedricki Meek, Field Col. Mus., Zool., V, 1904, 208, fig. 66,
Obispo, Vera Cruz, Mexico.
3366. *Astatheros octofasciatus* (Regan).
Atlantic slope, Isthmus of Tehuantepec to British Honduras.
Heros octofasciatus Regan, Ann. Mus. Hist. Nat. Geneva, Rev. Suisse,
Zool., 1903, 417, pl. 13, fig. 1, Mexico.
3367. *Astatheros granadense* (Meek).
Lake Nicaragua.
Cichlasoma granadense Meek, Field Mus. Nat. Hist., Zool., VII, 1907,
121, Lake Nicaragua, Nicaragua, Granada.

3368. *Astatheros citrinellus* (Günther).
Great Lakes of Nicaragua.
Heros citrinellus Gunther, Proc. Zool. Soc. Lond., 1864, 153, Lake Nicaragua.
3369. *Astatheros erythræum* (Günther).
Great lakes of Nicaragua.
Heros erythræus Gunther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 391, 457, pl. 75, fig. 2, Lake Managua
- Genus 1166. *AMPHILOPHUS* Agassiz.
Amphilophus Agassiz, Proc. Bost. Soc. Nat. Hist., VI, 1858, 401 (*A. fraebeli* Agassiz); a large-lipped species from Lake Nicaragua not otherwise described, probably *labiatus*; these species, *labiatus* and *lobochilus*, have been placed in *Astatheros*, but the name *Amphilophus* has priority
3370. *Amphilophus labiatus* (Günther).
Great lakes of Nicaragua.
Heros labiatum Gunther, Proc. Zool. Soc. Lond., 1864, 27, Pl. IV, fig. 1, Lake Nicaragua.
3371. *Amphilophus lobochilus* (Günther).
Lake Managua, Nicaragua.
Heros lobochilus Gunther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 391, 457, pl. 75, fig. 1, Lake Managua, Nicaragua.
- Genus 1167. *THORICHTHYS* Meek.
Thorichthys Meek, Field Col. Mus., Zool., V, 1904, 222 (*T. ellioti* Meek).
3372. *Thorichthys ellioti* Meek.
Atlantic slope of Vera Cruz to Guatemala.
Thorichthys ellioti Meek, Field Col. Mus., Zool., V, 1904, 223, Motzongongo, Vera Cruz, Mexico.
3373. *Thorichthys callolepis* (Regan).
Atlantic slope of Isthmus of Tehuantepec, Mexico.
Heros callolepis Regan, Ann. Mag. Nat. Hist., ser. 7, XIII. 1904, 258. San Domingo de Guzman, Mexico.
3374. *Thorichthys aureus* (Günther).
Atlantic slope, Vera Cruz, Mexico, to Guatemala.
Heros aureus Gunther, Cat., IV, 1862, 292, Lake Yzabal, Guatemala.
3375. *Thorichthys affinis* (Günther).
Lake Peten, Guatemala.
Heros affinis Gunther, Cat., IV, 1862, 292, Lake Peten.
3376. *Thorichthys acutus* (Miller).
Small tributaries of the Rio Motagua, Guatemala.
Cichlasoma acutum Miller, Bull. Amer. Mus. Nat. Hist., XXIII, 1907, 117, fig. 6, Rio Tenedores, Guatemala.
3377. *Thorichthys helleri* (Steindachner).
Lowland streams of Mexico emptying into the Gulf south of Vera Cruz.
Heros helleri Steindachner, Denks. Akad. Wiss. Wien, XXIII, 1864, 64, Pl. IV, fig. 1, Rio Teapa, Tabasco, Mexico.
Heros maculipinnis Steindachner, loc. cit., 69, Pl. IV, fig. 2, Rio Zanopa, Mexico.
- Genus 1168. *HERICHTHYS* Baird and Girard.
Herichthys Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 25 (*H. cyanoguttatus* Baird and Girard).
3378. *Herichthys cyanoguttatus* Baird and Girard.
East slope of Mexico from Rio Grande to Rio Panuco.
Herichthys cyanoguttatus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 25, Rio Grande, Brownsville, Tex.
Heros teporatus Fowler, loc. cit., LV, 1903, 321, Victoria, Tamaulipas.
3379. *Herichthys pavonaceus* (Garman).
Spring near Monclova, Coahuila, Mexico.
Heros pavonaceus Garman, Bull. Mus. Comp. Zool., VIII, 1881, 93, spring near Monclova, Coahuila, Mexico.

3380. *Herichthys bocourti* (Vaillant and Pellegrin).
Lake Isabel, Guatemala
Neotroplus bocourti Vaillant and Pellegrin, Bull. Mus. Hist. Nat., Paris, 1902, 85, Lake Isabel, Guatemala
3381. *Herichthys carpintis* (Jordan and Snyder).
Rio Panuco, Mexico, and its lowland tributaries.
Neotroplus carpintis Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 146, fig. 22, Laguna de Carpinte, Tampico, Mexico.
3382. *Herichthys troscheli* (Steindachner).
Mexico.
Heros troscheli Steindachner, Ichthyologische Notizen, IV, 1867, 12, Mexico.
- Genus 1169. **TOMOCICHLA** Regan.
Tomocichla Regan, Ann. Mag. Nat. Hist., II, ser. 8, 1908, 463 (*Herichthys underwoodi* Regan)
3383. *Tomocichla underwoodi* (Regan).
Costa Rica.
Herichthys underwoodi Regan, Biol. Central Amer., Pisces, 1908, 30, Costa Rica.
3384. *Tomocichla geddesi* (Regan).
Southern Mexico.
Herichthys geddesi Regan, Ann. Mag. Hist., XVI, ser. 7, 1905, 436, southern Mexico.
- Genus 1170. **PETENIA** Günther.
Petenia Günther, Cat., IV, 1862, 301 (*P. splendida* Günther).
3385. *Petenia splendida* Günther.
Lake Peten, Guatemala; Montecristo; Frontera.
Petenia splendida Günther, Cat., IV, 1862, 301, Lake Peten, Guatemala.
- Genus 1171. **PARANEETROPLUS** Regan.
Paraneetroplus Regan, Ann. Mag. Nat. Hist., XVI, ser. 7, 1905, 436 (*P. bulleri* Regan).
3386. *Paraneetroplus bulleri* Regan.
Rio de Sarabia, Mexico.
Paraneetroplus bulleri Regan, Ann. Mag. Nat. Hist., XVI, ser. 7, 1905, 437, Rio de Sarabia, Mexico.
- Genus 1172. **NEETROPLUS** Günther.
Neetroplus Günther, Trans. Zool. Soc. Lond. VI, 1866, 469 (*N. nematopus* Günther).
3387. *Neetroplus sieboldii* (Kner and Steindachner).
New Granada, and from the Pacific slope of Panama
Heros sieboldii Kner and Steindachner, Abhandl. Bayer Akad. Wiss., X, 1864, 13, pl. 2, fig. 2, Panama, Colombia.
3388. *Neetroplus nematopus* Günther.
Great lakes of Nicaragua.
Neetroplus nematopus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 392, 470, pl. 74, fig. 4, Lake Managua.
- Genus 1173. **HEROTILAPIA** Regan.
Herotilapia Regan, Ann. Mag. Nat. Hist., V, 38 (*Tilapia buettikoferi* Boulenger).
3389. *Herotilapia multispinosa* (Günther).
Great lakes of Nicaragua.
Heros multispinosus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 391, 453, pl. LXXIV, fig. 2, Lake Managua.
- Genus 1174. **ÆQUIDENS** Eigenmann and Bray.
Æquidens Eigenmann and Bray, Ann. N. Y. Acad. Sci., 1894, 616 (*Acara tetramerus* Heckel).
3390. *Æquidens cœruleopunctata* (Kner and Steindachner).
Panama to northwest Ecuador.
Acara cœruleopunctata Kner and Steindachner, Sitz. Bayer. Akad., 1863, 222, northwest Ecuador.

Genus 1175. *GEOPHAGUS* Heckel.

Geophagus Heckel, Ann. Mus. Wien, II, 1840, 383 (*G. magasema* Heckel = *Sparus surinamensis* Bloch)

Mesops Gunther, Cat., IV, 1862, 311 (*Geophagus cupido* Heckel).

Satanoperca Gunther, loc. cit., 312 (*Geophagus demon* Heckel).

Subgenus *SATANOPERCA* Günther.3391. *Geophagus crassilabris* Steindachner

Panama.

Geophagus (*Satanoperca*) *crassilabris* Steindachner, Sitz. Akad. Wiss. Wien, 1876, 74, 65, Pl. VII, Panama.

Order UU. PHARYNGOGNATHI

Family 251. LABRIDÆ

Genus 1176. *CENTROLABRUS* Gunther, *Rock cooks*.

Centrolabrus Gunther, Cat., IV, 1862, 92 (*Labrus exoletus* Linnæus).

3392. *Centrolabrus exoletus* (Linnæus). *Rock cook*.

Coast of northern Europe south to Cornwall; Greenland.

Labrus exoletus Linnæus, Syst. Nat., Ed. X, 1758, 287, Atlantic Ocean.

Labrus pentacanthus Lacépède, Hist. Nat. Poiss., III, 1803, 503; after Linnæus

Crenilabrus microstoma Couch, in Thompson, Proc. Zool. Soc. London, 1837, 55, Cornwall

Genus 1177. *TAUTOGOLABRUS* Gunther, *Cunners*

Tautogolabrus Günther, Cat., IV, 1862, 89 (*Ctenolabrus bergall* Günther = *Labrus adpersus* Walbaum)

3393. *Tautogolabrus adpersus* (Walbaum). *Cunner*; *Chogset*, *Blue perch*; *Bergall*; *Berg-gyll*; *Nibbler*; *Conner*; *Gall Bengal*.

Atlantic coasts of North America from Labrador to Sandy Hook.

Labrus adpersus Walbaum, Artedi Piscium, 1792, 254, no locality; after Bergall of Schöpf.

Tautoga niger Mitchill, Report, in part, in Fishes of New York, 1814, 23, New York.

Tautoga cerulea Mitchill, Report, loc. cit., 1814, 24, New York.

Labrus chogset Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 402, pl. 3, fig. 2, New York

Labrus chogset fulva Mitchill, loc. cit., 1815, 403, New York.

Ctenolabrus unnotatus Cuvier and Valenciennes, Hist. Nat. Poiss., XIII, 1839, 174 (239), New York.

Ctenolabrus bergall Günther, Cat., IV, 1862, 90, Canada.

Genus 1178. *TAUTOGA* Mitchill, *Tautogs*.

Hiatula Lacépède, Hist. Nat. Poiss., II, 1800, 522; name preoccupied.

Tautoga Mitchill, Rept. Fishes New York, 1814, 23 (*T. niger* Mitchill = *Labrus tautoga* Mitchill = *Labrus onitis* Linnæus).

3394. *Tautoga onitis* (Linnæus). *Tautog*; *Blackfish*; *Oysterfish*; *Tautoga*; *Smooth blackfish*; *Cub*; *Sea dog*.

Atlantic coast of United States; Cape Ann to Charleston.

Labrus onitis Linnæus, Syst. Nat., Ed. X, 1758, 286, locality unknown.

Labrus hiatula Linnæus, Syst. Nat., Ed. XII, 1766, 475, Carolinas

Labrus carolinus Bonnaterre, Tableau Encyclopéd. et Méthod., Ichthyologie, 1788, 113, Carolinas.

Labrus subfucus Walbaum, Artedi Pisc., 1792, 254; after Schöpf.

?*Labrus tassellatus* Bloch, Naturgesch. Ausl. Fische, VI, 1792, pl. 8, Norway.

Hiatula gardemana Lacépède, Hist. Nat. Poiss., II, 1800, 522, Carolinas.

Labrus americanus Bloch and Schneider, Syst. Ichth., 1801, 261; after Schöpf.

Labrus tautoga Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 398, Long Island; Rhode Island; Cape Cod; Sandy Circle.

Tautoga fusca Mitchill, Rept. Fish. N. Y., 1814, 23, New York.

Tautoga rubens Mitchill, loc. cit., 23, New York.

Tautoga atia Mitchill, loc. cit., 23, New York.

Tautoga niger Mitchill, loc. cit., 23, New York.

Genus 1179. *ACHNOLAIMUS* Cuvier and Valenciennes. *Capitaines*.
Lachnolaimus Cuvier and Valenciennes, Hist. Nat. Poiss., XIII, 1839,
 200 (274) (*L. aigula* Cuvier and Valenciennes=*Labrus maximus*
 Walbaum).
Lachnolæmus Günther; corrected spelling.

3395. *Lachnolaimus maximus* (Walbaum). *Capitaine*; *Perro perro*; *Hogfish*.
 West Indies, north to Key West and the Bermudas.
Suillus Catesby, Nat. Hist. Carol., II, 1743, 15, pl. 15; no locality;
 probably Bahamas.
Labrus maximus Walbaum, Artedi Pisc., 1792, 261, no locality; after
 Catesby.
Lachnolaimus suillus Cuvier, Règne Animal, ed. 2, II, 1829, 257; after
 Catesby.
Lachnolaimus aigula Cuvier and Valenciennes, Hist. Nat. Poiss., XIII,
 1839, 202 (277), pl. 378, St. Bartholomew.
Lachnolaimus dux Cuvier and Valenciennes, loc. cit., 208 (285), Mar-
 tinique.
Lachnolaimus caninus Cuvier and Valenciennes, loc. cit., 210 (288),
 St. Thomas; San Domingo.
Lachnolaimus psittacus Cuvier and Valenciennes, loc. cit., 212 (291);
 Porto Rico.

Genus 1180. *BODIANUS* Bloch.

- Bodianus* Bloch, Nat. Ausl. Fische, IV, 1790, 31, 33 (*B. bodianus*
 Bloch=*Labrus rufus* Linnæus); in part; type by tautonymy; not by
 first restriction to species of *Cephalopholis*.
Harpe Lacépède, Hist. Nat. Poiss., IV, 1803, 426 (*H. carulcoareum*
 Lacépède=*Labrus rufus* Linnæus).
Cossyphus Cuvier and Valenciennes, loc. cit., XIII, 1839, 102 (*Bodianus*
bodianus Bloch); name twice preoccupied.
Crenilabrus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839 (*Lutjanus*
verres Bloch=*Labrus rufus* Linnæus); name preoccupied.
3396. *Bodianus rufus* (Linnæus). *Ladyfish*; *Spanish ladyfish*; *Spanish hogfish*;
Pudiano; *Perro colorado*.
 West Indies, north to Key West, south to Rio de Janeiro.
Labrus rufus Linnæus, Syst. Nat., Ed. X, 1758, 284, America; after
 Catesby.
Pudiano vermelho Marcgrave, Hist. Bras., 1648, 145, Brazil.
Turdus flavus Catesby, Nat. Hist. Carolina, II, 1743, 11, pl. 11, fig. 1,
 Bahamas.
Perro colorado Parra, Descr. Dif. Piez. Hist. Nat. Cuba, 1787, 3, lam. 3,
 fig. 1, Habana.
 ?*Bodianus bodianus* Bloch, Naturgesch. Ausl. Fische, IV, 1790, 33,
 Brazil.
 ?*Lutjanus verres* Bloch, loc. cit., V, 1791, 9, locality uncertain
 ?*Sparus falcatus* Bloch, loc. cit., 20, Martinique; after Plumier.
Labrus semiruber Lacépède, Hist. Nat. Poiss., III, 1802, 428, Rio
 Janeiro.
Bodianus blochii Lacépède, loc. cit., IV, 1803, 279; after Bloch.
Harpe coeruleo-aureus Lacépède, loc. cit., 426, pl. 8, fig. 2, Martinique.
3397. *Bodianus diplotænia* (Gill).
 Pacific coast of tropical America; Cape San Lucas; Panama; Revilla-
 giedo Islands; Mazatlan.
Harpe diplotænia Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 140,
 Cape San Lucas; female.
Harpe pectoralis Gill, loc. cit., 141, Cape San Lucas; male.
3398. *Bodianus pulchellus* (Poey).
 Cuba.
Cossyphus pulchellus Poey, Memorias, II, 1860, 208, Habana.
- Genus 1181. *DECODON* Günther.
Decodon Günther, Cat., IV, 1862, 101 (*Cossyphus puellaris* Poey).
3399. *Decodon puellaris* (Poey).
 West Indies; Cuba; Barbados; Snapper Banks off Pensacola, Fla.
Cossyphus puellaris Poey, Memorias, II, 186, 2101, Habana.

Genus 1182. *PIMELOMETOPON* Gill. *Fatheads*.

Pimelometopon Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 58 (*Labrus pulcher* Ayres).

3400. *Pimelometopon pulcher* (Ayres). *California redfish; Fathead; Sheep-head*.

Southern California, from Point Conception to Ascension Island, Lower California.

Labrus pulcher Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 3, San Diego, Calif.

Genus 1183. *CLEPTICUS* Cuvier.

Clepticus Cuvier, Règne Anim., ed. 2, II, 1829, 261 (*Rabirubia genizara* Parra = *Brama parræ* Bloch and Schneider).

3401. *Clepticus parræ* (Bloch and Schneider). *Genizara; Janissary; Creole*. West Indies.

Rabirubia genizara Parra, Dif. Piezas de Hist. Nat. Cuba, 1787, 44, pl. 21, fig. 1, Habana; author not binomial.

Brama parræ Bloch and Schneider, Syst. Ichth., 1801, 100, Habana.

Family 252. *CORIDÆ*Genus 1184. *IRIDIO*⁸⁴ Jordan and Evermann. *Doncellas*.

Halichares Rupell, Neue Wirbelthiere, Fische, 1835, 17 (*bimaculatus*); not *Halichærus* Nilsson, 1820, a genus of seals.

Hermiulus Swainson, Nat. Hist. Classn. Fishes, II, 1839, 228, (*Labrus guttatus* Bloch); as restricted by Bonaparte (1841); restricted by Swain (1882) to *Cheilio auratus* Quoy and Gaimard.

Parajulis Bleeker, Ned. Tijdschr. Dierk., II, 1865, 251 (*Julis pæcilopecterus* Temminck and Schlegel).

Iridio Jordan and Evermann, Check List, 1895, 412 (*Julis dimidiatus* Agassiz = *Labrus cyanocephalus* Bloch).

3402. *Iridio radiata* (Linnæus). *Puddingwife; Doncella; Pudiano verde*.

West Indies, Brazil north to Florida Keys, Bermuda, and St. Paul Rocks.

Labrus radiatus Linnæus, Syst. Nat., Ed. X, 1758, 288, Bahamas; based on Catesby.

Pudiano verde Marcgrave, Hist. Pisc. Brasil., 1648, 146, Brazil.

Turdus oculo-radiatus Catesby, Nat. Hist. Carolina, II, 1743, 12, pl. 12, fig. 1, Bahamas.

Doncella Parra, Desc. Dif. Piez. Hist. Nat. Cuba, 1787, 95, lam. 37, fig. 1, Habana.

Labrus brasiliensis Bloch, Naturgesch. Ausl. Fische, V, 1791, 125, Brazil.

Julis crotaphus Cuvier, Règne Anim., ed. 2, II, 1829, 258; after Parra.

Julis cyanostigma Cuvier and Valenciennes, Hist. Nat. Poiss., XIII, 1839, 286 (391), Martinique.

Julis opalina Cuvier and Valenciennes, loc. cit., 287 (392), Martinique.

Julis patatus Cuvier and Valenciennes, loc. cit., 291 (398), Martinique; Cuba.

Julis principis Cuvier and Valenciennes, loc. cit., 294 (402), Bahia.

3403. *Iridio bivittata*⁸⁵ (Bloch). *Slimy dick; Doncella*.

West Indies, north to Beaufort, N. C., and south to Brazil.

Labrus bivittatus Bloch, Naturgesch. Ausl. Fische, V, 1791, 133, from a painting by Plumier made at Martinique.

Sparus radiatus Linnæus, Syst. Nat., ed. XII, 1766, 472, Carolinas.

Labrus psittaculus Lacépède, Hist. Nat. Poiss., III, 1800, 452, Martinique; after Plumier.

Julis humeralis Poey, Memorias, II, 1860, 212, Cuba.

Chærojulis grandisquamis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 206, Beaufort, N. C.

Chærojulis arangoi Poey, Enumeratio, 1876, 109, Habana.

Platyglossus florealis Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 287, Pensacola, Fla.

⁸⁴ This genus is very close to the Old World genus *Halichares*.

⁸⁵ Regarded by some authors as the young of *Iridio radiata*.

3404. *Iridio cyanocephalus* (Bloch).
West Indies, south to Brazil.
Labrus cyanocephalus Bloch, Naturgesch. Ausl. Fische, V, 1791, 139,
location unknown; Museum of Link
Julis dimidiatus Agassiz, in Spix, Pisc. Bras., 1829, 96, pl. 53, Brazil.
Julis internasalis Poey, Memorias, II, 1861, 421, Cuba.
3405. *Iridio maculipinna* (Müller and Troschel).
West Indies to Beaufort, N. C.
Julis maculipinna Müller and Troschel, in Schomburgk, Hist. Barbados,
1848, 674, Barbados.
3406. *Iridio garnoti* (Cuvier and Valenciennes).
West Indies, Cuba, Martinique, St. Croix.
Julis garnoti Cuvier and Valenciennes, Hist. Nat. Poiss., XIII, 1839,
285 (390), Martinique.
Julis cinctus Poey, Memorias, II, 1860, 211, Cuba.
Julis ruptus Poey, loc. cit., 212, Cuba.
3407. *Iridio kirschii* Jordan and Evermann.
Cuba to Brazil.
Julis crotaphus Cuvier and Valenciennes, Hist. Nat. Poiss., XIII,
1839, 289 (395), Bahia; not of Cuvier.
Iridio kirschii Jordan and Evermann, Check List, 1895, 413, Bahia,
Brazil.
3408. *Iridio poeyi* (Steindachner).
Surinam
PlatyGLOSSUS poeyi Steindachner, Ichth. Notizen, VI, 1867, 49, Surinam.
3409. *Iridio caudalis* (Poey).
Cuba.
Julis caudalis Poey, Memorias, II, 1860, 213, Habana.
3410. *Iridio pictus* (Poey).
West Indies, north to the Snapper Banks off Pensacola, Fla.
Julis pictus Poey, Memorias, II, 1860, 214, Cuba.
3411. *Iridio elegans* Bean.
Bermudas to Curacao.
Iridio elegans Bean, Field Mus. Nat. Hist., Zool., VII, 1906, 65, fig. 6,
Nonesuch Island, Bermudas.
3412. *Iridio similis* Nichols.
Bermudas.
Iridio similis Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 61,
Bermudas.
3413. *Iridio frenatus* Nichols.
Bermuda.
Iridio frenatus Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 61,
Bermuda.
3414. *Iridio decoratus* Bean.
Bermudas.
Iridio decoratus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 29, Nonesuch
Island, Bermudas.
3415. *Iridio meyeri* Bean.
Bermudas.
Iridio meyeri Bean, Proc. Biol. Soc. Wash., XIX, 1906, 29, Nonesuch
Island, Bermudas.
3416. *Iridio microstomus* Bean.
Bermudas.
Iridio microstomus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 30, None-
such Island, Bermudas.
3417. *Iridio semicinctus* (Ayres). *Kelpfish; Señorita*.
Southern California, Santa Barbara Islands to Cedros Island.
Julis semicinctus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 32,
Cerros [Cedros] Island, off coast of Lower California.
3418. *Iridio dispilus* (Günther).
Pacific coast of tropical America, Mazatlan to Panama.
PlatyGLOSSUS dispilus Günther, Proc. Zool. Soc. Lond., 1864, 25, Panama.

3419. *Iridio nicholsi* (Jordan and Gilbert).
Revillagigedo and Galapagos Islands.
PlatyGLOSSUS nicholsi Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 231, Braithwaite Bay, Socorro Island.
3420. *Iridio sellifer* (Gilbert).
Revillagigedo Archipelago
Halchæres sellifer Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 67, Clarion Island.
- Genus 1185. *OXYJULIS* Gill. *Señoritas*.
Oxyjulis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 330 (*Julis modestus* Girard).
3421. *Oxyjulis californicus* (Gunther) *Señorita*.
Coast of southern California; common from Monterey to Guadalupe Island.
Julis modestus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 151, San Diego, Monterey, San Miguel; preoccupied.
Halchæres californicus Gunther, Ann. Mag. Nat. Hist., ser. 3, VIII, 1861, 386, name only; substitute for *Julis modestus*, preoccupied.
- Genus 1186. *PSEUDOJULIS* Bleeker.
Pseudojulis Bleeker, Proc. Zool. Soc. Lond., 1861, 412 (*P. girardi* Bleeker).
- Subgenus *PSEUDOJULIS*.
3422. *Pseudojulis inornatus* Gilbert.
Pacific coast of Mexico; only the type known.
Pseudojulis inornatus Gilbert, Proc. U. S. Nat. Mus., XI II, 1890, 67, Pacific coast of Mexico.
3423. *Pseudojulis melanotis* Gilbert.
Pacific coast of Mexico; only the type known.
Pseudojulis melanotis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 66, Pacific coast of Mexico.
- Genus 1187. *EMMEEKIA* Jordan and Evermann.
Emmeekia Jordan and Evermann, Check List, 1895, 413 (*Pseudojulis venustus* Jenkins and Evermann).
3424. *Emmeekia venusta* (Jenkins and Evermann).
Gulf of California.
Pseudojulis venustus Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 145, Guaymas, Sonora, Mexico.
- Genus 1188. *JULIDIO* Jordan and Evermann.
Julidio Jordan and Evermann, Check List, 1895, 413 (*Pseudojulis adustus* Gilbert).
3425. *Julidio adustus* (Gilbert).
Revillagigedo Islands.
Pseudojulis adustus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 66, Socorro Island.
3426. *Julidio notospilus* (Günther).
Pacific coast of Mexico, Panama, and Clarion Island of Revillagigedo Archipelago.
Pseudojulis notospilus Günther, Proc. Zool. Soc. Lond., 1864, 26, Panama.
- Genus 1189. *THALASSOMA* Swainson.
Thalassoma Swainson, Nat. Hist. Class. Fishes, II, 1839, 224 (*Labrus purpureus* Rüppell).
Chlorichthys Swainson, loc. cit., 232 (*Labrus bifasciatus* Bloch).
Julis Günther, not of Cuvier and Valenciennes.

3427. *Thalassoma bifasciatum* (Bloch).
West Indies; Tortugas; Jamaica; Cuba; San Domingo; and Martinique.
? *Labrus bifasciatus* Bloch, Naturgesch. Ausl. Fische, V, 1791, 131,
West Indies.
Labrus bifasciatus torquatus Bloch and Schneider, Syst. Ichth., 1801,
243, Antilles; after Gronow.
Julis detorsor Cuvier and Valenciennes, Hist. Nat. Poiss., XIII, 1839,
298 (408), San Domingo.
Labrus ornatus Gronow, Syst. Nat., ed. Gray, 1854, 83, Antilles.
Julis gillmanus Poey, Memorias, II, 1860, 214, Cuba
? *Julis nitida* Gunther, Cat., IV, 1862, 190, Jamaica
3428. *Thalassoma socorroensis* Gilbert.
Revillagigedo Archipelago.
Thalassoma socorroense Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 69,
Socorro Island
3429. *Thalassoma nitidum* (Gunther).
West Indies; Jamaica; Tortugas
Julis nitida Gunther, Cat., IV, 1862, 190, Jamaica.
Julis nitidissima Goode, Am. Jour. Sci. Arts, XIV, 1877, 293, Bermuda.
3430. *Thalassoma grammaticum* Gilbert.
Revillagigedo Archipelago.
Thalassoma grammaticum Gilbert, Proc. U. S. Nat. Mus., XIII, 1890,
68, Socorro Island; Clarion Island.
3431. *Thalassoma virens* Gilbert
Revillagigedo Archipelago.
Thalassoma virens Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 68,
Socorro Island.
3432. *Thalassoma lucasanum* (Gill).
Gulf of California, Cape San Lucas to Mazatlan and Tres Marias
Islands; Panama Bay.
Julis lucasanus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 142,
Cape San Lucas.
3433. *Thalassoma steindachneri* (Jordan).
Acapulco, Pacific coast of Mexico.
Thalassoma steindachneri Jordan, Rept. U. S. Fish Com., XV, 1887,
654, Acapulco; after Steindachner.
Julis melanochir Steindachner, Ichth. Beitr., III, 1875, 63, Acapulco;
not of Bleeker
- Genus 1190. *DORATONOTUS* Günther.
Doratonotus Gunther, Cat., IV, 1862, 124 (*D. megalepis* Gunther).
3434. *Doratonotus megalepis* Gunther.
West Indies, north to Key West.
Doratonotus megalepis Günther, Cat., IV, 1862, 125, St. Christopher.
Doratonotus thalassinus Jordan and Gilbert, Proc. U. S. Nat. Mus.,
VII, 1884, 28, Key West, Fla.
3435. *Doratonotus decoris* Evermann and Marsh.
Porto Rico.
Doratonotus decoris Evermann and Marsh, Rept. U. S. Fish Com.,
XXV, 1899, 354, Ponce, Porto Rico.
3436. *Doratonotus boekei* Metzelaar
Caribbean Sea.
Doratonotus boekei Metzelaar, Trop. Atlant. Visschen, 1919, 107, fig. 31,
Curaçao, Bonaire.
- Genus 1191. *XYRULA* Jordan.
Xyrula Jordan, Rept. U. S. Fish Com., XV, 1887, 656 (*Xyrichthys jessiae* Jordan).
3437. *Xyrula jessiae* (Jordan).
Snapper Banks off Tampa, Fla.
Xyrichthys jessiae Jordan, Proc. U. S. Nat. Mus., X, 1887, 698, off
Tampa Bay, Fla.

Genus 1192. NOVACULICHTHYS Bleeker.

Novaculichthys Bleeker, Proc. Zool. Soc. Lond., 1861, 414 (*Labrus tænnurus* Bloch).

Dimalacocentrus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 223 (*Novacula kallasoma* Bleeker)

3438. *Novaculichthys martinicensis* (Cuvier and Valenciennes).

Martinique.

Xyrichthys martinicensis Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 37 (49), Martinique

Xyrichthys vitta Cuvier and Valenciennes, loc. cit., 38 (51), no locality.

3439. *Novaculichthys infirmus* (Bean)

Cozumel, Yucatan.

Xyrichthys infirmus Bean, Bull. U. S. Fish Com., VIII, 1888, 199, pl. 29, fig. 2, Cozumel

3440. *Novaculichthys rosipes* (Jordan and Gilbert).

Key West; Florida; Tortugas

Xyrichthys rosipes Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 27, Key West, Tortugas.

3441. *Novaculichthys ventralis* (Bean).

Cozumel Island, Yucatan.

Xyrichthys ventralis Bean, Bull. U. S. Fish Com., VIII, 1888, 198, pl. 29, fig. 1, Cozumel, Yucatan.

Genus 1193. XYRICHTHYS Cuvier. *Razorfishes*.

Xyrichthys Cuvier, Mém. Mus. d'Hist. Nat., I, 1815, 324, 329 (*X. novacula* Cuvier)

Novacula Cuvier, Règne Anim., Ed. I, 1817, 265 (*Coryphæna novacula* Linnæus).

Amorphocephalus Bowdich, Exc. Madeira, 1825, 238 (*A. granulatus* Bowdich).

3442. *Xyrichthys psittacus* (Linnæus). *Razorfish*.

West Indies, north to Pensacola and Charleston.

Coryphæna psittacus Linnæus, Syst. Nat., Ed. XII, 1766, 448, Charleston, S. C.

Coryphæna lineata Gmelin, Syst. Nat., 1788, 1195, Charleston, S. C.

Xyrichthys venustus Poey, Enumeratio, 1876, 110, Martinique.

Xyrichthys vermiculatus Poey, Memorias, II, 1860, 215, Cuba.

3443. *Xyrichthys mundiceps* Gill.

Cape San Lucas, Lower California.

Xyrichthys mundiceps Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 143, Cape San Lucas.

3444. *Xyrichthys modestus* Poey.

Cuba.

Xyrichthys modestus Poey, Repertorio, II, 1867, 238, Cuba.

3445. *Xyrichthys binghami* Mowbray.

Bahama Islands.

Xyrichthys binghami Mowbray, Marine Life, I, Apr. 2, 1925, 1, Cay Sal Bank, Bahamas.

Genus 1194. INIISTIUS Gill.

Iniistius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 143 (*Xyrichthys pavo* Cuvier and Valenciennes).

3446. *Iniistius pavoninus* (Cuvier and Valenciennes).

Hawaii and Pacific coast of Mexico.

Xyrichthys pavoninus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 63, Hawaii.

Iniistius mundicorpus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 145, Cape San Lucas.

Genus 1195. BERMUDICHTHYS Nichols.

Bermudichthys Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 62 (*B. subfurcatus* Nichols).

3447. *Bermudichthys subfurcatus* Nichols.

Bermuda.

Bermudichthys subfurcatus Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 62, Bermuda.

Family 253 SPARISOMIDÆ

Genus 1196. CRYPTOTOMUS Cope.

Cryptotomus Cope, Trans. Amer. Philos. Soc., XIV, 1870, 462 (*C. roseus* Cope).

Nicholsina Fowler, Copeia, No. 14, 1915, 3 (*Cryptotomus beryllinus* Jordan and Swain).

Subgenus CRYPTOTOMUS Cope.

3448. *Cryptotomus roseus* Cope

West Indies, south to Brazil.

Cryptotomus roseus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 462, St. Martins, West Indies.

3449. *Cryptotomus crassiceps* Bean.

Bermudas.

Cryptotomus crassiceps Bean, Proc. Biol. Soc. Wash., XIX, 1906, 32, Coopers Island, Bermudas.

Subgenus NICHOLSINA Fowler.

3450. *Cryptotomus beryllinus* Jordan and Swain. *Parrotfish*.

Florida Keys, north to New Jersey, south to Rio Janeiro.

Cryptotomus beryllinus Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 101, Habana.

Sparisoma sp. Bean, Bull. U. S. Fish Com., VII, 1887, 137, Somers Point, N. J.

3451. *Cryptotomus dentiens* (Poey).

Cuba.

Scarus dentiens Poey, Memorias, II, 1861, 422, Cuba.

3452. *Cryptotomus retractus* (Poey).

West Indies, north to Pensacola.

Calliodon retractus Poey, Synopsis, 1868, 245, Habana.

3453. *Cryptotomus ustus* (Cuvier and Valenciennes).

West Indies, north to Charleston, south to Bahia.

Calliodon ustus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 212 (286), pl. 405, Brazil.

3454. *Cryptotomus auropunctatus* (Cuvier and Valenciennes).

San Domingo.

Calliodon auro-punctatus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 215 (290), Santo Domingo.

Genus 1197. LEPTOSCARUS Swainson.

Leptoscarus Swainson, Nat. Hist. Classn. Fishes, 1839, 11 (*Scarus vaigiensis* Quoy and Gaimard = *Scarus spinidens* Cuvier and Valenciennes).

Calotomus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 70 (*C. xenodon* Gilbert).

3455. *Leptoscarus xenodon* (Gilbert).

Socorro Island, Revillagigedo Archipelago.

Calotomus xenodon Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 70, Socorro Island, Revillagigedo Islands.

Genus 1198. SPARISOMA Swainson. *Vezas*.

Sparisoma Swainson, Nat. Hist. Classn. Fishes, etc., II, 1839, 227 (*Scarus abildgardii* Bloch).

Euscarus Jordan and Evermann, Check List, 1895, 416 (*Labrus cretensis* Linnæus).

Subgenus SPARISOMA Swainson.

3456. *Sparisoma xystrodon* Jordan and Swain.

West Indies, north to Key West.

Sparisoma xystrodon Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 99, Key West; Habana.

3457. *Sparisoma atomarium* (Poey).

Cuba.

Scarus atomarius Poey, Memorias, II, 1861, 423, Habana.

3458. *Sparisoma radians* (Cuvier and Valenciennes).
West Indies, south to Brazil; Tortugas.
Scarus radians Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 153 (206), Brazil.
Scarus lacrimosus Poey, Memorias, II, 1861, 422, Cuba.
3459. *Sparisoma hoplomystax* (Cope)
West Indies, from Key West to Bahia.
Scarus hoplomystax Cope, Trans. Amer. Phil. Soc., XIV, 1871, 462, St. Martins, West Indies.
Sparisoma cyanolene Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 98, Key West, Fla.
3460. *Sparisoma niphobles* Jordan and Bollman
Bahamas; Tortugas; Cape Florida; Key West; Porto Rico; abundant about Porto Rico.
Sparisoma niphobles Jordan and Bollman, Proc. U. S. Nat. Mus., XI, 1888, 551, Green Turtle Cay, Bahamas.
3461. *Sparisoma aurofrenatum* (Cuvier and Valenciennes).
West Indies.
Scarus aurofrenatus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 142 (191), San Domingo.
Scarus miniofrenatus Poey, Memorias, II, 1860, 279, Cuba.
3462. *Sparisoma oxybrachium* (Poey).
Cuba.
Scarus oxybrachius Poey, Synopsis, 1868, 342, Habana.
3463. *Sparisoma abildgaardi* (Bloch). *Red parrotfish; Roseback parrot.*
West Indies south to Brazil; Tortugas; generally common.
Sparus abildgaardi Bloch, Nat. Ausl. Fische, V, 1791, 22, pl. 259, America.
Vieja Parra, Descr. Dif. Piezas Hist. Nat., 1787, 58, pl. 28, fig. 2, Cuba.
Scarus coccineus Bloch and Schneider, Syst. Ichth., 1801, 289, Cuba; after Parra.
Sparus aureo-ruber Lacépède, Hist. Nat. Poiss., IV, 1803, 55, 163, Martinique; on a drawing by Plumier.
Scarus amplius Ranzani, Nov. Comm. Ac. Scient. Inst. Bonon., V, 1842, 324, pl. 25, Brazil (fide Guichenot).
Scarus erythrinoides Guichenot, Mem. Soc. Sci. Nat. Cherbourg, XI, 1865, 10, San Domingo.
3464. *Sparisoma distinctum* (Poey).
Cuba.
Scarus distinctus Poey, Memorias, II, 1861, 423, Habana.
3465. *Sparisoma chrysopterus* (Bloch and Schneider). *Blue parrotfish; Vieja.*
West Indies, south to Bahia.
Scarus chrysopterus Bloch and Schneider, Syst. Ichth., 1801, 286, pl. 57, American seas
Vieja Parra, Descr. Dif. Piezas Hist. Nat., 1787, 59, pl. 28, fig. 4, Cuba.
Scarus chloris Bloch and Schneider, Syst. Ichth., 1801, 289, Cuba.
Scarus lateralis Poey, Memorias, II, 1860, 219, Cuba.
?Scarus spinidens Guichenot, Mem. Soc. Sci. Nat. Cherbourg, XI, 1865, 15, Bahia.
3466. *Sparisoma lorito* Jordan and Swain.
West Indies.
Sparisoma lorito Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 95, Habana.
3467. *Sparisoma viride* (Bonnaterre). *Green parrotfish.*
Tortugas to West Indies.
Scarus viridis Bonnaterre, Enc. Meth., 1788, 96, Bahamas; after Catesby.
Piscis viridis bahamensis Catesby, Nat. Hist. Car., II, 1743, 29, pl. 29, Bahamas.
Scarus melanotis Bleeker, Verslagen Koning. Akad. Wetenschappen, XIV, 1862, 126, St. Croix.
Scarus (Catesby) Lacépède, Hist. Nat. Poiss., IV, 1803, 16; after Catesby.

3468. *Sparisoma flavescens* (Bloch and Schneider). *Mud parrot*; *Vieja colorada*, *Vieja mager*, *Parrotfish*.
West Indies; Key West to Rio Janeiro; Woods Hole, Mass.
Scarus flavescens Bloch and Schneider, Syst. Ichth., 1801, 290, Cuba;
after Parra.
Vieja Parra, Descr. Dif. Piezas Hist. Nat., 1787, 59, pl. 28, fig. 4, Cuba.
Scarus squalidus Poey, Memorias, II, 1860, 218, Cuba.
3469. *Sparisoma rubripinne* (Cuvier and Valenciennes).
West Indies.
Scarus rubripinnus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV,
1839, 147 (199), San Domingo.
Scarus virens Cuvier and Valenciennes, loc. cit., 1839, 150 (203), Porto
Rico; Martinique.
Scarus circumnotatus Poey, Memorias, II, 1861, 423, Cuba.
Scarus truncatus Poey, Synopsis, 1868, 339, Cuba.
?Scarus emarginatum Poey, Synopsis, 1868, 340, Cuba.
3470. *Sparisoma strigatum* (Gunther).
Scarus strigatus Gunther, Cat., IV, 1862, 212, locality unknown, but
as the genus *Sparisoma* is chiefly confined to American waters, this
species is probably American.
3471. *Sparisoma frondosum* (Cuvier).
West Indies, south to Brazil.
Scarus frondosus Cuvier, in Agassiz, Spix, Pisc. Brasil, 1829, 98, Bahia.
Scarus aracanga Gunther, Cat., IV, 1862, 209, Jamaica.
3472. *Sparisoma brachiale* (Poey).
Cuba.
Scarus brachialis Poey, Memorias, II, 1861, 345, Cuba.
?Scarus humeralis Poey, loc. cit., 422, Habana.
3473. *Sparisoma maschalespilos* (Bleeker).
Surinam.
Scarus maschalespilos Bleeker, Verslagen Koning. Akad. Vet., XIV,
1862, 127, Surinam.

Family 254. SCARIDÆ

- Genus 1199. *SCARUS* Forskål.
Callyodon Gronow, Zoophyl., I, 1763, 72 (*Scarus croicensis* Bloch);
work not binomial.
Scarus Forskål, Descr. Anim., 1775, 25 (*Scarus croicensis* Bloch); not of
Gronow.
Hemistoma Swainson, Classn. Fish., etc., II, 1839, 226 (*Scarus pepo*
Bennett = *H. reticulata* Swainson).
Petronason Swainson, loc. cit., 226 (*Coryphæna psittacus* Linnæus).
Erythys Swainson, loc. cit., 226 (*Labrus croicensis* Bloch).
Chlorurus Swainson, loc. cit., 227 (*Scarus gibbus* Rüppell).
3474. *Scarus bollmani* Jordan and Evermann.
Gulf of Mexico in moderately deep water.
Scarus bollmani Jordan and Evermann, Proc. U. S. Nat. Mus., IX,
1886, 470, snapper banks off Tampa Bay, Fla.
3475. *Scarus punctulatus* Cuvier and Valenciennes.
West Indies; Tortugas.
Scarus punctulatus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV,
1839, 144 (195), Martinique.
Scarus diadema Cuvier and Valenciennes, loc. cit., 146 (196), Martinique.
Pseudoscarus tæniopterus Gunther, Cat., IV, 1862, 226, Trinidad.
3476. *Scarus tæniopterus* Desmarest.
West Indies.
Scarus tæniopterus Desmarest, Dict. Classique, XV, 1831, 244, pl. 12,
Cuba.
Scarus vetula Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839,
143 (193), St. Thomas; not of Bloch and Schneider.
Scarus virginalis Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884,
88, Habana.

3477. *Scarus trispinosus* Cuvier and Valenciennes
Brazil.
Scarus trispinosus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 135 (182), Brazil.
Scarus quadrispinosus Cuvier and Valenciennes, loc. cit., 146 (197), Martinique.
3478. *Scarus aracanga* Gunther.
West Indies.
Scarus aracanga Gunther, Cat., IV, 1862, 209, Jamaica.
Pseudoscarus aracanga Gunther, loc. cit., 227, Jamaica.
3479. *Scarus cuzamilæ* Bean.
Cozumel Island, Yucatan
Scarus cuzamilæ Bean, Bull. U. S. Fish. Com., VIII, 1888, 196, Cozumel, Yucatan
3480. *Scarus vetula* Bloch and Schneider. *Mudfish; Vieja; Oldwife.*
Tortugas; West Indies.
Vieja Parra. Dif. Piezas, etc., 1787, 58, pl. 28, fig. 1, Habana.
Scarus vetula Bloch and Schneider, Ichth., 1801, 289, Cuba.
Scarus superbus Poey, Memorias, II, 1860, 218, Cuba.
3481. *Scarus gnathodus* (Poey).
Cuba.
Pseudoscarus gnathodus Poey, Repertorio, II, 1867, 240, Habana.
Subgenus *CALLYODON* Gronow.
3482. *Scarus croicensis* Bloch *Bullon.*
West Indies to Key West and Woods Hole, Mass.
Scarus croicensis Bloch, Nat. Ausl. Fische, IV, 1790, 27, St. Croix.
Scarus insulæ-santæ-crucis Bloch and Schneider, Syst. Ichth., 1801, 285, St. Croix.
Scarus alternans Cuvier and Valenciennes, Hist. Nat. Poiss., 1839, 148 (200), Martinique
Pseudoscarus lneolatus Poey, Repertorio, II, 1867, 239, Cuba.
3483. *Scarus evermanni* Jordan.
Gulf of Mexico, in moderately deep water.
Scarus evermanni Jordan, Proc. U. S. Nat. Mus., IX, 1886, 469, Snapper Banks off Tampa Bay, in moderately deep water.
3484. *Scarus flavomarginatus* Cuvier and Valenciennes.
Martinique.
Scarus flavo-marginatus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 150 (202), Martinique
3485. *Scarus acutus* Poey.
Cuba.
Scarus acutus Poey, Memorias, II, 1861, 216, Habana.
3486. *Scarus cæruleus* (Bloch). *Blue parrotfish; Loro; Clamagore.*
West Indies; Chesapeake Bay.
Coryphæna cærulea Bloch, Naturgesch. Ausl. Fische, II, 1786, 120 (148), pl. 176, probably Martinique; in part; after Catesby.
Loro Parra, Descr. Dif. Piezas Hist. Nat., 1787, 57, pl. 27, fig. 1, Cuba.
Trompa Parra, loc. cit., fig. 2, Cuba
Scarus trilobatus Lacépède, Hist. Nat. Poiss., IV, 1803, 21, Martinique.
?Scarus holocyanæus Lacépède, loc. cit., 45, Martinique.
Scarus obtusus Poey, Memorias, II, 1861, 217, Cuba.
Scarus nuchalis Poey, loc. cit., 220, Cuba.
3487. *Scarus compressus* (Osburn and Nichols).
Gulf of California.
Callyodon compressus Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 170, fig. 12, Concepcion Bay, Gulf of California.
3488. *Scarus noyesi* Heller and Snodgrass.
Gulf of California to Galapagos Islands.
Scarus noyesi Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 206, Tagus Cove, Albemarle Island.

- 3489. *Scarus emblematicus* Jordan and Rutter.**
Jamaica.
Scarus emblematicus Jordan and Rutter, Proc. Acad. Nat. Sci. Phila., XLIX, 1897, 122, Jamaica.
- Genus 1200. PSEUDOSCARUS Bleeker.**
Pseudoscarus Bleeker, Versl. Akad. Wet. Amsterdam, Scaroiden, XII, 1861, 230 (*P. microrrhinos* Bleeker)
Loro Jordan and Evermann, Check List, 1895, 418 (*Scarus guacamara* Cuvier).
- Subgenus PSEUDOSCARUS Bleeker.**
- 3490. *Pseudoscarus caelestinus* (Cuvier and Valenciennes). *Loro*.**
West Indies.
Scarus caelestinus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 134 (180), St. Thomas.
- 3491. *Pseudoscarus microps* (Osburn and Nichols)**
Gulf of California.
Callyodon microps Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXV, 1916, 170, fig. 12, Santa Catalina Island, Gulf of California.
- 3492. *Pseudoscarus simplex* Poey.**
Cuba.
Pseudoscarus simplex Poey, Repertorio, I, 1867, 185, Habana.
- 3493. *Pseudoscarus pleianus* (Poey).**
St. Thomas; only the type known.
Scarus pleianus Poey, Memorias, II, 1861, 393, St. Thomas, West Indies.
- 3494. *Pseudoscarus guacamara* (Cuvier). *Guacamara*, Green parrotfish.**
West Indies, north to Florida Keys, south to Rio de Janeiro; straying to Argentine.
Guacamara Parra, Descr. Dif. Piezas Hist. Nat., 1787, p. 54, pl. 26, Cuba.
Scarus guacamara Cuvier, Règne Anim., ed. 2, II, 1829, 235; after Parra.
Scarus turchesus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 134 (181), Porto Rico
Scarus rostratus Poey, Memorias, II, 1861, 221, Cuba
- 3495. *Pseudoscarus perrico* (Jordan and Gilbert) *Perrico*.**
Pacific coast of Mexico, from La Paz to Mazatlan
Scarus perrico Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 357, Mazatlan.
- 3496. *Pseudoscarus plumbeus* Bean.**
Bermudas.
Pseudoscarus plumbeus Bean, Proc. Biol. Soc. Wash., XXV, 1912, 125, Bermudas.

Order VV. GOBOIDEA

Family 255. ELEOTRIDÆ. Sleepers

- Genus 1201. GOBIOMORUS Lacépède.**
Gobiomorus Lacépède, Hist. Nat. Poiss., II, 1800, 583 (*G. dormitor* Lacépède); first restriction to this species.
Philypnus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 255 (*Platycephalus dormitor* Bloch).
Lembus Günther, Cat., I, 1859, 505 (*L. maculatus* Günther).
- 3497. *Gobiomorus dormitor* Lacépède. *Sleeper*; *Guavina*.**
Streams of the West Indies and the Atlantic shores of Central America, Mexico and Surinam.
Gobiomorus dormitor Lacépède, Hist. Nat. Poiss., II, 1800, 583, Martinique; from a drawing by Plumier.
Guavina Parra, Descr. Dif. Piezas Hist. Nat. Cuba, 1787, 105, lam. 39, fig. 1, Habana
Platycephalus dormitor Bloch and Schneider, Ichth., 1801, 60, Martinique.
Eleotris longiceps Günther, Proc. Zool. Soc. Lond., 1864, 151, Nicaragua.
Eleotris dormitrix Cuvier, Règne Anim., ed. 2, II, 1829, 246, Antilles.

- 3498. *Gobiomorus lateralis* (Gill).** *Aboma de mar*
Streams of Pacific coast of Mexico and Central America; from Sonora to Panama.
Phylpnus lateralis Gill, Proc Acad. Nat. Sci. Phila., XII, 1860, 123, Cape San Lucas.
- Genus 1202. DORMITATOR** Gill.
Prochilus Cuvier, Règne Anim., ed. 1, II, 1817, 294 (*Scæna maculata* Bloch); name preoccupied.
Dormitator Gill, Proc Acad Nat Sci. Phila., XIV, 1862, 240 (*D. gundlachi* Poey).
- 3499. *Dormitator maculatus* (Bloch).** *Guavina*; *Mapo*. *Paneca*; *Mapiro*; *Striped sleeper*.
Both coasts of tropical America, ranging from South Carolina through the West Indies to Para, Cape San Lucas, and Panama
Scæna maculata Bloch, Ichth., 1785, pl. 299, fig. 2, West Indies.
Eleotris mugiloides Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 170 (226), Martinique; Surinam
Eleotris sima Cuvier and Valenciennes, loc. cit., 174 (232), Vera Cruz
Eleotris latifrons Richardson, Voy Sulphur, Fishes, 1844, 57, pl. 35, figs. 4 and 5, locality unknown, supposed to be from Pacific coast of Central America.
Eleotris grandisquama Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 173 (229), America.
Eleotris somnolentus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169, near mouth of Rio Grande.
Eleotris omocyaneus Poey, Memorias, II, 1861, 269, Habana
Dormitator micropthalmus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 170, Panama.
Dormitator gundlachi Poey, Synopsis, 1868, 396, Cuba.
Dormitator lineatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 271, Savannah.
- Genus 1203. GUAVINA** Bleeker.
Guavina Bleeker, Esquisse d'un Syst. Nat. Gobiod, in Archiv Neerl. Sci., IX, 1874 (*Eleotris guavina* Cuvier and Valenciennes).
- 3500. *Guavina guavina* (Cuvier and Valenciennes).** *Guarubaco*; *Guavina*.
East coast of tropical America, Cuba to Rio Janeiro.
Eleotris guavina Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 168 (223) Martinique.
- Genus 1204. LEPTOPHYLPNUS** Meek and Hildebrand.
Leptophilypnus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 361 (*L. fluvialis* Meek and Hildebrand).
- 3501. *Leptophilypnus fluviatilis* Meek and Hildebrand.**
Fresh waters of Panama.
Leptophilypnus fluviatilis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 361, Mindi, Canal Zone, Panama.
- Genus 1205. ELEOTRIS** Schneider.
Eleotris Schneider, Syst. Ichth., ed. Bloch, 1801, 65 (*Gobius pisonis*, etc., Gmelin).
Culius Bleeker, Esquisse, Gobioides, 1874, 304 (*C. fuscus* Bleeker).
- 3502. *Eleotris pisonis* (Gmelin).** *Guavina*; *Telard*; *Sleeper*.
Streams of the West Indies; southern Florida to Brazil.
Gobius pisonis Gmelin, Syst. Nat., 1789, 1206, Rio Almendares, Cuba; based on *Eleotris* of Gronow.
Amore pixuma Marcgrave, Hist. Brasil., IV, 1648, 166, Brazil.
Eleotris capite plagioplateo Gronow, Mus. Ichth., II, 1757, 168; after Marcgrave
Gobius amorea Walbaum, Artedi Pisc., III, 1792, 205; based on *Eleotris* of Gronow.
Eleotris gyrinus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 166 (220), pl. 356, Martinique; San Domingo; Surinam.
Eleotris (Culius) belizanus Sauvage, Bull. Soc. Philom. Paris, ser. 7, IV, 1879, 55, Belize, Cayenne.

3503. *Eleotris perniger* (Cope).
West Indies, south to Rio Janeiro.
Culius perniger Cope, Trans Amer Phil. Soc., XIV, 1870, 473, St. Martins, West Indies
3504. *Eleotris amblyopsis* (Cope).
Atlantic coast of America, from Charleston, S C, to Surinam; streams of West Indies.
Culius amblyopsis Cope, Trans Amer. Phil. Soc., XIV, 1870, 473, Surinam
3505. *Eleotris abacurus* Jordan and Gilbert.
Coast of South Carolina
Eleotris abacurus Jordan and Gilbert, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 228, Charleston, S. C.
3506. *Eleotris pictus* Kner and Steindachner. *Guayana*.
Streams about the Gulf of California, from Sonora south to Panama; Rio Presidio, Mazatlan.
Eleotris pictus Kner and Steindachner, Abh. Akad. Wiss. Wien, X, 1864, 18, pl 3, fig 1, Rio Bayano, near Panama.
Culius xquidens Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 461, Rio Presidio, near Mazatlan.
3507. *Eleotris tubularis* Heller and Snodgrass.
Cocos Island.
Eleotris tubularis Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 210, Pl. X, Cocos Island.
- Genus 1206. *ALEXURUS* Jordan.
Alexurus Jordan, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 511 (*A. armiger* Jordan and Richardson); perhaps inseparable from *Eleotris*.
3508. *Alexurus armiger* Jordan.
La Paz, Lower California, to Panama.
Alexurus armiger Jordan, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 511, pl. 48, La Paz, Lower California.
- Genus 1207. *HEMIELEOTRIS* Meek and Hildebrand.
Hemieleotris Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 364 (*Eleotris latifasciatus* Meek and Hildebrand).
3509. *Hemieleotris latifasciatus* (Meek and Hildebrand).
Streams of Panama.
Eleotris latifasciatus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1912, 63, Rio Cardenis, Panama
- Genus 1208. *EROTELIS* Poey. *Esmeraldas de Mar*.
Erotelis Poey, Memorias, II, 1861, 273 (*E. valenciennesi* Poey = *Eleotris smaragdus* Cuvier and Valenciennes).
3510. *Erotelis smaragdus* (Cuvier and Valenciennes). *Esmeralda negra*; *Esmeralda de mar*.
Key West and Cuba; coral shores among algae.
Eleotris smaragdus Cuvier and Valenciennes. Hist. Nat. Poiss., XII, 1837, 173 (231), Cuba.
Erotelis valenciennesi Poey, Memorias, II, 1861, 273, Cuba.
- Genus 1209. *MICROELEOTRIS* Meek and Hildebrand.
Microeleotris Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 362 (*M. panamensis* Meek and Hildebrand).
3511. *Microeleotris panamensis* Meek and Hildebrand.
Pacific slope of central Panama.
Microeleotris panamensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 363, Rio Juan Diaz, Panama.
3512. *Microeleotris mindii* Meek and Hildebrand.
Mindi, near Atlantic coast of Canal Zone.
Microeleotris mindii Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 364, brackish creek at Mindi, Canal Zone.

Genus 1210. EVIOTA Jenkins.

Eviota Jenkins, Bull. U. S. Fish Com., XXII, 1902, 501 (*Eviota epiphanes* Jenkins).

3513. *Eviota personata* Jordan and Thompson.

Coral reefs about the Tortugas, Fla.; Garden Key.

Eviota personata Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904, 251, fig. 3, Garden Key, Fla.

Genus 1211. IOGLOSSUS Bean.

Ioglossus Bean, in Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 297 (*I. callurus* Bean).

3514. *Ioglossus calliurus* Bean.

Gulf of Mexico; Snapper Banks off Pensacola, Fla.

Ioglossus calliurus Bean, in Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 297, Pensacola, Fla.

Genus 1212. GYMNELEOTRIS Bleeker.

Gymneleotris Bleeker, Esquisse d'un Syst. Nat. des Gobiod., in Archiv Néerl. Sci., IX, 1874, 304 (*Eleotris seminuda* Gunther).

3515. *Gymneleotris seminuda* (Günther).

Panama.

Eleotris seminudus Günther, Proc. Zool. Soc. Lond., 1864, 24, pl. 4, figs 2, 2a, Panama.

Genus 1213. PYCNOMMA Rutter.

Pycnomma Rutter, Proc. Calif. Acad. Sci., 3d ser., III, 1904, 252 (*P. semisquamatum* Rutter).

3516. *Pycnomma semisquamatum* Rutter.

Gulf of California.

Pycnomma semisquamatum Rutter, Proc. Calif. Acad. Sci., 3d ser., III, 1904, 252, Pl. XXIV, Gulf of California.

Genus 1214. CHRIOLEPIS Gilbert.

Chriolepis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 557 (*C. minutillus* Gilbert).

3517. *Chriolepis minutillus* Gilbert

Gulf of California.

Chriolepis minutillus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 558, Gulf of California, at Albatross station 2825.

Family 256. GOBIDÆ. Gobies**Genus 1215. LOPHOGOBIOUS Gill. Crested gobies.**

Lophogobius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 240 (*Gobius cristagalli* Valenciennes, in Cuvier and Valenciennes=*Gobius cyprinoides* Pallas).

3518. *Lophogobius cyprinoides* (Pallas).

West Indies; Cuba.

Gobius cyprinoides Pallas, Spic., Zool., VIII, 1770, 17, pl. 1, fig. 5, "Amboina."

Gobius crista-galli Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 98 (130), Habana.

Genus 1216. EVORTHODUS Gill.

Evorthodus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 195 (*E. breviceps* Gill).

3519. *Evorthodus breviceps* Gill.

Trinidad and Surinam

Evorthodus breviceps Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 195, Trinidad.

Genus 1217. CTENOGOBIOUS Gill.

Ctenogobius Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 374, 430 (*C. fasciatus* Gill).

3520. *Ctenogobius fasciatus* Gill.

West Indies.

Ctenogobius fasciatus Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 376, Trinidad.

- Genus 1218. RHINOGOBIUS** ⁹⁶ Gill.
Rhinogobius Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 145 (*R. similis* Gill).
Coryphopterus Gill, loc. cit., XV, 1863, 263 (*C. glaucofrænum* Gill).
- 3521. *Rhinogobius glaucofrænum* (Gill).**
 Florida Keys; Tortugas.
Coryphopterus glaucofrænum Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 263, coast of the State of Washington; evidently an error.
- 3522. *Rhinogobius manglicola* (Jordan and Starks).**
 Pacific coast of Mexico at Mazatlan
Gobius manglicola Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 495, Mazatlan, Mexico.
- 3523. *Rhinogobius quadriporus* (Cuvier and Valenciennes).**
 Surinam.
Gobius quadriporus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 66 (87), Surinam.
- 3524. *Rhinogobius stigmaturus* (Goode and Bean).**
 Florida Keys.
Gobius stigmaturus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 418, no type locality given, but specimens probably from Florida Keys.
- 3525. *Rhinogobius shufeldti* (Jordan and Eigenmann).**
 Gulf coast of United States.
Gobius shufeldti Jordan and Eigenmann, Proc. U. S. Nat. Mus., IX, 1886, 495, New Orleans.
- 3526. *Rhinogobius boleosoma* (Jordan and Gilbert).**
 Gulf of Mexico, Pensacola to Key West.
Gobius boleosoma Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 295, Laguna Grande, Pensacola, Fla.
- 3527. *Rhinogobius encæomus* (Jordan and Gilbert).**
 South Carolina to Key West.
Gobius encæomus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 611, Charleston, S. C.
- 3528. *Rhinogobius eigenmanni* (Garman).**
 Off Key West.
Gobius eigenmanni Garman, Bull. Lab. Nat. Sci. State Univ. Iowa, IV, No. 1, 1896, 88, pl. 3, fig. 1, off Key West in 60 fathoms.
- 3529. *Rhinogobius curacao* (Metzelaar).**
 Caribbean Sea.
Gobius curacao Metzelaar, Trop. Atl. Viss., 1919, 136, fig. 38, Antilles.
- 3530. *Rhinogobius mowbrayi* Bean.**
 Bermudas to Caribbean Sea.
Rhinogobius mowbrayi Bean, Field Mus. Nat. Hist., Zool., VII, 1906, 81, Challenger Bank, 28 fathoms, rock cavity, Bermudas.
- 3531. *Rhinogobius tortugæ* (Jordan).**
 Dry Tortugas, Garden Key, West Indies.
Ctenogobius tortugæ Jordan, Bull. U. S. Fish Com., XXII, 1902, 541, pl. 1, Garden Key, Tortugas.
- 3532. *Rhinogobius parvus* (Meek).**
 Rio San Francisco, La Antigua, Mexico.
Gobius parvus Meek, Field Col. Mus., Zool., III, 1902, 121, La Antigua.
- 3533. *Rhinogobius claytonii* (Meek).**
 Rio San Francisco, La Antigua, Mexico.
Gobius claytonii Meek, Field Col. Mus., Zool., III, 1902, 121, La Antigua.
- 3534. *Rhinogobius garmani* (Eigenmann and Eigenmann).**
 Dominica; Fort de France; Martinique; St. Kitts.
Gobius garmani Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 61, Dominica; Fort de France, Martinique, and St. Kitts.

* Some of the species named below may belong to *Ctenogobius*, from which this genus differs but slight y

Genus 1219. GOBIUS Linnæus.*Gobius* Linnæus, Syst. Nat., Ed. X, I, 1758, 263 (*G. niger* Linnæus)**3535. *Gobius translucens* Nichols.**

Porto Rico.

Gobius translucens Nichols, Bull. Amer. Mus. Nat. Hist., XXXIV, 1915, 145, fig. 2, near San Antonio Bridge, San Juan, Porto Rico**3536. *Gobius hipoliti* Metzelaar.**

West Indies.

Gobius hipoliti Metzelaar, Bijdr. Dierk. Amsterdam, XXII, 1922, 138, with figure, West Indies.**Genus 1220. BATHYGOBIUS Bleeker. *Mapos*.***Bathygobius*⁸⁷ Bleeker, Arch. Neerl. Acad. Sci., XIII, 1878, 54 (*Gobius nebulo-punctatus* Rüppell = *Gobius fuscus* Rüppell).*Mapo* Smitt, Oefv. Kong. Vet. Akad. Stockholm, 1899, 542 (*Gobius soporator* Cuvier and Valenciennes).**3537. *Bathygobius soporator* (Cuvier and Valenciennes). *Sleeper*, *Mapo*; *Caïman*; *Oopu*.**

Barbados, Pará, Itapuaná, Cuba, Bahamas, Orange Key, Bahía, Pernambuco, St. Thomas, Charleston, Tortugas, Florida Keys, Martinique, Sao Mathias, Curaçao, Rio de Janeiro, Rio Doce

Gobius soporator Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 42 (56), Martinique.? *Gobius albopunctatus* Cuvier and Valenciennes, loc. cit., 43 (57), Isle of France.? *Gobius padangensis* Bleeker, Nat. Tijds. Ned. Ind., I, 1850, 249, Padang.*Gobius catulus* Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169, St. Joseph Island, Tex.? *Gobius breviceps* Blyth, Proc. Asia. Soc. Beng., 1858, 271, Great Andaman Island (in Jour., XXVII, 1858).*Gobius mapo* Poey, Memorias, II, 1861, 277, Cuba.*Gobius lacertus* Poey, loc. cit., 278, Cuba.*Gobius carolinensis* Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 268, Charleston, S. C.*Gobius brunneus* Poey, Synopsis, 1868, 393, Habana.**3538. *Bathygobius fuscus* (Rüppell).**

Red Sea to Galapagos Islands.

Gobius fuscus Rüppell, Atl. Reis. Nord. Af. Fish., 1828, 137, Red Sea.*Gobius punctillatus* Rüppell, loc. cit., 138, Red Sea.*Gobius nebulo-punctatus* Rüppell, Neue Wirbelthiere, Fische, 1835, 139, Red Sea.*Gobius lineatus* Jenyns, Zool. Voy. *Beagle*, 1842, 95, pl. 19, fig. 2, Galapagos Islands.*Gobius homocyanus* Vaillant and Sauvage, Rev. Mag. Zool., III, 1875, 280, Sandwich [Hawaiian] Islands*Gobius sandwicensis* Günther, Rept. Shore Fishes, *Challenger*, I, 1880, pt. 6, 60, Honolulu.**Genus 1221. RHINOGOBIOPS Hubbs.***Rhinogobiops* Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 169, Feb. 17, 1926, p. 1 (*Gobius nicholsii* Bean).**3539. *Rhinogobiops nicholsii* (Bean).**

Coast of British Columbia, south to southern California.

Gobius nicholsii Bean, Proc. U. S. Nat. Mus., IV, 1881, 469, Departure Bay, British Columbia.**Genus 1222. GOBIONELLUS Girard.***Gobionellus* Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 168 (*G. hastatus* Girard)*Smaragdus* Poey, Memorias, II, 1860, 280 (*Gobius smaragdus* Cuvier and Valenciennes).⁸⁷ An ill-chosen name for these shallow-water gobies.

3540. *Gobionellus oceanicus* (Pallas) *Esmeralda; Endoimi; Emerande; Bacalhao sabara*.
Gulf coast of United States, southward through the West Indies.
Gobius oceanicus Pallas, Spic., VIII, 1770, 4, locality unknown; after Gronow.
Gobius lanceolatus Bloch, Fische Deutschlands, II, 1783, 8, pl. 38, fig. 1, Martinique.
Gobius bacalau Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 90 (119), Surinam; Cayenne.
3541. *Gobionellus hastatus* Girard. *Emerald fish; Sharp-tailed goby*.
Gulf of Mexico.
Gobionellus hastatus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 168, St. Joseph Island, Tex.
3542. *Gobionellus smaragdus* (Cuvier and Valenciennes). *Esmeralda*.
West Indies, south to Rio Janeiro, north to St. Augustine, Fla., and to Charleston, N. C.
Gobius smaragdus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 91 (120), Cuba.
Smaragdus valenciennesi Poey, Memorias, II, 1861, 280, Cuba; after Valenciennes.
3543. *Gobionellus sagittula* (Günther)
Gulf of California, south to Panama.
Eucenogobius sagittula Gunther, Proc. Zool. Soc. Lond., 1861, pt. 3, 371, 372, Pacific coast of Central America
Gobius longicaudus Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 146, Guaymas.
3544. *Gobionellus strigatus* (O'Shaughnessy).
Coast of Surinam.
Eucenogobius strigatus O'Shaughnessy, Ann. Mag. Nat. Hist., ser. IV, XV, 1875, 145, Surinam.
Gobius kraussi Steindachner, Ichth. Beitr., VIII, 1879, 16, Surinam
3545. *Gobionellus bayamonensis* (Evermann and Marsh).
Porto Rico.
Gobius bayamonensis Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 355, San Juan Market, Porto Rico.
3546. *Gobionellus lyricus* (Girard).
Gulf of Mexico, from Galveston to Cuba and the Lesser Antilles.
Gobius lyricus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169, Brazos Santiago, Tex.
Gobius wurdemanni Girard, loc. cit., 169, Brazos Santiago.
Smaragdus costalesi Poey, Memorias, II, 1861, 280, Habana.
3547. *Gobionellus stigmaticus* (Poey).
Coast of North Carolina, Florida Keys, West Indies, southward to Rio Janeiro; common at Habana.
Smaragdus stigmaticus Poey, Memorias, II, 1861, 281, Cuba.
3548. *Gobionellus microdon* (Gilbert).
San Juan Lagoon, west coast of Mexico.
Gobius microdon Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 554, San Juan Lagoon, north of Rio Ahomè, Mexico.
- Genus 1223. *ZONOGOBIUS* Bleeker.
Zonogobius Bleeker, Esquisse Syst. Nat. Gobioiden, 1874, 323 (*Gobius semifasciatus* Kner).
3549. *Zonogobius zebra* (Gilbert).
Pacific coast of Mexico and southern California.
Gobius zebra Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 73, Albatross station 2989, Pacific coast of Mexico.
- Genus 1224. *EUCTENOGOBIOUS* Gill
Eucenogobius Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1859, VII, 45 (*E. badius* Gill).

3550. *Euctenogobius badius* Gill
Mouth of Amazon; recorded also from Martinique.
Euctenogobius badius Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1859, 45, Amazon.
Gobius bosci Sauvage, Bull. Soc. Philom. Paris, IV, 1880, 44, Martinique; not *Gobius bosci* Lacépède.
3551. *Euctenogobius poeyi* (Steindachner).
Barbados
Gobius poeyi Steindachner, Ichth. Notizen, VI, 1867, 44, Barbados.
- Genus 1225. **LYTHRYPNUS** Jordan and Evermann.
Lythrypnus Jordan and Evermann, Check List, 1895, 458 (*Gobius dalli* Gilbert).
3552. *Lythrypnus dalli* (Gilbert)
Catalina Harbor, Calif.
Gobius dalli Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 73, Avalon, Catalina Harbor, Calif., in 35 fathoms.
- Genus 1226. **XENOGOBIUS** Metzelaar.
Xenogobius Metzelaar, Trop. Atlant. Vissch., 1919, 140 (*X. weberi* Metzelaar).
3553. *Xenogobius weberi* Metzelaar
Caribbean Sea.
Xenogobius weberi Metzelaar, Trop. Atlant. Vissch., 1919, 140, fig. 40, Curaçao.
- Genus 1227. **ELACATINUS** Jordan.
Elacatinus Jordan, Bull. U. S. Fish Com., XXII, 1902, 542 (*E. oceanops* Jordan).
3554. *Elacatinus oceanops* Jordan.
Garden Key, Tortugas.
Elacatinus oceanops Jordan, Bull. U. S. Fish Com., XXII, 1902, 542, pl. 2, Garden Key, Fla.
- Genus 1228. **CHONOPHORUS** Poey.
Chonophorus Poey, Memorias, II, 1860, 274 (*C. bucculentus* Poey = *Gobius banana* Cuvier and Valenciennes; published in July, 1860, and therefore presumably older in date than *Awaous*)
Awaous Steindachner, Ver. Mat. Phrs., XLII, 1860, 289 (*Gobius ocellaris* Cuvier and Valenciennes; in an article presented on July 12, 1860).
3555. *Chonophorus banana* (Cuvier and Valenciennes). *Guavina hoyera*; *Aboma de rio*.
West Indies; both coasts of Mexico, south to Brazil; Cuba; Sinaloa; common; Lower California to Panama.
? *Gobius tarasica* Lichtenstein, Abh. Akad. Wiss. Berlin, 1822, 273, Brazil; not *Tajaasica* Macgregor.
Gobius banana Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 78 (103), San Domingo.
Gobius martinicus Cuvier and Valenciennes, loc. cit., 79 (105), Martinique.
Chonophorus bucculentus Poey, Memorias, II, 1860, 275, Cuba.
Chonophorus contractus Poey, loc. cit., Cuba.
Gobius dolichocephalus Cope, Trans. Amer. Phil. Soc. Phila., XIII, 1869, 403, near Orizaba, Mexico.
Euctenogobius latus O'Shaughnessy, Ann. Mag. Nat. Hist., ser. 4, XV, 1875, 146, Bahia.
3556. *Chonophorus mexicanus* (Günther).
Fresh-water streams of Atlantic slope of Mexico.
Gobius mexicanus Günther, Cat., III, 1861, 61, Mexico.
3557. *Chonophorus nelsoni* (Evermann).
Streams of Sinaloa.
Awaous nelsoni Evermann, Proc. Biol. Soc. Wash., XII, 1898, 3, fresh-water pools at Rosario, Sinaloa, Mexico.
- Genus 1229. **GNATHOLEPIS** Bleeker.
Gnatholepis Bleeker, Esquisse Syst. Nat. Gobioiden, 1874, 318 (*Gobius anjerensis* Bleeker).

3558. *Gnatholepis thompsoni* Jordan.
Bush Key, Tortugas.
Gnatholepis thompsoni Jordan, Bull. U. S. Fish Com., XXII, 1902, 541,
pl. 1, Bush Key, Tortugas.
- Genus 1230. *BOLLMANNIA* Jordan.
Bollmannia Jordan, Proc. U. S. Nat. Mus., XII, 1889, 164 (*B. chlamydes*
Jordan).
3559. *Bollmannia chlamydes* Jordan.
Pacific Ocean off coast of Colombia.
Bollmannia chlamydes Jordan, Proc. U. S. Nat. Mus., XII, 1889, 164,
Pacific Ocean off coast of Colombia.
3560. *Bollmannia ocellata* Gilbert.
Gulf of California.
Bollmannia ocellata Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 555,
northern part of Gulf of California at *Albatross* stations 3031 and 3035
in 30 and 33 fathoms.
3561. *Bollmannia macropoma* Gilbert.
Gulf of California.
Bollmannia macropoma Gilbert, Proc. U. S. Nat. Mus., XIV, 1891,
556, near La Paz, Gulf of California, at *Albatross* station 2996 in
112 fathoms.
3562. *Bollmannia stigmatura* Gilbert.
Northern part of the Gulf of California.
Bollmannia stigmatura Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 556,
Gulf of California at *Albatross* stations 3016 and 3017 in 76 and 58
fathoms.
3563. *Bollmannia boqueronensis* Evermann and Marsh.
Porto Rico.
Bollmannia boqueronensis Evermann and Marsh, Rept. U. S. Fish Com.,
XXV, 1899 (Dec. 19), 356, Ensenada del Boqueron, Porto Rico.
- Genus 1231. *ABOMA* Jordan and Starks.
Aboma Jordan and Starks, Fishes of Sinaloa, Proc. Calif. Acad. Sci.,
2d ser., V, 1895, 497 (*A. etheostoma* Jordan and Starks).
3564. *Aboma etheostoma* Jordan and Starks.
Pacific coast of Mexico.
Aboma etheostoma Jordan and Starks, Fishes of Sinaloa, Proc. Calif.
Acad. Sci., 2d ser., V, 1895, 498, pl. 50, Mazatlan, Mexico.
3565. *Abomia lucretiæ* (Eigenmann and Eigenmann).
Pearl Island, Gulf of Panama.
Gobius lucretiæ Eigenmann and Eigenmann, Proc. Calif. Acad. Sci.,
2d ser., I, 1888, 57, Pearl Island, Gulf of Panama.
3566. *Aboma chiquita* (Jenkins and Evermann).
Gulf of California.
Gobius chiquita Jenkins and Evermann, Proc. U. S. Nat. Mus., XI,
1888, 146, Guaymas, Sonora, Mexico.
- Genus 1232. *MICROGOBIUS* Poey.
Microgobius Poey, Enumeratio, 126, 1876 (*M. signatus* Poey).
Zalypnus Jordan and Evermann, Check List, 1895, 459 (*Gobius em-*
blematicus Jordan and Gilbert).
3567. *Microgobius emblematicus* (Jordan and Gilbert).
Lower California to Panama.
Gobius emblematicus Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881,
330, Bay of Panama.
Microgobius cyclolepis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 74,
Albatross station 3020, Lower California.
3568. *Microgobius signatus* Poey.
West Indies; common about Cuba.
Microgobius signatus Poey, Enumeratio, 1876, 127, pl. 5, fig. 3, Cuba.
3569. *Microgobius gulosus* (Girard).
Coast of Florida to Texas; common north to Indian River, Fla.
Gobius gulosus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169,
Indianola, Tex.

3570. *Microgobius meeki* Evermann and Marsh.
Culebra and Vieques Islands, Porto Rico.
Microgobius meeki Evermann and Marsh, Rept. U. S. Fish Com., XXV,
1899 (Dec. 19), 356, between Vieques and Culebra Islands east of
Porto Rico
3571. *Microgobius thalassinus* (Jordan and Gilbert)
South Atlantic coast, Virginia to South Carolina.
Gobius thalassinus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
612, Charleston Harbor, S. C.
Microgobius euleps Eigenmann and Eigenmann, Proc. Calif. Acad. Sci.,
2d ser., I, 1888, 69, Fortress Monroe, Va.
3572. *Microgobius miraflorensis* Gilbert and Starks
Bay of Panama.
Microgobius miraflorensis Gilbert and Starks, Mem. Calif. Acad. Sci.,
IV, 1904, 176, Pl. XXIX, Bay of Panama.
- Genus 1233. **EUCYCLOGOBIUS** Gill.
Eucyclogobius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 279 (*Gobius
newberryi* Girard).
3573. *Eucyclogobius newberryi* (Girard).
Streams of California, in small brooks near the sea; locally common in
San Luis Obispo Creek.
Gobius newberryi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 136,
Tomaes Bay, Calif.
- Genus 1234. **LEPIDOGOBIUS** Gill.
Lepidogobius Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1859, 14 (*Gobius
lepidus* Girard).
Cyclogobius Steindachner, Sitzb. Kais. Akad. Wiss. Wien, XLII, 1860,
284 (*Gobius lepidus* Girard).
3574. *Lepidogobius lepidus* (Girard).
Pacific coast of North America, from Vancouver Island to Lower
California, in rather deep water off San Francisco Bay.
Gobius lepidus Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 127,
25a, figs. 5 and 6; substitute for *Gobius gracilis*, preoccupied.
Gobius gracilis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 134, San
Francisco.
- Genus 1235. **GILLICHTHYS** Cooper.
Gillichthys Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 109 (*G.
mirabilis* Cooper).
Gillia Gunther, Zool. Rec., 1864, 137 (*Gillichthys mirabilis* Cooper);
substitute name preoccupied.
Aprolepis Hubbs, Occ. Papers, Univ. Mich. Mus. Zool., I, 1921, 99
(*A. barbaræ* Hubbs); young of *Gillichthys mirabilis*.
3575. *Gillichthys mirabilis* Cooper. *Long-jawed goby*.
Pacific coast of North America, from San Francisco to Guaymas.
Gillichthys mirabilis Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863,
109, San Diego Bay.
Gobius townsendi Eigenmann and Eigenmann, Proc. U. S. Nat. Mus.,
XI, 1888, 463, San Diego, Calif.
Aprolepis barbaræ Hubbs, Occ. Papers Univ. Mich. Mus. Zool., I, 1921,
99, young.
3576. *Gillichthys detrusus* Gilbert and Scofield.
Horseshoe Bend, near mouth of Colorado River, in Mexico.
Gillichthys detrusus Gilbert and Scofield, Proc. U. S. Nat. Mus., XX,
1896, 498, near mouth of Colorado River.
- Genus 1236. **QUIETULA** Jordan and Evermann.
Quietula Jordan and Evermann, in Jordan and Starks, Proc. Calif.
Acad. Sci., 2d ser., V, 1895, 839 (*Gillichthys y-cauda* Jenkins and
Evermann).
3577. *Quietula y-cauda* (Jenkins and Evermann).
Pacific coast of North America from Guaymas to Vancouver Island.
Gillichthys y-cauda Jenkins and Evermann, Proc. U. S. Nat. Mus., XI,
1888, 147, Guaymas, Sonora.
Gillichthys guaymasiæ Jenkins and Evermann, loc. cit., 148, Guaymas,
Sonora.

Genus 1237. CLEVELANDIA Eigenmann and Eigenmann.

Clevelandia Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 73 (*C. longipinnis* Eigenmann and Eigenmann = *Clevelandia rosæ* Jordan and Evermann)

3578. *Clevelandia ios* (Jordan and Gilbert).

Puget Sound and neighboring waters.

Gobiosoma ios Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 437, Saanich Arm, Vancouver Island.

Clevelandia longipinnis Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 73; not of Steindachner.

Clevelandia rosæ Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 229, San Diego, Calif.

Genus 1238. ILYPNUS Jordan and Evermann.

Ilypnus Jordan and Evermann, Check List, 1895, 460 (*Lepidogobius gilberti* Eigenmann and Eigenmann).

3579. *Ilypnus gilberti* (Eigenmann and Eigenmann).

San Diego Bay and southward; abundant at Magdalena Bay, Conception Bay, and St. George's Bay in the Gulf of California.

Lepidogobius gilberti Eigenmann and Eigenmann, Proc. U. S. Nat. Mus., XI, 1888, 464, San Diego Bay, Calif.

Genus 1239. EVERMANNIA Jordan.

Evermannia Jordan, Proc. Calif. Acad. Sci., 2d ser., IV, 1896, 592 (*Gobiosoma zosterurum* Jordan and Gilbert).

3580. *Evermannia zosterura* (Jordan and Gilbert)

Pacific coast of Mexico at Mazatlan, Mexico

Gobiosoma zosterurum Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 361, Mazatlan.

3581. *Evermannia longipinnis* (Steindachner).

Gulf of California.

Gobiosoma longipinne Steindachner, Ichth. Beitr., VIII, 1879, 27, Las Animas Island, Gulf of California.

3582. *Evermannia panamensis* Gilbert and Starks.

Panama Bay.

Evermannia panamensis Gilbert and Starks, Mem. Calif. Acad. Sci., VI, 1904, 179, Pl. XXX, Panama Bay.

Genus 1240. GARMANNIA Jordan and Evermann. *Half-naked gobies.*

Garmannia Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 496, pl. 49 (*Gobius paradoxus* Gunther).

3583. *Garmannia paradoxa* (Günther).

Pacific coast of Mexico, Mazatlan to Panama.

Gobius paradoxus Gunther, Cat., III, 1861, 549, Pacific coast of Central America.

3584. *Garmannia hemigymna* (Eigenmann and Eigenmann)

West Indies.

Gobius hemigymnus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 66, dredged in the West Indies.

3585. *Garmannia rubra* Rosen.

Bahamas.

Garmannia rubra Rosen, Arsekr. Univ. Lund, VII, 1912, 50, Bahamas.

Genus 1241. ENYPNIAS Jordan and Evermann.

Enypnias Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2233 (*Gobius seminudus* Günther).

3586. *Enypnias seminudus* (Günther).

Pacific coast of Central America.

Gobius seminudus Günther, Cat., III, 1861, 554, Pacific coast of Central America.

Genus 1242. GOBIOSOMA Girard.

Gobiosoma Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169 (*Gobius alepidotus* Bloch and Schneider = *Gobius bosci* Lacépède).

3587. *Gobiosoma bosci* (Lacépède). *Naked goby*; *Clinging goby*; *Goby*, *Variegated goby*; *Mud-creeper*; *Oysterfish*.
Atlantic coast of United States, Cape Cod to Florida.
Gobius bosci Lacépède, Hist. Nat. Poiss., II, 1798, 555, pl. 16, fig. 1, Charleston, S. C.
Gobius alepidotus Bloch and Schneider, Syst. Ichth., 1801, 547; after Lacépède.
Gobius viridipallidus Mitchill, Trans. Lit. and Phil. Soc. N. Y., I, 1814, 379, pl. 1, fig. 8, New York.
3588. *Gobiosoma molestum* Girard.
Gulf coast of the United States from Key West to Texas, and south to Bahia.
Gobiosoma molestum Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169, Indianola, Tex.
Gobiosoma alepidotum Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 297, Laguna Grande, Pensacola, Fla.
3589. *Gobiosoma histrio* Jordan.
Known only from the Gulf of California at Guaymas.
Gobiosoma histrio Jordan, Proc. U. S. Nat. Mus., VII, 1884, 260, Guaymas, Mexico.
3590. *Gobiosoma crescentale* Gilbert.
Gulf of California.
Gobiosoma crescentalis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 557, Gulf of California, at Albatross station 2825.
3591. *Gobiosoma multifasciatum* Steindachner
West Indies; known from Cuba, St. Thomas, and the Lesser Antilles.
Gobiosoma multifasciatum Steindachner, Ichth. Beitr., V, 1876, 183, Lesser Antilles.
Gobius lineatus Pöcy, Memorias, II, 1861, 424, Cuba; not of Jenyns.
3592. *Gobiosoma horsti* Metzelaar.
Curaçao, Caribbean Sea.
Gobiosoma horsti Metzelaar, Marine Fishes Lesser Antilles, 1923, Curaçao.
3593. *Gobiosoma longum* Nichols.
West Indies, north to Florida.
Gobiosoma longum Nichols, Bull. Amer. Mus. Nat. Hist., XXXIII, 1914, 143, fig. 1, Florida.
- Genus 1243. *BARBULIFER* Eigenmann and Eigenmann.
Barbulifer Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser. I, 1888, 70 (*B. papillosus* Eigenmann and Eigenmann = *Gobiosoma cauthœcum* Jordan and Gilbert).
3594. *Barbulifer cauthœcus* (Jordan and Gilbert).
About Key West, Fla.
Gobiosoma cauthœcum Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 29, Key West, Fla.
Barbulifer papillosus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 70, Key West, Fla.
- Genus 1244. *EVERMANNICHTHYS* Metzelaar.
Evermannichthys Metzelaar, Trop. Atl. Vissch., 1919, 139 (*E. spongicola* Metzelaar).
Radcliffei Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 99, 1921, 2 (*Garmannia spongicola* Radcliffe).
3595. *Evermannichthys spongicola* (Radcliffe).
Off shore banks of North Carolina and Florida in cavities of sponges.
Garmannia spongicola Radcliffe, Proc. U. S. Nat. Mus., LII, 1917, 423, off Cape Hatteras, N. C., and near Tarpon Springs, west coast of Florida.

3596. *Evermannichthys metzelaari* Hubbs.
Caribbean sea, living in sponges.
Evermannichthys spongicola Metzelaar, Trop. Atl. Vissch., 1919, 189, figs. 39 and 40, 139, Curaçao, name preoccupied by *Evermannichthys spongicola* (Radcliffe).
Evermannichthys metzelaari Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 144, Dec. 21, 1923, pl. 1, Curaçao; substitute for *E. spongicola* Metzelaar, preoccupied.
- Genus 1245. **LETHOPS** Hubbs.
Lethops Hubbs, Occ. Papers Univ. Mich. Mus. Zool., 1925 (*Lethops connectens* Hubbs).
3597. *Lethops connectens* Hubbs.
Coast of California in tide pools.
Lethops connectens Hubbs, Occ. Papers Mus. Zool. Univ. Mich., 1925, Carmel Bay; Point Firmin, near Los Angeles.
- Genus 1246. **TYPHLOGOBIUS** Steindachner. *Blind gobies*.
Typhlogobius Steindachner, Ichth. Beitr., VIII, 1879, 24 (*T. californiensis* Steindachner).
Othonops Rosa Smith (Mrs. Eigenmann), Proc. U. S. Nat. Mus., II, 1881, 19 (*O. eos* Smith = *Typhlogobius californiensis* Steindachner).
3598. *Typhlogobius californiensis* Steindachner. *Blind goby of Point Loma; Pinkfish*
Coast of Lower California, from San Diego southward to Cedros Island; especially common at Point Loma.
Typhlogobius californiensis Steindachner, Ichth. Beitr., VIII, 1879, 24, False Bay, San Diego, Calif.
Othonops eos Rosa Smith, Proc. U. S. Nat. Mus., II, 1881, 19, Point Loma, San Diego, Calif.
- Genus 1247. **SICYDIUM** Cuvier and Valenciennes.
Sicydium Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 126 (168) (*Gobius plumieri* Bloch).
3599. *Sicydium plumieri* (Bloch). *Sirago*.
Fresh waters of the West Indies.
Gobius plumieri Bloch, Ichth., 1786, 125, pl. 178, fig. 3, Martinique; on a drawing by Plumier.
Sicydium siragus Poey, Memorias, II, 1861, 278, Santiago de Cuba.
3600. *Sicydium caguitæ* Evermann and Marsh.
Porto Rico.
Sicydium caguitæ Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 355, Rio de Caguitas at Caguas, Porto Rico.
3601. *Sicydium vincente* Jordan and Evermann.
Kingston, St. Vincent Island; Haiti.
Sicydium vincente Jordan and Evermann, Check List, 1895, 45, St. Vincent Island.
3602. *Sicydium antillarum* Ogilvie-Grant.
Barbados.
Sicydium antillarum Ogilvie-Grant, Proc. Zool. Soc. Lond., 1884, 157, Barbados.
3603. *Sicydium punctatum* Perugia.
Martinique.
Sicydium punctatum Perugia, Annali del Museo Civico di Storia Naturale di Genova, ser. 2, XVI, 1896, 18, Martinique.
3604. *Sicydium buscki* Evermann and Clark.
Santo Domingo.
Sicydium buscki Evermann and Clark, Proc. U. S. Nat. Mus., XXX, 1906, 854, fig. 3, San Francisco Mountain, Santo Domingo.
- Genus 1248. **COTYLOPUS** Guichenot.
Cotylopus Guichenot, Notes sur l'Isle de la Réunion, II, 1862, annex b., 9 (*C. acutipinnis*).
Sicya Jordan and Evermann, Check List, 1895, 456, (*Sicydium gymnogaster* Ogilvie-Grant).

3605. *Cotylopus salvini* (Ogilvie-Grant).
Streams near Panama.
Sicydium salvini Ogilvie-Grant, Proc. Zool. Soc. Lond., 1884, 159, pl. 12, fig. 2, Panama.
3606. *Cotylopus cocoensis* Heller and Snodgrass
Chatham Bay, Cocos Island, northwest of Panama.
Cotylopus cocoensis Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 211, Pl. XI, Cocos Island.
3607. *Cotylopus punctatus* Regan.
Tequesixtlan, Mexico.
Cotylopus punctatus Regan, Ann. Mag. Nat. Hist., ser. 7, XVI, 1905, 362, Tequesixtlan, Mexico.

Family 257. GOBIOIDIDÆ

Genus 1249. GOBIOIDES Lacépède.

Gobioides Lacépède, Hist. Nat. Poiss., II, 1800, 580 (*G. broussonnetii* Lacépède).

3608. *Gobioides broussonnetii* Lacépède.

West Indies to Brazil; once taken near New Orleans.

Gobioides broussonnetii Lacépède, Hist. Nat. Poiss., II, 1800, 580, probably from Surinam; "given by Holland to France."

Gobius brasiliensis Bloch and Schneider, Syst. Ichth., 1801, 69, Brazil.

Gobius oblongus Bloch and Schneider, loc. cit., 548; based on Lacépède.

Gobioides barreto Poey, Memorias, II, 1861, 282, Cuba.

Amblyopsis mexicanus O'Shaughnessy, Ann. Mag. Nat. Hist., ser. 4, XV, 1875, 147, Mexico.

Genus 1250. TYNTLASTES Günther.

Tyntlastes Günther, Proc. Zool. Soc. Lond., 1862, 193 (*Amblyopus sagitta* Günther).

3609. *Tyntlastes sagitta* (Günther).

Coast of Lower California.

Amblyopus sagitta Günther, Proc. Zool. Soc. Lond., 1862, 193, "California"; probably from Lower California.

3610. *Tyntlastes brevis* (Günther).

Panama.

Amblyopus brevis Günther, Proc. Zool. Soc. Lond., 1864, 151, Panama.

Genus 1251. CAYENNIA Sauvage.

Cayennia Sauvage, Bull. Soc. Philom., ser. 7, IV, 1880, 57 (*C. guichenoti* Sauvage).

3611. *Cayennia guichenoti* Sauvage.

Cayenne, French Guiana.

Cayennia guichenoti Sauvage, Bull. Soc. Philom., ser. 7, IV, 1880, 57, Cayenne.

Order WW. DISCOCEPHALI

Family 258. ECHENEIDÆ. Remoras

Genus 1252. ECHENEIS Linnæus.

Echeneis Linnæus, Syst. Nat., Ed. X, 1758, 261 (*E. naucrates* Linnæus, as first restricted by Gill, 1862).

Leptecheneis Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 60 (*Echeneis naucrates* Linnæus).

3612. *Echeneis naucrates* Linnæus. *Sharksucker; Pega; Pegador; Suckingfish; White-tailed remora; Indian remora*

Warm seas, universally distributed; common north to Cape Cod and San Francisco.

Echeneis naucrates (misprint for *naucrates*) Linnæus, Syst. Nat., Ed. X, 1758, 261, Pelago Indico.

Echeneis albacauda Mitchill, Amer. Month. Mag., II, 1818, 244, New York.

Echeneis lunata Bancroft, Proc. Comm. Zool. Soc., I, 1830, 124, Kingston, Jamaica.

- Echeneis vittata* Lowe, Proc. Zool. Soc. Lond., 1839, 89, Madeira.
Echeneis guarcan Poey, Memorias, II, 1861, 248, Cuba.
Echeneis verticalis Poey, loc. cit., 253, Cuba.
Echeneis metallica Poey, loc. cit., 252, Cuba.
Echeneis fusca Gronow, Cat. Fish., ed. Gray, 1854, 92; after Linnæus.
Echeneis neutratorides Zuieww, Nova Acta Acad. Sci. Imp. Petropol., IV, 1789, 279, no locality.
Echeneis lineata Holbrook, Ichth. South Carolina, 1855, 101, Charleston, S. C.; not of Menzies.
Echeneis holbrookii Gunther, Cat., II, 1860, 382, Jamaica.

Genus 1253. REMILEGIA Gill.

- Remilegia* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 239 (*Echeneis scutata* Gunther).

3613. Remilegia australis (Bennett).

- Tropical seas; recorded by Doctor Lutken from 10° N., 39° W.
Echeneis australis Bennett, Nar Whaling Voyage, II, 1840, 273, pls. 24-26.
Echeneis scutata Gunther, Ann. Mag. Nat. Hist., 1860, 401, pl. 10, fig. B, Ceylon.

Genus 1254. REMORA Forster.

- Remora* Forster, Cat. Anim. N. Amer., 1770, 20 (*Echeneis remora* Linnæus).

3614. Remora remora (Linnæus) Remora; Sucker, Pega.

- Warm seas, north to New York and San Francisco.
Echeneis remora Linnæus, Syst. Nat., Ed. X, 1758, 260, Pelago Indico.
Echeneis squalipeta Daldorf, Skrivt. Naturrh. Selsk., II, 1793, 157, Atlantic Ocean between the tropics.
Echeneis jacobæa Lowe, Proc. Zool. Soc. Lond., 1839, 89, Madeira.
Echeneis remoroides Bleeker, Natur. Tijd. Ned. Ind. (Batæ, IX, 1855, 70, Batæ).
Echeneis parva Gronow, Cat. Fish., ed. Gray, 1854, 92, no locality; after Linnæus.
Echeneis postica Poey, Memorias, II, 1861, 255, Habana.

Genus 1255. REMORINA Jordan and Evermann.

- Remorina* Jordan and Evermann, Check List, 1895, 490 (*Echeneis albescens* Temminck and Schlegel).

3615. Remorina albescens (Temminck and Schlegel).

- Tropical Pacific, northward to San Francisco and Japan; common at Galapagos Islands.
Echeneis albescens Temminck and Schlegel, Fauna Japonica, Poiss., 1845, 272, pl. 120, fig. 4, Nagasaki, Japan.

Genus 1256. REMOROPSIS Gill.

- Remoropsis* Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 60 (*Echeneis brachyptera* Lowe).

3616. Remoropsis brachyptera (Lowe). Swordfish sucker.

- Warm seas; occasionally north to Cape Cod.
Echeneis brachyptera Lowe, Proc. Zool. Soc. Lond., 1839, 89, Madeira.
Echeneis sexdecim-lamellata Eydoux and Gervais, Voy. Favorite, V, 1839, 77, pl. 31, Indian Ocean.
Echeneis quatuordecimlamnatus Storer, Rept. Fishes Mass., 1839, 155, Holmes Hole.
Echeneis pallida Temminck and Schlegel, Fauna Japonica, Poiss., 1845, 271, pl. 120, fig. 2, 3, Japan.
Echeneis neuhofti Bleeker, Sumatra, Naturk. Tijd. Neder. Indie, IV, 1853, 279, Sumatra, Priaman.

Genus 1257. PHTHEIRICHTHYS Gill.

- Phtheirichthys* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 239 (*Echeneis lineata* Menzies).

3617. Phtheirichthys tropicus (Euphrasen).

- Tropical seas, ranging north to South Carolina and Pensacola.
Echeneis tropica Euphrasen, Nya. Handl., XII, 1791, 315, Atlantic between the tropics.
Echeneis apicalis Poey, Memorias, II, 1861, 254, Cuba.
?Echeneis sphyænarum Poey, loc. cit., 255, Cuba, on barracudas.

Genus 1258. RHOMBOCHIRUS Gill.

Rhombochirus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 88 (*Echeneis osteochir* Cuvier).

3618. *Rhombochirus osteochir* (Cuvier). *Spearfish sucker*.

West Indies, north to Cape Cod.

Echeneis osteochir Cuvier, Règne Animal, ed. 2, II, 1829, 348, no locality given.

Echeneis terapleurorum Poey, Memorias, II, 1861, 256, Cuba.

Order XX. JUGULARES

Family 259. DRACONETTIDÆ

Genus 1259. DRACONETTA Jordan and Fowler.

Draconetta Jordan and Fowler, Proc. U. S. Nat. Mus., XXV, 1903, 939 (*D. zenica* Jordan and Fowler).

3619. *Draconetta acanthopoma* Regan.

Depths of north Atlantic

Draconetta acanthopoma Regan, Ann. Mag. Nat. Hist., May, 1904; north Atlantic, in deep water.

Family 260. CALLIONYMIDÆ. Dragonets

Genus 1260. CALLIONYMUS Linnæus.

Callionymus Linnæus, Syst. Nat., Ed. X, 1758, 249, (*C. lyra* Linnæus).

3620. *Callionymus bairdi* Jordan.

Snapper Banks, Gulf of Mexico, between Pensacola and Tampa, Fla.

Callionymus bairdi Jordan, Proc., U. S. Nat. Mus., X, 1887, 501, Snapper Banks off Pensacola, Fla.

3621. *Callionymus agassizii* Goode and Bean.

West Indies off Barbados and Santa Cruz

Callionymus agassizii Goode and Bean, Three Cruises of the *Blake*, II, 1888, 29, fig., Gulf of Mexico.

Callionymus himantophorus Goode and Bean, Oceanic Ichth., 1895, 296, figs. 268 A and B, off Barbados in 209 fathoms.

3622. *Callionymus calliurus* Eigenmann and Eigenmann.

Key West, Fla.

Callionymus calliurus Eigenmann and Eigenmann, Proc., Calif. Acad. Sci., 2d ser., I, 1888, 76, South Beach, Key West, Fla.

3623. *Callionymus pauciradiatus* Gill.

Matanzas, Cuba.

Callionymus pauciradiatus Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1867, 143, Matanzas, Cuba.

3624. *Callionymus bermudarum* Barbour.

Castle Island, Bermudas.

Callionymus bermudarum Barbour, Bull. Mus. Comp. Zool., XLVI, 1905, No. 7, 129, Castle Island, Bermudas.

3625. *Callionymus boekei* Metzelaar.

Caribbean Sea.

Callionymus boekei Metzelaar, Trop. Atl. Vissch., 1916, 149, fig. 47, Curaçao.

3626. *Callionymus sancti-eustatii* Metzelaar.

Callionymus sancti-eustatii Metzelaar, Trop. Atl. Vissch., 1916, 150, fig. 47, St. Eustatius.

3627. *Callionymus atrilabiatus* Garman.

Off Bay of Panama in deep water.

Callionymus atrilabiatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 122, off Bay of Panama.

Family 261. PTEROPSARIDÆ

Genus 1261. HYPsicOMETES Goode.

Hypsicometes Goode, Proc. U. S. Nat. Mus., III, 1880, 347 (*H. gobioides* Goode).

3628. *Hypsicometes gobioides* Goode.

Gulf Stream.

Hypsicometes gobioides Goode, Proc. U. S. Nat. Mus., III, 1880, 348, Gulf Stream at Albatross station 871, 40° 2' 54" N., 70° 23' 40" W., in 115 fathoms.

Family 262. AMMODYTIDÆ. Sand lances or sand launces

Genus 1262. AMMODYTES Linnæus. *Sand launces*; *Sand lances*.

Ammodytes Linnæus, Syst. Nat., Ed. X, 1758, 247 (*A. tobianus* Linnæus).

Argyrotaenia Gill, Fishes East Coast U. S., Suppl. to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 40 (*Ammodytes vittatus* DeKay; probably a deformed example).

3629. *Ammodytes americanus* DeKay. *Sand lance*; *Sand lance*; *Sand eel*;

Lant.

Newfoundland to Cape May and Cape Hatteras; abundant on sandy shores.

Ammodytes americanus DeKay, New York Fauna: Fishes, 1842, 317, Stratford, Conn.

Ammodytes vittatus DeKay, loc. cit., 318, pl. 60, fig. 197, New York.

3630. *Ammodytes dubius* Reinhardt.

Greenland to Cape Cod.

Ammodytes dubius Reinhardt, Dansk. Vidensk. Selsk. Afhandl., VII, 1838, 131, Greenland.

3631. *Ammodytes personatus* Girard. *Sand lance*; *Sand lance*.

Japan to Alaska, south to Monterey, Calif.

Ammodytes personatus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 137, Cape Flattery, Wash.

3632. *Ammodytes alascanus* Cope *Alaska sand lance*

North Pacific coast of North America, Sitka to Aleutian Islands.

Ammodytes alascanus Cope, Proc. Am. Phil. Soc., XIII, 1873, 30, Sitka, Alaska.

Family 263. BLEEKERIIDÆ

Genus 1263. RHYNCHIAS Gill.

Rhynchias Gill, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2841 (*Ammodytes septipinnis* Pallas); species and genus doubtful, not seen by recent authors.

3633. *Rhynchias septipinnis* (Pallas).

Kamchatka.

Ammodytes septipinnis Pallas, Rosso-Asiat., III, 1811, 227, Tab., XLVIII, fig. 3, Kamchatka.

Family 264. BATHYMASTERIDÆ. Ronquils

Genus 1264. BATHYMASTER Cope.

Bathymaster Cope, Proc. Amer. Phil. Soc., XIII, 1873, 31 (*B. signatus* Cope).

3634. *Bathymaster signatus* Cope.

Shores of Alaska.

Bathymaster signatus Cope, Proc. Amer. Phil. Soc. XIII, 1873, 31, near Sitka, Alaska.

3635. *Bathymaster cæruleofasciatus* Gilbert and Burke.

Southern Bering Sea.

Bathymaster cæruleofasciatus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 84, fig. 29, Agattu Island, Alaska.

Genus 1265. *RONQUILUS* Jordan and Starks.

Ronquilus Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 838 (*Bathymaster jordanii* Gilbert).

3636. *Ronquilus jordanii* (Gilbert). *Ronquil*.

Alaska to Puget Sound.

Rathbunella jordanii Gilbert, Proc. U. S. Nat. Mus., XI, 1888, 554, Elliott Bay at Seattle; Alaska.

Genus 1266. *RATHBUNELLA* Jordan and Evermann.

Rathbunella Jordan and Evermann, Check List, 1895, 463 (*Bathymaster hypoplectus* Gilbert).

3637. *Rathbunella hypoplecta* (Gilbert).

Coast of California about the Santa Barbara Islands.

Bathymaster hypoplectus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 97, coast of California, south of Point Conception, at Albatross station 2944.

3638. *Rathbunella alleni* Gilbert.

Monterey Bay, Calif.

Rathbunella alleni Gilbert, Proc. Calif. Acad. Sci., 3d ser., III, 1904, 267, Monterey Bay, Calif.

Family 265. *ZAPRORIDÆ*Genus 1267. *ZAPRORA* Jordan.

Zaprora Jordan, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 202 (*Z. silenus* Jordan).

3639. *Zaprora silenus* Jordan.

East coast of Vancouver Island; only four specimens known.

Zaprora silenus Jordan, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 203, pl. 20, Nanaimo, Vancouver Island.

Family 266. *CHIASMODONTIDÆ*. Black swallows.Genus 1268. *CHIASMODON* Johnson.

Chiasmodon Johnson, Proc. Zool. Soc. Lond., 1863, 408 (*C. niger* Johnson).

Chiasmodus Gunther, altered spelling

3640. *Chiasmodon niger* Johnson. *Black swallower*.

Deep waters of the Atlantic; taken at Madeira and off Massachusetts coast.

Chiasmodon niger Johnson, Proc. Zool. Soc. Lond., 1863, 408, Madeira.

3641. *Chiasmodon subniger* Garman.

Pacific coast of Mexico, off Lower California, in deep water.

Chiasmodon subniger Garman, Mem., Mus. Comp. Zool., XXIV, 1899, 73, off Gulf of California at Albatross station 3429.

Genus 1269. *PSEUDOSCOPELUS* Lütken.

Pseudoscopus Lütken, Spolia Atlantica, Scopelini, 64, in Vid. Selsk. Skr. (6), VII, 1892, 285 (*P. scriptus* Lütken).

3642. *Pseudoscopus scriptus* Lütken.

Old Bahama Straits, in deep water.

Pseudoscopus scriptus Lütken, Spolia Atlantica, Scopelini, 64, in Vid. Selsk. Skr. (6), VII, 1892, 285, Old Bahama Straits.

Family 267. *OPISTHOGNATHIDÆ*. JawfishesGenus 1270. *OPISTHOGNATHUS* Cuvier.

Opisthognathus Cuvier, Règne Anim., Ed. I, 1817, 252 (*O. sonnerati* Cuvier).

3643. *Opisthognathus macrognathus* Poey.

Florida Keys to Cuba.

Opisthognathus macrognathus Poey, Memorias, II, 1861, 284, Cuba.

Opisthognathus megastoma Gunther, Cat., I, 1860, 255, Gulf of Mexico.

Opisthognathus scaphiurus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 417, Garden Key, Fla.

3644. *Opisthognathus lonchurum* Jordan and Gilbert.
Gulf of Mexico, in deep water; two specimens known, taken from the stomach of a red snapper at Pensacola, Fla.
Opisthognathus lonchurus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 290, Snapper Banks off Pensacola, Fla.
3645. *Opisthognathus punctatum* Peters.
West coast of Mexico; Mazatlan; Magdalena Bay.
Opisthognathus punctatus Peters, Berl. Monatsber., 1869, 708, Mazatlan, Mexico.
3646. *Opisthognathus ommatum* Jenkins and Evermann
Bay of Guaymas, Gulf of California.
Opisthognathus ommata Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 153, Guaymas, Sonora.
- Genus 1271. GNATHYPOPS^{ss} Gill.
Gnathypops Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 241 (*Opisthognathus maxillosus* Poey).
3647. *Gnathypops maxillosa* (Poey).
Cuba, north to Garden Key, Fla.
Opisthognathus maxillosus Poey, Memorias, II, 1861, 286, Cuba.
3648. *Gnathypops macrops* (Poey).
Coast of Cuba.
Opisthognathus macrops Poey, Memorias, II, 1861, 287, coast of Cuba.
3649. *Gnathypops cuvieri* (Cuvier and Valenciennes).
Bahia.
Opisthognathus cuvieri Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 504, Bahia (may be identical with *macrops*).
3650. *Gnathypops rhomalea* (Jordan and Gilbert).
Gulf of California.
Opisthognathus rhomaleus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 276, Santa Maria Cove, Lower California.
3651. *Gnathypops scopis* Jenkins and Evermann.
Gulf of California at Guaymas.
Gnathypops scopis Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 152, Guaymas, Sonora, Mexico.
3652. *Gnathypops snyderi* Jordan and Evermann.
Gulf of California.
Gnathypops snyderi Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2285, San Luis Gonzales Bay, Gulf of California.
3653. *Gnathypops mystacina* Jordan.
Gulf of Mexico; known from the Pensacola "Snapper Banks," from the stomach of a red snapper (*Neomænis blackfordi*).
Gnathypops mystacinus Jordan, Proc. U. S. Nat. Mus., VII, 1884, 37, Snapper Banks off Pensacola, Fla.
3654. *Gnathypops aurifrons* Jordan and Thompson.
Garden Key, Fla.
Gnathypops aurifrons Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904, 252, fig. 4, Garden Key, Fla.
- Genus 1272. LONCHOPISTHUS Gill.
Lonchopisthus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 241 (*Opisthognathus micrognathus* Poey).
3655. *Lonchopisthus micrognathus* (Poey).
Cuba.
Opisthognathus micrognathus Poey, Memorias, II, 1861, 287, Cuba.

^{ss} It is possible that some or all of the short-jawed forms are the females of species of *Opisthognathus*

Family 268. URANOSCOPIDÆ. Stargazers

Genus 1273. *ASTROSCOPUS* Brevoort. *Stargazers*.*Astroscoptes* Brevoort, in Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 20 (*Uranoscoptes anoplos* Cuvier and Valenciennes = *Uranoscoptes y-græcum* Cuvier and Valenciennes).

3656. *Astroscoptes y-græcum* (Cuvier and Valenciennes). *Stargazer*.
South Atlantic coast, from Cape Hatteras to the Caribbean Sea.
Uranoscoptes y-græcum Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 229 (308), locality unknown.
Uranoscoptes anoplos Cuvier and Valenciennes, loc. cit., VIII, 1831, 363 (493), Atlantic coast of North America.
3657. *Astroscoptes guttatus* Abbot. *Spotted stargazer*.
Atlantic coast of the United States, from Long Island to Virginia; not known south of Cape Hatteras.
Astroscoptes guttatus Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 365, pl. 7, Cape May, N. J.
3658. *Astroscoptes zephyreus* Gilbert and Starks.
Magdalena Bay, Lower California.
Astroscoptes zephyreus Jordan and Starks, Proc. U. S. Nat. Mus., XIX, 1896, 453, Magdalena Bay, Lower California.

Genus 1274. *KATHETOSTOMA* Günther.*Kathetostoma* Günther, Cat., II, 1860, 231 (*Uranoscoptes lævis* Bloch and Schneider).

3659. *Kathetostoma averruncus* Jordan and Bollman.
Pacific Ocean off coast of Colombia, and Bay of Panama, in deep water.
Kathetostoma averruncus Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 163, off coast of Colombia, 8° 57' N, 79° 31' 30" W., at *Albatross* station 2800 in 7 fathoms.
3660. *Kathetostoma albigutta* (Bean).
Gulf of Mexico.
Kathetostoma albigutta Bean, Proc. U. S. Nat. Mus., XVIII, 1892, 121, Gulf of Mexico, at *Albatross* station 2403, 28° 42' 30" N., 85° 29' W.
- Genus 1275. *EXCESTIDES* Jordan and Thompson.
Excestides Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904, 253, (*Excestides egregius* Jordan and Thompson).
3661. *Excestides egregius* Jordan and Thompson.
Garden Key, Fla.
Excestides egregius Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904, 253, fig. 5, Garden Key, Fla.

Family 269. DACTYLOSCOPIDÆ. Sand stargazers

Genus 1276. *DACTYLOSCOPUS* Gill.*Dactyloscopus* Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 132 (*D. tridigitatus* Gill).*Congrammus* Fowler, loc. cit., LVIII, 1906, 105 (*C. moorei* Fowler).

3662. *Dactyloscopus tridigitatus* Gill. *Sand conger*.
West Indies, north to Key West.
Dactyloscopus tridigitatus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 132, Barbados.
Congrammus moorei Fowler, loc. cit., LVIII, 1906, 105, Hailer's Rock, Florida Keys.
3663. *Dactyloscopus pectoralis* Gill.
Cape San Lucas, Lower California.
Dactyloscopus pectoralis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 267, Cape San Lucas, Lower California.
3664. *Dactyloscopus cinctus* Osburn and Nichols.
Gulf of California.
Dactyloscopus cinctus Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 176, fig. 14, Cape San Lucas.

- 3665. *Dactyloscopus poeyi* Gill.**
Cuba
Dactyloscopus poeyi Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 266, Cuba.
- 3666. *Dactyloscopus lunaticus* Gilbert.**
Gulf of California.
Dactyloscopus lunaticus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 99, Gulf of California.
- Genus 1277. GILLELLUS Gilbert.**
Gillellus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 98 (*G. semicinctus* Gilbert).
- 3667. *Gillellus semicinctus* Gilbert.**
Gulf of California; also off the Florida coast.
Gillellus semicinctus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 98, near north end of Gulf of California, at *Albatross* stations 2827, 24° 11' 45" N., 109° 55' W., and 2829, 22° 52' N., 109° 55' W., in 10 and 31 fathoms.
- 3668. *Gillellus arenicola* Gilbert.**
Cape San Lucas, Lower California.
Gillellus arenicola Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 99, Cape San Lucas
- 3669. *Gillellus ornatus* Gilbert.**
Gulf of California.
Gillellus ornatus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 558, Gulf of California.
- Genus 1278. ESLOSCOPUS Jordan and Evermann.**
Esloscopus Jordan and Evermann, Check List, 1895, 465 (*Dactyloscopus zelotes* Jordan and Gilbert).
- 3670. *Esloscopus zelotes* (Jordan and Gilbert).**
Panama.
Dactyloscopus zelotes Jordan and Gilbert, in Jordan and Evermann, Check List, 1895, 464, Panama.
- Genus 1279. DACTYLAGNUS Gill.**
Dactylagnus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 505 (*D. mundus* Gill).
- 3671. *Dactylagnus mundus* Gill.**
Carmen Island, Gulf of California.
Dactylagnus mundus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 505, Carmen Island, Gulf of California.
- Genus 1280. MYXODAGNUS Gill.**
Myxodagnus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 269 (*M. opercularis* Gill).
- 3672. *Myxodagnus opercularis* Gill.**
Cape San Lucas, Lower California.
Myxodagnus opercularis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 270, Cape San Lucas, Lower California.

Family 270. CLINIDÆ

- Genus 1281. MNIERPES Jordan and Evermann.**
Mnierpes Jordan and Evermann, Check List, 1895, 468 (*Clinus macrocephalus* Günther).
- 3673. *Mnierpes macrocephalus* (Günther).**
Pacific coast of Central America; Panama.
Clinus macrocephalus Günther, Cat., III, 1861, 266; Pacific coast of Central America.
- Genus 1282. GOBIOCLINUS Gill.**
Gobioclinus Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 103 (*Clinus gobio* Cuvier and Valenciennes).

3674. *Gobioclinus gobio* (Cuvier and Valenciennes).
Lesser Antilles.
Clinus gobio Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 292 (395), Lesser Antilles.
- Genus 1283. **HISTIOCLINUS** Metzelaar.
Histioclinus Metzelaar, Trop. Atl. Vissch., 1919, 157 (*H. veliger* Metzelaar).
3675. *Histioclinus veliger* Metzelaar.
Caribbean Sea.
Histioclinus veliger Metzelaar, Trop. Atl. Vissch., 1919, 157, Bonaire.
- Genus 1284. **STARKSIA** Jordan and Evermann.
Starksia Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 231, (*Labrosomus cremnobates* Gilbert).
3676. *Starksia cremnobates* (Gilbert).
Gulf of California.
Labrosomus cremnobates Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 100, Gulf of California.
- Genus 1285. **ALLOCLINUS** Hubbs.
Alloclinus Hubbs, Papers, Mich. Acad. Sci. Arts and Letters, VII, 1926 (1927), 351 (*Starksia holderi* Lauderbach).
3677. *Alloclinus holderi* (Lauderbach)
Santa Catalina, Calif.
Starksia holderi Lauderbach, Proc. U. S. Nat. Mus., XXXII, 1907, 74, fig. 6, Santa Catalina Island.
- Genus 1286. **CRYPTOTREMA** Gilbert.
Cryptotrema Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 101 (*C. corallinum* Gilbert).
3678. *Chryptotrema corallinum* Gilbert
Coast of California, south of Point Conception.
Cryptotrema corallinum Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 101, coast of California, south of Point Conception.
- Genus 1287. **EXERPES** Jordan and Evermann.
Ezerpes Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 232 (*Auchenopterus asper* Jenkins and Evermann).
3679. *Ezerpes asper* (Jenkins and Evermann).
Guaymas, Mexico.
Auchenopterus asper Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 154, Guaymas, Mexico.
- Genus 1288. **CORALLICOLA** Jordan and Evermann.
Corallicola Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2369 (*Cremnobates marmoratus* Steindachner).
3680. *Corallicola marmoratum* (Steindachner). *Rockfish*.
Florida Keys to Cuba; Key West.
Cremnobates marmoratus Steindachner, Ichth. Beitr., V, 1876, 174, pl. 12, fig. 6, near Cuba.
3681. *Corallicola nigripinne* (Steindachner).
Barbados.
Clinus nigripinnis Steindachner, Ichth. Notizen, VI, 1867, 45, Barbados.
3682. *Corallicola altivelis* (Lockington).
La Paz, Lower California.
Cremnobates altivelis Lockington, Proc. Acad. Nat. Sci. Phila., XXXIII, 1881, 116, La Paz, Lower California.
- Genus 1289. **AUCHENOPTERUS** Günther.
Auchenopterus Günther, Cat., III, 1861, 275 (*A. monophthalmus* Günther; not *Auchenipterus* Cuvier and Valenciennes).
Cremnobates Günther, Proc. Zool. Soc. Lond., 1861, 374 (same type; substitute name).
Tekla Nichols, Copeia, No. 110, 1922, 69 (*Cremnobates fasciatus* Steindachner).

3683. *Auchenopterus monophthalmus* Gunther.
Gulf of California to Panama.
Auchenopterus monophthalmus Gunther, Cat., III, 1861, 275, Panama.
3684. *Auchenopterus integripinnis* (Rosa Smith).
Coast of California and southward to Todos Santos, Lower California.
Cremnobates integripinnis Rosa Smith, Proc. U. S. Nat. Mus., III, 1880, 147, La Jolla, near San Diego.
3685. *Auchenopterus mexicanus* Gilbert
La Paz to Mazatlan, Mexico.
Auchenopterus mexicanus Gilbert, Proc. Calif. Acad. Sci., 3d ser., III, 1904, 269, Pl. XXIX, La Paz, Lower California.
3686. *Auchenopterus affinis* (Steindachner).
West Indies; Key West; St. Thomas.
Cremnobates affinis Steindachner, Ichth. Beitr., V, 1876, 178, St. Thomas.
3687. *Auchenopterus fasciatus* (Steindachner)
Florida Straits; north to Key West; Porto Rico
Cremnobates fasciatus Steindachner, Ichth. Beitr., V, 1876, 176, Florida Straits.
3688. *Auchenopterus nox* (Jordan and Gilbert).
Key West, Fla.
Cremnobates nox Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 30, Key West.
3689. *Auchenopterus albicaudus* Evermann and Marsh.
Porto Rico.
Auchenopterus albicaudus Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 360, Arroyo, Porto Rico.
3690. *Auchenopterus fajardo* Evermann and Marsh.
Porto Rico.
Auchenopterus fajardo Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 361, Fajardo, Porto Rico.
3691. *Auchenopterus rubescens* Evermann and Marsh.
Porto Rico.
Auchenopterus rubescens Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 360, Puerto Real, Porto Rico.
3692. *Auchenopterus cingulatus* Evermann and Marsh.
Porto Rico.
Auchenopterus cingulatus Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 361, Ponce, Porto Rico,
3693. *Auchenopterus grandicomus* Rosen.
Bahamas.
Auchenopterus grandicomus Rosen, Fauna Bahamas, 1911, 69, Andros Island.
- Genus 1290. *AUCHENISTIUS* Evermann and Marsh.
Auchenistius Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 359 (*A. stahli* Evermann and Marsh).
3694. *Auchenistius stahli* Evermann and Marsh.
Porto Rico.
Auchenistius stahli Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 359, Ponce, Porto Rico.
- Genus 1291. *PARACLINUS* Mocquard.
Acanthoclinus Mocquard, Bull. Soc. Philom. Paris, X, 1885, 18 (*A. chaperi* Mocquard); name preoccupied.
Paraclinus Mocquard, loc. cit., I, 1888, 41 (*Acanthoclinus chaperi* Mocquard).
3695. *Paraclinus chaperi* (Mocquard).
Bay of Guanta, near Barcelona, in Venezuela.
Acanthoclinus chaperi Mocquard, Bull. Soc. Philom. Paris, X, 1885, 18, Bay of Guanta, Venezuela.

- Genus 1292. **ENNEAPTERYGIUS** Rüppell.
Enneapterygius Rüppell, Neue Wirbelthiere, Fische, 1835, 2 (*E. pusillus* Rüppell).
Enneanectes Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 501 (*Tripterygium carminale* Jordan and Gilbert).
3696. **Enneapterygius carminalis** (Jordan and Gilbert).
 Mazatlan, Mexico, and Caribbean Sea.
Tripterygium carminale Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 362, Mazatlan, Mexico.
- Genus 1293. **GILLIAS** Evermann and Marsh.
Gillias Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 357 (*G. jordani* Evermann and Marsh).
3697. **Gillias jordani** Evermann and Marsh.
 Porto Rico.
Gillias jordani Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899, (Dec. 19), 357, Cardona Lighthouse Reef, Ponce, Porto Rico.
- Genus 1294. **GIBBONSIA** Cooper.
Gibbonsia Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 109 (*Myxodes elegans* Cooper).
Blakea Steindachner, Ichth. Beitr., V, 1876, 148 (*Myxodes elegans* Cooper).
3698. **Gibbonsia elegans** (Cooper). *Spotted kelpfish; Señorita*.
 Coast of southern California; Point Conception to Todos Santos.
Myxodes elegans Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 109, San Diego; Santa Barbara.
Clinus ocellifer Mocquard, Bull. Soc. Philom. Paris, 1886, 44, California.
Clinus evdes Jordan and Gilbert, Synopsis, 1883, 763, Monterey, Calif.
3699. **Gibbonsia elegans montereyensis** Hubbs.
 Coast of southern California.
Gibbonsia elegans montereyensis Hubbs, Papers Mich. Acad. Sci. Arts and Letters, VII, 1926, 354, reef at Pacific Grove, Calif.
3700. **Gibbonsia metzi** Hubbs.
 California coast, chiefly in tidepools.
Gibbonsia metzi Hubbs, Papers Mich. Acad. Sci. Arts and Letters, VII, 1926, 360 Pacific Grove, Calif.
- Genus 1295. **HETEROSTICHUS** Girard.
Heterostichus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 143 (*H. rostratus* Girard).
3701. **Heterostichus rostratus** Girard. *Kelpfish*.
 Coast of California from San Francisco to San Diego.
Heterostichus rostratus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 143, San Diego, Calif.
- Genus 1296. **NEOCLINUS** Girard.
Neoclinus Girard, U. S. Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 114 (*N. blanchardi* Girard).
3702. **Neoclinus blanchardi** Girard.
 Coast of California, Monterey to San Diego.
Neoclinus blanchardi Girard, U. S. Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 114, off San Diego, Calif.
- Genus 1297. **PTEROGNATHUS** Jordan and Evermann.
Pterognathus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 57 (*Neoclinus satiricus* Girard); name only suggested but not adopted.
Pterognathus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2354 (*Neoclinus satiricus* Girard).
3703. **Pterognathus satiricus** (Girard).
 Coast of California, from Monterey to Santa Barbara.
Neoclinus satiricus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 57, Monterey Bay.

Genus 1298. LABRISOMUS Swainson.

- Labrisomus* Swainson, Nat. Hist. Classn. Fishes, II, 1839, 277 (*Clinus pectinifer* Cuvier and Valenciennes=*Clinus nuchipinnis* Quoy and Gaimard).
- Lepisoma* DeKay, N. Y. Fauna: Fishes, 1842, 41, *L. cirrhosum* DeKay=*Clinus nuchipinnis* Quoy and Gaimard).
- 3704. *Labrisomus nuchipinnis* (Quoy and Gaimard).**
West Indies; South Atlantic coast.
Clinus nuchipinnis Quoy and Gaimard, Voy. *Uranié*, Zool., 1824, 255, Bay of Rio de Janeiro.
Clinus pectinifer Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 276 (374), Bahia.
Clinus capillatus Cuvier and Valenciennes, loc. cit., 278 (377), Martinique.
Clinus canariensis Valenciennes, in Webb and Berthelot, Poiss. Îles Canaries, 1839, 60, Pl. XVII, fig. 3, Canary Islands.
Lepisoma cirrhosum DeKay, N. Y. Fauna: Fishes, 41, 1842, Florida.
- 3705. *Labrisomus heilneri* Nichols.**
Bahamas.
Labrisomus heilneri Nichols, Amer. Mus. Novitates, XXVI, 1921, p. 2, fig. 2, Bahamas.
- 3706. *Labrisomus xanti* Gill.**
Pacific coast of Mexico.
Labrisomus xanti Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 107, Cerro Blanco, Cape San Lucas.
- 3707. *Labrisomus herminier* (LeSueur).**
St. Bartholomew, West Indies.
Blennius herminier LeSueur, Jour. Acad. Nat. Sci. Phila., IV, 1824, 361, St. Bartholomew, West Indies.
- 3708. *Labrisomus bucciferus* Poey.**
Cuba.
Labrosomus bucciferus Poey, Synopsis, 1868, 399, Cuba.
- 3709. *Labrisomus lentiginosus* Bean.**
Paget Parish, Bermudas.
Labrisomus lentiginosus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 31, tide pool at Does Rock, Paget Parish, Bermudas.
- 3710-3711. *Labrisomus microlepidotus* Poey.**
Cuba.
Labrosomus microlepidotus Poey, Anal. Soc. Española Hist. Nat., XIX, 1880, 246, Pl. VIII, fig. 2, Cuba.
- Genus 1299-1300. MALACOTENUS Gill.**
Malacotenus Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 103 (*Clinus delalandi* Cuvier and Valenciennes).
- 3712. *Malacotenus delalandi* (Cuvier and Valenciennes).**
Coast of Brazil and the Pacific coast of Mexico.
Clinus delalandii Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 279 (378), Brazil.
Clinus zonifer Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 361, Mazatlan.
- 3713. *Malacotenus ocellatus* (Steindachner).**
Bahama Islands.
Clinus ocellatus Steindachner, Ichth. Beitr., V, 1876, 182, Bahama Islands.
- 3714. *Malacotenus varius* (Poey).**
Cuba.
Myxodes varius Poey, Enumeratio, 1876, 132, pl. 5, fig. 2, Habana.

3715. *Malacoctenus macropus* (Poey).
Cuba.
Myxodes macropus Poey, Synopsis, III, 1868, 399, Habana.
3716. *Malacoctenus lugubris* (Poey).
Cuba.
Myxodes lugubris Poey, Enumeratio, 1876, 131, Cuba.
3717. *Malacoctenus gilli* (Steindachner).
Barbados.
Clinus gilli Steindachner, Ichth. Notizen, VI, 1867, 46, Barbados.
3718. *Malacoctenus bimaculatus* (Steindachner).
Small rocky islands north coast of Cuba.
Clinus bimaculatus Steindachner, Ichth. Beitr., V, 1876, 180, islands off Cuba.
3719. *Malacoctenus puertoricensis* Evermann and Marsh
Porto Rico.
Malacoctenus puertoricensis Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 358, Hucare, Porto Rico.
3720. *Malacoctenus versicolor* (Poey).
Cuba.
Myxodes versicolor Poey, Enumeratio, 1876, 131, pl. 5, fig. 1, Cuba.
3721. *Malacoctenus biguttatus* (Cope).
New Providence, Bahamas.
Labrisomus biguttatus Cope, Trans. Amer. Phil. Soc. Phila., XIV, 1871, 473, New Providence, Bahamas.
- Genus 1301. *ERICTEIS* Jordan.
Erieteis Jordan, Bull. U. S. Fish Com., XXII, 1902, 543 (*E. kalisheræ* Jordan).
3722. *Erieteis kalisheræ* Jordan.
Bush Key, Tortugas.
Erieteis kalisheræ Jordan, Bull. U. S. Fish Com., XXII, 1902, 543, pl. 2, Bush Key, Fla.
- Genus 1302. *ACTEIS* Jordan.
Acteis Jordan, Bull. U. S. Fish Com., XXII, 1902, 543 (*Malacoctenus moorei* Evermann and Marsh).
3723. *Acteis moorei* (Evermann and Marsh).
Porto Rico.
Malacoctenus moorei Evermann and Marsh, Fishes of Porto Rico, in Bull. U. S. Fish Com., XX, 1900, 309, fig. 97, Culebra Island, Porto Rico.
3724. *Acteis culebræ* (Evermann and Marsh).
Porto Rico.
Malacoctenus culebræ Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 357, Culebra Island.

Family 271. *BLENNIIDÆ*

- Genus 1303. *BLENNIUS* Linnæus. *Blennies*.
Blennius Linnæus, Syst. Nat., ed. X, 1758, 256 (*B. galerita* Linnæus).
Salaria Forskål, Descr. Anim., 1777, 22 (*Blennius galerita* Linnæus).
Adonis Gronow, Cat. Fish., Ed. Gray, 1854, 93 (*A. pavoninus* Gronow = *Blennius ocellaris* Linnæus).
3725. *Blennius fabbri* Nichols.
Florida.
Blennius fabbri Nichols, Bull. Amer. Mus. Nat. Hist., XXVIII, 1910, 161, Miami, Fla.; perhaps young of *Chasmodes bosquianus*.
3726. *Blennius stearnsi* Jordan and Gilbert.
Gulf of Mexico.
Blennius stearnsi Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 300, Pensacola snapper banks.

3727. *Blennius fucorum* Cuvier and Valenciennes. *Seaweed blenny*.
Off the Azores; also off New York.
Blennius fucorum Cuvier and Valenciennes, Hist. Nat. Poiss., XI,
1836, 194 (263), pl. 324, 240 miles south of the Azores.
Blennius oceanicus Cuvier and Valenciennes, loc. cit., 195 (265), open
sea.
3728. *Blennius cristatus* Linnæus. *Rockfish*.
Tropical Atlantic.
Blennius cristatus Linnæus, Syst. Nat., Ed. X, 1758, 256, Indies; after
Gronow.
Blennius crinitus Cuvier and Valenciennes, Hist. Nat. Poiss., XI,
1836, 175 (237), La Rochelle, France.
Blennius nuchifilis Cuvier and Valenciennes, loc. cit., 186 (253), Isle of
Ascension.
Blennius asterias Goode and Bean, Proc. U. S. Nat. Mus., V, 1882,
416, Garden Key, Fla.
3729. *Blennius favosus* Goode and Bean.
Gulf of Mexico; Garden Key, Fla.
Blennius favosus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882,
416, Garden Key, Fla.
3730. *Blennius pilicornis* Cuvier.
Coast of Brazil and north to the West Indies; recorded from Rio
Janeiro, Bahia, the Tortugas, and off the coast of Florida
Blennius pilicornis Cuvier, Règne Anim., ed. 2, II, 1829, 237, coasts of
Europe, Rio Janeiro.
3731. *Blennius marmoreus* Poey.
Cuba.
Blennius marmoreus Poey, Enumeratio, 1876, 130, Cuba.
3732. *Blennius truncatus* Poey.
Cuba.
Blennius truncatus Poey, Memorias, II, 1861, 424, Cuba.
3733. *Blennius vinctus* Poey.
Cuba.
Blennius vinctus Poey, Repertorio, II, 1867, 243, Habana.
- Genus 1304. *LIPOPHRYIS* Gill.
Pholis Fleming, Brit. Anim., 1828, 207 (*Blennius lævis* = *Blennius
pholis* Linnæus); name preoccupied.
Lipophrys Gill, Amer. Nat., XXX, June, 1896, 498 (*Blennius pholis*
Linnæus).
3734. *Lipophrys carolinus* (Cuvier and Valenciennes).
South Carolina.
Pholis carolinus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836,
204 (276), Carolina.
- Genus 1305. *SCARTELLA* Jordan.
Scartella Jordan, Proc. U. S. Nat. Mus., IX, 1886, 50 (*Blennius micro-
stomus* Poey).
Blennioides Jordan and Evermann, Fishes North and Mid. Amer.,
1898, 2386 (*Blennius brevipinnis* Günther).
3735. *Scartella microstoma* (Poey).
Cuba.
Blennius microstomus Poey, Memorias, II, 1861, 288, Cuba.
3736. *Scartella brevipinnis* (Günther).
Pacific coast of Mexico, from Mazatlan to Panama.
Blennius brevipinnis Günther, Cat., III, 1861, 226, Pacific coast of
Central America.
- Genus 1306. *HYPLEUROCHILUS* Gill.
Hypleurochilus Gill, Cat. Fishes E. Coast of N. Amer., Suppl. Proc.
Acad. Nat. Sci. Phila., XIII, 1861, 44 (*Blennius multifilis* Girard =
Blennius geminatus Wood).

- 3737. *Hypleurochilus geminatus* (Wood). Blenny.**
 South Atlantic and Gulf coasts of United States.
Blennius geminatus Wood, Jour. Acad. Nat. Sci. Phila., IV, 1824, 278,
 Charleston, S. C.
Blennius multifilis Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169,
 St. Joseph's Island, Tex.
- Genus 1307. HYPSOBLENNIUS Gill.**
Hypsoblennius Gill, Cat. Fishes E. Coast N. Amer., Suppl. Proc. Acad.
 Nat. Sci. Phila., XIII, 1861, 44 (*Blennius hantz* LeSueur).
Isesthes Jordan and Gilbert, Synopsis, 1883, 757 (*Blennius gentilis*
 Girard).
- 3738. *Hypsoblennius hantz* (LeSueur).**
 Coasts of North and South Carolina, south of Indian River, Fla.;
 locally common.
Blennius hantz LeSueur, Jour. Acad. Nat. Sci. Phila., 1st ser., IV, 1825,
 363, Charleston, S. C.
Blennius punctatus Wood, loc. cit., 278, Charleston, S. C.; not of
 Fabricius, 1780.
- 3739. *Hypsoblennius gilberti* (Jordan).**
 California, from Point Conception southward to Todos Santos; Santa
 Barbara Islands, among rocks.
Isesthes gilberti Jordan, Proc. U. S. Nat. Mus., V, 1882, 349, Santa
 Barbara, Calif.
- 3740. *Hypsoblennius gentilis* (Girard).**
 Coast of California from Monterey to Cape San Lucas.
Blennius gentilis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 149,
 Monterey, Calif.
- 3741. *Hypsoblennius piersoni* Gilbert and Starks.**
 Bay of Panama.
Hypsoblennius piersoni Gilbert and Starks, Mem. Calif. Acad. Sci., IV,
 1904, 191, Pl. XXXII, Bay of Panama.
- 3742. *Hypsoblennius striatus* (Steindachner).**
 Panama.
Blennius striatus Steindachner, Ichth. Beitr., V, 1876, 15, Panama.
- 3743. *Hypsoblennius ionthas* (Jordan and Gilbert).**
 Pensacola, Fla.
Isesthes ionthas Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
 299, Pensacola.
Isesthes scrutator Jordan and Gilbert, loc. cit., 300, Pensacola; Galveston.
- Genus 1308. CHASMODES Cuvier and Valenciennes.**
Chasmodes Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 218
 (295) (*Blennius bosquianus* Lacépède).
Blenitrachus Swainson, Classn. Fish., II, 1839, 78, 274 (*Pholis quadri-*
faciatus Wood).
- 3744. *Chasmodes bosquianus* (Lacépède). Banded blenny; Bosc's shanny;**
Four-banded blenny.
 New York to Florida; common southward in shallow water.
Blennius bosquianus Lacépède, Hist. Nat. Poiss., II, 1800, 493, South
 Carolina; female.
- 3745. *Chasmodes quadrifasciatus* (Wood).**
 Probably south Atlantic coast of United States.
Pholis quadrifasciatus Wood, Jour. Acad. Nat. Sci. Phila., IV, 1824,
 282, locality unknown; probably South Carolina.
- 3746. *Chasmodes saburræ* Jordan and Gilbert.**
 Pensacola Bay, Fla.
Chasmodes saburræ Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
 298, Pensacola, Fla.
- 3747. *Chasmodes novemlineatus* (Wood). Nine-lined shanny.**
 South Atlantic coast of the United States, South Carolina to Florida.
Pholis novemlineatus Wood, Jour. Acad. Nat. Sci. Phila., IV, 1825, 280,
 Charleston Harbor, S. C.

3748. *Chasmodes jenkinsi* Jordan and Evermann.
Guaymas, Sonora, Mexico.
Chasmodes jenkinsi Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 232, pl. 39, Guaymas, Mexico.
- Genus 1309. *HOMESTHES* Gilbert.
Homesthes Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2394 (*H. caulopus* Gilbert).
3749. *Homesthes caulopus* Gilbert
Panama Bay.
Homesthes caulopus Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2394, Panama.
- Genus 1310. *SCARTICHTHYS* Jordan and Evermann.
Scartes Jordan and Evermann, Check List, 1895, 471 (*Salarias salarias rubropunctatus* Cuvier and Valenciennes); name preoccupied.
Scartichthys Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2395 (*Salarias rubro-punctatus* Cuvier and Valenciennes).
3750. *Scartichthys rubropunctatus* (Cuvier and Valenciennes).
Coast of Peru and Chile, north to Panama.
Salarias rubro-punctatus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 257 (348), Juan Fernandez, off Chile.
- Genus 1311. *RUPISCARTES* Swainson.
Alicus Commerson, in Lacépède, Hist. Nat. Poiss., II, 1800, 458 (*A. saltatorius* Commerson); not binomial and not accepted by Lacépède.
Alicus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 337; casual notice; the name not accepted as that of a genus (*Blennius saliens*⁸⁹ = *Salarias alicus* Cuvier and Valenciennes).
Rupiscartes Swainson, Class. Fish., II, 1839, 275 (*Salarias atlanticus* Cuvier and Valenciennes).
3751. *Rupicartes atlanticus* (Cuvier and Valenciennes).
Tropical America on both coasts, north to West Indies and to Todos Santos, Lower California; abundant.
Salarias atlanticus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 238 (321), Atlantic Ocean.
- Genus 1312. *ENTOMACRODUS* Gill.
Entomacrodus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 168 (*E. nigricans* Gill).
3752. *Entomacrodus chiostrictus* (Jordan and Gilbert).
Pacific coast of Mexico.
Salarias chiostrictus Jordan and Gilbert, Synopsis, 1883, 363, Mazatlan, Mexico.
3753. *Entomacrodus margaritaceus* (Poey).
Cuba.
Salarias margaritaceus Poey, Memorias, II, 1861, 289, Cuba.
3754. *Entomacrodus decoratus* Poey.
Cuba.
Entomacrodus decoratus Poey, Synopsis, 1868, 398, Cuba.
3755. *Entomacrodus nigricans* Gill.
West Indies.
Entomacrodus nigricans Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 168, Barbados.
3756. *Entomacrodus cruentatus* Garman.
Off Cocos Island in deep water.
Entomacrodus cruentatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 123, Pl. L; off Cocos Island, Panama.
- Genus 1313. *SALARICHTHYS* Guichenot.
Salarichthys Guichenot, Mém., Soc. Sci. Nat. Cherbourg, XIII, 1868, 96 (*Salarias vomerinus* Cuvier and Valenciennes = *Salarias textilis* Cuvier and Valenciennes).

⁸⁹ "Un petit *salarias* qui nous paraît être celui, la même pour lequel Commerson a établi son genre *Alicus*."

3757. *Salarichthys textilis* Quoy and Gaimard.
West Indies, from Bermudas to Brazil.
Salaris textilis Quoy and Gaimard, in Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 227 (307), Ascension Island.
Salaris vomerinus Cuvier and Valenciennes, loc. cit., 258 (349), Bahia.
- Genus 1314. **BRANNERELLA** Gilbert.
Brannerella Gilbert, Proc. Wash. Acad. Sci., II, 1900, 180 (*B. brasiliensis* Gilbert).
3758. *Brannerella brasiliensis* Gilbert
Brazil, north to Curaçao, in the Caribbean Sea.
Brannerella brasiliensis Gilbert, Proc. Wash. Acad. Sci., II, 1900, 180, Maceio, Brazil.
3759. *Brannerella sluiteri* Metzelaar.
Caribbean Sea.
Brannerella sluiteri Metzelaar, Trop. Atl. Vissch., 1919, 155, Bonaire.
- Genus 1315. **OPHIOBLENNIUS** Gill.
Blennophis Valenciennes, in Webb and Barthelot, Îles Canaries, Poiss., 1844, 60 (*B. webbi* Valenciennes), name preoccupied.
Ophioblennius Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 103 (*B. webbi* Valenciennes); substitute for *Blennophis* Valenciennes, not *Blennophis* Swainson, a genus of *Clinidæ*.
3760. *Ophioblennius webbi* (Valenciennes).
West Indies and Canary Islands.
Blennophis webbi Valenciennes, in Webb and Barthelot, Îles Canaries, Poiss., 1839, 60, pl. 20, fig. 1, Fort Aventura, Canary Islands.
3761. *Ophioblennius steindachneri* Jordan and Evermann.
Pacific coast of Mexico.
Ophioblennius steindachneri Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2401, Tres Marias Islands; after Steindachner.
- Genus 1316. **CORALLIOZETUS** Evermann and Marsh.
Coralliozetus Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 362 (*C. cardonæ* Evermann and Marsh).
3762. *Coralliozetus cardonæ* Evermann and Marsh.
Porto Rico.
Coralliozetus cardonæ Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 362, reef at Cardona Light House, Playa de Ponce, Porto Rico.

Family 272. EMBLEMARIIDÆ

- Genus 1317. **EMBLEMARIA** Jordan and Gilbert.
Emblemaria Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 627 (*E. nivipes* Jordan and Gilbert).
3763. *Emblemaria atlantica* Jordan and Evermann.
Snapper Banks off Pensacola; Gulf of Mexico.
Emblemaria atlantica Jordan and Evermann, Fishes North and Mid. Amer., 1908, 2402, snapper banks off Pensacola, Fla.
3764. *Emblemaria nivipes* Jordan and Gilbert.
Pearl Island near Panama.
Emblemaria nivipes Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 627, Pearl Island, near Panama.
3765. *Emblemaria oculocirris* Jordan.
Gulf of California.
Emblemaria oculocirris Jordan, in Gilbert, Proc. U. S. Nat. Mus., XIX, 1896, 456, La Paz.
3766. *Emblemaria pandionis* Evermann and Marsh.
Porto Rico.
Emblemaria pandionis Evermann and Marsh, Bull. U. S. Fish Com., XX, 1900, 318, between Vieques and Culebra Islands.

3767. Emblemaria marki Mowbray.

Bermudas.

Emblemaria marki Mowbray, in Bean, Proc. Biol. Soc. Wash., XX V, 1912, 126, Hamilton Harbor, Bermudas.**Genus 1318. ACANTHEMBLEMARIA** Metzelaar.*Acanthemblemaria* Metzelaar, Trop. Atl. Vissch., 1919, 159 (*A. spinosa* Metzelaar).**3768. Acanthemblemaria spinosa** Metzelaar.

Caribbean Sea.

Acanthemblemaria spinosa Metzelaar, Trop. Atl. Vissch., 1919, 159, fig. 53, Curaçao.**Family 273. ATOPOCLINIDÆ****Genus 1319. ATOPOCLINIS** Vaillant.*Atopoclinus* Vaillant, Bull. Soc. Philom. Paris, ser. 8, VI, 1894, 73 (*A. ringens* Vaillant).**3769. Atopoclinus ringens** Vaillant.

Gulf of California.

Atopoclinus ringens Vaillant, Bull. Soc. Philom., ser. 8, VI, 1894, 74, Gulf of California.**Family 274. CHÆNOPSIDÆ****Genus 1320. CHÆNOPSIS** Gill.*Chænopsis* Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1863, 141, Pl. 3, fig. 3 (*C. ocellatus* Poey).*Lucioblennius* Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 103 (*L. alepidotus* Gilbert).**3770. Chænopsis ocellata** Poey.

Matanzas, Cuba.

Chænopsis ocellatus Poey, in Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1863, 143, Matanzas, Cuba.**3771. Chænopsis alepidota** (Gilbert).

Gulf of California; Avalon Bay, Calif.

Lucioblennius alepidotus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 103, Lower California.**3772. Chænopsis lucius** (Osburn and Nichols).

Gulf of California.

Lucioblennius lucius Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 179, fig. 15, San Jose Island, Gulf of California.**Genus 1321. PHOLIDICHTHYS** Bleeker.*Pholidichthys* Bleeker, Boerce, in Natur. Tijds. Neder. Indie, XI, 1856, 406 (*P. leucotænia* Bleeker).**3773. Pholidichthys anguilliformis** Lockington.

Gulf of California.

Pholidichthys anguilliformis Lockington, Proc. Acad. Nat. Sci. Phila., XXXIII, 1881, 118, San Jose Island, Lower California.**Genus 1322. PSEDNOBLENNIUS** Jenkins and Evermann.*Pseudoblennius* Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 156 (*P. hypacanthus* Jenkins and Evermann).**3774. Pseudoblennius hypacanthus** Jenkins and Evermann.

Gulf of California at Guaymas, Sonora, Mexico.

Pseudoblennius hypacanthus Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 156, Guaymas, Mexico.**Genus 1323. STATHMONOTUS** Bean.*Stathmonotus* Bean, Proc. U. S. Nat. Mus., VIII, 1885, 191 (*S. hemphillii* Bean).**3775. Stathmonotus hemphillii** Bean.

Key West, Fla.

Stathmonotus hemphillii Bean, Proc. U. S. Nat. Mus., VIII, 1885, 191, pl. 13, Key West, Fla.

3776. *Stathmonotus tekla* Nichols.

Key West, Fla.

Stathmonotus tekla Nichols, Bull. Amer. Mus. Nat. Hist., XXVIII, 1910, 161, fig. 1c. XXX, Pl. XI, fig. 2, Key West, Fla.**Family 275. CEBIDICHTHYIDÆ****Genus 1324. CEBIDICHTHYS Ayres.***Cebidichthys* Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 59 (*Apodichthys violaceus* Girard).**3777. *Cebidichthys violaceus* (Girard).**

Coast of California from Crescent City to Carpenteria.

Apodichthys violaceus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 150, Monterey, Calif.*Cebidichthys cristagalli* Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 58, San Francisco.**Family 276. CHIROLOPHIDÆ****Genus 1325. BRYOSTEMMA Jordan and Starks.***Bryostemma* Jordan and Starks, Proc. Calif. Acad. Sci. 2d ser, V, 1895, 841 (*Blennius polyactocephalus* Pallas).**3778. *Bryostemma polyactocephalum* (Pallas).**

Northern Japan and Kamchatka to Puget Sound.

Blennius polyactocephalus Pallas, Zool. Rosso-Asiat, III, 1811, 179, Kamchatka.*Charolophus japonicus* Herzenstein, Mélanges Biologiques Soc. Sci. Petersb., XIII, 1890, 123, Yezo, Japan.**3779. *Bryostemma nugator* Jordan and Williams.**

Puget Sound.

Bryostemma nugator Jordan and Williams, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 843, pl. 101, Seattle, Wash.**3780. *Bryostemma decoratum* Jordan and Snyder.**

Point Orchard, Wash.

Bryostemma decoratum Jordan and Snyder, Proc. U. S. Nat. Mus., XXV, 1903, 615, Point Orchard, near Seattle, Wash.**3781. *Bryostemma tarsodes* Jordan and Snyder.**

Unalaska.

Bryostemma tarsodes Jordan and Snyder, Proc. U. S. Nat. Mus., XXV, 1903, 614, Unalaska.**Genus 1326. BRYOLOPHUS Jordan and Snyder.***Bryolophus* Jordan and Snyder, Proc. U. S. Nat. Mus., XXV, 1903, 617 (*B. lysimus* Jordan and Snyder).**3782. *Bryolophus lysimus* Jordan and Snyder.**

Unalaska.

Bryolophus lysimus Jordan and Snyder, Proc. U. S. Nat. Mus., XXV, 1903, 617, Unalaska.**Genus 1327. GYMNOCLINUS Gilbert and Burke.***Gymnoclinus* Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 86 (*G. cristulatus* Gilbert and Burke).**3783. *Gymnoclinus cristulatus* Gilbert and Burke.**

Bering Island.

Gymnoclinus cristulatus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 86, fig. 30, Nikolski, Bering Island.**Family 277. PHOLIDÆ****Genus 1328. APODICHTHYS Girard.***Apodichthys* Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 150 (*A. flavidus* Girard).

3784. *Apodichthys flavidus* Girard.
Point Conception to Vancouver Island.
Apodichthys flavidus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 150, Presidio (San Francisco), Calif.
Apodichthys vrescens Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 55 (54), San Francisco.
Apodichthys sanguineus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 279, California.
Apodichthys mornatus Gill, loc. cit., 279, Puget Sound.
3785. *Apodichthys univittatus* Lockington.
Lower California.
Apodichthys univittatus Lockington, Proc. Acad. Nat. Sci. Phila., XXXIII, 1881, 119, Lower California, probably from the Gulf.
- Genus 1329. *XERERPES* Jordan and Gilbert.
Xererpes Jordan and Gilbert, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 846 (*Apodichthys fucorum* Jordan and Gilbert).
3786. *Xererpes fucorum* (Jordan and Gilbert).
Coast of California.
Apodichthys fucorum Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 139, Point Pinos, near Monterey, Calif.
- Genus 1330. *ULVICOLA* Gilbert and Starks.
Ulvicola Gilbert and Starks, Proc. U. S. Nat. Mus., XIX, 1896, 455 (*U. sanctæ-rosæ* Gilbert and Starks).
3787. *Ulvicola sanctæ-rosæ* Gilbert and Starks.
Santa Rosa Island, coast of California.
Ulvicola sanctæ-rosæ Gilbert and Starks, Proc. U. S. Nat. Mus., XIX, 1896, 453, Santa Rosa Island, Calif.
- Genus 1331. *PHOLIS* Gronow.
Pholis Gronow, Acta Helvet., IV, 1760, 259 (*Blennius gunnellus* Linnæus).
Pholis Scopoli, Introd. Hist. Nat., 1777, 456 (*Blennius gunnellus* Linnæus).
Murænoides Lacépède, Hist. Nat. Poiss., II, 1800, 324 (*Murænoides sujef* Lacépède).
Centronotus Schneider, Systema, ed. Bloch, 1801, 165 (*Centronotus fasciatus* Schneider).
Dactyleptus Rafinesque, Anal. Nat., 1815, 82; substitute for *Murænoides*.
Gunnellus Fleming, Brit. Anim., 1828, 207 (*G. vulgaris* Fleming = *Blennius gunnellus* Linnæus).
Centronotus Cuvier, Règne Anim., ed. 2, II, 1829, 239 (*Blennius gunnellus* Linnæus).
Ophisomus Swainson, Classn. Anim., II, 1839, 279 (*Blennius gunnellus* Linnæus).
Urocentrus Kner, Sitzber. K. Akad. Wiss. Wien, LVIII, 1868, 51 (*U. pictus* Kner).
3788. *Pholis gunnellus* (Linnæus). *Gunnel*; *Butterfish*; *Rockfish*.
North-Atlantic, south to Cape Cod.
Blennius gunnellus Linnæus, Syst. Nat., Ed. X, 1758, 257, Atlantic Ocean.
Ophidion imberbe Linnæus, loc. cit., 259, Europe.
Blennius europæus Olafsen, Reise Island, I, 1772, 81, Iceland.
Blennius murænoides Sujef, Act. Petrop., II, 1779, 195, no locality, probably Baltic.
Ophidium mucronatum Mitchill, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 361, Pl. I, fig. 1, New York.
Gunnellus vulgaris Fleming, British Anim., 1842, 207, England.

3789. *Pholis fasciatus* (Bloch and Schneider).
Greenland to Alaska and Japan.
Centronotus fasciatus Bloch and Schneider, Syst. Ichth., 1801, 165, pl. 37, fig. 1, Tranquebar.
Gunnellus grænlandicus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 326 (442), pl. 340, Greenland; after Bloch and Schneider.
Gunnellus murænoïdes Valenciennes, in Cuvier, Règne Anim., Poiss., ed. 2, 1829, pl. 78, fig. 2, 916; after Bloch and Schneider.
Blennius lænia Pallas, Zoogr. Rosso-Asiat., III, 1811, 178, Kuril Islands.
Murænoïdes maxillaris Bean, Proc. U. S. Nat. Mus., IV, 1881, 147, St. Paul Island, Alaska.
3790. *Pholis ornatus* (Girard).
San Francisco to Alaska.
Gunnellus ornatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 149, Presidio (San Francisco), Calif.
Centronotus lætus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 27, Sitka or Unalaska.
3791. *Pholis gilli* Evermann and Goldsborough.
Bering Sea.
Pholis gilli Evermann and Goldsborough, Bull. U. S. Bur. Fish., XXVI, 1906, 337, fig. 110, Bering Sea.
- Genus 1332. RHODYMENICHTHYS Jordan and Evermann.
Rhodymenichthys Jordan and Evermann, Check List, 1895, 474 (*Gunnellus ruberrimus* Cuvier and Valenciennes = *Blennius dolichogaster* Pallas).
3792. *Rhodymenichthys dolichogaster* (Pallas). *Butterfish*.
Okhotsk Sea to San Francisco
Blennius dolichogaster Pallas, Zoogr. Rosso-Asiat., III, 1811, 175, Kamchatka.
Gunnellus ruberrimus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 324 (440), Kuril Islands; after Pallas.
- Genus 1333. ASTERNOPTERYX Ruppell.
Asternopteryx Ruppell, in Günther, Cat., III, 1861, 288 (*A. gunelliformis* Ruppell).
3793. *Asternopteryx gunelliformis* Ruppell.
Greenland.
Asternopteryx gunelliformis Ruppell, in Günther, Cat., III, 1861, 288, Greenland; type in Senckenberg Museum, Frankfurt; a second specimen from Greenland examined by us.
- Genus 1334. ANOPLARCHUS Gill.
Anoplarchus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 261 (*A. purpureus* Gill).
3794. *Anoplarchus cristagalli* (Günther).
Alaska to San Francisco.
Centronotus cristagalli Günther, Cat., III, July, 1861, 289, Vancouver Island.
Anoplarchus purpureus Gill, Proc. Acad. Nat. Sci. Phila., XIII, Sept., 1861, 262, Washington Territory.
3795. *Anoplarchus cristagalli archolepis* Hubbs.
Coast of central California.
Anoplarchus purpureus archolepis Hubbs, Papers Mich. Acad. Sci., Arts and Letters, VII, 1926, 375, Reef at Carmel, Calif.
3796. *Anoplarchus insignis* Gilbert and Burke.
Attu Island.
Anoplarchus insignis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 88, fig. 32, Attu Island, Bering Sea.
- Genus 1335. PLECTOBRANCHUS Gilbert.
Plectobranchnus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 102 (*P. evides* Gilbert).

3797. *Plectobranchnus evides* Gilbert.
Coasts of Oregon and Washington.
Plectobranchnus evides Gilbert, Proc. U. S. Nat. Mus., XIII, 1898, 102,
coasts of Oregon and Washington.

Family 278. XIPHISTERIDÆ

Genus 1336. XIPHISTER Jordan and Gilbert.

- Xiphidion* Girard, Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 119 (*X mucosum* Girard); name preoccupied in *Orthoptera*.
Xiphister Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 135
(*Xiphidion mucosum* Girard).

3798. *Xiphister mucosus* (Girard).

- Monterey to Alaska.
Xiphidion mucosum Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858,
119, South Farallon Island, off California.
Xiphidium cruoreum Cope, Proc. Amer. Phil. Soc., XIII, 1873, 27,
Sitka, Alaska.

Genus 1337. EPIGEICHTHYS Hubbs.

- Epigeichthys* Hubbs, Papers Mich. Acad. Sci. Arts and Letters, VII,
1926 (*Xiphister rupestris* Jordan and Gilbert).

3799. *Epigeichthys atropurpureus* (Kittlitz).

- Pacific coast of America from Alaska to Monterey.
Ophidium atropurpureum Kittlitz, Denkwurd, einer Reise Russ.-Amer.,
I, 1858, 225, Alaska.
Xiphister rupestris Jordan and Gilbert, Proc. U. S. Nat. Mus., III,
1880, 137, Monterey, Calif.

Genus 1338. PHYTICHTHYS Hubbs.

- Xiphistes* Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895,
846 (*Xiphistes chirus* Jordan and Gilbert); name preoccupied in
insects.
Phytichthys Hubbs, in Jordan, Class. Fishes, 1923, 234 (*Xiphister chirus*
Jordan and Gilbert).

3800. *Phytichthys chirus* (Jordan and Gilbert).

- Monterey to Alaska.
Xiphister chirus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880,
135, Point Pinos, near Monterey, Calif.
Xiphistes ulvæ Jordan and Starks, Fishes Puget Sound, Proc. Calif.
Acad. Sci., 2d ser., V, 1895, 847, pl. 102, Waadda Island, Neah Bay.

3801. *Phytichthys versicolor* (Gilbert and Burke).

- Bering Sea.
Xiphistes versicolor Gilbert and Burke, Bull. U. S. Bur. Fish., XXX,
1910, 88, fig. 33; Attu; Agattu, Bering Sea.

Family 279. STICHÆIDÆ

Genus 1339. STICHÆUS Reinhardt.

- Stichæus* Reinhardt, K. Dansk. Vidensk. Selsk. Skrift Natur. og
Math. Afhandl., VI, 1837, CX (*Blennius punctatus* Fabricius).
Notogrammus Bean, Proc. U. S. Nat. Mus., IV, 1881, 147 (*N. rothrocki*
Bean); young example.

3802. *Stichæus punctatus* (Fabricius). *Spotted snake blenny*.

- Arctic seas from Greenland to northern Siberia, south to Bristol Bay
and Newfoundland.
Blennius punctatus Fabricius, Fauna Groenlandica, 1780, 153, Greenland.
Notogrammus rothrocki Bean, Proc. U. S. Nat. Mus., IV, 1881, 146,
Plover Bay and Cape Lisburne, Siberia.

Genus 1340. PLAGIOGRAMMUS Bean.

- Plagiogrammus* Bean, Proc. U. S. Nat. Mus., XVI, 1893, 699 (*P.*
hopkinsi Bean).

3803. *Plagiogrammus hopkinsi* Bean.

- Monterey Bay, Calif.
Plagiogrammus hopkinsi Bean, Proc. U. S. Nat. Mus., XVI, 1893, 700,
with fig. on p. 699; Monterey, Calif.

Genus 1341. ULVARIA Jordan and Evermann.

Ulvaria Jordan and Evermann, Check List, 1895, 475 (*Pholis subbifurcatus* Storer).

3804. *Ulvaria subbifurcata* (Storer). *Radiated shanny*.

North Atlantic, south of Cape Cod.

Pholis subbifurcatus Storer, Rept. Fish. Mass., 1839, 63, Nahant, Mass.

Genus 1342. EUMESOGRAMMUS Gill.

Eumesogrammus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 210 (*Clinus præcisus* Krøyer).

3805. *Eumesogrammus præcisus* (Krøyer).

Greenland.

?*Clinus præcisus* Krøyer, Naturh. Tidsskr., I, 1837, 25, Greenland.

Clinus unimaculatus Reinhardt, Dansk. Vidensk. Selsk., VII, 1838, 114, 121, Greenland.

Family 280. LUMPENIDÆ**Genus 1343. LEPTOCLINUS** Gill.

Ctenodon Nilsson, Scand. Fauna, IV, 1853, 190 (*Clinus maculatus* Fries); name preoccupied.

Leptoclinus Gill, Cat. Fish. E. coast N. Amer., Suppl., Proc. Acad. Nat. Sci. Phila., XIII, 1861, 45 (*Clinus aculeatus* Reinhardt=*Clinus maculatus* Fries).

3806. *Leptoclinus maculatus* (Fries). *Langbarn*.

North Atlantic coasts of Europe; this or some closely similar species is recorded from Bering Sea, Robben Island, Aleutian Islands, and Hood Canal, Puget Sound.

Clinus maculatus Fries, Kgl. Vet. Ak. Handl., 1837, 51, Bohuslaan, Sweden.

Clinus aculeatus Reinhardt, K. Dansk. Vid. Selsk. Natur. Afh., VII, 1838, 114, 122, Spitzbergen.

Genus 1344. POROCLINUS Bean.

Poroclinus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 40 (*P. rothrocki* Bean).

3807. *Poroclinus rothrocki* Bean.

Between Nagai and Big Koniushi Islands, Alaska.

Poroclinus rothrocki Bean, Proc. U. S. Nat. Mus., XIII, 1890, 40, Albatross station 2952, lat. 55° 15' N., long. 159° 37' W., in 58 fathoms.

Genus 1345. ANISARCHUS Gill.

Anisarchus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 209 (*Clinus medius* Reinhardt).

3808. *Anisarchus medius* (Reinhardt).

Greenland to Norway and Spitzbergen; this or a similar species on coasts of southern and central Alaska.

Clinus medius Reinhardt, K. Dansk. Vidensk. Selsk. Skrift. Afh., VII, 1838, 114, 121, Greenland.

Genus 1346. LUMPENUS Reinhardt.

Lumpenus Reinhardt, Dansk. Vidensk. Selsk. Natur., VI, 1837, CX (*Blennius lumpenus* Fabricius=*Gunnellus fabricii* Cuvier and Valenciennes).

Leptogunnellus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 26 (*L. gracilis* Ayres=*Blennius anguillaris* Pallas).

Centroblennius Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 209 (*Lumpenus nubilus* Richardson=*Gunnellus fabricii* Cuvier and Valenciennes).

Leptoblennius Gill, loc. cit., 1864, 209 (*Blennius serpentinus* Storer=*Blennius lampetæformis* Walbaum).

3809. *Lumpenus gracilis* (Ayres).

San Francisco to Alaska.

Blennius anguillaris Pallas, Zoogr. Rosso-Asiat., III, 1811, 176, about Kamchatka; no definite locality; not of Peck.

Leptogunnellus gracilis Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 26, San Francisco.

3810. *Lumpenus mackayi* Gilbert.
Nushagak River, Alaska.
Lumpenus mackayi Gilbert, Rept. U. S. Fish Com., XIX, 1893, 450,
pl. 32, Nushagak River, Alaska.
3811. *Lumpenus fabricii* (Cuvier and Valenciennes).
Arctic seas; Greenland, Gulf of St. Lawrence, Wellington Sound,
Bristol Bay, and other localities in Bering Sea.
Gunellus fabricii Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836,
317 (431), Greenland.
Lumpenus nubilus Richardson, Last Arctic Voyage, Fishes, 1855, 259,
pl. 28, Wellington Sound.
3812. *Lumpenus lampetræformis* (Walbaum). *Snake blenny; Tangbrosme;*
Eel blenny; Snakefish.
North Atlantic and Arctic, on both shores; south to Sweden and Green-
land; rarely to Cape Cod.
Blennius lampetræformis Walbaum, Artedi Pisc., III, 1792, 184, Iceland.
Centronotus islandicus Schneider, ed. Bloch, Syst. Ichth., 1801, 167,
Iceland.
? *Chnus nebulosus* Fries, Vet. Akad. Handl., 1837, 55, Bohuslän, Sweden.
? *Chnus mohr* Krøyer, Naturh. Tidsskr., I, 1837, R, 32, Iceland.
Blennius gracilis Stuvitz, Nye Mag. for Naturvid., I, 1838, 406, west
coast of Norway.
Blennius serpentinus Storer, Proc. Bost. Soc. Nat. Hist., III, 1848, 30,
Massachusetts Bay
- Genus 1347. **LUMPENELLA** Hubbs.
Lumpenella Hubbs, Papers Mich. Acad. Sci. Arts and Letters, VII,
1926, 378 (*Lumpenus longirostris* Evermann and Goldsborough).
3813. *Lumpenella longirostris* (Evermann and Goldsborough).
Southeastern Alaska.
Lumpenus longirostris Evermann and Goldsborough, Bull. U. S. Bur.
Fish., XXVI, 1906, 340, fig. 115, Lynn Canal, Alaska.

Family 281. PTILICHTHYIDÆ

- Genus 1348. **PTILICHTHYS** Bean.
Ptilichthys Bean, Proc. U. S. Nat. Mus., IV, 1881, 157 (*P. goodei* Bean).
3814. *Ptilichthys goodei* Bean.
Aleutian Islands.
Ptilichthys goodei Bean, Proc. U. S. Nat. Mus., IV, 1881, 157, Port
Levasheff, Unalaska.

Family 282. CRYPTACANTHODIDÆ

- Genus 1349. **CRYPTACANTHODES** Storer.
Cryptacanthodes Storer, Rept. Fish. Mass., 1839, 27 (*C. maculatus*
Storer).
3815. *Cryptacanthodes maculatus* Storer. *Wrymouth; Ghostfish*
North Atlantic, south to Cape Cod.
Cryptacanthodes maculatus Storer, Rept. Fish. Mass., 1839, 28, Massa-
chusetts.
Fraseria borealis DeKay, New York Fauna Fishes, 1842, 316, New
York; after Peck.
Cryptacanthodes mornatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863,
332, coast of Massachusetts.
- Genus 1350. **LYCONECTES** Gilbert.
Lyconectes Gilbert, Rept. U. S. Fish Com., XIX, 1893, 452 (*L. aleutensis*
Gilbert).
3816. *Lyconectes aleutensis* Gilbert.
Aleutian Islands.
Lyconectes aleutensis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 452,
pl. 34, near Unalaska.
- Genus 1351. **DELOLEPIS** Bean.
Delolepis Bean, Proc. U. S. Nat. Mus., IV, 1881, 465 (*D. virgatus* Bean).

3817. *Delolepis giganteus* Kittlitz
Coasts of British Columbia and southern Alaska, south to Puget Sound.
Delolepis giganteus Kittlitz, Denk. Reise Russ. Amer., 226, 1858, Siberia.
Delolepis virgatus Bean, Proc. U. S. Nat. Mus., IV, 1881, 466, King-combe Inlet, British Columbia, and Port Wrangel, Alaska.

Family 283. ANARHICADIDÆ

Genus 1352. ANARHICHAS Linnæus.

Anarhichas Linnæus, Syst. Nat., Ed. X, 1758, 247 (*A. lupus* Linnæus).

3818. *Anarhichas lupus* Linnæus *Wolfish*
North Atlantic, south to Cape Cod and France: rather common in America and Europe.
Anarhichas lupus Linnæus, Syst. Nat., Ed. X, 1758, 247, no definite locality.
Anarhichas strigosus Gmelin, Syst. Nat., Tome I, Pt. III, 1789, 1144, British seas.
Anarhichas vomerinus Agassiz, in Störck, Hist. Fish. Mass., 1867, 265, pl. 18, fig. 1, Cusk Rocks between Boston and Cape Ann.
3819. *Anarhichas orientalis* Pallas.
Alaska.
Anarhichas orientalis Pallas, Zoogr. Rosso-Asiat., III, 1811, 77, Kamchatka.
Anarhichas lepturus Bean, Proc. U. S. Nat. Mus., II, 1879, 212, St. Michaels, Alaska.

Genus 1353. LYCICHTHYS Gill.

Lycichthys Gill, Ann. Rec. Indust. & Sci., 1876, CLXVII (*Anarrhichas latifrons* Steenstrup and Hallgrímsson).

3820. *Lycichthys latifrons* (Steenstrup and Hallgrímsson).
North Atlantic, on both coasts; chiefly north of the Arctic Circle.
Anarrhichas latifrons Steenstrup and Hallgrímsson, Förh. Skand. Naturf., 3 die Møte, 1842, 647; name only, no description; Viden-skabelige Meddeleker Natur Forening Kjøbenhavn, 1877-1878, 109-114.
3821. *Lycichthys minor* (Olafsen).
North Atlantic, on both coasts, chiefly north of the Arctic Circle.
Anarrhichas minor Olafsen, Reise i. Island, sec. 682b, 1774, tab. 42, Iceland.
Anarrhichas karra Bonnaterre, Encycl. Ichth., 1788, 38, Iceland, after Olafsen.
Anarrhichas maculatus Bloch and Schneider, Syst. Ichth., 1801, 496, Iceland; after Olafsen.
Anarrhichas leopardus Agassiz, in Spix, Pisc. Brasil., 1829, 92, tab. 51, "Atlantic Ocean."

Family 284. ANARRHICHTHYIDÆ

Genus 1354. ANARRHICHTHYS Ayres.

Anarrhichthys Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 31 (*A. ocellatus* Ayres = *A. felis* Girard).

3822. *Anarrhichthys felis* (Girard). *Wolf eel*.
Pacific coast, from Monterey north to Puget Sound; common.
Anarrhichas felis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 150, San Francisco; brief notice.
Anarrhichthys ocellatus Ayres, Proc. Calif. Acad. Nat. Sc., I, 1855, 31, San Francisco.

Family 285. CERDALIDÆ

Genus 1355. MICRODESMUS Günther.

Microdesmus Günther, Proc. Zool. Soc. Lond., 1864, 26 (*M. dipus* Günther).

3823. *Microdesmus dipus* Gunther.
Pacific coast of tropical America.
Microdesmus dipus Gunther, Proc. Zool. Soc. Lond., Jan. 26, 1864,
26, pl. 3, fig. 2, Central America.
3824. *Microdesmus retropinnis* Jordan and Gilbert.
Panama.
Microdesmus retropinnis Jordan and Gilbert, Bull. U. S. Fish Com., I,
1881, 331, Panama.
- Genus 1356. CERDALE Jordan and Gilbert.
Cerdale Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 332 (*C.
ionthas* Jordan and Gilbert).
3825. *Cerdale ionthas* Jordan and Gilbert.
Panama.
Cerdale ionthas Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881,
332, Panama.
- Genus 1357. LEPTOCERDALE Weymouth.
Leptocerdale Weymouth, Proc. U. S. Nat. Mus., XXXVIII, 1910,
142 (*L. longipinnis* Weymouth).
3826. *Leptocerdale longipinnis* Weymouth.
Coast of Louisiana.
Leptocerdale longipinnis Weymouth, Proc. U. S. Nat. Mus., XXXVIII,
1910, 142, Cameron, La.

Family 286. SCYTALINIDÆ

- Genus 1358. SCYTALINA Jordan and Gilbert.
Scytalina Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 266
(*S. cerdale* Jordan and Gilbert).
Scytaliscus Jordan and Gilbert, Proc. U. S. Nat. Mus., VI, 1883, III;
same type, there being already a genus *Scytalinus*.
3827. *Scytalina cerdale* Jordan and Gilbert.
Straits of Juan de Fuca, Waadda Island, near Cape Flattery; also in
Aleutian Islands burrowing deep in gravel and sand.
Scytalina cerdale Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880,
267, Waadda Island, Neah Bay.

Family 287. ZOARCIDÆ

- Genus 1359. MACROZOARCES Gill.
Macrozoarces Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 258 (*Blennius
anguillaris* Peck).
3828. *Macrozoarces americanus* (Bloch and Schneider). *Eelpout*; *Muttonfish*;
Mother-of-eels; *Lang*; *Lamper eel*; *Conger eel*.
Delaware to Labrador; rather common northward.
? *Enchelyopus* Gronow, Zoophyl., No. 266, 1763, 77, America.
? *Blennius americanus* Bloch and Schneider, Syst. Ichth., 1801, 171,
America; after Gronow.
Blennius anguillaris Peck, Mem. Amer. Acad. Sci., II, 1804, 46, fig. 3,
New Hampshire.
? *Blennius fimbriatus* Mitchill, Trans. Lit. and Phil. Soc. N. Y., 1815,
374, pl. 1, fig. 6, New York.
Blennius ciliatus Mitchill, loc. cit., 374, pl. 1, fig. 6, New York.
Zoarces labrosus Cuvier, Règne Anim., ed. 2, II, 1829, 240, America.
Zoarces gronovi Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836,
246 (469); after Gronow.
- Genus 1360. EMBRYX Jordan and Evermann.
Embryx Jordan and Evermann, Fishes North and Mid. Amer., 1898,
2458 (*Lycodopsis crotalinus* Gilbert).
3829. *Embryx crotalinus* (Gilbert).
Northern Pacific and off Santa Barbara Islands.
Lycodopsis crotalinus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 105,
off Santa Barbara Islands at Albatross station 2980 in 603 fathoms.

3830. *Embryx crassilabris* (Gilbert).
Coast of southern California.
Lycodopsis crassilabris Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 106,
off southern California at Albatross station 2839.
3831. *Embryx parallelus* Gilbert.
Monterey Bay.
Embryx parallelus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 360,
Pl. X, IX, fig. 16, Monterey Bay, 394 to 406 fathoms.
- Genus 1361. *LYCOPSIS* Collett.
Lycopsis Collett, Proc. Zool. Soc. Lond., 1879, 381 (*Lycodes pacificus*
Collett).
Leurynnus Lockington, Proc. U. S. Nat. Mus., II, 1879, 326 (*L. paucidens*
Lockington = *Lycodes pacificus* Collett).
3832. *Lycodopsis pacificus* (Collett).
San Francisco to central Alaska; Afognak Bay.
Lycodes (Lycodopsis) pacificus Collett, Proc. Zool. Soc. Lond., 1879,
381; said to be from Japan; doubtless an error.
Leurynnus paucidens Lockington, Proc. U. S. Nat. Mus., II, 1879, 326,
off San Francisco, Calif.
3833. *Lycodopsis scaurus* Garman.
Off Bay of Panama in deep water.
Lycodopsis scaurus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
132, Pl. XXXII, off Bay of Panama.
- Genus 1362. *APRODON* Gilbert.
Aprodon Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 106 (*A. corteziana*
Gilbert).
3834. *Aprodon cortezianus* Gilbert.
Cortez Banks, near San Diego, Calif.
Aprodon corteziana Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 107,
Cortez Banks, off San Diego, at Albatross stations 2925 and 2948 in
339 and 266 fathoms.
- Genus 1363. *LYCODES* Reinhardt.
Lycodes Reinhardt, Königl. Danske. Vidensk. Selsk. Naturv. Math.
Afh., VII, 1838, 153 (*L. vahli* Reinhardt).
3835. *Lycodes vahli* Reinhardt. *Vahl's lycodes; Misarkornak*.
Coast of Greenland.
Lycodes vahli Reinhardt, Königl. Danske. Vidensk. Selsk. Nat. Math.
Afh., VII, 1838, 153, pl. 5, Greenland.
3836. *Lycodes frigidus* Collett.
North Atlantic, off New England coast; Spitzbergen.
Lycodes frigidus Collett, Forh. Vid. Selsk. Christ., 1878, Nos. 14, 45,
near Spitzbergen.
3837. *Lycodes terræ-novæ* Collett.
Newfoundland Banks.
Lycodes terræ-novæ Collett, Campagnes Scientifiques *L'Hirondelle*,
Poiss., 1896, 55, Newfoundland Banks in 25 fathoms.
3838. *Lycodes reticulatus* Reinhardt. *Akulliakitsoq; Spotted lycodes*.
North Atlantic from Greenland south to Narragansett Bay; also
northern Europe.
Lycodes reticulatus Reinhardt, Königl. Danske Vid. Selsk. Afh., VII,
1838, 167, Greenland.
Lycodes perspicillum Krøyer, Dansk. Vidensk. Selsk. Afhandl., XI,
1845, p. cccxxiii, Newfoundland.
3839. *Lycodes esmarkii* Collett.
North Atlantic; Gulf Stream, in about latitude 40°, to Finmark and
Spitzbergen.
Lycodes esmarkii Collett, Norges Fiske, 1875, 95; American specimens
from the Gulf Stream in about latitude 40°.
3840. *Lycodes jordani* Evermann and Goldsborough.
Puget Sound and southeastern Alaska.
Lycodes jordani Evermann and Goldsborough, Bull. U. S. Bur. Fish.,
XXVI, 1906, 343, fig. 120, Puget Sound.

3841. *Lycodes gracilis* Sars
Atlantic Ocean off Nova Scotia.
Lycodes gracilis Sars, Christ. Vid. Selsk. Forh., 1866, 40-45, Drøbak, Norway.
Lycodes zoarchus Goode and Bean, Oceanic Ichth., 1896, 308, off Nova Scotia.
3842. *Lycodes concolor* Gill and Townsend.
Bering Sea.
Lycodes concolor Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 233, Bering Sea in 276 fathoms.
3843. *Lycodes digitatus* Gill and Townsend.
Bering Sea.
Lycodes digitatus Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 232, Bering Sea in 49 fathoms.
3844. *Lycodes palearis* Gilbert.
Bristol Bay, Alaska.
Lycodes palearis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 454, Bristol Bay, Alaska, at *Albatross* stations 3253 and 3254 in 36 and 46 fathoms.
3845. *Lycodes brevipes* Bean.
Aleutian Islands to Puget Sound.
Lycodes brevipes Bean, Proc. U. S. Nat. Mus., XIII, 1890, 38, between Unga and Nagai islands in 110 fathoms.
3846. *Lycodes anguis* Garman.
Off Lower California and Panama in deep water.
Lycodes anguis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 133, off Bay of Panama.
3847. *Lycodes serpens* Garman.
Off Lower California in deep water.
Lycodes serpens Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 134, off Gulf of California.
3848. *Lycodes incisus* Garman.
Off Bay of Panama in deep water.
Lycodes incisus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 135, Pl. XXX, off Bay of Panama.
3849. *Lycodes cicatrifer* Garman.
Off Bay of Panama in deep water.
Lycodes cicatrifer Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 136, Pl. XXXI, off Bay of Panama.
- Genus 1364. *LYCIAS* Jordan and Evermann.
Lycias Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2461 (*Lycodes seminudus* Reinhardt).
3850. *Lycias nebulosus* (Krøyer). *Cloudy lycodes*.
Greenland.
Lycodes nebulosus Krøyer, Kongl. Danske Vidensk. Selsk. Afhand., XI, 1845, p. ccxxxii, Greenland.
3851. *Lycias seminudus* (Reinhardt). *Half-naked lycodes*.
North Atlantic, from Greenland to Spitzbergen.
Lycodes seminudus Reinhardt, Kongl. Danske Selsk. Afhand., VII, 1838, 221, Omenak, Greenland.
- Genus 1365. *LYCENCHELYS* Gill.
Lycenchelys Gill, Proc. Acad. Nat. Sci. Phila., XXXVI, 1884, 180 (*Lycodes muræna* Collett).
3852. *Lycenchelys verrillii* (Goode and Bean).
Coast of Massachusetts and northward.
Lycodes verrillii Goode and Bean, Amer. Jour. Sci. Arts, XIV, 1877, 474; 27 miles south and west of Chebucto Head near Halifax.

3853. *Lycenchelys paxillus* (Goode and Bean).

Gulf Stream.

Lycodes paxillus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 44, between La Have and Sable Island Banks.*Lycodes paxilloides* Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 207, off Newfoundland.**3854. *Lycenchelys porifer* (Gilbert)**

Off Lower California.

Lycodes porifer Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 104, off Lower California at Albatross station 3009 in 857 fathoms.**Genus 1366. *LYCODALEPIS* Bleeker.***Lycodalepis* Bleeker, Versl. Akad. Amst., ed. 2, VIII, 1874, 369 (*Lycodes muosus* Richardson).**3855. *Lycodalepis polaris* (Sabine). *Arctic lycodes*.**

Gulf of Georgia and Alaska.

Blennius polaris Sabine, Parry's Jour., Voyage 1819-20, App. CCXII, north Georgia.*Lycodes turneri* Bean, Proc. U. S. Nat. Mus., I, 1878, 463, St. Michaels, Alaska.*Lycodes coccineus* Bean, Proc. U. S. Nat. Mus., IV, 1881, 144, Big Diomed Island, Bering Strait.**3856. *Lycodalepis mucosus* (Richardson).**

Arctic seas.

Lycodes mucosus Richardson, Last Arctic Voyage, 1855, 362, pl. 26, Northumberland Sound.**Genus 1367. *FURCIMANUS* Jordan and Evermann.***Furcella* Jordan and Evermann, Fishes North and Mid. Amer., 1896, 480 (*Lycodes diapterus* Gilbert); name preoccupied.*Furcimanus* Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2472 (*Lycodes diapterus* Gilbert).**3857. *Furcimanus diapterus* (Gilbert).**

Coasts of California and Oregon.

Lycodes diapterus Gilbert, Proc., U. S. Nat. Mus., XIV, 1891, 564, off coast of Oregon at Albatross stations 2892, 2896, and others in 82 to 376 fathoms.**Genus 1368. *LYCODONUS* Goode and Bean.***Lycodonus* Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 208 (*L. mirabilis* Goode and Bean).**3858. *Lycodonus mirabilis* Goode and Bean.**

Off the New England coast.

Lycodonus mirabilis Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 208, lat. 38° 20' 8" N., long. 73° 23' 20" W., in 740 fathoms.**Genus 1369. *LYCONEMA* Gilbert.***Lyconema* Gilbert, Rept. U. S. Fish Com., XIX, 1893, 471 (*L. barbatum* Gilbert).**3859. *Lyconema barbatum* Gilbert.**

Coast of Alaska.

Lyconema barbatum Gilbert, Rept. U. S. Fish Com., XIX, 1893, 471, pl. 35, coast of Alaska, at Albatross station 3129 in 204 fathoms.**Genus 1370. *LYCOGRAMMA* Gilbert.***Lycogramma* Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 364, (*Maynea brunnea* Bean).**3860. *Lycogramma brunnea* (Bean).**

Coast of California to Alaska.

Maynea brunnea Bean, Proc. U. S. Nat. Mus., XIII, 1890, 38, off Queen Charlotte Island in 876 fathoms.**Genus 1371. *MAYNEA* Cunningham.***Maynea* Cunningham, Trans. Linn. Soc. Lond., XXVII, 1871, 471 (*M. patagonica* Cunningham).

3861. *Maynea californica* Gilbert
Santa Barbara Islands.
Maynea californica Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 362,
pl. 19, fig. 17, San Nicolas Island off southern California.
3862. *Maynea bulbiceps* Garman.
Off Bay of Panama in deep water.
Maynea bulbiceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 140,
Pl. E, off Bay of Panama.
- Genus 1372. *BOTHROCARA* Bean.
Bothrocara Bean, Proc. U. S. Nat. Mus., XIII, 1890, 38 (*B. mollis* Bean).
3863. *Bothrocara pusilla* (Bean).
Coast of Alaska north to Unalaska.
Maynea pusilla Bean, Proc. U. S. Nat. Mus., XIII, 1890, 39, Aleutian
Islands, lat. 55° 10' N., long. 160° 18' W., in 110 fathoms.
3864. *Bothrocara mollis* Bean.
North Pacific to southern California.
Bothrocara mollis Bean, Proc. U. S. Nat. Mus., XIII, 1890, 39, off
Queen Charlotte Islands in 876 fathoms.
3865. *Bothrocara remigera* Gilbert.
Monterey Bay.
Bothrocara remigera Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 366,
pl. 20, fig. 19, Monterey Bay in 718 to 756 fathoms.
- Genus 1373. *BOTHROCAROPSIS* Garman.
Bothrocaropsis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 127
(*B. alalonga* Garman).
3866. *Bothrocaropsis alalonga* Garman.
Off Gulf of California in deep water.
Bothrocaropsis alalonga Garman, Mem. Mus. Comp. Zool., XXIV,
1899, 127, XXXII, off Gulf of California.
3867. *Bothrocaropsis rictolata* Garman.
Off Bay of Panama in deep water.
Bothrocaropsis rictolata Garman, Mem. Mus. Comp. Zool., XXIV,
1899, 128, Pl. XXIII, off Bay of Panama.
3868. *Bothrocaropsis elongata* Garman.
Off Bay of Panama in deep water.
Bothrocaropsis elongata Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 129,
Pl. XXXIII, off Bay of Panama.
- Genus 1374. *GYMNELIS* Reinhardt.
Gymnelis Reinhardt, Dansk. Vidensk. Selsk. Afhandl., VII, 1838, 116,
131 (*Ophidium viridis* Fabricius).
Cepolopolis Kaup, Archiv für Natur., 1856, 96 (*Ophidium viridis*
Fabricius).
3869. *Gymnelis viridis* (Fabricius). *Unernak*.
Arctic seas south to Alaska and Nova Scotia.
Ophidium viridis Fabricius, Fauna Grœn., 1780, 141, Greenland.
Ophidium unernak Lacépède, Hist. Nat. Poiss., II, 1800, 280, Greenland.
? *Gymnelis pictus* Günther, Cat., IV, 1862, 324, no locality.
3870. *Gymnelis stigma* (Lay and Bennett).
Kotzebue Sound, Alaska.
Ophidium stigma Lay and Bennett, Zool. Beechey's Voy., 1839, 67,
pl. 20, fig. 1, Kotzebue Sound.
Gymnelis viridis unimaculatus Richardson, Last Arctic Voyage, 1855,
371, Northumberland Sound.
3871. *Gymnelis conorhynchus* Garman.
Off Bay of Panama in deep water.
Gymnelis conorhynchus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
131, Pl. XXXI, off Bay of Panama.

Genus 1375. LYCOCARA Gill.

Uroneles Günther, Cat., IV, 1862, 325 (*Ophidium parrii* Ross); name preoccupied.

Lycocara Gill, Proc. Acad. Nat. Sci. Phila., XXXVI, 1884, 180 (*Ophidium parrii* Ross).

3872. *Lycocara parrii* (Ross).

Baffins Bay.

Ophidium parrii Ross, in Parry's Third Voyage, App., 1826, 109, Baffins Bay.

Genus 1376. MELANOSTIGMA Günther.

Melanostigma Günther, Proc. Zool. Soc. Lond., 1881, 20 (*M. gelatinosum* Günther).

3873. *Melanostigma gelatinosum* Günther.

Deep waters of the western Atlantic; originally known from the Straits of Magellan, but since obtained in various localities from Cape Cod to the West Indies.

Melanostigma gelatinosum Günther, Proc. Zool. Soc. Lond., 1881, 21, pl. 2, fig. A, Straits of Magellan.

3874. *Melanostigma pammelas* Gilbert.

Coast of Alaska.

Melanostigma pammelas Gilbert, Rept. U. S. Fish Com., XIX, 1893, 472, pl. 35, coast of Alaska at Albatross station 3202, lat. 36° 46' 10" N., long. 121° 58' 45" W., in 382 fathoms.

Genus 1377. PHYCOCÆTES Jenyns.

Phycocætes Jenyns, Zool. Beagle, 1842, 168 (*P. latitans* Jenyns).

3875. *Phycocætes suspectus* Garman.

Off Gulf of California in deep water.

Phycocætes suspectus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 137, Pl. XXX, off Gulf of California.

Family 288. LYCODAPODIDÆ**Genus 1378. LYCODAPUS Gilbert.**

Lycodapus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 107 (*L. fierasfer* Gilbert).

3876. *Lycodapus fierasfer* Gilbert.

Pacific coast of America, Lower California to southeastern Alaska.

Lycodapus fierasfer Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 108, off Lower California and Washington at Albatross stations 2980, 3010, and 3072 in 610 to 1005 fathoms.

3877. *Lycodapus dermatinus* Gilbert.

Coast of California in deep water.

Lycodapus dermatinus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 471, pl. 35, coast of California at Albatross station 3162, 37° 54' 10" N., 123° 30' W., in 552 fathoms.

3878. *Lycodapus parviceps* Gilbert.

Bering Sea, north of Unalaska Island.

Lycodapus parviceps Gilbert, Rept. U. S. Fish Com., XIX, 1893, 455, Bering Sea north of Unalaska Island at Albatross station 3324 in 109 fathoms.

3879. *Lycodapus extensus* Gilbert.

Bering Sea north of Unalaska Island.

Lycodapus extensus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 455, Bering Sea north of Unalaska Island at Albatross station 3324 in 109 fathoms.

3880. *Lycodapus lycodon* Gilbert.

Monterey Bay.

Lycodapus lycodon Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 371, Monterey Bay, 152 to 268 fathoms.

3881. *Lycodapus grossidens* Gilbert.
Bering Sea.
Lycodapus grossidens Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915,
373, Funter Bay, southeast Alaska, in 350 fathoms.
3882. *Lycodapus attenuatus* Gilbert.
Monterey Bay.
Lycodapus attenuatus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915,
372, pl. 21, Monterey Bay, in 285 to 357 fathoms.
3883. *Lycodapus mandibularis* Gilbert.
Monterey Bay.
Lycodapus mandibularis Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915,
369, pl. 20, fig. 20, Monterey Bay, in 144 to 293 fathoms.

Family 289. DEREPODICHTHYIDÆ

Genus 1379. DEREPODICHTHYS Gilbert.

Derepodichthys Gilbert, Rept. U. S. Fish Com., XIX, 1893, 456 (*D. alepidotus* Gilbert).

3884. *Derepodichthys alepidotus* Gilbert.
Coast of British Columbia, off Queen Charlotte Island.
Derepodichthys alepidotus Gilbert, Rept. U. S. Fish Com., XIX, 1893,
456, Queen Charlotte Island at Albatross station 3342 in 1,588 fathoms.

Family 290. BROTULIDÆ

Genus 1380. BROTULA Cuvier.

Brotula Cuvier, Règne Anim., ed. 2, II, 1829, 335 (*Enchelyopus barbatus* Bloch and Schneider).

3885. *Brotula barbata* (Bloch and Schneider). *Brotula*.
West Indies.
Enchelyopus barbatus Bloch and Schneider, Syst. Ichth., 1801, 52,
Habana; after Parra.
Brotula Parra, Dif. Piezas Hist. Nat., 1787, 70, lam. 31, fig. 2, Habana.

Genus 1381. STYGICOLA Gill.

Stygicola Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 252 (*Lucifuga dentatus* Poey).

3886. *Stygicola dentatus* (Poey).
Caves in southern Cuba.
Lucifuga dentatus Poey, Memorias, II, 1861, 102, cave of Cajío, Cuba.

Genus 1382. LUCIFUGA Poey.

Lucifuga Poey, Memorias, II, 1860, 95 (*L. subterraneus* Poey).

3887. *Lucifuga subterranea* Poey. *Pez ciego*.
Cuba, in caves.
Lucifuga subterraneus Poey, Memorias, II, 1861, 96, caves of San Antonio in southern Cuba.

Genus 1383. BROSMOPHYCIS Gill.

Brosomphycis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 168
(*Brosmius marginatus* Ayres).
Halias Ayres, Proc. Calif. Acad. Nat. Sci., II, 1861, 52 (*Brosmius marginatus* Ayres).

3888. *Brosomphycis marginatus* (Ayres).
San Francisco to Puget Sound; scarce, recently taken at San Pedro,
Calif.
Brosmius marginatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 13,
Bay of San Francisco.

Genus 1384. OGILBIA Jordan and Evermann.

Ogilbia Jordan and Evermann, in Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 132 (*O. cayorum* Evermann and Kendall).

3889. *Ogilbia cayorum* Evermann and Kendall.
Key West.
Ogilbia cayorum Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 132, pl. 9, fig. 14, Key West, Fla.
3890. *Ogilbia verrillii* (Garman).
Bermudas.
Brosomphycis verrilli Garman, Trans. Conn. Acad. Sci., X, 1900, 511, Bailey Bay, Bermuda Islands.
3891. *Ogilbia ventralis* (Gill).
Lower California, about Cape San Lucas.
Brosomphycis ventralis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 253, Cape San Lucas.
- Genus 1385. BYTHITES Reinhardt.
Bythites Reinhardt, Kongl. Dansk. Vidensk. Selsk. Skrift. Afhandl., VII, 1837, 178 (*B. fuscus* Reinhardt).
3892. *Bythites fuscus* Reinhardt.
The only specimen known, now in the Museum at Copenhagen, was obtained in Greenland nearly a century ago.
Bythites fuscus Reinhardt, Kongl. Dansk. Vidensk. Selsk. Skrift. Afhandl., VII, 1838, 179, Greenland.
- Genus 1386. CATÆTYX Günther.
Catætyx Günther, Challenger Rept., XXII, 1887, 104 (*Siremba messieri* Günther).
3893. *Catætyx rubrirostris* Gilbert.
Coast of California south of Point Conception, in deep water.
Catætyx rubrirostris Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 111, coast of California, at Albatross stations 2909, 2925, and 2936 in 205 and 356 fathoms.
3894. *Catætyx simus* Garman.
Off Galapagos Islands to Bay of Panama, in deep water.
Catætyx simus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, Pl. E 1, 168, off Bay of Panama.
- Genus 1387. DICROMITA Goode and Bean.
Dicromita Goode and Bean, Oceanic Ichth., 1895, 319 (*D. agassizii* Goode and Bean).
3895. *Dicromita agassizii* Goode and Bean.
Granada and Gulf of Mexico.
Dicromita agassizii Goode and Bean, Oceanic Ichth., 1896, 319, fig. 285, Gulf of Mexico in 26 fathoms.
- Genus 1388. MONOMITOPUS Alcock.
Monomitopus Alcock, Ann. Mag. Nat. Hist., ser. 6, VI, 1890, 297 (*Siremba nigripinnus* Alcock).
3896. *Monomitopus torvus* Garman.
Off Bay of Panama in deep water.
Monomitopus torvus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 157, Pl. XL, off Bay of Panama.
- Genus 1389. MONOMEROPUS Garman.
Monomeropus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 157 (*Monomeropus malispinosus* Garman).
3897. *Monomeropus malispinosus* Garman.
Off Bay of Panama in deep water.
Monomeropus malispinosus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 158, Pl. XL, off Bay of Panama.
- Genus 1390. BASSOZETUS Gill.
Bathynectes Günther, Ann. Nat. Hist., II, 1878, 20 (*B. compressus* Günther); name preoccupied.
Bassozetus Gill, Proc. U. S. Nat. Mus., VI, 1883, 259 (*B. normalis* Gill).
Bathyonus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1886, 603 (*Bassozetus catena* Goode and Bean).

3898. *Bassozetes compressus* (Günther).
Gulf of Mexico and elsewhere in deep water.
Bathynectes compressus Günther, Ann. Mag. Nat. Hist., II, 1878, 20,
southeast of New Guinea in 1,400 fathoms.
3899. *Bassozetes normalis* Gill.
West Indies, off Dominica, to 40° north latitude, in region of Gulf
Stream.
Bassozetes normalis Gill, Proc. U. S. Nat. Mus., VI, 1883, 259, Gulf
Stream, at Albatross station 2042, 39° 33' N., 68° 26' 45" W., in
1,555 fathoms.
3900. *Bassozetes catena* (Goode and Bean).
Gulf of Mexico in deep water.
Bathyonus catena Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885,
603, at Albatross station 2379, 28° 00' 15" N., 87° 42' W., in 1,467
fathoms.
3901. *Bassozetes tænia* (Günther).
Mid-Atlantic.
Bathyonus tænia Günther, Challenger Rept., XXII, 1887, 110, pl. 23,
fig. A, Mid-Atlantic.
3902. *Bassozetes nasus* Garman.
Off Gulf of Panama and Pacific coast of southern Mexico in deep water.
Bassozetes nasus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 159,
Pl. LXXVII, off Pacific coast of southern Mexico.
- Genus 1391. *ERETMICHTHYS* Garman.
Eretmichthys Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 164
(*E. pinnatus* Garman).
3903. *Eretmichthys pinnatus* Garman.
Galapagos Islands to Bay of Panama in deep water.
Eretmichthys pinnatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
165, Pl. XXXV, off Bay of Panama.
3904. *Eretmichthys ocella* Garman.
Off Bay of Panama in deep water.
Eretmichthys ocella Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
166, Pl. XXVII, off Bay of Panama.
- Genus 1392. *MÆBIA* Goode and Bean.
Mæbia Goode and Bean, Oceanic Ichth., 1895, 331 (*Bathynectes gracilis*
Günther).
3905. *Mæbia promelas* (Gilbert).
Gulf of California.
Porogadus promelas Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 546,
Gulf of California in 1,005 fathoms.
- Genus 1393. *NEOBYTHITES* Goode and Bean.
Neobythites Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 600
(*N. gillii* Goode and Bean).
3906. *Neobythites gillii* Goode and Bean.
Gulf of Mexico to Brazil, in deep water.
Neobythites gillii Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885,
601, Gulf of Mexico, at Albatross station 2402, 28° 36' N., 85° 33' W.,
in 111 fathoms
Neobythites ocellatus Günther, Challenger Rept., XXII, 1887, 103, pl. 21,
fig. B, off Pernambuco in 350 fathoms.
3907. *Neobythites marginatus* Goode and Bean.
Off Barbados.
Neobythites marginatus Goode and Bean, Bull. Mus. Comp. Zool., XII,
1886, off Barbados in 209 fathoms.
- Genus 1394. *HOLCOMYCTERONUS* Garman.
Holcomycteronus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 162
(*H. digitatus* Garman).

- 3908. *Holcomycteronus digittatus* Garman.**
Gulf of California to Colombia in deep water.
Holcomycteronus digittatus Garman, Mem. Mus. Comp. Zool., XXXVI, 1899, 163, Gulf of California.
- Genus 1395. BENTHOCOMETES Goode and Bean.**
Benthocometes Goode and Bean, Oceanic Ichth., 1895, 327 (*Neobythites robustus* Goode and Bean).
- 3909. *Benthocometes robustus* (Goode and Bean).**
West Indies to lat. 29° N. in Gulf Stream.
Neobythites robustus Goode and Bean, Bull. Mus. Comp. Zool., XII, 1883, 161, off Morro Castle, Cuba.
- Genus 1396. BASSOGIGAS Gill.**
Bassogigas Gill, in Goode and Bean, Oceanic Ichth., 1895, 328 (*B. gillii* Goode and Bean).
- 3910. *Bassogigas gillii* Goode and Bean.**
Off Cape Henlopen, Del.
Bassogigas gillii Goode and Bean, Oceanic Ichth., 1895, 328, fig. 291, off Cape Henlopen, Del., at Albatross station 2684 in 1,106 fathoms.
- 3911. *Bassogigas stelliferoides* (Gilbert)**
Off Bay of Panama in deep water, and Gulf of California.
Neobythites stelliferoides Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 112, off Lower California at Albatross station 2996 in 112 fathoms.
- Genus 1397. BARATHRODEMUS Goode and Bean.**
Barathrodemus Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882, 200 (*B. manatinus* Goode and Bean).
- 3912. *Barathrodemus manatinus* Goode and Bean.**
Gulf Stream, north to Bermudas.
Barathrodemus manatinus Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882, 200, lat. 33° 51' 20" N., long. 76° W., in 647 fathoms.
- Genus 1398. NEMATONUS Günther.**
Nematonus Günther, *Challenger* Rept., XXII, 1887, 114 (*Bathyonus pectoralis* Goode and Bean).
- 3913. *Nematonus pectoralis* (Goode and Bean).**
Gulf of Mexico and off Dominica.
Bathyonus pectoralis Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 604, Albatross station 2380, lat. 28° 02' 30" N., long. 87° 43' 45" W., in 1,430 fathoms.
- Genus 1399. POROGADUS Goode and Bean.**
Porogadus Goode and Bean Proc. U. S. Nat. Mus., VIII, 1885, 602 (*P. miles* Goode and Bean).
- 3914. *Porogadus miles* Goode and Bean.**
Atlantic coast, off Delaware Bay, in deep water.
Porogadus miles Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 602, at Albatross station 2230, lat. 38° 27' N., long. 73° 2' W., in 1,168 fathoms.
- 3915. *Porogadus longiceps* Garman.**
Off Cocos Islands to Bay of Panama in deep water.
Porogadus longiceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 153, Pl. F, off Bay of Panama.
- 3916. *Porogadus atripectus* Garman.**
Off Bay of Panama and Pacific coast of Mexico in deep water.
Porogadus atripectus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 154, Pl. XXXVII, off Bay of Panama.
- 3917. *Porogadus breviceps* Garman.**
Gulf of California in deep water.
Porogadus breviceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 155, Gulf of California.
- Genus 1400. PENOPUS Goode and Bean.**
Penopus Goode and Bean, Oceanic Ichth., 1895, 335 (*P. macdonaldi* Goode and Bean).

3918. *Penopus macdonaldi* Goode and Bean.

Gulf Stream.

Penopus macdonaldi Goode and Bean, Oceanic Ichth., 1895, 336, fig. 293, Gulf Stream at Albatross station 2716, lat. 38° 29' 30" N., long. 70° 57' W., in 1,631 fathoms.**Genus 1401. *DICROLENE* Goode and Bean.***Dicrolene* Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882, 202 (*D. intronigra* Goode and Bean).**3919. *Dicrolene intronigra* Goode and Bean.**

Gulf Stream, from lat. 15° to 40° N.

Dicrolene intronigra Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882, 202, Gulf Stream, lat. 34° N.**3920. *Dicrolene pullata* Garman.**

Off Gulf of Panama in deep water.

Dicrolene pullata Garman, Mem. Comp. Zool., XXIV, 1899, 152, off Gulf of Panama.**3921. *Dicrolene nigra* Garman.**

Off Galapagos Islands to Bay of Panama in deep water.

Dicrolene nigra Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 150, Pl. XXXVII, off Galapagos Islands.**3922. *Dicrolene filamentosa* Garman.**

Off Bay of Panama and Pacific coast of Mexico in deep water.

Dicrolene filamentosa Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 149, Pl. F, off Bay of Panama.**Genus 1402. *MIXONUS* Günther.***Mixonus* Günther, Challenger Rept., XXII, 1887, 108 (*Bathynectes laticeps* Günther).**3923. *Mixonus laticeps* (Günther).**

Mid-Atlantic in deep water.

Bathynectes laticeps Günther, Ann. Mag. Nat. Hist., II, 1878, 20, Mid-Atlantic at Challenger station 104 in 2,500 fathoms.*Siremo guentheri* Vaillant, Expd. Sci. Travailleur et Talisman, 1888, 268, Pl. XXIV, fig. 5, off the Cape Verde Islands at a depth of 3,200 meters.**3924. *Mixonus caudalis* Garman.**

Between Galapagos and Cocos Islands in deep water.

Mixonus caudalis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 148, Pl. XXXVI, off Cocos Islands.**Genus 1403. *ACANTHONUS* Günther.***Acanthonus* Günther, Ann. Mag. Nat. Hist., II, 1878, 22 (*A. armatus* Günther).**3925. *Acanthonus spinifer* Garman.**

Off Bay of Panama in deep water.

Acanthonus spinifer Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 170, Pl. F, off Bay of Panama.**Genus 1404. *LEUCICORUS* Garman.***Leucicorus* Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 146 (*L. lusciosus* Garman).**3926. *Leucicorus lusciosus* Garman.**

Off Pacific coast of Mexico in deep water.

Leucicorus lusciosus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 146, Pl. XXXVIII, off Pacific coast of Mexico.**Genus 1405. *SCIADONUS* Garman.***Sciadonus* Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 171 (*S. pedicellaris* Garman).**3927. *Sciadonus pedicellaris* Garman.**

Off Cocos Island in deep water.

Sciadonus pedicellaris Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 172, Pl. F, off Cocos Island.

- Genus 1406. **LAMPROGRAMMUS** Alcock.
Lamprogrammus Alcock, Ann. Mag. Nat. Hist., VIII, 1891, 32 (*L. niger* Alcock).
3928. **Lamprogrammus illustris** Garman.
 Off Bay of Panama.
Lamprogrammus illustris Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 174, Pl. XXXIV, off Bay of Panama.
- Genus 1407. **BARATHRONUS** Goode and Bean.
Barathronus Goode and Bean, Mem. Mus. Comp. Zool., XII, 1886, 164 (*B. bicolor* Goode and Bean).
3929. **Barathronus bicolor** Goode and Bean.
 Off Guadalupe Island in deep water.
Barathronus bicolor Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, 164, off Guadalupe at Blake station 71 in 769 fathoms.
- Genus 1408. **APHYONUS** Günther.
Aphyonus Günther, Ann. Mag. Nat. Hist., II, 1878, 22 (*A. gelatinosus* Günther).
3930. **Aphyonus mollis** Goode and Bean.
 Gulf of Mexico in deep water.
Aphyonus mollis Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, 163, Gulf of Mexico at Blake station 221, lat. 24° 36' N, long. 84° 5' W., in 955 fathoms.
- Genus 1409. **ALLECTOR** Heller and Snodgrass.
Allector Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 228 (*A. chelonix* Heller and Snodgrass).
3931. **Allector chelonix** Heller and Snodgrass.
 Between Clipperton Island and Galapagos Archipelago.
Allector chelonix Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 228, fig. 1, between Clipperton Island and the Galapagos.
- Family 291. **OPHIDIIDÆ**
- Genus 1410. **LEPOPHIDIUM** Gill.
Lepophidium Gill, Am. Nat., XXIX, 1895, 16; substitute for *Leptophidium* Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 210 (*L. profundorum* Gill); preoccupied in serpents by *Leptophidium* Halliwell, 1860.
3932. **Lepophidium marmoratum** (Goode and Bean).
 Gulf Stream.
Lepophidium marmoratum Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 423, lat. 23° 10' 39" N., long. 82° 20' 21" W., in 213 fathoms.
3933. **Lepophidium profundorum** (Gill).
 Gulf Stream, off coast of Florida.
Leptophidium profundorum Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 211, Gulf Stream, off coast of Florida.
3934. **Lepophidium cervinum** (Goode and Bean).
 Gulf Stream.
Lepophidium cervinum Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 422, lat. 40° 01' N., long. 69° 56' W., depth 76 fathoms.
3935. **Lepophidium brevibarbe** (Cuvier).
 West Indies and Brazil.
Ophidium brevibarbe Cuvier, Règne Anim., ed. 2, II, 1829, 359, Brazil.
3936. **Lepophidium prorates** (Jordan and Bollman).
 Panama and south of Panama.
Leptophidium prorates Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 172, Panama and south of Panama.
3937. **Lepophidium emmelas** (Gilbert).
 Coast of Lower California.
Leptophidium emmelas Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 110, coast of Lower California at Albatross stations 3007 and 3008 in 362 and 306 fathoms.

3938. *Lepophidium pardale* (Gilbert).
Lower California.
Leptophidium pardale Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 108,
off Lower California.
3939. *Lepophidium microlepis* (Gilbert).
Off Lower California
Leptophidium microlepis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890,
109, off Lower California.
3940. *Lepophidium stigmatistium* (Gilbert).
Coast of Lower California.
Leptophidium stigmatistium Gilbert, Proc. U. S. Nat. Mus., XIII, 1890,
109, off Lower California at Albatross station 2996 in 112 fathoms.
- Genus 1411. *OPHIDION* Linnæus. *Cusk eels*.
Ophidion Linnæus, Syst. Nat., Ed. X, 1758, 259 (*Ophidion barbatum*
Linnæus).
Ophidium of most authors; revised spelling.
3941. *Ophidion graëllsi* Poey.
Cuba.
Ophidion graëllsi Poey, Memorias, II, 1861, 425, Habana
3942. *Ophidion beani* Jordan and Gilbert.
Gulf of Mexico; Snapper Banks off Pensacola, Fla.
Ophidion beani Jordan and Gilbert, Proc. U. S. Nat. Mus., VI, 1883, 143,
Snapper Banks off Pensacola.
3943. *Ophidion holbrooki* (Putnam).
Key West, Fla.; Gulf of Mexico.
Ophidium holbrooki Putnam, Proc. Bost. Soc. Nat. Hist., XVI, 1874,
342, Key West, Fla.
- Genus 1412. *RISSOLA* Jordan and Evermann.
Rissola Jordan and Evermann, Check List, 1895, 483 (*Ophidium mar-*
ginatum DeKay).
3944. *Rissola marginata* (DeKay). *Sand eel; Slippery dick*.
South Atlantic coast of United States from New York south to Pensa-
cola.
Ophidium marginatum DeKay, New York Fauna: Fishes, 1842, 315,
New York Harbor.
? *Ophidium josephi* Girard, U. S. and Mex. Bound. Surv., Zool., 1859,
29, Saint Joseph Island, Tex.
- Genus 1413. *OTOPHIDIUM* Gill.
Otophidium Gill, in Jordan, Cat. Fish. N. Amer., 1885, 914 (*Genypterus*
omostigma Jordan and Gilbert).
Chilaria Jordan and Evermann, Check List, 1895, 482 (*Ophidium taylori*
Girard).
3945. *Otophidium taylori* (Girard).
Coast of California, from Monterey to San Diego.
Ophidium taylori Girard, Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 138,
Monterey, Calif.
3946. *Otophidium omostigma* (Jordan and Gilbert).
Gulf of Mexico, Pensacola snapper banks.
Genypterus omostigma Jordan and Gilbert, Proc. U. S. Nat. Mus., V,
1882, 301, Pensacola snapper banks.
3947. *Otophidium scrippsii* Hubbs.
Long Beach to Cedros Island, Calif.
Ophidium scrippsii Hubbs, Univ. Calif. Publ. Zool., XVI, 1916, 166,
pl. 20, Cedros Island, off coast of Lower California.
3948. *Otophidium galeoides* (Gilbert).
Coast of Lower California.
Otophodon galeoides Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 110,
off Lower California.

3949. *Otophidium welshi* Nichols and Breder.

Coast of Texas.

Otophidium welshi Nichols and Breder, Proc. Biol. Soc. Wash., XXXV, 1922, 15, fig. 3, Texas, *Grampus* station 10478, southwest from Brayes Light.**Family 292. CARAPIDÆ****Genus 1414. CARAPUS Rafinesque.***Carapus* Rafinesque, Indice, 1810, 57, restricted by the International Commission of Nomenclature to *Ophidium imberbe*⁹⁰ Linnæus.*Fierasfer* Cuvier, Règne Anim., ed. 1, II, 1817, 239 (*Ophidium imberbe* Linnæus).*Echadon* Thompson, Proc. Zool. Soc. Lond., 1837, 55 (*E. drummondi* Thompson).*Diaphasia* Lowe, loc. cit., 1843, 92 (*Gymnotus acus* Brunnich).*Oxybeles* Richardson, Voy. *Erebus* and *Terror*, Fishes, 1844-48, 73 (*O. homei* Richardson).*Porobronchus* Kaup, Ann. Mag. Nat. Hist., 1860, 272 (*P. linearis* Kaup; larva of *Fierasfer acus*).*Vexillifer* Gasco, Bull. Assoc. Nat. Med. Napoli, 1870, 59 (*V. dephillippi* Gasco; larva of *Fierasfer acutus*).*Lefroyia* Jones, Zoologist, IX, 1874, 3838 (*L. bermudensis* Jones).**3950. *Carapus affinis* (Günther). Pearlfish.**

Lower California to Panama; abundant about the Pearl Islands off Panama, north to Point Loma, San Diego, Calif.

Fierasfer affinis Günther, Cat., IV, 1862, 381, no locality given.*Fierasfer dubius* Putnam, Proc. Bost. Soc. Nat. Hist., XVI, 1874, 344, Pearl Islands.**3951. *Carapus arenicola* (Jordan and Gilbert).**

Pacific coast of Mexico

Fierasfer arenicola Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 363, Mazatlan, Mexico.**3952. *Carapus bermudensis* (Jones).**

West Indies; recorded from Bermuda, Habana, Key West, and St. Thomas.

Lefroyia bermudensis Jones, Zoologist, IX, 1874, 3837, Bermuda.**Genus 1415. LEPTOFIERASFER Meek and Hildebrand.***Leptofierasfer* Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1928, 964 (*L. macrurus* Meek and Hildebrand).**3953. *Leptofierasfer macrurus* Meek and Hildebrand.***Leptofierasfer macrurus* Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1928, 964, Pl. XCIX, fig. 1, Chame Point, Panama.**Suborder HAPLODOCI****Family 293. BATRACHOIDIDÆ****Genus 1416. BATRACHOIDES Lacépède.***Batrachoides* Lacépède, Hist. Nat. Poiss., II, 1800, 451 (*B. tau* Lacépède); not *Gadus tau* Linnæus = *Batrachus surinamensis* Schneider.*Batrachus* Schneider, ed. Bloch, Syst. Ichth., 1801, 42 (*B. surinamensis* Schneider); substitute for *Batrachoides*.*Batrictius* Rafinesque, Anal. Nat., 1815, 82; substitute for *Batrachoides*.**3954. *Batrachoides surinamensis* (Schneider). Sapo.**

Coast of Guiana and Brazil.

Batrachus surinamensis Schneider, ed. Bloch, Syst. Ichth., 1801, 43, Surinam; from a specimen in the museum of Paris.⁹⁰ With this decision we do not willingly concur, as the name *Carapus* is clearly cognate with *Gymnotus carapo*, a species of Gymnotidae included but not named in the original account.

3955. *Batrachoides goldmani* Evermann and Goldsborough.
Rio Usumacinta at Montecristo, Tabasco, Mexico.
Batrachoides goldmani Evermann and Goldsborough, Bull. U. S. Fish
Com., XXI, 1901, 159, fig. 8, Rio Usumacinta, Mexico.
3956. *Batrachoides boulengeri* Gilbert and Starks.
Bay of Panama.
Batrachoides boulengeri Gilbert and Starks, Mem. Calif. Acad. Sci.,
IV, 1904, 182, Pl. XXXI, Bay of Panama.
3957. *Batrachoides pacifici* (Günther).
Panama.
Batrachus pacifici Günther, Cat., III, 1861, 173, Panama.
- Genus 1417. OPSANUS Rafinesque. *Toadfishes*.
Opsanus Rafinesque, Amer. Month. Mag., II, 1818, 203 (*O. cerapalus*
Rafinesque=*Gadus tau* Linnæus).
Batrachus of authors, not of Schneider.
3958. *Opsanus tau* (Linnæus). *Toadfish*; *Sapo*; *Slimer*; *Oysterfish*; *Scorpion*.
Atlantic coast, Massachusetts to the West Indies.
Gadus tau Linnæus, Syst. Nat., ed. XII, 1766, 439, Charleston, S. C.
Cottus glaber Schöpfung, Schrift. Naturf. Freunde, VII, 1788, 146, Long
Island.
Cottus chætodon Bloch and Schneider, Syst. Ichth., 1801, 62, New York;
after Schöpfung.
Lophius bufo Mitchell, Trans. Lit. Phil. Soc., I, 1815, 463, New York.
Opsanus cerapalus Rafinesque, Amer. Month. Mag., II, 1818, 204,
south coast of Long Island.
Batrachoides vernullas LeSueur, Mém. Mus. Hist. Nat., V, 1819, 157,
pl. 17, coast of Rhode Island.
Batrachoides variegata LeSueur, Jour. Acad. Nat. Sci. Phila., III, pt. 2,
1824, 398, Egg Harbor, N. J.
Batrachus celatus DeKay, New York Fauna: Fishes, 1842, 170, pl. 50,
fig. 161, New York.
3959. *Opsanus hildebrandi* Breder.
Colon Market, Panama.
Opsanus hildebrandi Breder, Amer. Mus. Novit., No. 188, Sept. 28,
1925, I, Colon Market, Panama; place of collection unknown.
3960. *Opsanus pardus* (Goode and Bean). *Sapo*.
Gulf of Mexico in deep water.
Batrachus tau pardus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879,
336, Pensacola snapper banks
- Genus 1418. PORICHTHYS Girard. *Midshipmen*.
Porichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 141 (*P.*
notatus Girard).
3961. *Porichthys notatus* Girard. *Singingfish*; *Midshipman*; *Cabezon*; *Sapo*.
Pacific coast from Puget Sound to Panama.
Porichthys notatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 141,
San Francisco.
3962. *Porichthys margaritatus* (Richardson).
Pacific coast of tropical America.
Batrachus margaritatus Richardson, Voy. Sulphur, Fishes, 1844-1843,
67, Pacific coast of Central America.
Porichthys nautopædium Jordan and Bollman, Proc. U. S. Nat. Mus.,
XII, 1889, 171, Pacific Ocean off coast of Colombia in 16 fathoms.
3963. *Porichthys greeniei* Gilbert and Starks.
Panama Bay.
Porichthys greeniei Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904,
184, Panama Bay.
- Genus 1419. NAUTOPÆDIUM Jordan.
Nautopædium Jordan, Proc. Acad. Nat. Sci. Phila., 1919, 342 (*Batrachus*
porosissimus Cuvier and Valenciennes).

3964. *Nautopædium porosissimum* (Cuvier and Valenciennes). *Bagre sapo*. South Carolina to Montevideo; Pensacola.
Batrachus porosissimus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 373 (501), Surinam, Cayenne, Rio Janeiro, and St. Catherine.
Porichthys plectrodon Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 291, Galveston, Tex.

Genus 1420. **THALASSOPHRYNE** Gunther. *Poison toadfishes*.
Thalassophryne Gunther, Cat., III, 1861, 175 (*T. maculosa* Gunther).
Dæctor Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2325 (*Thalassophryne dowi* Jordan and Gilbert).

Subgenus **THALASSOPHRYNE**.

3965. *Thalassophryne maculosa* Gunther.
 Caribbean Sea.
Thalassophryne maculosa Gunther, Cat., III, 1861, 175, Puerto Cabello near Aspinwall.

3966. *Thalassophryne reticulata* Gunther.
 Panama.
Thalassophryne reticulata Gunther, Proc. Zool. Soc. Lond., 1864, 150, 155, Panama.

Subgenus **DÆCTOR** Jordan and Evermann.

3967. *Thalassophryne dowi* (Jordan and Gilbert).
 Pacific coast of North America from Punta Arenas to Panama.
Thalassophryne dowi Jordan and Gilbert, Proc. U. S. Nat. Mus., X, 1887, 388, Punta Arenas.

Order YY. XENOPTERYGII

Family 294. GOBIESOCIDÆ

Genus 1421. **CAULARCHUS** Gill.
Caularchus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Lepadogaster mæandricus* Girard=*Lepadogaster reticulatus* Girard); name preoccupied.

3968. *Caularchus mæandricus* (Girard). *Suckfish*.
 Pacific coast of United States, from Vancouver Island to Monterey.
Lepadogaster reticulatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 155, San Luis Obispo, Calif.; name preoccupied.
Lepadogaster mæandricus Girard, Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 130, San Luis Obispo, Calif., and the Farallons.

Genus 1422. **BRYSSETÆRES** Jordan and Evermann.
Bryssætæres Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 230 (*Gobiesox pinniger* Gilbert).

3969. *Bryssætæres pinniger* (Gilbert).
 Gulf of California; known from Puerto Refugio (Angel Island), San Luis Gonzales Bay, and La Paz.
Gobiesox pinniger Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 94, Puerto Refugio, Gulf of California.

Genus 1423. **CAULISTIUS** Jordan and Evermann.
Caulistius Jordan and Gilbert, Check List, 1895, 491 (*Gobiesox papillifer* Gilbert).
Bryssophilus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2329 (*Gobiesox papillifer* Gilbert); accidental duplication overlooked in proof reading.

3970. *Caulistius papillifer* (Gilbert).
 Magdalena Bay, Lower California.
Gobiesox papillifer Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 96, Magdalena Bay, Lower California.

Genus 1424. **GOBIESOX** Lacépède. *Clingfishes*.
Gobiesox Lacépède, Hist. Nat. Poiss., II, 1800, 595 (*G. cephalus* Lacépède).
Megaphalus Rafinesque, Anal. Nat., 1815, 86 (substitute for *Gobiesox* Lacépède).
Sicyogaster Brisout de Barneville, Rev. Zool., 1846, 144 (*Gobiesox marmoratus* Jenyns).

3971. *Gobiesox nigripinnis* (Peters).
Puerto Cabello, near Aspinwall, Colon.
Cotylin nigripinnis Peters, Monatsber. Akad. Wiss. Berlin, 1859, 412,
Puerto Cabello, near Aspinwall, Colon.
3972. *Gobiesox macrophthalmus* Günther.
Probably West Indies.
Gobiesox macrophthalmus Gunther, Cat., III, 1861, 502, locality
unknown.
3973. *Gobiesox androsiensis* Rosen.
Bahamas
Gobiesox androsiensis Rosen, Arsskr. Univ. Lund., VII, 1912, 50,
Bahamas.
3974. *Gobiesox costaricensis* Meek.
Costa Rica.
Gobiesox costaricensis Meek, Field Mus. Nat. Hist., Zool., X, 1912, 74,
Zent, Costa Rica.
3975. *Gobiesox gyrinus* Jordan and Evermann.
West Indies.
Gobiesox gyrinus Jordan and Evermann, Fishes North and Mid Amer.,
1898, 2331, Cordova; name only in Check List, 1895, 491.
3976. *Gobiesox cephalus* Lacépède. *Têtard; Testar*.
West Indies.
Gobiesox cephalus Lacépède, Hist. Nat. Poiss., II, 1800, 595, Mar-
tinique.
Lepadogaster testar Bloch and Schneider, Syst. Ichth., 1801, 4, Mar-
tinique; after Plumier.
3977. *Gobiesox tudes* Richardson.
Probably West Indies.
Gobiesox tudes Richardson, Voy. Sulphur, Fishes, 1845, 103, pl. 46,
figs. 1-3; habitat unknown, erroneously supposed to be China.
3978. *Gobiesox strumosus* Cope.
Maryland to Florida.
Gobiesox strumosus Cope, Proc. Acad. Nat. Sci. Phila., XXII, 1870,
121, Hilton Head, S. C.
3979. *Gobiesox virgatulus* Jordan and Gilbert.
From Pensacola Bay north to Charleston, S. C.
Gobiesox virgatulus Jordan and Gilbert, Proc. U. S. Nat. Mus., V,
1882, 293, Pensacola, Fla.
3980. *Gobiesox adustus* Jordan and Gilbert.
Pacific coast of Mexico.
Gobiesox adustus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
360, Mazatlan, Mexico.
3981. *Gobiesox funebris* Gilbert.
Gulf of California; Puerto Refugio (Angel Island) and La Paz.
Gobiesox funebris Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 95, Puerto
Refugio, Gulf of California.
3982. *Gobiesox rhodospilus* Günther.
Panama.
Gobiesox rhodospilus Gunther, Proc. Zool. Soc. Lond., 1864, 25, Panama.
3983. *Gobiesox cerasinus* Cope.
St. Martins, West Indies.
? *Cyclopterus nudus* Linnæus, Syst. Nat., ed. X, 1758, 260, "India";
not identifiable.
Gobiesox cerasinus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 473, St.
Martins, West Indies.
3984. *Gobiesox vittatus* Metzelaar.
Caribbean Sea.
Gobiesox vittatus Metzelaar, Marine Fish. Lesser Antilles, Bijdr.
Dierk., 1922, 140, with fig., Curaçao.

3985. *Gobiesox sancti-martini* Metzelaar.
West Indies.
Gobiesox sancti-martini Metzelaar, Trop. Atl. Vissch., 1919, 154, fig. 48,
St. Martin's Island.
3986. *Gobiesox yuma* Nichols.
Coast of Florida.
Gobiesox yuma Nichols, Bull. Amer. Mus. Nat. Hist., XXXVII, Dec.
29, 1917, 876, fig. 1, south of Sanibel Light, Florida west coast.
- Genus 1425. *SICYASES* Müller and Troschel.
Sicyases Muller and Troschel, Wiegmann's Archiv, 1843, 298, *S.*
sanguineus.
Tomocodon Brisout de Barneville, Rev. Zool., 1846, 144 (*T. chilensis*
Barneville).
3987. *Sicyases erythroptus* (Jordan and Gilbert).
Mazatlan; also Tres Marias Islands.
Gobiesox erythroptus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV,
1881, 360, Mazatlan, Mexico.
3988. *Sicyases rubiginosus* Poey.
Matanzas, Cuba
Sicyases rubiginosus Poey, Synop., 1868, 391, wharves of Palmasola,
Matanzas, Cuba.
3989. *Sicyases carneus* Poey.
Matanzas, Cuba.
Sicyases carneus Poey, Synop., 1868, 392, wharves of Palmasola,
Matanzas, Cuba.
3990. *Sicyases hæres* (Jordan and Bollman).
Green Turtle Cay, Bahamas.
Gobiesox hæres Jordan and Bollman, Proc. U. S. Nat. Mus., XI, 1888,
552, Green Turtle Cay, Bahamas.
3991. *Sicyases punctulatus* Poey.
Cuba.
Sicyases punctulatus Poey, Enumeratio, 1876, 124, Habana.
3992. *Sicyases fasciatus* Peters.
Puerto Cabello near Aspinwall.
Sicyases fasciatus Peters, Monatsber. Akad. Wiss. Berlin, 1859, 412,
Puerto Cabello.
- Genus 1426. *RIMICOLA* Jordan and Evermann.
Rimicola Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI,
1896, 231 (*Gobiesox muscarum* Meek and Pierson).
3993. *Rimicola eigenmanni* (Gilbert).
Point Loma, near San Diego, to Monterey Bay, Calif.
Gobiesox eigenmanni Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 96,
Point Loma near San Diego, Calif.
Gobiesox muscarum Meek and Pierson, Proc. Calif. Acad. Sci., 2d ser.,
V, 1895, 571, pl. 71, Monterey Bay, Calif.
3994. *Rimicola beryllinus* (Hildebrand and Ginsberg).
Key West, Fla.
Gobiesox (Rimicola) beryllinus Hildebrand and Ginsberg, Bull. U. S.
Bur. Fish., XLII, 1926, 213, fig. 5, Boca China, Key West, Fla.
- Genus 1427. *ARBACIOSA* Jordan and Evermann.
Arbaciosa Jordan and Everman, Proc. Calif. Acad. Sci., 2d ser., VI,
1896, 230 (*Gobiesox humeralis* Gilbert).
3995. *Abraciosa rhessodon* (Rosa Smith).
San Diego to the northern part of the Gulf of California.
Gobiesox rhessodon Rosa Smith (Mrs. Eigenmann), Proc. U. S. Nat.
Mus., IV, 1881, 140, San Diego, Calif.
3996. *Abraciosa humeralis* (Gilbert).
Gulf of California; Puerto Refugio, Angel Island; La Paz.
Gobiesox humeralis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 95,
Puerto Refugio, Gulf of California.

3997. *Arbacia rupestris* (Poey).
Coral reefs of Cuba.
Gobiesox rupestris Poey, Memorias, II, 1861, 283, Cuba.
3998. *Arbacia zebra* (Jordan and Gilbert).
Mazatlan, Mexico.
Gobiesox zebra Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 359, Mazatlan, Mexico.
3999. *Arbacia eos* (Jordan and Gilbert).
Pacific coast of Mexico.
Gobiesox eos Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 360, Mazatlan, Mexico.

Order ZZ. PLECTOGNATHI

Suborder SCLERODERMI

Family 295. TRIACANTHIDÆ

- Genus 1428. *HOLLARDIA* Poey.
Hollardia Poey, Memorias, II, 1861, 348 (*H. hollardi* Poey).

4000. *Hollardia hollardi* Poey.
West Indies.
Hollardia hollardi Poey, Memorias, II, 1861, 348, pl. 18, fig. 11, Cuba

Family 296. BALISTIDÆ

- Genus 1429. *BALISTES* Linnæus.
Balistes Linnæus, Syst. Nat., ed. X, 1758, 327 (*B. vetula* Linnæus).
Capriscus Rafinesque, Indice, 1810, 41, 58 (*C. porcus* Rafinesque = *Balistes capriscus* Gmelin).

4001. *Balistes naufragium* Jordan and Starks. *Pez puerco de piedra*.
Mazatlan; Panama Bay.
Balistes naufragium Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 488, Mazatlan.
4002. *Balistes capriscus* Gmelin. *Leatherjacket; Cucuyo; Turbo; Triggerfish; Filefish*.
Tropical parts of the Atlantic; Gulf Stream; Mediterranean.
Balistes carolinensis Gmelin, Syst. Nat., I, 1788, 1468, Carolinas.
Balistes capriscus Gmelin, loc. cit., 1789, 1471, Indian and American oceans; after Gronow (this name preferred by first reviewer).
Balistes bumwa Lacépède, Hist. Nat. Poiss., V, 1798, 669, pl. 21, fig. 1, Nice.
Balistes caprinus Valenciennes, Ichth. Îles Canaries, 1836, 94, pl. 16, Canaries.
Balistes fuliginosus DeKay, N. Y. Fauna: Fishes, 1842, 339, pl. 57, fig. 188, New York.
Balistes tænopterus Poey, Memorias, II, 1861, 326, Cuba.
4003. *Balistes vetula* Linnæus. *Oldwife; Old wench; Cochino; Blue-striped triggerfish; Bessy cerka*.
Tropical parts of the Atlantic; West Indies; Gulf Stream to Woods Hole.
Guaperva Marcgrave, Hist. Bras., 1648, 163, Brazil.
Balistes vetula Linnæus, Syst. Nat., Ed. X, 1758, 329, Ascension Island; after *Balistes vetula* of Osbeck, Iter Chinensis, 294, 1757.
Balistes bellus Walbaum, Artedi Pisc., III, 1792, 467, West Indies; after Krøyer.
Balistes equestris Gronow, Cat. Fishes, ed. Gray, 1854, 31, American seas.

4004. *Balistes forcipatus* Gmelin.

West coast of Africa to Rhode Island and Brazil.

Balistes forcipatus Gmelin, Syst Nat, I, 1789, 1472, Brazil; after *Guaperva lata* of Lista

Balistes punctatus Gmelin, loc. cit., 1472; after Stipvisch of Willoughby.

Balistes spilopterygius Walbaum, Artdi Pisc., III, 1792, 455, Brazil.

Balistes guttatus Walbaum, loc cit, 467, St. Vincent.

Balistes ciliaris Bloch and Schneider, Syst. Ichth., 1801, 471, Brazil.

Balistes lberiensis Steindachner Ichth. Notizen, IV, 1867, 9, Monrovia in Liberia.

?*Balistes powelli* Cope, Proc. Acad. Nat. Sci. Phila., XXII, 1870, 120, Newport, R. I.

Balistes moribundus Cope, Trans Amer. Phil. Soc., XIV, 1870, 478, St. Martins.

Genus 1430. *SUFFLAMEN* Jordan.

Pachynathus Swainson, Classn Fishes, II, 1839, 326 (*Balistes triangularis*=*Balistes capistratus* Shaw); not *Pachygnatha* nor *Pachygnathus*; both these names earlier used for genera of spiders, the name *Pachygnathus* regarded as a misprint and not valid according to decision of the International Commission of Nomenclature.

Sufflamen Jordan, Copeia, April 12, 1916, 27 (*Balistes capistratus* Shaw).

4005. *Sufflamen verres* (Gilbert and Starks). *Pez puerco*.

Pacific coast of tropical America, from Magdalena Bay to the Galapagos.

Balistes capistratus of authors; not of Shaw, Gen. Zool., V, 1804, 417; after *Balistes bridé* Lacépède, a species of the South Seas and East Indies.

Balistes verres Gilbert and Starks, Mem Calif Acad. Sci., V, 1904, 153, Pl. XXVI, fig 49, Panama and Mazatlan.

Genus 1431. *VERRUNCULUS* Jordan.

Verrunculus Jordan, Copeia, Sept, 1924, 82 (*Balistes polylepis* Steindachner).

4006. *Verrunculus polylepis* (Steindachner) *Pez puerco*.

San Diego to Panama; common southward.

Balistes polylepis Steindachner, Ichth. Beitr., V, 1876, 21, Magdalena Bay; Mazatlan; Acapulco.

Genus 1432. *CANTHIDERMIS* Swainson. *Sobacos*.

Canthidermis Swainson, Nat. Hist. Classn. Anim., II, 1839, 325, (*Balistes angulosus* Quoy and Gaimard).

4007. *Canthidermis angulosus* (Quoy and Gaimard).

Hawaii to Cocos Island; rare.

Balistes angulosus Quoy and Gaimard, Voy. Uranie, Zool., 1824, 210, Sandwich Islands (Hawaiian Islands).

4008. *Canthidermis sobaco* (Poey). *Sobaco*.

West Indies.

Balistes sobaco Poey, Memorias, II, 1861, 324, Habana.

Balistes asperrimus Cope, Trans. Amer. Phil. Soc., XIV, 1870, 478, St. Martins.

4009. *Canthidermis sufflamen* (Mitchill). *Sobaco*.

West Indies; Habana.

Sobaco Parra, Dif. Piezas, Hist. Cuba, 1787, 17, fig. 10, Habana.

Balistes sufflamen Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 467, locality unknown, said by DeKay to be from the south Atlantic.

Balistes macrops Poey, Memorias, II, 1861, 326, Habana.

4010. *Canthidermis maculatus* (Bloch) *Ocean turbot*.
West Indies.
Balistes maculatus Bloch, Nat. Ausl. Fisch., II, 1786, 25, West Indies.
Balistes americanus Gmelin, Syst. Nat., I, 1789, 1473, America; after Gronow.
Balistes macropterus Walbaum, Arted. Pisc., III, 1792, 465, America.
Balistes rufus Gronow, Cat. Fishes, ed. Gray, 1854, 36, American oceans.
Balistes longus Gronow, loc. cit., 1854, American oceans.
Balistes melanopterus Cope, Trans. Amer. Phil. Soc., XIV, 1870, 478, Darien.
4011. *Canthidermis willughbeii* (Lay and Bennett).
Pacific coast of Mexico
Balistes willughbeii Lay and Bennett, Zoology Beechey's Voyage, 1839, 68, pl. 21, fig. 2, Acapulco, Mexico.
- Genus 1433. **XANTHICHTHYS** Kaup.
Xanthichthys Kaup, in Richardson, Enc. Brit., ed. XII, 1856, 313
Balistes curassavicus Gmelin).
4012. *Xanthichthys ringens* (Linnæus). *Cocuyo*.
West Indies and southward, recorded from Mauritius.
Balistes ringens Linnæus, Syst. Nat., Ed. X, 1758, 329, no locality.
Cucuyo Parra, Dif. Piezas, etc., Cuba, 1787, 19, Habana, after Linnæus.
Balistes curassavicus Gmelin, Syst. Nat., I, 1788, 1472, Curaçao.
Balistes notatus Gronow, Cat. Fishes, ed. Gray, 1854, 36, both Indies.
Balistes nudus Gronow, loc. cit., 36, American oceans.
Balistes cicatricosus Poey, Memorias, II, 1861, 327, Cuba.
Balistes heckelii von Müller, Reisen in Vereinigten Staaten, Canada und Mexico, I, 1864, 182, Mexico.
4013. *Xanthichthys lineopunctatus* (Hollard).
Panama; Pacific coast of Mexico to East Indies, taken at San Diego.
Balistes lineopunctatus Hollard, Ann. Sci. Nat., ser. 4, I, 1854, 65, Bourbon Island.
Balistes mento Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 228
Clarion Island of the Revillagigedo group.
- Genus 1434. **MELICHTHYS** Swainson.
Melichthys Swainson, Class. Anim., II, 1839, 325 (*Balistes ringens* Bloch = *Balistes buniva* Lacépède).
4014. *Melichthys piceus* (Poey). *Galafate*; *Black oldwife*; *Calafate*.
West Indies and southward.
? *Balistes nigra* Osbeck, Iter Chinensis, 1757, 295, Ascension Island; pre-Linnæan.
Galafate Parra, Dif. Piezas, etc., 1787, 18, Habana.
Balistes piceus Poey, Proc. Acad. Nat. Sci. Phila., XV, 1863, 180, Cuba.
4015. *Melichthys radula* (Solander).
East Indies, Clarion and Socorro Islands; Cocos Island.
Balistes radula Solander, in Richardson, Voy. H. M. S. *Samarang*, Fishes, 1848, 21.
Melichthys bispinosus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 125, Clarion and Socorro Islands of the Revillagigedo group.

Family 297. **MONACANTHIDÆ**. FilefishesGenus 1435. **CANTHERINES** Swainson.

- Cantherines* Swainson, Classn. Fishes, II, 1839, 327 (*Cantherines nasutus* Swainson = *Balistes sandwichensis* Quoy and Gaimard).
Pseudomonacanthus Bleeker, loc. cit., 12 (*Monacanthus macrurus* Bleeker).
Liomonacanthus Bleeker, Ned. Tijdschr., III, 1866, 13 (*Monacanthus pardalis* Rüppell).

- 4016. *Cantherines pullus* (Ranzani). *Liza colorado*.**
 West Indies and coast of Brazil; southern Florida.
Liza colorado Parra, Dif. Piezas, etc., 1787, pl. 23, p. 49, Cuba.
Monacanthus pullus Ranzani, Nov. Comm. Act. Sci. Inst. Bonon., V, 1842, 4, pl. 1, Brazil.
Monacanthus macrocerus Holland, Ann Sci. Nat., II, 1854, 327, pl. 12, fig. 1, Bahia.
Monacanthus ruppelii Castelnau, Anim. Nouv. Amer. Sud., Poiss., 1855, 97, pl. 47, fig. 2, Bahia.
Monacanthus irroratus Poey, Memorias, II, 1861, 330, Cuba.
Monacanthus striatus Poey, loc. cit., 329, Cuba.
Monacanthus parraianus Poey, Proc. Acad. Nat. Sci. Phila., XV, 1863, 185, Cuba; after Parra.
Monacanthus punctatus Poey, Synopsis, 1868, 437, Cuba.
- 4017. *Cantherines carolæ* (Jordan and McGregor).**
 Islands off the Pacific coast of Mexico; Revillagigedo Islands.
 ?*Balistes sandwichensis* Quoy and Gaimard, Voy. l'*Uranie*, Zool., 1824, 214, Sandwich [Hawaiian] Islands.
 ?*Cantherines nasutus* Swainson, Nat. Hist. Class. Fishes, II, 1839, 327; substitute for *B. sandwichensis*.
Cantherines carolæ Jordan and McGregor, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1713 (Clarion Island by error), Socorro Island.
- 4018. *Cantherines amphioxys* (Cope).**
 St. Martins Island, West Indies.
Monacanthus amphioxys Cope, Trans. Amer. Phil. Soc., XIV, 1870, 477, St. Martins Island.
- Subgenus **MONACANTHUS**.
- Genus **1436. MONACANTHUS** Cuvier.
Monacanthus Cuvier, Règne Anim., Ed. I, 1817, 152 (*Balistes chinensis* Bloch).
Trichoderma Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 328 (*Balistes scapus* Lacépède).
Stephanolepis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 78 (*Monacanthus setifer* Bennett).
- 4019. *Monacanthus ciliatus* (Mitchill). *Leatherfish*; *Liza*; *Flapmango*.**
 West Indies and Florida; Florida Keys.
Balistes ciliatus Mitchill, Amer. Month. Mag. and Crit. Rev., II, March, 1818, 326, Bahama Straits.
Monacanthus piraaca Kner, Reise Fregatte *Novara* um Erde, Zool., Fische, 1867, 396, Rio Janeiro.
Monacanthus occidentalis Günther, Cat., VIII, 1870, 237, Puerto Caballo.
Monacanthus davidsonii Cope, Trans. Amer. Phil. Soc. Phila., XIV, 1870 (1871), 476, Florida Reef.
- Subgenus **STEPHANOLEPIS** Gill.
- 4020. *Monacanthus hispidus* (Linnæus). *Foofish*; *Filefish*; *Leatherfish*; *Horny cony*; *Liza*; *Thread filefish*; *Mingo*.**
 Cape Cod to Cuba; south Atlantic coast and Florida Keys; West Indies to Brazil.
Balistes hispidus Linnæus, Syst. Nat., Ed. XII, 1766, 405, Carolinas.
Balistes broccus Mitchill, Trans. Lit. Phil. Soc., I, 1815, 467, New York.
Monacanthus filamentosus Valenciennes, Îles Canaries Ichth. Atl., 1836, 95, pl. 16, fig. 1, Canaries.
Monacanthus gallinula Valenciennes, loc. cit., 95, Canaries.
Monacanthus varius Ranzani, Novi Comment. Ac. Sci. Inst. Bonon., V, 1842, 6, Pl. II, Brazil.
Monacanthus massachusettsensis DeKay, N. Y. Fauna: Fishes, 1842, 336, pl. 57, fig. 187, Massachusetts Bay.
Monacanthus setifer DeKay, loc. cit., 337, pl. 59, fig. 194, New York.
Monacanthus signifer Storer, Synop. Fish. N. Amer., 1846, 497, Massachusetts.
Monacanthus auriga Lowe, Proc. Zool. Soc. Lond., 1850, 253, Madeira.

4021. *Monacanthus oppositus* Poey.
Cuba.
Monacanthus oppositus Poey, *Memorias*, II, 1861, 331, Habana.
4022. *Monacanthus spilonotus* Cope.
Gulf of Mexico.
Monacanthus spilonotus Cope, *Trans. Amer. Phil. Soc. Phila.*, XIV, 1870, 476, Gulf of Mexico.
4023. *Monacanthus tuckeri* Bean.
West Indies, north to Bermudas.
Monacanthus tuckeri Bean, *Field Mus. Nat. Hist., Zool.*, VII, 1906, 78, fig. 11, Bermudas.
- Genus 1437. *CERATACANTHUS* Gill.
Ceratacanthus Gill, *Fish. E. Coast N. Amer.*, *Proc. Acad. Nat. Sci. Phila.*, XIII, 1861, 57 (*Balistes auranthacus* Mitchill = *Balistes schæpfi* Walbaum).
4024. *Ceratacanthus schæpfi* (Walbaum). *Filefish; Orange filefish; Laja; Foolfish; Unicornfish; Sunfish.*
Cape Cod to the Carolinas and Florida.
Balistes schæpfi Walbaum, *Artedi Pisc.*, 1792, 461, Long Island; after Schopf, *Berlin Ges. Naturf.*, VIII, 1788, 186.
Balistes auranthacus Mitchill, *Trans. Lit. Phil. Soc. N. Y.*, I, 1815, 468, New York.
Aluterus cuspidata DeKay, *New York Fauna: Fishes*, 1842, 338, New York.
Aluterus holbrooki Hollard, *Ann. Sci. Nat.*, IV, 1855, 7, North America.
Aluterus cultrifrons Hollard, *Ann. Sci. Nat.*, ser. 4, IV, 1855, 8, pl. 1, fig. 2, New York and Bahia.
- Genus 1438. *DAVIDIA* Miranda Ribeiro.
Davidia Miranda Ribeiro, *Archiv Mus. Nac. Rio Janeiro*, XVII, 1915, 9 (*Alutera punctata* Agassiz).
4025. *Davidia punctata* (Agassiz). *Long mingo.*
Jamaica to Bahia.
Alutera punctata Agassiz, *Pisc. Brasil.*, 1829, 134, pl. 76, Brazil.
- Genus 1439. *OSBECKIA* Jordan and Evermann.
Osbeckia Jordan and Evermann, *Check List*, 1895, 424 (*Balistes scripta* Osbeck).
4026. *Osbeckia scripta*¹ (Osbeck). *Unicornfish; Lija trompa; Olilepa; Ohua.*
Tropical seas; West Indies; North Carolina; off west coast of Mexico; Clarion Island, Revillagigedo Islands.
Balistes scriptus Osbeck, *Iter Chinensis*, I, 1757, 144, Canaries.
Unicornis piscis bahamensis Catesby, *Nat. Hist. Carolina*, etc., II, 1743, 19, pl. 19, Bahamas.
Balistes laevis Bloch, *Naturgesch. Ausl. Fische*, IV, 1795, 82, pl. 414, Morocco.
Monacanthum proboscideum Ranzani, *Nov. Comm. Ac. Sc. Inst. Bonon.*, V, 1842, 8, Brazil.
Aluterus venosus Hollard, *Ann. Sc. Nat.*, ser. 4, IV, 1855, 14, pl. 1, fig. 3, New Ireland.
Alutera picturata Poey, *Proc. Acad. Nat. Sci. Phila.*, XV, 1863, 183, Cuba.
- Genus 1440. *ALUTERA* Cuvier.
Les Alutères Cuvier, *Règne Anim.*, ed. 1, 1817, 153 (*Balistes monoceros* Osbeck).
Alutera Oken, *Isis*, 1817, 153 ("Les Alutères" Cuvier; *Balistes monoceros* Linnæus).

¹ Whether *Osbeckia venosa* Hollard, East Indies to Hawaii and the Revillagigedos, is distinct from *O. scripta* of the Atlantic is a matter not yet determined. The distinction of *Osbeckia* and *Ceratacanthus* from *Alutera* is questionable.

4027. *Alutera monoceros*⁹² Linnæus. *Laja barbuda*; *Loulu*.
Pacific Ocean; East Indies; Japan; Hawaii.
Capriscus minima etc, Klein, *Miasus*, II, 1742, 25 (no locality; figure very bad).
Balistes monoceros Osbeck, *Iter Chinensis*, 1757, 110, Asia, Linnæus, *Syst. Nat.*, ed. X, 1758, 327, Asia and America.
Laja barbuda Parra, *Dif. Piezas Hist. Nat.*, 1787, 48, pl. 22, fig. 2, Habana.
Balistes kleini Gmelin, *Syst. Nat.*, 1788, 1472, Indian seas; after Klein.
Balistes barbatus Walbaum, *Artedi Pisc.*, III, 1792, 464; after Klein.
Balistes monoceros unicolor Bloch and Schneider, *Syst. Ichth.*, 1801, 463, after Gmelin.
Aluteres berardi Lesson, *Voy. Coquille*, Zool., 1830, 108, pl. 7, New Guinea.
Alutera cinerea Temminck and Schlegel, *Fauna Japon.*, Poiss., 1850, 292, pl. 131, fig. 1, Japan.
Alutarius macracanthus Bleeker, *Verh. Bat. Gen.*, Balistini, XXIV, Art. 11, 1852, 22, pl. 3, fig. 7, East Indies.
Alutarius amphacanthoides Bleeker, loc. cit., 23, pl. 2, fig. 5, East Indies.
Alutarius obliteratus Cantor, *Malayan Fishes*, 1850, 352, Pinang.
Balistes linguatula Gronow, *Cat.*, ed. Gray, 1854, 35, Indian seas.
Aluterus anginosus Hollard, *Ann. Sci. Nat.*, IV, 1855, 11, East Indies.
Balistes uncornu Basilewsky, *Nouv. Mém. Soc. Nat. Moscow*, X, 1855, 263, China.
4028. *Alutera guntheriana* Poey. *Liza barbuda*.
West Indies; once taken at Woods Hole (1898).
Alutera guntheriana Poey, *Proc. Acad. Nat. Sci. Phila.*, XV, 1863, 184, Habana.

Suborder OSTRACODERMI

Family 298. OSTRACIIDÆ. Trunkfishes

- Genus 1441. OSTRACION Linnæus.
Ostracion Linnæus, *Syst. Nat.*, Ed. X, 1758, 330 (*O. cubicus* Linnæus).
Cibotus Kaup, *Wiegmann's Archiv*, XXI, 1855, 215 (*Ostracion cubicus* Linnæus).
Ostracium of authors; improved orthography.
4029. *Ostracion punctatum* Bloch and Schneider.
Tropical Pacific in general; Clipperton Island, Galapagos Islands, Hawaiian Islands.
Ostracion lentiginosum Bloch and Schneider, *Syst. Ichth.*, 1801, 501, Ile de France.
Ostracion punctatus Bloch and Schneider, loc. cit., 501, Honolulu.
4030. *Ostracion clippertonense* Snodgrass and Heller.
Clipperton Island.
Ostracion clippertonense Snodgrass and Heller, *Shore Fishes*, Galapagos Islands, in *Proc. Wash. Acad. Sci.*, VI, 1904, 410, Clipperton Island.
- Genus 1442. RHINESOMUS Swainson.
Rhinesomus Swainson, *Nat. Hist. Classn. Fishes*, II, 1839, 194, 324 (*Ostracion triquetus* Linnæus)
4031. *Rhinesomus triquetus* (Linnæus). *Trunkfish*; *Rock shellfish*; *Drunkenfish*; *Chapin*; *Platefish*.
West Indies, north to the Bermudas; Key West and Pensacola; Woods Hole, Mass.
Ostracion triquetus Linnæus, *Syst. Nat.* Ed. X, 1758, 330, "India."
Ostracion concatenatus Bloch, *Naturgesch. Ausl. Fische*, I, 1785, 101, Martinique.
- Genus 1443. CHAPINUS Jordan and Evermann.
Chapinus Jordan and Evermann, *Check List*, 1895, 424 (*Ostracion bicaudalis* Linnæus).

⁹² Whether the West Indian *Alutera guntheriana* is distinguishable from *Alutera monoceros* of the Pacific is uncertain.

- 4032. *Chapinus bicaudalis* (Linnæus).** *Chapin; Spotted trunkfish; Pesce toro; Bullfish.*
West Indies, from Cuba to Ascension Island; Tortugas.
Ostracion bicaudalis Linnæus, Syst. Nat., Ed. X, 1753, 330, "India"; after Artedi.
- Genus 1444. LACTOPHRYS Swainson.** *Three-angled trunkfishes.*
Lactophrys Swainson, Nat. Hist. Classn. Fishes, II, 1839, 194, 324 (*Ostracion trigonus* Linnæus, etc.).
- 4033. *Lactophrys trigonus* (Linnæus).** *Common trunkfish; Chapin; Shellfish; Cuckold; Boxfish.*
West Indies, north to Bermuda and Key West; occasionally northward in Gulf Stream to Holmes Hole, Mass., and Chesapeake Bay.
Ostracion trigonus Linnæus, Syst. Nat., Ed. X, 1753, 330, India.
Ostracion yalei Storer, Bost. Jour. Nat. Hist., I, 1836, 353, Holmes Hole and Marthas Vineyard.
Ostracion undulatus Poey, Synopsis, 1868, 441, Habana.
Ostracium expansum Cope, Trans. Amer. Phil. Soc., XIV, 1870, 474, figs. 9-10, locality unknown.
- Genus 1445. ACANTHOSTRACION Bleeker.**
Acanthostracion Bleeker, Atlas Ichth., V, 1865, 27, (*Ostracion quadricornis* Linnæus = *Ostracion tricornis* Linnæus).
- 4034. *Acanthostracion quadricornis* (Linnæus).** *Cuckold, Toro; Cowfish; Horned trunkfish.*
Tropical seas; common from Carolina to Brazil; Marquesas Islands; Florida Keys; occasionally at Pensacola and Galveston and north in the Gulf Stream to Charleston and Chesapeake Bay.
Ostracion tricornis Linnæus, Syst. Nat., Ed. X, 1753, 331, "India"; after Artedi.
Ostracion quadricornis Linnæus, loc. cit., 331, "India"; after Artedi; (this name preferred to the preceding by the first reviser).
Ostracion hster Lacépède, Hist. Nat. Poiss., I, 1798, 468; after Willoughby.
Ostracion sexcornutus Mitchill, Amer. Month. Mag., II, 1818, 328, mouth of Mississippi River.
Ostracion guineensis Bleeker, Ned Tydskr. Dierk., II, 1865?, 298, Guinea.
Ostracion gronovi Bleeker, loc. cit., II, 298, Guinea.
Acanthostracion polygonius Poey, Enumeratio, 1876, 175, Cuba.

Suborder GYMNOdontes

Family 299. TETRAODONTIDÆ. Puffers

- Genus 1446. LAGOCEPHALUS**⁹³ Swainson.
Lagocephalus Swainson, Nat. Hist. Classn. Fishes, II, 1839, 194, 328 (*Tetraodon stellatus* Donovan).
- 4035. *Lagocephalus lævigatus* (Linnæus).** *Smooth puffer; Rabbitfish; Smooth swellfish; Tambor.*
Cape Cod to Brazil; rare north of Cape Hatteras.
Tetraodon lævigatus Linnæus, Syst. Nat., Ed. XII, 1766, 411, Charleston, S. C.
Tetraodon curvus Mitchill, Trans. Lit. Phil. Soc., I, 1815, 472, New York.
Tetraodon mathematicus Mitchill, loc. cit., I, 1815, 474, New York.
Holocanthus melanothos Gronow, Syst. Nat., ed. Gray, 1854, 24, Carolinas.
Tetraodon lineolatus Poey, Synopsis, 1868, 432, Cuba.
- 4036. *Lagocephalus pachycephalus* (Ranzani).** *Jugfish.*
Jamaica to Brazil.
Tetraodon pachycephalus Ranzani, Nov. Comm. Ac. Sci. Inst. Bonon., IV, 1840, 73, Pl. X, fig. 2, Brazil.

⁹³ The American species of *Lagocephalus* are very well separated from *Sphaeroides* but various Japanese species of the former genus grade off into the latter.

Genus 1447. SPHÆROIDES Lacépède. *Swellfishes*.

- Les *Sphéroides* Lacépède, Hist. Nat. Poiss., II, 1798, 22, Pl. I, fig. 1
(*La Sphéroïde tuberculée*, a front view of *Tetraodon spengleri* Bloch).
Sphæroides, Allegem. Lit. Zeit. Berlin, Sept. 24, 1798, unsigned review,
adopts this spelling as a technical name for the genus.
Sphæroides Duméril, Zool. Anal., 1806, 342; after Lacépède.
Orbidus Rafinesque, Analyse de la Nature, 1815, 90 (substitute for Les
Sphéroides Lacépède).
Crayracion Klein, Gesellschaft Schauplatz, IV, 1777, 788 (*Tetraodon*
spengleri Bloch; not binomial); after Klein, 1742.
Holocanthus Gronow, Cat. Fish., 1854 (*H. leionothus* Gronow).
Anchisomus (Kaup) Richardson, Zool. H. M. S. Herald, 1854, 156
(*Tetraodon spengleri* Bloch).
Crayracion Bleeker, Atlas Ichth., 1865, 65 (*Tetraodon spengleri* Bloch).
Stenometopus Bibron, Rev. Mag. Zool., VIII, 1855, 279 (*Tetraodon*
téstudineus Linneus).
- 4037. *Sphæroides maculatus* (Bloch and Schneider). *Puffer; Swelltoad; Blower; Swellbelly; Swellfish; Toadfish*.**
Atlantic coast of the United States from Cape Ann to St. Johns River
and Biscayne Bay, Fla.
Tetrodon hispidus var. *maculatus* Bloch and Schneider, Syst. Ichth.,
1801, 504, Rhode Island; after Schopf.
Tetrodon turgidus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 473,
pl. 5, fig. 5, New York.
- 4038. *Sphæroides spengleri* (Bloch). *Southern puffer; Swelltoad; Tambor*.**
West Indies; coasts of Texas and Florida, south to Rio Janeiro and to
the Madeiras and Canaries; rarely north to Woods Hole.
Tetrodon spengleri Bloch, Ichth., I, 1782, 135, pl. 144, East Indies.
Le *Tetrodon* Plumier Hist. Nat. Poiss., I, 1797, 504, Martinique.
Le *Sphéroïde tuberculé* Lacépède, Hist. Nat. Poiss., II, 1798, 22, Mar-
tinique.
Sphæroides tuberculatus Pillot edition of Lacépède, VI, 1831, 279,
Martinique.
- 4039. *Sphæroides nephelus* (Goode and Bean). *Puffer; Swellingfish*.**
Florida.
Tetrodon nephelus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882,
412, Indian River and Pensacola, Fla.
- 4040. *Sphæroides marmoratus* (Ranzani). *Spinyback blowfish; Swellfish*.**
West Indies to Brazil; Jamaica.
Tetrodon marmoratus Ranzani, Nov. Comm. Ac. Sci. Inst. Bonon.,
IV, 1840, 72, pl. 10, fig. 1, Brazil.
- 4041. *Sphæroides harperi* Nichols.**
West Indies (St. Eustatius) north to Florida.
Sphæroides harperi Nichols, Bull. Am. Mus. Nat. Hist., XXXIII,
1914, 81, Cape Sable, Fla.
- 4042. *Sphæroides asterias* (Blosser).**
St. Croix.
Sphæroides asterias Blosser, Ann. Carnegie Mus., VI, 300, 1909, St.
Croix.
- 4043. *Sphæroides eulepidotus* (Metzelaar).**
Caribbean Sea.
Tetrodon (Sphæroides) eulepidotus Metzelaar, Trop. Atl. Visschen, 1919,
170, Curaçao.
- 4044. *Sphæroides lobatus* (Steindachner). *Botete*.**
Gulf of California to Galapagos Islands; Mazatlan; Panama.
Canthogaster? lobatus Steindachner, Ichth. Notizen, X, 1870, 18, pl. 5,
fig. 3, Altata.
- 4045. *Sphæroides angusticeps* (Jenyns).**
Pacific coast Lower California (La Paz), to the Galapagos Islands.
Tetrodon angusticeps Jenyns, Voyage Beagle, Fishes, 1842, 154, 28,
Galapagos Islands.

Genus 1448. CHEILICHTHYS Müller.

Cheilichthys Müller, Abhandl. Akad. Wiss. Berl., 1839, 252 (*Tetraodon testudineus* Linnæus).

Apsicephalus Hollard, Ann. Sci. Nat., Zool., VIII, 1857, 275 (*Tetraodon fluviatilis* Hamilton).

4046. *Cheilichthys testudineus* (Linnæus). Tambor; Glovefish; Blowfish; Swellfish; Puffer.

West Indies; Gulf Stream, occasionally to Newport, R. I.

Orbis lævis variegatus Catesby, Nat. Hist. Carolinas, II, 1743, 28, pl. 28, Virginia.

Tetraodon testudineus Linnæus, Syst. Nat., Ed. X, 1758, 332, India; based on Balk and Artedi.

?*Tetraodon punctatus* Bloch and Schneider, Syst. Ichth., 1801, 506, Brazil.

Tetraodon geometricus Bloch and Schneider, loc. cit., 508, Virginia; after Catesby.

Tetraodon ammocryptus Gosse, Nat. Sojourn Jamaica, 1851, 287, Jamaica.

Holocanthus leionothos Gronow, Cat. Fishes, ed. Gray, 1854, 24, American oceans.

4047. *Cheilichthys annulatus* (Jenyns).

Pacific coast of tropical America; Galapagos Islands.

Tetraodon annulatus Jenyns, Zool. Beagle, 1842, 153, Galapagos Islands.

Anchisomus geometricus (Kaup) Richardson, Voyage Herald, 1854, 156, pl. 30, Galapagos Islands.

Tetraodon heraldi Gunther, Cat., VIII, 1870, 283, Galapagos Islands and Panama.

4048. *Cheilichthys politus* (Girard).

Pacific coast of tropical America, San Diego to Mazatlan.

Tetraodon politus Girard, Pacific R. R. Surv., X, 1858, 340, San Diego, Calif.

4049. *Cheilichthys fürthi* (Steindachner).

Panama.

Tetraodon furthii Steindachner, Ichth. Beitr., V, 1876, 22, Panama.

4050. *Cheilichthys trichocephalus* (Cope) Harry blowfish.

Gulf Stream off Newport, R. I.; only the type known.

Tetraodon trichocephalus Cope, Proc. Acad. Nat. Sci. Phila., XXII, 1870, 120, Gulf Stream off Newport, R. I.

4051. *Cheilichthys pachygaster* (Müller and Troschel).

Barbados.

Tetraodon (Cheilichthys) pachygaster Müller and Troschel, in Schomburgk's Hist. Barbados, 1848, 677, Barbados.

Genus 1449. TETRAODON Linnæus.

Tetraodon Linnæus, Syst. Nat., Ed. X, 1758, 333 (*T. lineatus* Linnæus).

Les Ovoides Lacépède, Hist. Nat. Poiss., I, 1797, 256 (fascé, face view).

Ovoides Cuvier, Leçons Anat. Comp., I, 1800, tab. 1 (*Ovoides fasciatus* Lacépède = *Tetraodon stellatus* Linnæus).

Ovum Bloch and Schneider, Syst. Ichth., 1801, 530 (*O. commersoni* Bloch and Schneider = "l'Ovoide Fascé" Lacépède, 1797); preoccupied in mollusks.

Oonidus Rafinesque, Anal. Nat., 1815, 90, for "les Ovoides" of Lacépède.

Arothron Müller, Abh. Akad. Wies. Berlin, 1839, 252 (*A. testudinatus* Müller = *Tetraodon reticularis* Bloch and Schneider).

Dilobomycter Bibron, Rev. Mag. Zool., VIII, 1855, 279 (*Tetraodon reticularis* Bloch and Schneider).

Dichotomycter Bibron, loc. cit., 279 (*Tetraodon pluviatilis* Hamilton).

4052. *Tetraodon hispidus* Linnæus *Maki-maki*; *Kete*.
Panama and neighboring islands; Hawaii; Samoa; to China and the Red Sea.
Tetraodon hispidus Linnæus, Syst. Nat., Ed. X, 1758, 333, India.
Tetraodon perspicillaris Ruppell, Atlas, Reise Nord Africa, 1828, 63, Red Sea.
Tetraodon implutus Jenyns, Voy. *Beagle*, Fish., 1842, 152, Keeling Island.
Tetraodon latera Richardson, Voy. *Sulphur*, Zool., 1844, 124, pl. 61, fig. 2, Canton, China.
*Arothron erethizon*⁹⁴ Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 631, Panama.

Subgenus *OVOIDES* Cuvier.

4053. *Tetraodon setosus* Smith.
West coast of Mexico; Revillagigedo, Clipperton, Cocos, and Galapagos Islands.
Tetraodon setosus Smith, Bull. Calif. Acad. Sci., 2d ser., VI, Nov. 13, 1886, Mexico.

Genus 1450. *GUENTHERIDIA* Gilbert and Starks.

Guentheridia Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 158 (*Tetraodon formosus* Günther).

4054. *Guentheridia formosa* (Günther).

Panama.

Tetraodon formosus Günther, Cat., VIII, 1870, 233, "South America."

Genus 1451. *COLOMESUS* Gill.

Colomesus Gill, Proc. U. S. Nat. Mus., VII, 1884, 422 (*Tetraodon psittacus* Bloch and Schneider)

4055. *Colomesus psittacus* (Bloch and Schneider).

River mouths, Guiana and northern Brazil; Rio Branco, Essequibo, Marañon, and Capin at Para; West Indies in fresh waters only.

Tetraodon psittacus Bloch and Schneider, Syst. Ichth., 1801, 505, "Malabar."

Cheilichthys asellus Muller and Troschel, in Schomburgk, British Guiana, III, 1848, 641, Guiana.

Family 300. *CANTHIGASTERIDÆ*. Sharp-nosed puffers

Genus 1452. *CANTHIGASTER* Swainson.

Canthigaster Swainson, Nat. Hist. Fish., Amphib., etc., II, 1835, 194 (*Tetraodon rostratus* Bloch, called *Psilonotus* later in the text).

Psilonotus Swainson, loc. cit., 1835, 328, same type.

Prilonotus (Kaup) Muller, Richardson, Voyage *Herald*, 1854, 162 (*Tetraodon rostratus* Bloch).

Tropidichthys Bleeker, Nat. Tijdschr. Neder-Indie, VI, 501 (*T. margaritatus* Bleeker).

Anosmius Peters, Monatsber. Akad. Wiss. Berlin, XXI, 1855, 274 (*T. lænatus* Peters).

Eumycteris Jenkins, Bull. U. S. Fish Com., XIX, 1899, 399 (*E. biteniatus* Jenkins).

4056. *Canthigaster rostratus* (Bloch).

West Indian fauna; Pensacola; Madeiras and Bermudas.

Tetraodon rostratus Bloch, Ichth., I, 1782, pl. 146, India.

Tetraodon capistratus Lowe, Proc. Zool. Soc. Lond., 1839, 90, Madeira.

Prilonotus (*Anchisomus*) *caudacinctus* Richardson, Voyage *Herald*, 1854, 162, pl. 30, figs 1-3, locality unknown.

Tetraodon ornatus Poey, Synopsis, 1868, 433, Cuba.

4057. *Canthigaster punctatissimus* (Günther).

Pacific coast of America, Gulf of California to Panama.

Tetraodon punctatissimus Günther, Cat., VIII, 1870, 302, Panama.

Tetraodon oxyrhynchus Lockington, Proc. Acad. Nat. Sci. Phila., XXXIII, 1881, 116, Gulf of California

⁹⁴ This American form shows no distinction from the widespread *Tetraodon hispidus*.

Family 301. DIODONTIDÆ

Genus 1453. DIODON Linnæus. *Porcupinefishes*.

Diodon Linnæus, Syst. Nat., Ed. X, 1758, 335 (*D. hystrix* Linnæus).

Paradiodon Bleeker, Neder. Tijdschr. Dierk., II, 1865, 271 (*Diodon hystrix* Linnæus).

4058. *Diodon hystrix* Linnæus. *Porcupinefish*; *Erizo*; *Puerco espino*; *Soursap*. Tropical seas, north to Lower California, Hawaii, and Florida.

Diodon hystrix Linnæus, Syst. Nat., Ed. X, 1758, 335, India; after Artedi.

Guamajacu guará Marcgrave, Hist. Nat. Bras., 1648, 159, Brazil.

Erizo Parra, Desc. Dif. Piezas Hist. Nat. Cuba, 1787, 60, pl. 29, fig. 1, Habana.

Le Diodon plumier Lacépède, Hist. Nat. Poiss., II, 1798, 1, pl. 3, fig. 3, Martinique.

Diodon brachiatum Bloch and Schneider, Syst. Ichth., 1801, 513, Cuba.

Diodon punctatus Cuvier, Mém. du Mus. d'Hist. Nat., IV, 1818, 132, no locality.

Diodon echinus Rafinesque, in Bonaparte, Cat. Met. Pisc. Eur., 1846, 87, Mediterranean Sea.

4059. *Diodon holacanthus* Linnæus. *Balloonfish*

All warm seas; north to Florida Keys, Lower California, and Hawaiian Islands.

Diodon holacanthus Linnæus, Syst. Nat., ed. X, 1758, 335, India; based on Artedi (misprint).

Erizo guanabana Parra, Desc. Dif. Piezas Hist. Nat. Cuba, 1787, 62, pl. 29, fig. 2, Habana.

Le Diodon tacheé Lacépède, Hist. Nat. Poiss., II, 1798, 13, New Cytherea.

Diodon liturosus Shaw, Gen. Zool., V, 1804, 436; after Lacépède.

Diodon spinosissimus Cuvier, Mém. Mus. Hist. Nat., IV, 1818, 134, pl. 7, no locality.

Diodon novem-maculatus Cuvier, loc. cit., 136, pl. 6, no locality.

Diodon sex-maculatus Cuvier, loc. cit., 136, pl. 7, no locality.

Diodon multimaculatus Cuvier, loc. cit., 137, no locality.

Diodon 4-maculatus Cuvier, loc. cit., 137, pl. 6, Otaïti.

Diodon maculatus var. a, Gunther, Cat., VIII, 1870, 307, St. Croix; Jamaica; Panama; South America, etc.; after Lacépède.

Diodon melanopsis Kaup, Wiegmann's Archiv, 1855, 228.

4060. *Diodon maculifer* Kaup.

Cape of Good Hope; recorded doubtfully from Cuba.

Diodon maculifer Kaup, Wiegmann's Archiv, 1855, 229, Cape of Good Hope.

Genus 1454. TRICHODIODON Bleeker.

Trichodiodon Bleeker, Atlas Ichth., Gymnodontes, V, 1867, 49 (*Diodon pilosus* Mitchell).

Trichocycilus Günther, Cat., VIII, 316 (*T. erinaceus* Günther).

4061. *Trichodiodon pilosus* (Mitchell). *Balloonfish*; *Hairy porcupinefish*; *Hairy boxfish*.

New York.

Diodon pilosus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 471; supposed to be from New York Harbor.

Diodon asper Cuvier, Mém. Mus. Hist. Nat., IV, 1818, 138; no locality.

?*Trichocycilus erinaceus* Günther, Cat., VII, 1870, 316; no locality.

Genus 1455. CYCLICHTHYS Kaup. *Swelltoads*.

Cychchihys Kaup, Wiegmann's Archiv, XXI, 1855, 231 (*Diodon orbiculatus* Bloch).

4062. *Cyclichthys schœpfi* (Walbaum). *Burrfish; Cucumberfish; Spotted balloonfish; Balloonfish; Rabbitfish; Swelltoad; Spiny boxfish.*
Cape Cod to West Indies and occasionally northward to Woods Hole, Mass.
Diodon schœpfi Walbaum, Artedi Pisc., 1792, 601, Long Island; after Schöpf.
? *Diodon meulenii* Walbaum, loc. cit., 602; no locality.
Diodon geometricus lineatus Bloch and Schneider, Syst. Ichth., 1801, 513, New York.
Diodon maculato striatus Mitchill, Fish. N. Y., 470, pl. 56, fig. 3, 814, New York.
Diodon rivulatus Cuvier, Mém. Mus. Hist. Nat., IV, 1818, 129, pl. 6, New York.
Diodon nigrolineatus Ayres, Proc. Bost. Soc. Nat. Hist., I, 1842, 68, Brookhaven, Long Island.
Diodon verrucosus DeKay, N. Y. Fauna: Fishes, 324, 1842, pl. 56, fig. 184, New York.
4063. *Cyclichthys spinosus* (Linnæus).
West Indies and coast of Brazil.
Guamancu atinga Marcgrave, Hist. Brasil., 1648, 168, Brazil.
Diodon spinosus Linnæus, Syst. Nat., Ed. X, 1758, 335, India.
Le Diodon orbe Lacépède, Hist. Nat. Poiss., II, 1793, 16, Rio Janeiro.
Diodon geometricus Bloch and Schneider, Ichth., 1801, 513, pl. 96, America.
Cyclichthys cornutus Kaup, Wiegmann's Archiv, 1855, 231, Bahia.
4064. *Cyclichthys fuliginosus* (DeKay). *Black burrfish; Unspotted balloonfish.*
Atlantic coast from Cape Cod southward; not common.
Diodon fuliginosus DeKay, N. Y. Fauna: Fishes, 1842, 324, pl. 55, fig. 181, New York Harbor.
4065. *Cyclichthys antillarum* (Jordan and Rutter).
Cuba and Jamaica.
Chilomycterus antillarum Jordan and Rutter, Proc. Acad. Nat. Sci. Phila., XLIX, 1897, 131, Kingston, Jamaica.
4066. *Cyclichthys briareos* (Metzelaar).
Lesser Antilles.
Chilomycterus briareos Metzelaar, Trop. Atl. Vissch., 1919, 173, fig. 55, St. Eustatius.
4067. *Cyclichthys antennatus* (Cuvier).
West Indies and southward; St. Croix; Jamaica; Porto Rico; Cape of Good Hope.
Diodon antennatus Cuvier, Mém. Mus. Hist. Nat., IV, 1818, 131; no locality.
? *Chilomycterus puncticulatus* Poey, Anal. Soc. Española Hist. Nat., X, 1881, 346, Porto Rico.
- Genus 1456. *CHILOMYCTERUS* Bibron. *Burrfishes.*
Chilomycterus Bibron, in Barneville, Rev. Zool., 1846, 40 (*Diodon reticulatus* Linnæus = *Diodon atinga* Linnæus).
Cyanichthys Kaup, Wiegmann's Archiv, IV, 1855, 231 (*C. cæruleus* Kaup; not of Quoy and Gaimard).
Diodon Bleeker, Atlas Ind. Gymnod., 1865, 55 (*D. atinga*); not of authors, who, after Kaup, restrict *Diodon* to *D. hystrix*.
- Subgenus *CHILOMYCTERUS* Bibron.
4068. *Chilomycterus atinga* (Linnæus). *Atinga.*
West Indies to Bermuda and Florida Keys.
Diodon atinga (misprinted *atinga*) Linnæus, Syst. Nat., Ed. X, 1758, 334, India.
Diodon reticulatus Linnæus, loc. cit., 334, India; after Artedi.
4069. *Chilomycterus affinis* Günther.
Galapagos; coast of southern California; Hawaii, Japan; scarce.
Chilomycterus affinis Günther, Cat., VIII, 1870, 314, no locality; California?
Chilomycterus californiensis Eigenmann, Amer. Nat., V, 1891, 1133, San Pedro, Calif.

Genus 1457. LYOSPHERA Evermann and Kendall.

Lyosphæra Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 131 (*L. globosa* Evermann and Kendall).

4070. Lyosphæra globosa Evermann and Kendall. *Marblefish*.

Atlantic coast of the United States; known from the mouth of the Rappahannock River and from Biscayne Bay, Fla.

Lyosphæra globosa Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 131, pl. 9, figs. 11 and 12, Rappahannock River near mouth of Windmill Creek, Va.

Family 302. MOLIDÆ. Headfishes**Genus 1458. MOLA** Kœlreuter. *Headfishes*.

Mola Kœlreuter, Nova Comm. Act. Petropol., VII, 1770, 337 (*Diodon mola* Kœlreuter=*Diodon mola* Linnæus).

Mola Cuvier, Tableau Elem. Hist. Nat. Anim., 1798, 323 (*M. rotunda* Cuvier=*Tetrodon mola* Linnæus).

Orthogoriscus Schneider, ed. Bloch, Syst. Ichth., 1801, 510 (*Diodon mola* Bloch).

Cephalus Shaw, Gen. Zool., V, 1804, 432 (*Diodon mola* Linnæus).

Orthragus Rafinesque, Caratteri, 1810, 17 (*Tetrodon mola* Linnæus).

Diplanchias Rafinesque, loc. cit., 17 (*D. nasus* Rafinesque).

Tympanomium Ranzani, Novi Comm. Acad. Sci. Inst. Bonon., III, 1839, 81 (*T. planci* Ranzani).

Ozodura Ranzani, loc. cit., 81 (*O. orsini* Ranzani=*Mola mola* Linnæus).

Trematopsis Ranzani, loc. cit., 81 (*T. willughbei* Ranzani).

Pedalion Swainson, Nat. Hist. Fish. Amphib. etc., I, 1838, 199, II, 195, 329 (*Tetrodon mola* Linnæus).

Molacanthus Swainson, loc. cit., II, 1839, 329 (*M. pallasi* Swainson=*Diodon mola* Pallas, young).

Pallasia Nardo, Ann. Sci. Lombardo-Veneto, X, 1839, V, 10, 112 (*P. pallasi* Nardo).

Acanthosoma DeKay, Nat. Hist. N. Y., 1842, 350 (*A. carnatum* DeKay).

Centaurus Kaup, Wiegemann's Archiv Naturg., XXI, 1855, 221 (*Ostracion boops* Richardson; larval).

Aledon Castelnau, Mém. Poiss., Afr. Aust., VII, 1861, 75 (*A. storeri* Castelnau).

4071. Mola mola (Linnæus). *Ocean sunfish; Globe-fish; Sunfish; Headfish; Mola; Pez luna; Pez mola*.

Temperate and tropical seas, northward to England, Cape Cod, and San Francisco; rare in the West Indies.

Tetrodon mola Linnæus, Syst. Nat., Ed. X, 1758, 334, 412, Mediterranean.

Diodon carinatus Mitchell, Ann. Lyc. Nat. Hist. New York, II, 1815, 264, pl. 5, fig. 1, New York.

Ozodura orsini Ranzani, Nov. Comm. Ac. Sci. Inst. Bonon., III, 1839, 82, Mediterranean Sea.

Tympanomium planci Ranzani, loc. cit., table, Adriatic Sea.

Diplanchias nasus Ranzani, loc. cit., table, "in marei siculo" (Sicily).

Trematopsis willughbei Ranzani, loc. cit., table, "in oceano".

Orthragoriscus retzii Ranzani, loc. cit., table, no locality.

Orthragoriscus ghini Ranzani, loc. cit., table, Mediterranean Sea.

Orthragoriscus rondeletii Ranzani, loc. cit., Mediterranean Sea.

Orthragoriscus blochii Ranzani, loc. cit., "in mari oceano."

Orthragoriscus alexandrini Ranzani, loc. cit., Adriatic Sea.

Orthragoriscus redi Ranzani, loc. cit., Mediterranean Sea.

Orthragoriscus oculatus Ranzani, loc. cit., no locality.

Orthragoriscus elegans Ranzani, loc. cit., Atlantic Ocean.

Orthragoriscus battaræ Ranzani, loc. cit., Adriatic Sea.

Orthragoriscus lunaris Gronow, Cat. Fishes, ed. Gray, 1854, 165, Mediterranean.

Orthragoriscus solaris Gronow, loc. cit., 165, Mediterranean.

Orthragoriscus analis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 31, fig. 54, San Francisco.

Ostracion boops Richardson, Voy, *Erebus* and *Terror*, Ichth., 1844, 52, South Atlantic (larva).

Genus 1459. *RANZANIA* Nardo. *King of the mackerels.*

Ranzania Nardo, Ann. Sci. Regn. Lombard. Venet., X, 1840, 105
(*Tetrodon truncatus* Retzius).

4072. *Ranzania truncata* (Retzius)

Pelagic; occasionally on the Atlantic coast; Bermudas; Martinique.
Tetrodon truncatus Retzius, Vet. Ak. Nya. Handl., VI, 1785, 2, 121,
Brazil.

4073. *Ranzania makua* Jenkins.

Open Pacific, Hawaii to Japan; very rare, once found on Pismo Beach
near Oceano, San Luis Obispo County, Calif.
Ranzania makua Jenkins, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 779,
with colored plate (frontispiece), Honolulu; probably not different
from *Ranzania truncata*; see Snyder, Proc. U. S. Nat. Mus., XLIV,
1913, 455.

Order AAA. PEDICULATI

Family 303. LOPHIIDÆ. Fishing frogs

Genus 1460. *LOPHIUS* Linnæus.

Lophius Linnæus Syst. Nat., Ed. X, 1758, 236 (*Lophius piscatorius*
Linnæus); after Artedi.

4074. *Lophius piscatorius* Linnæus. *Fishing frog; Monkfish; Goosefish;*
Angler; Allmouth; Bellowsfish.

North Atlantic, on both coasts.
Lophius piscatorius Linnæus, Syst. Nat., Ed. XII, 1766, 402, Seas of
Europe.

Lophius americanus Cuvier and Valenciennes, Hist. Nat. Poiss., XII,
1837, 380, Philadelphia.

Genus 1461. *LOPHIOMUS* Gill.

Lophiomus Gill, Proc. U. S. Nat. Mus., V, 1882, 552 (*Lophius setigerus*
Vahl.)

4075. *Lophiomus setigerus* (Vahl).

Pacific Ocean; not uncommon in rather deep water off coasts of China
and Japan; one specimen from Albatross station 2805, southwest of
Panama.

Lophius setigerus Vahl, Skrivt. Naturh., IV, 1797, 214, tab. 3, figs. 5
and 6, China Sea.

Lophius viviparus Bloch and Schneider, Syst. Ichth., 142, 1801, tab.
32, China Sea; after Vahl.

Family 304. ANTENNARIDÆ. Frogfishes

Genus 1462. *HISTRIO* Fischer.

*Histrio*⁸⁵ Fischer, Zoog., Tab. Syn. Illust., Ed. III, I, 1813 (*Lophius*
histrio Linnæus).

Pterophryne Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 90 (*Chironectes*
lævigatus Cuvier; name preoccupied).

Pterophrynoides Gill, Proc. U. S. Nat. Mus., I, 1878, 216 (*Chironectes*
lævigatus Cuvier); substitute for *Pterophryne*).

4076. *Histrio pictus* (Cuvier and Valenciennes). *Mousefish; Sargassumfish.*

Tropical parts of the Atlantic; abundant on our Gulf coast and occa-
sional to Cape Hatteras and beyond.

Lophius tumidus Osbeck, Iter Chinensis, 1757, 400, open sea; pre-
Linnæan.

Chironectes pictus Cuvier and Valenciennes, Hist. Nat. Poiss., XII,
1837, 393, Surinam.

Chironectes tumidus Cuvier and Valenciennes, loc. cit., 397, Sargasso Sea.

Chironectes arcticus Duben and Koren, K. Vet. Akad. Stockholm, 1844,
72, Vadsø, Norway, carried northward on the Gulf Stream.

⁸⁵ Name questionable on account of error in the published diagnosis, the description interchanged with
that of *Lophius*.

4077. **Histrio gibba** (Mitchill).
West Indies, north to Key West and the Tortugas.
Lophius gibbus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 1, pl. 4,
fig. 9, off St. Croix, lat. 22° N., long. 64° W.
Chironectes længatus Cuvier, Mém. Mus., III, 1817, 423, pl. 16, fig. 1;
South Carolina.
4078. **Histrio jagua** Nichols
Bermuda.
Histrio jagua Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 62,
Bermuda.
- Genus 1463. **ANTENNARIUS** ⁹⁸ Lacépède.
Antennarius Lacépède, Hist. Nat. Poiss., I, 1798, 421 (*Antennarius
bivertex totus ater puncto mediorum, laterum albo* Commerson (in foot-
note) = *Lophius commersonianus* Lacépède).
Chironectes Cuvier, Mém. Mus., III, 1817, 418 (*Lophius commersonianus*
Lacépède).
Capellaria Gistel, Thierreich, VIII, 1848 (*Lophius commersonianus*
Lacépède).
4079. **Antennarius scaber** (Cuvier).
Caribbean Sea; Port Castries; St. Lucia.
Chironectes scaber Cuvier, Mém. Mus., III, 1817, 425, pl. 16, fig. 2,
Martinique.
Lophius spectrum Gronow, Cat. Fish., ed. Gray, 1854, 49, Antilles;
after *Lophius acute scabra* Gronow, Zoophyl., 1781, 210.
4080. **Antennarius inops** Poey.
Porto Rico.
Antennarius inops Poey, Anal. Soc. Esp. Hist. Nat., X, 1881, 340 Porto
Rico.
4081. **Antennarius principis** (Cuvier and Valenciennes).
West Indies to Brazil.
Chironectes principis Cuvier and Valenciennes, Hist. Nat. Poiss., XII,
1837, 416, Brazil; on a drawing by Prince Maurice.
? *Chironectes mentzeli* Cuvier and Valenciennes, loc. cit., 416, Brazil; on
a drawing.
4082. **Antennarius tenebrosus** (Poey).
Cuba.
Chironectes tenebrosus Poey, Memorias, I, 1853, 219, pl. 17, fig. 1, Cuba.
4083. **Antennarius reticularis** Gilbert.
Gulf of California; only the type known.
Antennarius reticularis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 566;
Gulf of California at Albatross station 2825.
4084. **Antennarius strigatus** Gill.
Cape San Lucas; Panama.
Antennarius strigatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 92;
Cape San Lucas
Antennarius tenuifilis Gunther, Fish. Cent. Amer., 1869, 440, Panama;
young.
4085. **Antennarius sanguineus** Gill.
Pacific coast of tropical America, Cape San Lucas to Panama; scarce.
Antennarius sanguineus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863,
91; Cape San Lucas.
Antennarius leopardinus Gunther, Proc. Zool. Soc. Lond., 1864, 151;
Panama.
4086. **Antennarius ocellatus** (Bloch and Schneider).
West Indies, north to Florida.
Pescador Parra, Dif. Piezas, Hist. Nat., 1780, 1, tab. 1, Cuba.
Lophius histrio, var. *ocellatus* Bloch and Schneider, Syst. Ichth., 142,
180; after Parra.
Antennarius pleurophthalmus Gill, Proc. Acad. Nat. Sci. Phila., XV,
1863, 92; Key West.

⁹⁸ These fishes vary nominately in color and more or less in form. It is likely that the number of species should be reduced.

4087. *Antennarius tigris* (Poey).
Cuba
Chironectes tigris Poey, Memorias, I, 1853, 217, pl. 17, fig. 2; Cuba.
4088. *Antennarius nuttingii* Garman.
Great Bahama Banks.
?Chironectes mentzelii Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 417; Brazil; on a drawing by Prince Maurice.
Antennarius nuttingii Garman, Bull. Iowa Lab. Nat. Hist., 1896, 83, Pl. II; Great Bahama Banks.
4089. *Antennarius multiocellatus* (Cuvier and Valenciennes). *Martin pescador*.
Garden Key, Fla., to Martinique.
Chironectes multiocellatus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1837, 422, Martinique.
Antennarius annulatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 91, Garden Key, Fla.
Antennarius corallinus Poey, Repertorio, I, 1865, 188, Cuba.
4090. *Antennarius radiosus* Garman.
Key West.
Antennarius radiosus Garman, Bull. Lab. Nat. Hist. Iowa Univ., 1896, 85, pl. 1, off Key West in 50 fathoms.
4091. *Antennarius sonntagii* (Baron Müller).
Gulf of Mexico.
Chironectes sonntagii Baron J. W. von Müller, Reisen in den Verein. Staat., etc., Band I, 1864, 180, in floating seaweed, no exact locality stated.
4092. *Antennarius avalonis* Jordan and Starks.
Santa Barbara Islands.
Antennarius avalonis Jordan and Starks, Proc. U. S. Nat. Mus., XXXII, 1907, 77, fig. 8, Santa Catalina Island at Avalon, Calif.
4093. *Antennarius verrucosus* Bean.
Bermudas.
Antennarius verrucosus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 31, St. George Island, Bermuda Islands.
4094. *Antennarius astroscopus* Nichols.
Barbados.
Antennarius astroscopus Nichols, Bull. Amer. Mus. Nat. Hist., XXXI, 1912, 109, Barbados.

Family 305. CHAUNACIDÆ

Genus 1464. CHAUNAX Lowe.

- Chaunax* Lowe, Proc. Zool. Soc. Lond., XIV, 1846, 339 (*C. pictus* Lowe).
4095. *Chaunax pictus* Lowe.
Madeira; taken off the coast of Rhode Island in the Gulf Stream.
Chaunax pictus Lowe, Trans. Zool. Soc. Lond., 1846, 339, Camera de Lobos, Madeira.
4096. *Chaunax nuttingi* Garman.
Florida keys,
Chaunax nuttingi Garman, Bull. Nat. Hist. Iowa Univ., 1896, 86, Sand Key Light, Fla.

Family 306. OGCOCEPHALIDÆ. Batfishes

Genus 1465. OGCOCEPHALUS Fischer. *Seabats*.

- Ogcocephalus* Fischer, Zoognosia, ed. 3, I, 1813, 70, 78 (*Lophius vespertilio* Linnæus).
Oncocephalus Gill, corrected spelling.
Malthe Cuvier, Règne Anim., ed. I, II, 1817, 311 (*Lophius vespertilio* Linnæus).
Malithæa of authors, modified spelling.

4097. *Ogcocephalus vespertilio* (Linnæus). *Batfish; Diablo*.
West Indies, north to Carolina.
Lophius vespertilio Linnæus, Syst. Nat., Ed. X, I, 1758, 236, American seas; after *Lophius frontii unicorn* of Artedi.
4098. *Ogcocephalus nasutus* (Cuvier and Valenciennes).
West Indies.
Malthæa nasuta Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 452, Martinique.
Malthæa notata Cuvier and Valenciennes, loc. cit., 453, Surinam; snout 15 in length; body spotted.
4099. *Ogcocephalus radiatus* (Mitchill).
Coast of Florida and neighboring waters; very common in shallow bays among weeds, especially about the Florida keys.
Lophius radiatus Mitchill, Amer. Month. Mag., March, 1818, 326, Straits of the Bahamas.
Malihe cubifrons Richardson, Fauna Bor.-Amer., III, 1836, 103; said to be from Labrador, which is certainly an error.
? *Malthæa truncata* Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 454, America; snout wholly obsolete; perhaps a species of *Zalieutes*.
- Genus 1466. **ZALIEUTES** Jordan and Evermann.
Zalieutes Jordan and Evermann, Check List, 1895, 511 (*Malihe elater* Jordan and Gilbert).
4100. *Zalieutes elater* (Jordan and Gilbert).
Pacific coast of Mexico, south to Panama, in water of moderate depth; very rare near the shore but obtained in abundance by the *Albatross* at stations 2794 and 2795 near Panama.
Malihe elater Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 365, Mazatlan.
- Genus 1467. **MALTHOPSIS** Alcock.
Malthopsis Alcock, Deep Sea Fish. Invest., Ann. Mag. Nat. Hist., VIII, 1891, 26 (*M. luteus* Alcock).
4101. *Malthopsis spinosa* Garman.
Off Bay of Panama in deep water.
Malthopsis spinosa Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 104, Pl. XXII, *Albatross* stations 3392 and 3393, off Panama, in 1,020 to 1,270 fathoms.
4102. *Malthopsis spinulosa* Garman.
Off Bay of Panama, in deep water.
Malthopsis spinulosa Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 106, Pls. XXI, XXVI, *Albatross* station 3394, lat. 7° 21' N., long. 79° 35' W. in 511 fathoms.
- Genus 1468. **HALIEUTICHTHYS** Poey.
Halieutichthys Poey, Proc. Acad. Nat. Sci. Phila., XV, 1863, 90 (*H. reticulatus* Poey = *Lophius aculeatus* Mitchill).
4103. *Halieutichthys aculeatus* (Mitchill).
Cuba to southern Florida.
Lophius aculeatus Mitchill, Amer. Month. Mag., II, 1818, 325, Straits of Bahamas.
Halieutichthys reticulatus Poey, in Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 91, Cuba.
4104. *Halieutichthys caribbæus* Garman.
Jamaica to Barbados, in depths of 70 to 150 fathoms or more.
Halieutichthys caribbæus Garman, Bull. Nat. Hist. Iowa Univ., 1896, 87, pl. 4, fig. 2, Jamaica to Barbados.
4105. *Halieutichthys smithii* Evermann and Marsh.
Porto Rico.
Halieutichthys smithii Evermann and Marsh, Bull. U. S. Fish Com., XX, 1900, 339, fig. 112, Mayaguez Harbor, Porto Rico.
- Genus 1469. **HALIEUTÆA** Cuvier and Valenciennes.
Halieutæa Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 455 (*H. stellata* Cuvier and Valenciennes).

4106. *Halieutæa spongiosa* Gilbert.
Pacific Ocean, off Lower California.
Halieutæa spongiosa Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 124,
west of Revillagigedo Islands, at *Albatross* station 2992, lat. $18^{\circ} 17'$
 $30''$ N., long. $114^{\circ} 43' 15''$ W., in 460 fathoms.
- Genus 1470. **HALIEUTELLA** Goode and Bean.
Halieutella Goode and Bean, Proc. Biol. Soc. Wash., II, 1885, 88 (*H. lappa* Goode and Bean).
4107. *Halieutella lappa* Goode and Bean.
Gulf Stream.
Halieutella lappa Goode and Bean, Proc. Biol. Soc. Wash., II, 1882, 88,
Gulf Stream at *Fish Hawk* station 1151, lat. $39^{\circ} 58' 30''$ N., long.
 $70^{\circ} 37'$ W., in 125 fathoms.
- Genus 1471. **DIBRANCHUS** Peters.
Dibranchus Peters, Monatsber. Akad. Wiss. Berlin, 1875, 736 (*D. atlanticus* Peters).
4108. *Dibranchus atlanticus* Peters.
Gulf Stream.
Dibranchus atlanticus Peters, Monatsber. K. Akad. Wiss. Berlin, 1875,
736, with plate; West Africa, lat. 10° N. long. 17° W., in 360 fathoms.
- Family 307. **CERATHIDÆ.** Sea devils
- Genus 1472. **CERATIAS** Krøyer.
Ceratias Krøyer, Naturhist. Tidsskr., I, 1845, 639 (*C. hollbolli* Krøyer).
4109. *Ceratias hollbolli* Krøyer.
Greenland to Nova Scotia; four specimens known.
Ceratias hollbolli Krøyer, Naturh. Tidsskr., 1844, 639, Greenland.
- Genus 1473. **MANCALIAS** Gill.
Mancalias Gill, Proc. U. S. Nat. Mus., I, 1878, 227 (*Ceratias uranoscopus* Murray).
Typhlosaras Gill, Forest and Stream, 1883, Nov. 8, 284 (*T. shufeldti* Gill).
4110. *Mancalias uranoscopus* (Murray).
Deep seas; taken at Madeira and off the coast of southern New England.
Ceratias uranoscopus Murray, in Wyville Thompson, The Atlantic, II,
1878, 67, off Madeira.
4111. *Mancalias shufeldti* (Gill).
Western Atlantic.
Typhlosaras shufeldti Gill, Forest and Stream, November 8, 1883, 284,
western Atlantic.
- Genus 1474. **CRYPTOSARAS** Gill.
Cryptosaras Gill, Forest and Stream, Nov. 8, 1883 (*C. couesi* Gill).
4112. *Cryptosaras couesi* Gill.
Gulf Stream.
Cryptosaras couesi Gill, Forest and Stream, Nov. 8, 1883, 284, Gulf
Stream off New England.
- Genus 1475. **EDRIOLYCHNUS** Regan.
Edriolychnus Regan, Proc. Roy. Soc. Lond., ser. B, XCVII, 1925, 398
(*E. schmidt* Regan).
4113. *Edriolychnus schmidt* Regan.
Deep waters of the western Atlantic.
Edriolychnus schmidt Regan, Proc. Royal Soc. Lond., ser. B, XCVII,
1925, 398, western Atlantic; *Dana* expedition.
- Genus 1476. **MONOCERATIAS** Gilbert.
Monoceratias Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 379 (*M. acanthias* Gilbert).
4114. *Monoceratias acanthias* Gilbert.
Off Santa Barbara Islands, Calif.
Monoceratias acanthias Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915,
379, pl. 22, fig. 24, off Santa Cruz Island, 764 to 891 fathoms.

Genus 1477. BOROPHRYNE Regan.

Borophryne Regan, Ann. Mag. Nat. Hist., ser. 9, XV, 1925, 564 (*B. apogon* Regan).

4115. *Borophryne apogon* Regan.

Gulf of Panama.

Borophryne apogon Regan, Ann. Mag. Nat. Hist., ser. 9, XV, 1925, 564, Gulf of Panama.

Genus 1478. PHOTOCORYNUS Regan.

Photocorynus Regan, Proc. Roy. Soc. Lond., ser. B, XCVII, 1925, 393 (*P. spiniceps* Regan).

4116. *Photocorynus spiniceps* Regan.

Deep waters of Panama.

Photocorynus spiniceps Regan, Proc. Roy. Soc. Lond., ser. B, XCVII, 1925, 393, figs. 6-7, deep waters off Panama.

Family 308. HIMANTOLOPHIDÆ**Genus 1479. ONEIRODES Lütken.**

Oneiroides Lutken, Overs. Dansk. Vid. Selsk. Kjøbenhavn, 1871, 56 (*O. eschrichtii* Lütken).

4117. *Oneiroides eschrichtii* Lutken.

Deep sea off Greenland.

Oneiroides eschrichtii Lütken, Overs. Dansk. Vid. Selsk. Forhandl., 1871, 56, 9-18, pl. 2, deep sea off Greenland.

Genus 1480. HIMANTOLOPHUS Reinhardt.

Himantolophus Reinhardt, Dansk. Vid. Selsk. Afh. Kjøbenhavn, VII, 1838, 74 (*H. grænlandicus* Reinhardt).

4118. *Himantolophus grænlandicus* Reinhardt.

Greenland.

Himantolophus grænlandicus Reinhardt, Dansk. Vid. Selsk. Nat. Math. Afh., 1837, 74, Greenland.

Genus 1481. LOPHODOLUS Lloyd.

Lophodolus Lloyd, Mem. Indian Mus., II, 1909, 167 (*L. indicus* Lloyd).

4119. *Lophodolus acanthognathus* Regan.

North Atlantic.

Lophodolus acanthognathus Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 563, north Atlantic.

Genus 1482. CORYNOLOPHUS Gill.

Corynolophus Gill, Proc. U. S. Nat. Mus., I, 1878, 219 (*Himantolophus reinhardtii* Lütken).

4120. *Corynolophus reinhardtii* (Lütken).

Greenland; one specimen known.

Himantolophus reinhardtii Lütken, Kong. Dansk. Vid. Selsk., 1878, 321, Greenland. Perhaps a species of *Himantolophus*.

Genus 1483. DOLOPICHTHYS Garman.

Dolopichthys Garman in Mem. Mus. Comp. Zool., XXIV, 1899, 81, (*D. allector* Garman).

4121. *Dolopichthys gracilispinis* Regan.

North Atlantic.

Dolopichthys gracilispinis Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 563, north Atlantic.

4122. *Dolopichthys microlophus* Regan.

North Atlantic.

Dolopichthys microlophus Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 563, north Atlantic.

4123. *Dolopichthys anisacanthus* Regan.

North Atlantic.

Dolopichthys anisacanthus Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May 1925, 562, north Atlantic.

4124. *Dolopichthys luetkeni* Regan.
Gulf of Panama.
Dolopichthys luetkeni Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 562, Gulf of Panama.
4125. *Dolopichthys heteracanthus* Regan.
Gulf of Panama.
Dolopichthys heteracanthus Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 563, Gulf of Panama.
- Genus 1484. *LASIOGNATHUS* Regan.
Lasiognathus Regan, Ann. Mag. Nat. Hist., 9 ser., XV, 1925, 563 (*L. saccoctoma* Regan).
4126. *Lasiognathus saccoctoma* Regan.
Caribbean Sea.
Lasiognathus saccoctoma Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 563, Caribbean Sea.
- Genus 1485. *LINOPHRYNE* Collett.
Linophryne Collett, Proc. Zool. Soc. Lond., 1886, 138 (*L. lucifer* Collett).
4127. *Linophryne lucifer* Collett.
Open Atlantic.
Linophryne lucifer Collett, Proc. Zool. Soc. Lond., 1886, 138, pl. 15, off Madeira, lat. 36° N., long. 20° W.
4128. *Linophryne polypogon* Regan.
North Atlantic
Linophryne polypogon Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 565, north Atlantic.
4129. *Linophryne arborifer* Regan.
North Atlantic.
Linophryne arborifer Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 564, north Atlantic.
4130. *Linophryne macrodon* Regan.
Gulf of Panama.
Linophryne macrodon Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1895, 564, Gulf of Panama.
- Genus 1486. *CHÆNOPHRYNE* Regan.
Chænophryne Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 564 (*C. longiceps* Regan).
4131. *Chænophryne longiceps* Regan.
North Atlantic and Gulf of Panama.
Chænophryne longiceps Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 564, north Atlantic.

Family 309. MELANOCETIDÆ

- Genus 1487. *MELANOCETUS* Günther.
Melanocetus Günther, Proc. Zool. Soc. Lond., 1864, 301 (*M. johnsoni* Günther).
4132. *Melanocetus polyactis* Regan.
Gulf of Panama.
Melanocetus polyactis Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 565, Gulf of Panama.
4133. *Melanocetus niger* Regan.
Gulf of Panama.
Melanocetus niger Regan, Ann. Mag. Nat. Hist., ser. 9, XV, May, 1925, 565, Gulf of Panama.
- Genus 1488. *LYOCETUS* Günther.
Lyocetus Günther, *Challenger* Rept., XXII, 1887, 57 (*L. murrayi* Günther).

4134. *Lyocetus murrayi* (Günther).Mid-Atlantic at *Challenger* stations 106 and 348.*Melanocetus (Liocetus) murrayi* Günther, *Challenger* Rept., XXII, 1887, 57, pl. 11, fig. A, mid-Atlantic.**Family 310. CAULOPHRYNIDÆ****Genus 1489. CAULOPHRYNE Goode and Bean.***Caulophryne* Goode and Bean, *Oceanic Ichth.*, 1895, 496 (*Caulophryne jordani* Goode and Bean).**4135. *Caulophryne jordani* Goode and Bean.**

Gulf Stream.

Caulophryne jordani Goode and Bean, *Oceanic Ichth.*, 1895, 496, pl. 21, fig. 409, Gulf Stream off Carolinas in 1,276 fathoms.**Family 311. ACERATIIDÆ****Genus 1490. RHYNCHOCERATIAS Regan.***Rhynchoceratias* Regan, *Ann. Mag. Nat. Hist.*, 9 ser., XV, May, 1925, 566 (*R. brevirostris* Regan).**4136. *Rhynchoceratias leucorhinus* Regan.**

North Atlantic and Gulf of Panama.

Rhynchoceratias leucorhinus Regan, *Ann. Mag. Nat. Hist.*, 9 ser., XV, May, 1925, 566, north Atlantic.**4137. *Rhynchoceratias rostratus* Regan.**

Gulf of Panama.

Rhynchoceratias rostratus Regan, *Ann. Mag. Nat. Hist.*, 9 ser., XV, May, 1925, 567, Gulf of Panama.

INDEX

LIST OF COMMON NAMES OF FISHES OCCURRING IN CHECK LIST

Page	Page	Page
abadejo..... 315	anged..... 40	barracuda..... 255
aboma de mar..... 436	angel, black..... 360	California..... 255
aboma de rio..... 442	angelfish..... 22, 358, 361	European..... 255
aburabozu..... 373	angel shark, California..... 22	great..... 255
acedia..... 231	angel sharks..... 22	northern..... 255
achigan grand bouche..... 297	angler..... 504	short..... 255
achigan noir..... 297, 298	añil..... 321	ballacudas..... 255
achigan petite bouche..... 298	antonino..... 270	barred marlin..... 263
acoupa..... 354	aguaji..... 314	barred pickarel..... 173
aguaji..... 314	arctic charr..... 61	bashaw..... 155
agua vina..... 318	arctic flounder..... 227	basking shark..... 20
agua de casta..... 196, 263	arctic grayling..... 66	East Indian..... 18
agujon..... 196, 197	arctic lake herring..... 62	basking sharks..... 20
aku..... 258	arctic lamprey..... 10	bass..... 298, 348
akule..... 271	arctic lycodes..... 476	banklick..... 305
akullakitsok..... 474	arctic salmon, Ross's..... 60	bayou..... 297
Alabama darter..... 292	arctic sculpin..... 386	big-mouthed black..... 297
Alabama shad..... 43	argentino, sheppy..... 73	black..... 298, 317
Alaska blackfish..... 172	argentines..... 68	fresh-water..... 298
Alaska codfish..... 210	arnillo..... 325	of the Huron..... 297
Alaska greenfish..... 374	arrow-toothed halibut..... 221	black sea..... 317
Alaska pollack..... 209	Athabaska herring..... 62	blue..... 298
Alaska sand lance..... 451	Athabaska minnow..... 115	brown..... 298
albacore..... 200	atunga..... 502	buffalo..... 302
West Indies long-finned..... 260	atka fish..... 374	calico..... 305
yellow-finned..... 260	atka mackerel..... 374	channel..... 348
albacores..... 259	atka mackerels..... 374	common rock..... 303
long-finned..... 260	Atlantic salmon, common..... 50	cow..... 298
yellow-finned..... 260	atule..... 271	dwarf..... 298
aleky trout..... 212	Aubeaubee darter..... 291	fresh-water..... 303
alewife..... 41	aurora trout..... 60	gold..... 298
alewives..... 41	awa..... 39, 40	grass..... 305
añil..... 248		gray..... 297
alfone..... 411		green..... 297, 298
alfoncino..... 295	bacalao..... 315	gulf sea..... 318
alfonsin a casta..... 234	bacalhao sabara..... 441	hog..... 298
alfonsin a casta larga..... 234	bachelor..... 304, 305	kelp..... 317
alfonsinos..... 234	bachelor perch..... 304	Kentucky black..... 298
ahlonghi..... 260	bagré..... 151, 157	lake..... 297, 303, 305
alligatorfish, common..... 394	bagré sapo..... 488	Lake Erie..... 305
alligatorfishes..... 393	bagrés de rio..... 156	Lake Erie..... 304
alligator gar, Cuban..... 38	Bahama amphioxus..... 7	little..... 297
Mississippi..... 38	Bahama lancelet..... 7	little..... 298
alligator gars..... 38	bejonado..... 336	marsh..... 297
Allison's tuna..... 266	balao..... 199	minny..... 298
allmouth..... 504	blue-tailed..... 199	moss..... 297
alma seca..... 98	red-tailed..... 199	mountain..... 305
amberfish, great..... 279	balloonfish..... 501, 502	mud..... 297, 303
amberjack..... 279	spotted..... 502	Oswego..... 297
American anchovies..... 47	unspotted..... 502	pale river..... 297
American carp..... 103	bananafish..... 39	red..... 348
American eel..... 77	banded blenny..... 462	river..... 297, 298
American hake..... 214	banded darter..... 289	rock..... 297, 298, 317
American perch..... 281	banded drum..... 352	rock sea..... 318
American pike..... 173	banded larmus..... 344	round..... 304
American pike perches..... 282	banded mackerel..... 279	sea..... 317, 348
American saibling..... 60	banded rudderfish..... 279	shining..... 304
American seahorse, com- mon..... 244	banded sunfish..... 304	silver..... 305, 307
American shad..... 42	bang..... 42	slough..... 297
American smelt..... 67	banklick bass..... 305	small-mouthed black..... 298
American sole..... 229	banner pompano..... 276	southern rock..... 303
amphioxus, Bahama..... 7	barracuda..... 255	speckled..... 305
California..... 7	barb..... 350	spotted..... 297, 348
anchoa pelada..... 45	barbero..... 362, 363	spotted river..... 298
anchovies..... 47	barbero negro..... 363	stone..... 308
American..... 47	barbiers..... 322	straw..... 297, 305
anchovy, bay..... 47	barbu..... 256	strawberry..... 302, 305
California..... 47	barbudo..... 157, 234, 256	streaked-cheeked river..... 298
fy..... 48	barbudos..... 234, 256	striped..... 297, 307
Porto Rican..... 51	barfish..... 305	sun..... 302
striped..... 48	barndoor skate..... 25	sunfish..... 303
	barracuta..... 255	swago..... 298

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
bass, trout.....	298	black cod.....	373	blueback mullet.....	253
white.....	297, 303, 307	black crappie.....	305	blueback salmon.....	55
white lake.....	307	black croaker.....	348	blueback trout.....	61
white sea, of California.....	356	black drum.....	352	of Lake Crescent.....	57
wood.....	298	black-eared pondfish.....	299, 301	blue bass.....	298
yellow.....	297, 298, 307	blackears.....	300	blue bream.....	301
bass hogfish.....	298	blackeyes.....	298	blue-breasted darter.....	290
bass killifish.....	176	blackeye sunfish.....	298	blue cat.....	152
bass sunfish.....	303	blackfin.....	126	blue cod.....	374
basses, black.....	297	blackfin of Lake Michigan.....	63	blue darter.....	291
rock.....	303	blackfin snapper.....	326	blue dog.....	13
round.....	304	Alaska.....	112, 287, 317, 424	bluefish.....	290, 293, 317
sea.....	303	smooth.....	172	California.....	339, 356
bastard halibut.....	223	black fresh-water bass.....	293	bluefishes.....	280
bastard halibuts.....	222	black grouper.....	312, 314	bluegill.....	301
bastard margaret.....	329	black groupers.....	312	bluehead sucker.....	104
bastard weakfish.....	355	black grunt.....	323, 329	blue herring.....	42
batfish.....	31, 403, 507	black guatavere.....	309	blue Joe.....	301
batfishes.....	508	black Harry.....	317	blue Johnny.....	291
baya.....	314	blackhead minnow.....	145	blue-lined minnow.....	132
bay anchovy.....	47	blackhorse.....	102	blue-mouthed sunfish.....	301
bay mackerel.....	257	black huron.....	297	blue mullet.....	110
bay mare.....	105	black jewfish.....	312	blue parrotfish.....	432, 434
bay shark.....	15	black lamprey, small.....	9	blue perch.....	301, 424
bayou bass.....	297	black marlin.....	263	blue pike.....	282
Bear Lake herring, great.....	62	giant.....	263	blue runner.....	272
Bear Lake whitefish.....	65	black minnow.....	139	blue shark, great.....	17
beau gregoire.....	413	91, 92.....	91, 92	small.....	16
becuna.....	255	black moray.....	105	blue smelt, California.....	252
bellowsfish.....	246, 504	black mullet.....	139, 140	blue-spotted sunfish.....	298, 304
bellyfish.....	191	black-nosed dace.....	493	blue-striped triggerfish.....	401
bergal.....	424	black oldwife.....	493	blue sucker.....	146
berg-gylt.....	424	black perch.....	298, 317, 323	blue sunfish.....	298, 301
Bermuda catfish.....	261	black pilot.....	413	blue surgeon.....	362
Bermuda chub.....	340	black rockfish.....	365	blue-tailed halao.....	199
Bermuda lancelet.....	7	black rudderfish.....	268	blue tang.....	362
Bermuda silk snapper.....	328	black ruddersfishes.....	267	blunthead.....	176
berrugate.....	324	black ruffe.....	267	blunt-nosed minnow.....	145
besan.....	362	black ruffs.....	267	blunt-nosed river carp.....	103
beshow.....	373	black sculpin.....	387	blunt-nosed shiner.....	275
bessy cerka.....	491	black sea bass.....	317	boar grunt.....	330
besugo.....	337	black-sided darters.....	263	bobo.....	255
blajaiba.....	327	blacksmith.....	316	bob-tailed skate.....	25
blajaiba de lo alto.....	303	black soapfish.....	263	bocaccio.....	364
bielaya ryba.....	54	black spearfish.....	263	bocce dulce.....	13
big drum.....	362	black-spotted Spanish.....	257	boccon.....	48, 51
big-eared sunfish.....	300	mackerel.....	108, 107	boccone.....	354
big-eye.....	322, 323	black sucker.....	302	bodieron.....	373
big-eyed herring.....	39, 42	black sunfish.....	411	boga.....	340
big-eyed scad.....	271	black surffish.....	452	bonaci.....	314
bigeye minnow.....	134	black swallower.....	452	bonaci arara.....	314
big-eyes.....	322	black swallows.....	300	bonaci cardenal.....	313
big-headed darter.....	284	black-tailed sunfish.....	302	bonaci de Piedra.....	313
big-headed gurnard.....	403	black warmouth.....	317	bonaci gato.....	315
biglip sucker.....	106	black will.....	109	bone-dog.....	19, 21
bigmouth.....	302	black winter sucker.....	298	bone-eater.....	259
bigmouth buffalo.....	102	blanco perch.....	32	bonefish.....	39
big-mouthed black bass.....	297	blanketfish.....	357	bone-jack.....	259
big-mouthed buffaloish.....	102	blanquillo.....	357	bone shark.....	20
big-mouthed cat.....	153	blanquillos.....	460	bonita of Martinique.....	260
big-mouthed sunfish.....	302	blennies.....	462	bonito.....	258, 259, 279, 281
bigmouth surffish.....	411	banded.....	462	California.....	259
big-nose sunfish.....	299	eel.....	471	oceanic.....	258
big skate of California.....	26	four-banded.....	462	Pacific.....	258
big-toothed redhorse.....	111	seaweed.....	461	bonitos.....	259
bullfish.....	36, 196, 197, 242	snake.....	471	bonnethead shark.....	17
billy gar.....	36	spotted snake.....	499	Bonneville whitefish.....	65
bishop ray.....	30	blind catfishes.....	154	bonyfish.....	39, 44
Bissell's herring.....	63	blindfishes, cave.....	194	bonyfishes.....	39
bitterhead.....	115, 305	blindfish, Mammoth Cave.....	195	bony-tail.....	114
black and yellow rock cod.....	369	blindfish, small.....	195	bootleg sucker.....	107
black angel.....	360	blind gobies.....	447	boregat.....	373
black-banded rock cod.....	370	blind goby of Point Loma.....	447	borer.....	8
black-banded sunfish.....	303	blind simon.....	287	borders.....	8
black bass.....	293, 317	bloater.....	63	borlase.....	267
big-mouthed.....	297	of Lake Michigan.....	63	Bosc's shanny.....	462
Kentucky.....	293	blob.....	385	Boston mackerel.....	257
largemouth.....	297	bloody sunfish.....	300	botete.....	498
large-mouthed.....	297	blower.....	498	boucanello.....	326
of the Huron.....	297	blowfish.....	499	bout de tabac.....	319
small-mouthed.....	298	hairy.....	499	bowfin.....	38
black basses.....	297	spinyback.....	499	bowfins.....	38
black-belly.....	42	blue-and-green sunfish.....	298	boxfish.....	497
black buffalo.....	102	blue angel.....	361	hairy.....	501
black bullhead.....	154	blueback.....	42, 62	spiny.....	502
black burrfish.....	502			bramble shark.....	22

List of common names of fishes occurring in check list—Continued

Page	Page	Page
bramble sharks..... 22	bullhead shark..... 10	capitaines..... 425
branch herring..... 42	little..... 22	caranx, yellow..... 273
branch perch..... 302	bullhead sharks..... 10	carbonero..... 271
brassy sculpin..... 386	bullion..... 434	caronetta..... 16
breasm..... 115, 298-303, 336,	bull redfish..... 245	cardenal..... 295
blue..... 301	bull trout..... 61	cardinalfishes..... 295
calico..... 305	bumper..... 276	cardonniera..... 370
copper-headed..... 301	burbot..... 212	Carolina pompano..... 277
copper-nosed..... 301	burbot..... 212	Carolina robin..... 407
little..... 304	burfish..... 502	Carolina sea robin..... 407
moremouth..... 302	black..... 502	Carolina whiting..... 350
perchmouth..... 302	burfishes..... 502	carp..... 108
red-bellied..... 300	burrito..... 333, 334	American..... 108
redbreast..... 299	burritos..... 333	blunt-nosed river..... 108
red-eyed..... 303	burro..... 334	common river..... 108
red-headed..... 299	burro blanco..... 333	lake..... 103
red-sided..... 113	burros..... 333	silver..... 103
red-tailed..... 299	butter chub..... 146	carp sucker..... 103
sea..... 285	butterfish..... 267, 277, 309, 467,	carp suckers..... 103
spotted..... 299	lemon-yellow..... 309	casabe..... 276
warmouth..... 302	butterflyfishes..... 359	casabes..... 276
yellow..... 302	butterfly ray..... 29	catagnole..... 265
breams, sea..... 285		catalina..... 332
bride perch..... 297	caballa..... 257	catalineta..... 332, 361
bridge perch..... 304	caballito del mar..... 244	catalineta..... 361
bridled minnow..... 135	cabelellote..... 325	catalufa..... 322, 323
brier ray..... 25	caazon..... 344, 352, 387	catalufa de lo alto..... 323
brilliant sunfish..... 300	smooth..... 390	catalufas..... 322
bristle herring..... 246	cabezones..... 382	deep-water..... 323
broadbill..... 264	caberote..... 246	cat, big-mouthed..... 153
broadhead, grubber..... 47, 50	cabra mora..... 310	blue..... 152
broad shad..... 341	cabrilla..... 311, 317	brown..... 154
broad soles..... 229	spotted..... 317	channel..... 153
brochet de mer..... 305	cabrilla calamaria..... 314	of the Potomac..... 153
bronze-backer..... 298	cabrilla de astillero..... 314	duckbill..... 36
bronzed centrarchus..... 303	cabrilla de raizero..... 313	eel..... 152
bronze minnow..... 121	cabrilla pinta..... 310	Fulton..... 152
brook lamprey..... 9, 10	cabrilla pirita..... 315	great fork-tailed..... 152
brook lampreys..... 10	cabrillas verdes..... 317	long-tailed..... 212
brook minnow..... 140	cachucho..... 328	Mississippi..... 152
brook silverside..... 251	Caesar..... 331	mud..... 154, 155
brook stickleback..... 238	Caesar grunt..... 330	Russian..... 155
brook sucker..... 105	caffa bota..... 11	Sacramento..... 154
brook trout..... 60	cagon de lo alto..... 328	Schuylkill..... 154
eastern..... 60	cailieu..... 43	sea..... 149
of western Oregon to	cailieu tassart..... 44	spoonbill..... 36
southern Alaska..... 58	cailieu..... 440	stone..... 155
brotnula..... 479	calf..... 325	white..... 152, 153
brown bass..... 298	calafate..... 493	willow..... 152
brown buffalo..... 107	calico bass..... 305	yellow..... 153, 155
brown cat..... 154	calico bream..... 305	catfish..... 176, 256
brown hind..... 308	calico salmon..... 54	sea..... 148, 149
brown rock cod..... 369	California amphioxus..... 7	small..... 154
brown rockfish..... 368	California anchovy..... 47	catfishes, blind..... 154
brown trout..... 298	California angel shark..... 22	cave..... 159
brown-winged sea robin..... 407	California barracuda..... 255	gafttopsail..... 147
bubbler..... 352	California bluefish..... 339, 356	mailed..... 160
buffalo..... 103	California blue smelt..... 252	sea..... 147
bigmouth..... 102	California bonito..... 259	South American..... 156
black..... 102	California common skate..... 26	cats, channel..... 151
brown..... 102	California dogfish..... 21	stone..... 155
gourdseed..... 102	California flounder..... 226	tadpole..... 155
mongrel..... 102	California garfish..... 196	cat shark..... 14
quillback..... 102	California hagfish..... 8	cat sharks..... 11
razor-backed..... 102	California herring..... 41	cavalla..... 257, 273
round..... 102	California jewfish..... 308	cavalias..... 269
smallmouth..... 102	California lancelet..... 7	cave blindfishes..... 194
suckermouth..... 102	California pampano..... 266	cave catfishes..... 159
buffalo bass..... 302	California redfish..... 426	cavefish..... 194
buffalo cod..... 374	California sardine..... 43	Missouri..... 195
buffalohead, big-mouthed..... 102	California sole..... 222	cazon de playa shark..... 16
common..... 102	California stingray..... 31	céfalo..... 253
redmouth..... 102	California tomcod..... 210	centrarchus, bronzed..... 303
buffaloheadfishes..... 102	California torpedo..... 27	carnier..... 308
buffalo sculpin..... 381	California whiting..... 350	cero..... 257
buffalo sunfish..... 298	Califaver mullet..... 253	spotted..... 257
bugara..... 412	cambellite..... 304	chagrin..... 18
bugfish..... 44	candil..... 235	chamside..... 301
bullfish..... 497	candlefish..... 66	chan-sided perch..... 301
bullhead..... 385, 386	cane de mare..... 19	chan..... 40
black..... 154	capellin..... 66	channel bass..... 348
common..... 154	capellins..... 66	channel cat..... 152
prickly..... 383	Cape May goody..... 349	of the Potomac..... 153
Rocky Mountain..... 384	capidoli..... 20	channel cats..... 151
bullhead minnow..... 145	capitaine..... 425	chapin..... 499, 497

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
chappaul	114	coachman	320	common weakfish	355
characins	95	coalfish	209, 373	common whitefish	65
charoteer	320	coast-range trout	58	compressed redfin	126
charma ryba	172	cobbler	175	conejo	261
charr, Arctic	61	common	176	conejos	261
Greenland	60	cobblerfish	274, 277	coney	308
Oregon	61	cobblerfishes	274	conger, sand	454
charrs	60	Cobessicentic smelt	67	conger eel	54, 79, 473
chauffe soleil	413	cobia	281	of California	91
Cheney's darter	286	cobinito	363	conger eels	79
cherna americana	312	cobino	491	Connecticut River shad	42
cherna criolla	311	cocinera	272	conner	424
cherna de lo alto	310	cocinero	272	constantino	306
cherna de viveiro	312	cocinero dorado	272	constantino de las aletas	
chicharro	271	cock and hen paddle	397	prietas	306
chickwick	355	cockeye pilot	413	Copeland's darter	285
chicolar	261	cockfishes	274	copper-headed bream	301
chigh	113	cocuyo	493	copper-nosed bream	301
chimeras	10, 32	cod, black	373	coralfishes	359
chimeroids	32	blue	374	corbineta	346
chumook salmon	55	buffalo	374	corneta	245
chunquapin perch	299, 304, 305	cultus	374	cornetfishes	245
churivita	360	green	209	coronado	279
churivitas	360	Pacific	210	corporal	117
chuselmouth	112	wachna	210	corsair	367
chuselmouth jack	66	codfish, Alaska	210	corvalos	351
chuselmouths	112	common	210	corvina	344, 349, 355
chivin	117	Greenland	210	corvina de las aletas am-	
chivo	344	codfishes	209, 210	arillas	356
chogset	424	codfish shark	21	couchu	161
chopa	339	codling	213	coua	108
chopa amarilla	339	spotted	213	Coulter's whitefish	65
chopa blanca	340	codlines	212	courpata	269
chopa spina	337	codorniz	350	cow bass	297
chopper	102	cods, cultus	374	cowfish	31, 497
choupiquel	38	cod shark	21	cow-nosed ray	31
chub	63, 64, 109, 115, 117, 137, 297, 302, 340	coelho	261	cow pilot	415
Bermuda	340	coho salmon	54	cow shark	11
butter	146	cojinera	272	cow sharks	11
Columbia River	112	Colorado River squawfish	114	crab-eater	281
creek	117	Colorado River trout	57	crab-eaters	281
flat-headed	136	Columbia River chub	112	crabra	370
greased	146	Columbia River salmon	55	Cragin's darter	292
Indian	137	Columbia River sucker	106	Craig fluke	228
Kawah	113	Columbia River trout	56	crampfish	27
lake	136	common alligatorfish	394	crapet	304, 305
millpond	297	common American seahorse	244	carpet jaune	302
mud	117, 303	common Atlantic salmon	56	carpet mondoux	303
nigger	146	common bullhead	102	carpet noir	303
of the Rio Grande	120	common codfish	154	carpet vert	303
river	137	common cobbler	176	crappie	304, 305
Sacramento	120	common codfish	210	black	305
silver	117, 137	common dolphin	264	pole	304
southern	297	common eastern stickleback	238	timber	304
steel-backed	146	common flatfish	227	white	304
Storer's	137	common gar pike	36	crappies	304
chub mackerel	258	common gray shark	15	cravo	216
chub minnow	136	common grunt	330	crawl-a-bottom	107, 111, 283
chub sucker	108, 109	common gurnard	407	creek chub	117
chuchin	158	common halfbeak	198	creekfish	109
chum	54	common halibut	221	creek sucker	109
chuss	213	common herring	41	creole	426
cibi	271	common killifish	175	crolefish	320
cibi amarilla	272	common lake herring	63	crested gobies	438
cibi mancho	271	common mackerel	257	crevallé	272, 273
cigarfish	270	common mullet	253	horse	273
curstomes	7	common pampano	277	crimson darter	294
cisco, Lake Michigan	63	common pipefish	242	croaker	349, 352
mooneye	61	common rat-tail	205	black	348
of Green Lake	64	common redhorse	110	white	353
of Lake Bonneville	63	common river carp	103	yellow-tailed	350
of Lake Michigan	64	common rock bass	303	croakers	344
of Lake Ontario	63	common sawfish	23	cuatro ojos	194
of Lake Superior	63	common sculpin	386	cub	424
of Lake Tippecanoe	62	common scup	336	Cuban alligator gar	88
ciscos	61	common shad	42	cubera	325
cuvetta de Mer	409	common skate	25, 26	Cub shark	15
clamagore	434	common spotted moray	91	ruck old	497
clam-cracker	28	common stickleback	237	cucumferfish	502
clear-nosed skate	25	common sturgeon	34	cucuyo	491
clear ray	25	common sucker	105	cultus cod	374
clingfishes	498	common sunfish	300, 302	cultus cods	374
clinging goby	446	common surgeon	362	Cumberland darter	293
cloudy lycoodes	475	common swordfish	264	cunprida	234
		common trunkfish	497	cunner	424

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
cunners.....	424	dogfish.....	13, 19, 21, 174, 176	eels.....	54, 77
cusk.....	213, 214	California.....	21	conger.....	79
European.....	214	horned.....	19	cusk.....	485
rock.....	385	picked.....	19, 21	duckbill.....	82
cusk eels.....	485	smooth.....	13	long-necked.....	77
cutlassfish.....	262	spiny.....	21	ooze.....	78
cutlassfishes.....	262	dogfishes.....	21	snake.....	84
cut-lips.....	112, 146	dog salmon.....	54	snipe.....	83
cut-throat trout.....	56	dog shark.....	13	snub-nosed.....	78
of Utah Lake.....	56	dog snapper.....	325	spiny.....	75
cybium, spotted.....	257	dollardee.....	301	sym branchoid.....	76
		dollarfish.....	267, 275	thread.....	83
dab.....	222	dollfish.....	360	true.....	77
sand.....	222	Dolly Varden.....	305	worm.....	82
smear.....	228	Dolly Varden trout.....	61	eighteen-spined sculpin.....	386
dabs, mud.....	226	dolphin, common.....	264	eight-threaded threadfish.....	256
smear.....	228	small.....	265	electricfish.....	27
dace.....	115, 117, 128, 140	dolphins.....	264	electric rays.....	27
black-nosed.....	139, 140	dómine.....	261	elephantfish.....	33
horned.....	117	doncella.....	426	elephantfishes.....	32
long-nosed.....	139	doncellas.....	426	elephant shark.....	20
red-bellied.....	113	dorado.....	264	ellwife.....	42
rosy.....	121	dorados.....	264	emeraldfish.....	441
striped.....	140	dormeur.....	323	emerald minnow.....	123
daces.....	112	dory.....	282	emerald trout.....	58
daddy sculpin.....	387	hair-finned.....	275	emerande.....	441
daddy sculpins.....	386	ocellated.....	232	endormi.....	441
darter, Alabama.....	292	spinous.....	276	enjambe.....	308
Aubeenaubee.....	291	dotted paintedtail.....	297	entemedor.....	27
banded.....	289	dotted scad.....	270	Erie great herring.....	63
big-headed.....	284	double-bearded flyingfish.....	201	Erie herring.....	63
black-sided.....	283	doughbelly.....	146	erizo.....	501
blue.....	291	dourade.....	264	escolar.....	261
blue-breasted.....	290	dovetailfish.....	415	escolar chimo.....	297, 328
Cheney's.....	286	dragonets.....	450	escolar de natura.....	269
Copeland's.....	285	drum.....	103, 352	escolares.....	260, 261
Cragin's.....	282	banded.....	352	escribano.....	198, 199
crimson.....	284	big.....	352	esmeralda.....	441
Cumberland.....	283	black.....	352	esmeralda de mar.....	437
Evermann's.....	284	fresh-water.....	352	esmeralda negra.....	437
fan-tailed.....	293	gray.....	352	espada.....	264
fasiform.....	294	red.....	348	espadón emperador.....	264
Gilbert's.....	286	drumfish.....	352	eulachon.....	66
Grand Oranchoe.....	287	drummer, Jewsharp.....	351	European barracuda.....	255
green-sided.....	286	mongol.....	355	European cusk.....	214
Gunter's.....	283	whitemouth.....	348	European hake.....	214
Jessie's.....	291	drums, red.....	348	European sculpin.....	386
Johnny's.....	287	river.....	352	Everglades pugnny sunfish.....	305
Jordan's.....	290	sea.....	352	Evermann's darter.....	284
Julia's.....	292	drunkenfish.....	496	Evermann's trout.....	59
least.....	294	Dublin Pond trout.....	60		
manitou.....	283	duckbill cat.....	36	faguto.....	191
Maryland.....	284	duckbill eels.....	82	fair maid.....	336
Maxinkuckee.....	284	dusky shark.....	16	fallfish.....	117
Meadie's.....	286	dwarf bass.....	298	red.....	127
Niangua.....	285			fallfishes.....	117
northern.....	291	Eagle Lake trout.....	58	fall herring.....	42
ouachita.....	284	eagle rays.....	30, 31	fanegal.....	370
Putnam's.....	285	eared sunfish.....	300	fantail darters.....	293
rainbow.....	291	Easter mackerel.....	258	fan-tailed darters.....	293
sand.....	287, 288	eastern brook trout.....	60	fantail mullet.....	253, 254
shielded.....	283	eastern mud minnow.....	174	fatback.....	44, 280
Shumard's.....	285	eastern pickerel.....	173	fathead.....	426
Susan's.....	288	eastern rehorse.....	111	fathead minnow.....	145
tassellated.....	287	eastern stickleback, com- mon.....	238	fatheads.....	145, 428
tessellated.....	290	East Indian basking shark.....	18	fatherlasher.....	386
Tuscumbia.....	293	eel.....	91	fat priest.....	373
Whipple's.....	292	American.....	77	fat priests.....	373
darters.....	282	conger.....	54, 79, 473	female perch.....	302
black-sided.....	283	of California.....	91	fiatolas.....	266
fantail.....	293	fresh-water.....	77	fiddlefish.....	22
sand.....	289	lamper.....	473	fiddler.....	152
tessellated.....	287	lamprey.....	8	fiddlerfish.....	23
dealfish.....	216	moray.....	93	filefish.....	491, 494, 495
deceptive sunfish.....	299	sand.....	451, 485	orange.....	495
deep-water catalufus.....	323	sea.....	88	thread.....	494
démousselle.....	415	snipe.....	83	filefishes.....	493
dentuda.....	19	whipsnake.....	88	fine-scaled mullet.....	105
devilfish.....	28, 32	wolf.....	472	fine-scaled sucker.....	105
small.....	32	eel-back flounder.....	227	fishes, bony.....	39
diablo.....	507	eel blenny.....	471	soft-rayed.....	39
diamond flounder.....	225	eel cat.....	152	true.....	33
doctorfish.....	362	eel pout.....	212, 473	fishng-frog.....	504
dog, blue.....	18			fishng-frogs.....	504

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
fish of Dismal Swamp.....	194	gabian.....	31	golet.....	61
fish of the rice ditches of		gafftopsail.....	148, 276	goodies.....	348
South Atlantic States.....	194	gafftopsail catfishes.....	147	goody.....	349
fish shark.....	18	galafate.....	493	Cape May.....	349
five-bearded rockling.....	208	gall bengal.....	424	goosefish.....	504
five-bearded rocklings.....	208	gallwasp.....	163	gorbuscha.....	54
flamenco.....	326	gar.....	195	goujon.....	155
flannel-mouthed sucker.....	107	billy.....	36	gourami.....	314
flapmango.....	494	Cuban alligator.....	38	gourdseed buffalo.....	102
flasher.....	323	green.....	196	gourdseed sucker.....	102
flashers.....	323	long-nosed.....	36	grande écaille.....	39
flatfish.....	299, 302, 360	Mississippi alligator.....	38	Grand Oranchoe darter.....	287
common.....	227	Ohio.....	36	grass bass.....	305
flatfishes.....	217	salt-water.....	196	grass pike.....	173
flat-headed chub.....	136	silver.....	196	grass porgy.....	337
flesh-colored rock cod.....	369	silvery.....	196	grass rockfish.....	369
filer.....	304	garfish.....	196	grassy ground rockfish.....	320
floma.....	365	California.....	196	grayback.....	63
Florida lancelet.....	7	Ohio.....	36	gray bass.....	297
Florida sailfish.....	263	garibaldi.....	415	gray drum.....	352
flounder.....	269, 302	garibaldi.....	415	gray grunt.....	329
arctic.....	227	garlopa.....	314	grayling, arctic.....	66
California.....	226	gar pike, common.....	36	Michigan.....	66
diamond.....	225	gar pikes.....	36	Montana.....	66
eel-back.....	227	garrupa.....	365	graylings.....	65
four-spotted.....	224	garrupas.....	308	gray pike.....	282
gulf.....	224	gars, alligator.....	37	gray shark.....	15
lemon-yellow.....	227	short-nosed.....	37	common.....	15
peacock.....	217	gascon.....	270	gray sharks.....	13
pole.....	223	gaspereau.....	42	gray snapper.....	325
southern.....	224	gaspergou.....	352	gray sole.....	228
summer.....	224	gata.....	12	gray sucker.....	106
winter.....	227	gato.....	13	gray trout.....	355
flounders.....	217	genizara.....	426	greased chub.....	146
great.....	223	Georgian Bay herring.....	62	great amberfish.....	279
starry.....	226	ghostfish.....	471	great barraeuda.....	255
winter.....	227	ghost smelt.....	68	great Bear Lake herring.....	62
fluke.....	224	ghost smelts.....	68	great blue shark.....	17
Craig.....	228	giant black marlin.....	263	great flounders.....	226
flutemouth.....	245	Gila trout.....	114	great flyingfish.....	202
flyfish.....	367	Gilbert's darter.....	286	great fork-tailed cat.....	152
flyingfish.....	201, 407	gilded sunfish.....	301	great herring, Erie.....	63
double-bearded.....	201	gizzard shad.....	46	Great Lakes trout.....	59
great.....	202	gizzard shads.....	46	great northern pike.....	173
sharp-finned.....	200	glancefish.....	216	great pampano.....	277
single-bearded.....	201	glass-eye.....	282	great pike.....	172
flyingfishes.....	199, 200	glassfish.....	251	great pipefish.....	240
flying gurnard.....	409	globefish.....	503	great sculpin.....	387
flying gurnards.....	408	glovefish.....	499	great sea lamprey.....	8
flying perch.....	304	glut herring.....	42	great torpedo.....	27
flying robin.....	409	goatfish.....	343	great trout of Lake Tahoe.....	57
foolfish.....	494, 495	red.....	343	great tunnies.....	269
fork-beard.....	213	yellow.....	343	great white sharks.....	20
four-banded blenny.....	462	goatfishes.....	343	greenback trout.....	57
four-bearded rockling.....	209	gobies.....	438	green bass.....	297, 298
four-bearded rocklings.....	209	blind.....	447	green cod.....	209
four-eyed fish.....	194	crested.....	438	greenfish.....	280, 339
four-eyed fishes.....	194	half-naked.....	445	Alaska.....	374
four-spined stickleback.....	239	goby.....	446	green gar.....	196
four-spotted flounder.....	224	blind, of Point Loma.....	447	green jack.....	271
fox shark.....	18	clunging.....	446	green lamprey.....	8
Fraser River salmon.....	55	long-jawed.....	444	Greenland charr.....	60
French grunt.....	330	naked.....	446	Greenland codfish.....	210
frère jacques.....	235	sharp-tailed.....	441	Greenland halibut.....	221
fresh-water bass.....	303	variegated.....	446	greenling.....	374
black.....	298	gog.....	314	greenlings.....	373
fresh-water drum.....	352	goggle-eye.....	302, 305	green parrotfish.....	432, 435
fresh-water eel.....	77	goggle-eye jack.....	271	green perch.....	297, 298
fresh-water shark.....	16	goggler.....	271	green pike.....	173
fresh-water silverside.....	247	gold bass.....	298	green pipefish.....	242
fresh-water sunfish.....	299	golden angel.....	361	green-sided darter.....	286
fresh-water trout.....	297	golden trout of south fork		green sturgeon.....	34
triar.....	247	of Kern River.....	59	green sunfish.....	298
truars.....	246	golden trout of Sunapee		small.....	300
frigate mackerel.....	259	Lake.....	60	green trout.....	257, 298
frogfishes.....	504	golden trout of Volcano		grenadier.....	207
frostfish.....	210	Creek.....	59	grenadiers.....	202
frostfishes.....	262	golden trout of western		ground drummer.....	346
fry, anchovy.....	48	tributaries of Kern River.....	59	ground rockfish, grassy.....	320
hogmouth.....	48	goldeye.....	41	ground spearing.....	163
tarpon.....	51	goldeyes.....	41	grouper.....	311
fuernita.....	295	goldfish, sea.....	415	black.....	312, 314
Fulton cat.....	152	gold perch.....	301	Nassau.....	311
funal.....	370	goldring.....	304	poison.....	370, 371
fusiform darter.....	294	goldthread shiner.....	116	red.....	312

Last of common names of fishes occurring in check list—Continued

	Page		Page		Page
grouper, snowy.....	311	hairy boxfish.....	501	herrings, lake.....	41, 61
spotted.....	311	hairy porcupinefish.....	501	round.....	45
yellow.....	314	hake.....	213, 350	hickory jack.....	42
yellow-finned.....	311, 313	American.....	214	hickory shad.....	42, 46
groupers.....	308, 310	European.....	214	hind, brown.....	308
black.....	312	New England.....	214	red.....	312
growler.....	297	Old English.....	213	rock.....	308, 310
salmon-formed.....	297	Old England.....	214	speckled.....	312
grubber broadhead.....	47, 50	red.....	213	hitch.....	113
grubby.....	386	silver.....	214	boe.....	19
grunion.....	251	spotted.....	213	boe-mother.....	20
grunt, black.....	323, 329	squirrel.....	213	hog bass.....	298
boar.....	330	white.....	213	hogchoker.....	229
Cæsar.....	330	hakes.....	214	hogfish.....	283, 318, 335, 425
common.....	330	halfbeak, common.....	198	bass.....	298
French.....	330	slender.....	198	Spanish.....	425
gray.....	329	halfbeaks.....	198	hogmolly.....	107, 283
humpback.....	330	halfmoon.....	358	hogmouth fry.....	48
margaret.....	329	halfmoonfishes.....	358	hog sucker.....	107
open-mouthed.....	330	half-naked gobies.....	445	Holbrook's top minnow.....	185
redmouth.....	331	half-naked lycodes.....	475	holia.....	54
red-mouthed.....	335	halibut.....	221	hoopid salmon.....	54
striped.....	329, 332	arrow-toothed.....	221	hoosier.....	155
white.....	331	bastard.....	223	horned dace.....	117
yellow.....	330	common.....	221	horned dogfish.....	19
young.....	331	Greenland.....	221	horned-eared sunfish.....	299
grunts.....	329	Monterey.....	223	horned pout.....	153, 154
striped.....	331	halibuts.....	221	horned pouts.....	151
guabin, small.....	180	bastard.....	222	horned trunkfish.....	497
guacamara.....	435	hamlet.....	91, 311	horny cony.....	494
guaquanche pelón.....	255	hammerhead.....	18, 107	hornyhead.....	137
guajaco.....	185	hammerhead shark.....	18	horse crevallé.....	273
guacacones.....	184	hammerheaded sharks.....	17	horse-eyed jack.....	271
guajaca.....	191	handsawfish.....	171	horse-eye jack.....	272
guauvanche.....	255	handsawfishes.....	171	horsefish.....	244, 275
guapena.....	353	hannahill.....	317	horsehead.....	275
guardfish.....	156	harder.....	268	horse mackerel.....	270, 280
guarubaco.....	436	bardhead.....	199, 349	hound, smooth.....	13
guasa.....	310, 312	hardheads.....	199, 246	houndfish.....	196, 197
guasas.....	312	hardmouth.....	112	houndfishes.....	195
guatiwere.....	309	hard-mouthed suckers.....	105	humpback grunt.....	330
black.....	309	hardtall.....	272	humpback salmon.....	54
red.....	309	harelip sucker.....	112	humpback sucker.....	108
guatiwere amarilla.....	309	harlequin roach.....	302	humpback whitefish.....	65
guavina.....	354, 435, 436, 437	harvestfish.....	266, 267	huron.....	297
guavina hoyera.....	442	harvest pike.....	196	black.....	297
gudgeon.....	115, 132, 143	hayko.....	54	icefish.....	67
Niagara.....	139	headfish.....	503	inconnu.....	61
gudlar.....	216	headfishes.....	503	Indian chub.....	137
guitarfish.....	24	hechudo.....	50	Indianfish.....	302, 360
guitarfishes.....	23	hedghog ray.....	25	Indian remora.....	448
guitarro.....	23, 24	hemdurgan.....	363	Irish lord.....	331
gulf flounder.....	224	herring, arctic lake.....	62	Irish pompano.....	342
gulf menhaden.....	44	Athabaska.....	62	isabelita.....	361
gulf sea bass.....	318	big-eyed.....	42	isabelita de la alto.....	360
gulpers.....	95	Bessell's.....	63	iverkssoak.....	212
gunnel.....	467	blue.....	42	jabon.....	316
Gunther's darter.....	283	branch.....	42	jaboncillo.....	316
guppy.....	191	bristle.....	246	jack.....	173, 273, 364
gurnard, big-headed.....	408	California.....	41	chiselmouth.....	66
common.....	407	common.....	63	goggle-eye.....	271
flying.....	409	common lake.....	63	green.....	271
northern striped.....	407	Erne.....	63	hickory.....	42
red-winged.....	407	Erne great.....	63	horse-eye.....	272
southern striped.....	407	fall.....	42	horse-eyed.....	271
spinous.....	407	Georgian Bay.....	62	yellow.....	272
gurnards.....	407	glut.....	42	jackfish.....	279
flying.....	408	great Bear Lake.....	62	jack salmon.....	282
hackleback.....	35	jumbo.....	63	jacome.....	319
hacklehead.....	386	king of the.....	216	jallao.....	329
haddo.....	54	lake.....	63	janissary.....	426
haddock.....	211	Lake Huron.....	62	jaqueta.....	415
Jerusalem.....	216	least lake.....	62	jawfishes.....	452
Norway.....	363	Mackenzie River.....	62	jeniguan.....	330
hagfish.....	8	Michigan.....	62	jeniguan.....	331
round.....	8	rainbow.....	67	Jenny Lind.....	279
California.....	8	Rawson Lake.....	63	jerker.....	137
hagfishes.....	8	round.....	46	Jerusalem haddock.....	216
hair-finned dory.....	275	Saginaw Bay.....	62	Jessie's darter.....	291
hair-finned silverfish.....	275	Seneca Lake.....	62	jewish, black.....	312
hairtail.....	262	southern toothed.....	41	California.....	308
silvery.....	262	summer.....	42	spotted.....	312
hairtails.....	262	tailor.....	42	jewishfishes.....	308
hairy blowfish.....	499	thread.....	44	jewishsharp drummer.....	351
		toothed.....	40		
		wall-eyed.....	42		

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
iguagua	273	lake herrings	61	lisa blanca	253
iod	325	lake Huon herring	62	lisa cabezuda	253
John A. Grundle	38	lake lamprey	9	lisa francesca	39
John demon	304	lake lawyer	212	lisita	254
John dories	232	lake Michigan cisco	63	little bass	298
John Mariggle	39	lake minnow	123, 137	little bream	304
John paw	312	lake perch	281	little bullhead shark	22
Johnny	390	lake sculpin	389	littlehead porgy	336
Johnny darter	287	lake sheepshead	352	little killifish	175
Johnny verde	317	lake sturgeon	35	littlemouth porgy	337
loft-head porgy	336	Lake Superior whitefish	64	little pickerel	173
lopatan	335	Lake Tahoe trout	57	little red-eye	298
Jordan's darter	290	lama	15	little roncador	349
Jordan's trout	57	lamper eel	473	little skate	25
Jorobado	275	lamperina	8	little smelt	252
joturo	254	lamprey, arctic	10	little stargazer	385
joturo	497	brook	9, 10	little sunfish	304
joturo	302	great sea	8	little tunny	258
Julia's darter	292	green	8	liza	253
jumbo herring	63	lake	9	liza ojo de perdiz	254
jumper	297, 298	Mexican	9	lizardfish	163
jumping mullet	111	mud	9	lizardfishes	163
jumprocks	111	river	8	lodde	66
June sucker	108	shad	8	logfish	268
jurel	272, 278	silver	9	log perch	283
		small black	9	log perches	282
kale	363	three-toothed	10	long-eared sunfish	299, 300
kalog	387	western	10	long-finned albacore, West	
Kamchatka salmon trout	56	lamprey eel	8	Indies	260
Kamloops trout	58	lampreys	8	long-finned albacores	260
kawakawa	259	brook	8	long-finned sole	228
kawah chub	113	river	9	long-finned sunfish	304
kelp bass	317	lancelet, Bahama	7	long-headed trout of Cres-	
kelpfish	427, 458	Bermuda	7	cent Lake	57
speckled	458	California	7	longjaw	196
K. L. G. G.	65	Florida	7	Centro	63
K. L. G. G.	298	VIRGINIA	7	longjaw	64
K. R. G. G.	58	West Indian	7	longjaw	444
kete	500	lancelets	7	longjaws	195
killie	176	lancefish	171, 362	long mingo	495
killifish	176, 181	lancefish	56	long-mouthed minnow	121
common	175	lancefish	327	long-necked eels	77
little	175	langbarn	470	long-nosed dace	139
killifishes	174	lant	451	long-nosed gar	36
killy	176	lanternfishes	166, 169	long-nosed sucker	106
bass	176	laron	371	long-scaled sunfish	300
kingfish	257, 349, 350	laron	41	long-spined sunfish	386
kingfishes	350	laron	304	long-spined sunfish	212
king mullet	343	laron	297	long-spined sunfish	31
king of the herring	216	laron	297	long-spined sunfish	18
king of the mackerels	501	laron	110	longue	59
king of the mullets	295	laron	344	lookdown	275
king of the salmon	216	laron	62	loro	434, 435
king salmon	55	lavarets	64	Lost River sucker	108
kingston	22	lawyer	38, 325	loul	496
kisutch	54	lake	212	Lower California trout	59
kit	28	leaping tuna	259	lucky proach	386
kiver	299, 302	least darter	294	lumpfish	397
kiyi	64	least lake herring	62	spinous	398
Kodiak smelt	67	leathercoat	277	lumpsucker	397, 399
krashnaya ryba	55	leather-ear	299	lycodes, arctic	476
kuroka	263	leatherfish	494	cloudy	475
kurokajiki	263	leatherjacket	277, 491	half-naked	475
kweewee	108	leather	277	stott	474
kyack	42	leather	277	stott	474
		leatherwing	299	macabi	39
Labrador whitefish	64	lebis	28	macho	253
Lac de Mabre trout	61	lebranco	253	machuelo	44
lacefin	129	le kai salmon	5	machuelo	253
laccro	163	lemonfin	129	machuelo	62
ladfish	39, 425	lemon sole	228	mackerel, Atka	374
Spanish	425	lemon-yellow butterfish	309	banded	279
ladfishes	39	lemon-yellow flounder	227	bay	257
lady ray	30	leopard shark	14	black spotted Spanish	257
lafayette	267, 349	lcorne de Mer	397	Boston	257
lagarto	163	liza	494, 495	chub	258
lake bass	297, 303, 305	liza barbuda	496	common	257
white	307	liza colorada	494	Easter	558
lake carp	103	liza trompa	495	frigate	259
lake chub	136	lined sea robin	407	horse	270, 280
Lake Erie bass	305	lung	212, 213, 374, 473	Monterey Spanish	257
Lake Erie whitefish	65	thumble-eyed	213	skip	280
Lake herring	63	lungs	212	snake	260
arctic	62	honfish	371	snapp	280
common	63	lisa	253	snapping	280
least	62				

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
mackerel, Spanish	257	menhaden	44	Mississippi cat	152
of England	258	gulf	44	Missouri cavefish	195
spotted	257	short-tailed	44	Missouri sucker	102
thumb-eyed	258	menhagens	44	missuckee-kequock	307
tinker	258	Menominee whitefish	65	mojarra	343, 416
yellow	272, 273	mero	312	Mexican	343
mackerel midge	208	merco de lo alto	312	mojana almejero	329
mackerel porbeagle	19	merout	310, 364	mojarra blanca	341
mackerel scad	270	metalote	103	mojarra cantilena	341
mackerel scads	270	Mexican lamprey	9	mojarra cardinal	236
mackerel shark	19	Mexican mojarra	343	mojarra china	342
mackerel sharks	19	Mexican saifish	263	mojarra de Casta	341
mackerels, Atka	374	Mexican snapper	326	mojarra de las aletas anaril-	
king of the	504	Mexican sole	230	las	342
snake	261	mo	18	mojarra de las piedras	361
Mackinaw trout	59	Michigan herring	60	mojarra de ley	341
mademoiselle	345	midshipman	487	mojarra dorada	273
mademoiselles	345	midshipmen	487	mojarra garabata	337
madregal	279, 280	milkfish	40	mojarra negra	416
mad tom	156	milkfishes	40	mojarra plateada	416
mad toms	155	millers thumb	385	mojarra prieta	330
mahogany snapper	327	northern	386	mojarra rajado	415
maid	29	southern	385	mojarra verde	419
mailed catfishes	160	millers thumb	383	mojarra	340
maire d'Ampora	166	milhous	191	mojarra blanca	341
makajiki	263	millpond chub	297	mojarra de rio	416
maki-maki	500	millpond fluer	305	mojarritas	340
Malma trout	61	millpond perch	304	mojarron	332
malolo	109	mingo	494	mola	503
Mammoth Cave blindfish	195	minnie	133, 135	monda	278
mammy	146	minnow	116, 117, 133, 175	mongolar dummie	355
man-eaters	19	Athabaska	115	mongrel buffalo	102
man-eater sharks	20	bigeye	134	monkeyfish	22
mangrove minnow	175	black	139	monkfish	22, 504
mangrove snapper	313, 325	blackhead	145	Montana grayling	66
Manitoba whitefish	64	blue-lined	132	Monterey halibut	223
Manitou darter	283	blunt-nosed	145	Monterey roach	122
Manitoulin tullibee	64	bridled	135	Monterey Spanish mack-	
manjua	48	bronze	121	erel	257
manjuri	38	brook	140	moo-gad ee	107
manjuda	248	bullhead	145	mooneye	40, 64
man-of-war fish, Portuguese	268	chub	136	southern	41
man-of-war fishes	268	eastern mud	174	mooneye cisco	64
manta	32	emerald	123	mooneyes	40
manta raya	29	fathead	145	moonfish	193, 216, 275, 358
manta raya colorado	29	Holbrook's top	185	moonfishes	275
many-spined sunfish	304	lake	123, 137	moorish idol	362
mapiro	436, 440	leather-sided	119	moorish idols	362
mapo	440	long-mouthed	121	Moose Lake minnow	136
mapos	440	mangrove	175	moray, black	91, 92
marbled rockfish	314	Moose Lake	136	common spotted	91
marblefish	503	mud	174	spotted	93
margaret grunt	329	northern straw-colored	134	moray eel	93
margatefish	329	pearl	116	morays	90, 91
maria molle	413	pike	175	moremouth bream	302
marian	286	pin	109	morena pinta	90
maria prieta	332	plumbeus	136	morena pintita	92
mariposa	216	pursy	181	morena verde	92
marlin, barred	263	river	176	mosquitofish	185
black	163	rosy	123	moss bass	297
giant black	163	rosy-faced	124	mossbunker	44
striped	163	rotgut	146	mother-of-eels	473
marlinpikefish	263	sheepshead	181	mother-of-eels	305
marlinpikefishes	263	silver-mouthed	142	mother-of-eels	254
marsh bass	297	silvery	132, 143	mother-of-eels	254
Marston trout	61	smelt	143	mother-of-eels	104
Martin pascador	506	spot-tailed	132	mountain suckers	104, 105
Maryland darter	284	spotted	145	mountain trout	298
masamachus	59	spotted-tail	129	mousefish	504
maskinongy	172	star-headed	178	mud bass	297, 308
matajuelo	235	steel-colored	129	mud blower	212
matajuelo blanco	356	straw-colored	133, 134	mud cat	154, 155
matajuelo real	39	striped mud	174	mud chub	117, 302
matajuelos	356	sucker-mouthed	142	mud creeper	446
mattowacea	42	top	177, 178, 185	mud dabbler	173
Maxinkuckee darter	284	variegated	181	mud dabs	220
Mayfish	176	minnows	112	mudfish	38, 192, 434
May sucker	112	mud	174	mud lamprey	9
mazaara	263	silvery	123	mud minnow	174, 175
McCloud River rainbow		top	184	eastern	174
trout	58	minny bass	298	striped	174
Meadie's darter	288	misarkornak	474	mud minnows	174
medialuna	358	mishcuppaug	336	mud parrot	438
medium red salmon	54	Mississippi alligator gar	38	mud perch	215, 303
				mud sucker	107

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
mud sunfish.....	303	obispo.....	30	parrotfish.....	431, 433, 434
muffejaw.....	385	oceanic bonito.....	258	blue.....	432
muffejaws.....	385	ocean pipefish.....	243	green.....	432, 435
mukikichi.....	56	ocean sunfish.....	503	red.....	432
mullet.....	109, 110, 111	ocean tang.....	363	paru.....	360
black.....	105	ocean turbot.....	492	pascadillos del rey.....	252
blue.....	110	ocellated dory.....	232	pastor.....	268
blackback.....	253	ogac.....	210	patao.....	342
Califaver.....	253	Ohio gar.....	36	peacockfishes.....	280
common.....	253	Ohio garfish.....	36	peacock flounder.....	217
fantail.....	253, 254	Ohio shad.....	42	pealip sucker.....	112
fine-scaled.....	105	Ohio sturgeon.....	35	pearlfish.....	486
jumping.....	111	ohua.....	495	pearl minnow.....	116
king.....	343	oilfish.....	20, 261	pearlsides.....	73
mountain.....	254	oil shark.....	14	peche-peche.....	99
of Utah Lake.....	105	ojanco.....	327	peche pretre.....	365
red.....	111, 343	old England hake.....	214	peck-nosed skate.....	25
redeye.....	254	old English hake.....	213	pega.....	448, 449
sea.....	350	old wench.....	491	pegador.....	448
snipnose.....	268	oldwife.....	276, 349, 434, 491	peixe agulha.....	195
striped.....	253	black.....	493	peixe carago.....	12
trench.....	253	ohlepa.....	495	peixe fonda.....	331
whirligig.....	253	omilu.....	273	peixe rey.....	246, 252
white.....	109, 253	ono.....	256	pelerin.....	20
mullets.....	252	Ontario longjaw.....	63	Pensacola snapper.....	326
king of the.....	295	oolachan.....	66	pepesca.....	248
mountain.....	254	oouu.....	440	perch.....	297-302, 304, 409
striped.....	252	ooze eels.....	78	American.....	281
white.....	253	opah.....	216	backelor.....	304
mummichog.....	175, 176, 181	open-mouthed grunt.....	330	black.....	298, 317, 323
mufeca.....	360	Oquassa trout.....	61	blanco.....	298
murcielago.....	409	orange filefish.....	495	blue.....	301, 424
muscalonge.....	172	orange rockfish.....	365	branch.....	302
muskallunge.....	172	orange-spotted sunfish.....	300	bride.....	297
muskellunge.....	172	Oregon charr.....	61	bridge.....	304
Musquaw River whitefish.....	64	Oregon sturgeon.....	34	chain-sided.....	301
muttunfish.....	326, 342, 473	orelle noire.....	326	chinguapin.....	299, 304, 305
mykiss.....	56	orqueta.....	276	female.....	302
naccaysh.....	41	Oswego bass.....	297	flying.....	304
naulrod.....	42	ouachita darter.....	254	gold.....	301
naked goby.....	446	ouananiche.....	66	green.....	297, 298
namaycush.....	59	outilibi.....	309	lake.....	281
namaycush salmon.....	59	outilibi espagnol.....	308	log.....	283
Nassau grouper.....	311	ovate pampano.....	276	millpond.....	304
Navarro roach.....	121	oysterfish.....	424, 446, 487	mud.....	215, 303
needlefish.....	195, 196			pike.....	282
needlefishes.....	195	Pacific bonito.....	258	pirate.....	215
nee-kaio-kalung.....	392	Pacific cod.....	210	pond.....	302
negre.....	312	Pacific salmon.....	54	racoon.....	281
negrofish.....	309	Pacific volador.....	263	red.....	299, 300
nejorpallujak.....	214	paddle.....	397	red-bellied robin.....	299
Nelson's trout.....	59	paddlefish.....	36	redeye.....	303
Nelson's whitefish.....	65	paddlefishes.....	36	ringed.....	281
nerka.....	55	paintedtail.....	297	river.....	281
New England hake.....	214	dotted.....	297	robin.....	302
Newfoundland shark.....	14	pajarito.....	198	Sacramento.....	303
newlight.....	304	pale crappie.....	304	sand.....	302, 305
Niagara gudgeon.....	139	pale river bass.....	297	sea.....	307
Niangua darter.....	285	palometa.....	267, 276, 277	silver.....	304, 305, 307, 340, 345
nibbler.....	424	pampanito.....	276	speckled.....	304
nigger chub.....	146	pampano.....	274, 276, 277	spotted.....	305
nigger dick.....	146	California.....	266	strawberry.....	304
niggerfish.....	146, 309	common.....	277	Suckley.....	304
niggerhead.....	107	great.....	277	sun.....	299, 300, 303, 305
nigrone.....	16	round.....	276, 277	tin.....	304
nine-eyes.....	8	ovate.....	276	trout.....	214, 215, 288
nine-lined shanny.....	462	pampanos.....	276	viviparous.....	409
nine-spined sticklebacks.....	238, 239	paneca.....	276	warmouth.....	302
nissue trout.....	58	papagalio.....	438	white.....	304, 305, 307, 352, 410
northern barracuda.....	255	papagallos.....	280	yawmouth.....	302
northern darter.....	291	papermouth.....	280	yellow.....	281, 289, 299
northern miller's thumb.....	386	parche.....	350, 360	perchmouth bream.....	302
northern pomotis.....	302	pargo.....	324, 321	perches.....	281
northern roach.....	122	pargo amarillo.....	328	American pike.....	282
northern straw-colored minnow.....	134	pargo colorado.....	328, 337	log.....	282
northern striped gurnard.....	407	pargo criollo.....	326	river.....	281
northern sucker.....	106	pargo de lo alto.....	326	viviparous.....	409
northern whiting.....	350	pargo de raizero.....	327	permit.....	276, 277
North River shad.....	42	pargo guachenarigo.....	326	perrio.....	435
Norway haddock.....	363	pargo moreno.....	325	perro colorado.....	425
no-shee trout.....	58	pargo negro.....	325	perro perro.....	425
numbfish.....	27	pargo prieto.....	325	pesca blanca.....	137
nurse shark.....	12, 22	pargos.....	336	pesca pavone.....	18
nurse sharks.....	12	parrot, mud.....	433	pesca vermiglia.....	367
		roseback.....	432	pescadillo del rey.....	252, 354
				pescadito.....	120

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
pescado azul de dos colores	414	pipefish, ocean	243	puffer	498, 499
pescado blanco de chapala	251	West Indian	241	smooth	497
pescado blanco de chalo	250	pipefishes	240	southern	498
pescado colorado	348	pipper	199	puffers	497
pescado del rey	252	pipon	98	sharp-nosed	500
pescados blancos	250	piquer	362	Puget Sound pollack	209
pescados del rey	246	pirate	215	pug-nosed shiner	275
pesce rey	252	pirate perch	215	pumpkinseed	302
pesce tondo	19	pirate perches	215	pumpkinseeds	297
pesce toro	497	plance	226	purague	23
petite gueule	341	plain leatherjacket	277	pursy minnow	181
petite jaquette	414	plantalet	261	Putnam's darter	285
petite soie	333	plateado	98	Pyramid Lake trout	57
petitnègre	321	platefish	496	quasky	61
peto	256	pluma	337	queenfish	353
petos	256	plumbeus minnow	136	queenfishes	353
pez ciego	479	Plummer's pigfoot	370	querman	263
pez de espada	23	pogge	395	quiaquia	270
pez de gallo	280	pogy	44	quiebra	277
pez del rey	247, 252	po-he-wa	119	quillback	103
pez de pluma	336	poison grouper	370, 371	quillback buffalo	102
pez luna	303	poison toadfishes	488	quinnat salmon	55
pez mola	303	poisson bleu	65, 152	quusutch	54
pez puerco	492	poisson de marais	38	quiver	299, 302
pez puerco de piedra	491	poisson lune	216	quwwee	108
pez sierra	23	pole crappie	304		
pez zorro	18	pole flounder	228		
philosophie	363	pollack	209	rabbitfish	261, 497, 502
phoebe	319	Alaska	209	rabbitmouth sucker	112
picoconu	111	Puget Sound	209	rabrubia	327
picked dogfish	19, 21	wall-eyed	209	rabrubia de lo alto	320
pickeral	282	pollacks	209	rabrubias	327
barred	173	pomfret	265	racon perch	281
eastern	173	pomotis, northern	302	radiated shanny	470
little	173	red-tailed	299	ragfish	269
pickering	252	pompano, Carolina	277	ragfishes	266
picuda	255	Irish	342	rainbow darter	291
picudilla	255	pompon	332	rainbowfish	191
pigeon stingray	28	ponce prieto	329	rainbow herring	67
pigfish	335, 407	pondfish, black-eared	299, 301	rainbow trout	58
rock	407	pond perch	302	McCloud River	58
sand	407	yellow	297	of fish-culturists	58
pigfishes	335	pond smelt	68	rainfish	180
pigfoot, Plummer's	370	popeye	205	rainwaterfish	175
pigmy sculpin	386	poppyfish	296	rascera	365
pigmy sunfish	305	porbeagle	19	rasher	365
Everglades	305	mackerel	19	ratfish	33
pike, American	173	porbeagles	19	raton	256
blue	282	porcupinefish	501	rat-tail, common	205
common gar	36	hairy	501	rat-tails	202
grass	173	porcupinefishes	501	raven, sea	393
gray	282	porkfish	332	Rawson Lake herring	63
great	172	porgies	336	ray, bishop	30
great northern	173	porgy	336, 349	brier	25
green	173	grass	337	butterfly	29
harvest	196	jolt-head	336	clear	25
Sacramento	114	littlehead	337	cow-nosed	31
sand	282	littlemouth	337	hedgehog	25
sea	196	red	337	lady	30
wall-eyed	282	sauceyeye	336	possum	25
yellow	282	shad	337	shark	22
pike minnow	175	sheepshead	337	sharn-nosed	31
pike perch	282	southern	336	small electric	27
pike perches, American	282	speckled	337	rays	10, 23
pike shiner	121	three-tailed	358	eagle	30, 31
pikes	172	whitebone	337	electric	27
gar	36	porte enseigne	362	thick-tailed	23
pilot, black	413	Porto Rican anchovy	51	whip-tailed	28
cockeye	413	portugais	360	razorback	305
cow	415	Portuguese butterfly	359	razor-backed buffalo	102
shark	279	Portuguese man-of-war fish	268	razorback sucker	108
pilotfish	65, 278, 279	possum ray	25	razorback suckers	108
pilotfishes	339	postcroaker	349	razorfish	430
pincoo	43	Potomac shad	42	razorfishes	430
pinfish	337	pout, horned	153, 154	real swordfish	263
pinkfish	447	pouts, horned	151	red bass	348
pink salmon	54	prickfish	166	red-bellied bream	299, 300
pin minnow	109	prickly bullhead	383	red-bellied dace	113
pinous dory	276	prickly skate	25	red-bellied robin perch	299
pintado	257	pristfish	365	redbelly	299, 300, 302
pintano	415	puddingwife	426	redbreast	299
pintanos	414	pudiano	425	redbreast bream	299
pipefish	241	pudiano verde	426	red drum	348
common	242	puerco espino	501	red drums	348
great	240			redeye	298, 302, 303
green	242			little	298

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
red-eyed bream.....	303	robalito de las aletas ama-		rose trout.....	59
red-eyed sunfish.....	300, 303	rillas.....	306	Ross's arctic salmon.....	60
red-eye mullet.....	254	robalito de las aletas prietas.....	306	rosy dace.....	121
red-eye perch.....	303	robalo.....	305, 306	rosy-faced minnow.....	124
red fallfish.....	127	robalo prieto.....	306	rosy minnow.....	123
redfin.....	128, 131	robalos.....	305	rotgut minnow.....	146
compressed.....	126	robin.....	299, 302	roughback.....	227
redfish.....	55, 318, 348, 363, 370	Carolina.....	407	rough silverside.....	248
bull.....	348	flying.....	409	round bass.....	304
California.....	426	robin perch.....	299, 302	round basses.....	304
red goatfish.....	343	1ed-bellied.....	299	round buffalo.....	102
red grouper.....	312	rock.....	307	round herring.....	46
red guativera.....	309	rock bass.....	297, 298, 317	round herrings.....	45
red hake.....	213	common.....	303	round pampano.....	276, 277
red-headed bream.....	299	southern.....	303	round robin.....	270
red hind.....	311	rock basses.....	303	round stingray.....	29
redhorse.....	109	rock beauty.....	361	round sunfish.....	304
big-toothed.....	111	rock cod, black and yellow.....	369	roundtail.....	114
common.....	110	black-banded.....	370	round whitefish.....	65
eastern.....	111	brown.....	369	rousettes.....	11
short-headed.....	111	flesh-colored.....	369	rovalate.....	248
Texas.....	110	yellow-backed.....	369	rovetto.....	261
redmouth buffaloish.....	102	yellow-spotted.....	369	royal silver trout.....	56
redmouth grunt.....	331	rock cook.....	424	rubio volador.....	407
red-mouthed grunt.....	335	rock cooks.....	424	rudderfish.....	266, 268, 270, 340
red mullet.....	111, 343	rock cusk.....	385	banded.....	279
red parrotfish.....	432	rockfish.....	176, 283, 305, 307, 311, 313, 315, 456, 461, 467	black.....	268
red perch.....	290, 300	black.....	365	rudderfishes.....	267
red porgy.....	337	brown.....	368	black.....	267
red rockfish.....	368	grass.....	369	ruff.....	302
red rock trout.....	374	grassy ground.....	320	runner.....	272, 277, 279
red roncador.....	348	marbled.....	314	blue.....	272
red roncadors.....	348	orange.....	365	Russian cat.....	155
red salmon, medium.....	54	red.....	368	ruvotto.....	261
red sculpin.....	381	spotted.....	367	sabalito.....	99
red senfish.....	363	yellowtail.....	364	sabalo.....	39, 40
red-sided bream.....	118	rockfishes.....	303, 364	sabina.....	28
red-sided shiner.....	120	rock hind.....	308, 309, 310	sable.....	262
red snapper.....	326	lockling.....	203	sablefish.....	373
red-spotted sunfish.....	300	five-bearded.....	208	sablefishes.....	372, 303
red-spotted trout.....	61	four-bearded.....	209	sac-a-lait.....	177, 302, 304
red sturgeon.....	35	silvery.....	208	Sacramento cat.....	154
red sucker.....	106	three-bearded.....	208	Sacramento chub.....	120
red sunfish.....	299	rocklings.....	208	Sacramento perch.....	303
red-tailed balao.....	199	five-bearded.....	208	Sacramento pike.....	114
red-tailed bream.....	299	four-bearded.....	209	Sacramento River salmon.....	55
red-tailed pomotis.....	299	three-bearded.....	208	Sacramento squawfish.....	114
red-tail snappers.....	327	rock pigfish.....	407	Sacramento sturgeon.....	34
red-tail snappers.....	407	rock salmon.....	280	Sacramento sucker.....	106
red-tail snappers.....	407	rock sea bass.....	318	Sacramento sucker.....	106
raina.....	368	rock shellfish.....	496	Sacramento sucker.....	106
remora.....	449	rock sunfish.....	304	saibling, American.....	62
Indian.....	448	rock trout.....	373	salmeur.....	362
white-tailed.....	448	red.....	374	salmon.....	20, 103
remoras.....	448	Rocky Mountain bullhead.....	384	Florida.....	263
requiem.....	15	Rocky Mountain whitefish.....	65	Mexican.....	263
requiem.....	15	romero.....	278	salmon.....	263
rescacio.....	370	roncador.....	349	salmon sucker.....	103
rex sole.....	228	little.....	348	sallorfish.....	103
ribbonfish.....	262, 353	red.....	348	sallors' choice.....	329, 335, 337
ringed perch.....	281	yellow-finned.....	350	salma.....	338, 339
Rio Grande trout.....	57	roncador raado.....	330, 351	salmichi.....	132
river bass.....	297, 298	roncadores.....	344	salmon.....	54, 55
pale.....	297	red.....	348	blueback.....	55
spotted.....	298	ronco.....	346	calico.....	54
streaked-checked.....	298	ronco amarillo.....	330	chinook.....	55
river carp, blunt-nosed.....	103	ronco arará.....	330	coho.....	54
common.....	103	ronco blanco.....	330	Columbia River.....	55
river chub.....	137	ronco carbonero.....	330	common Atlantic.....	56
river drums.....	352	ronco condenado.....	330	dog.....	54
river lamprey.....	8	roncodina.....	349	Fraser River.....	55
river lampreys.....	9	ronco grande.....	330	hoopid.....	54
river minnow.....	176	ronco prieto.....	330	humpback.....	54
river perch.....	281	ronco ronco.....	330	jack.....	282
river perches.....	281	roncos.....	330	king.....	55
river sucker.....	111	rondanin.....	285	king of the.....	216
river sunfish.....	299	ronquil.....	452	landlocked.....	56
river trout.....	282	ronquils.....	451	le kai.....	54
roach.....	113, 299, 302, 305	Roosevelt trout.....	59	medium red.....	54
harlequin.....	302	roosterfish.....	280	Namaycush.....	59
Monterey.....	122	rooter.....	102	Pacific.....	54
Navarro.....	121	roseback parrot.....	432	pink.....	54
northern.....	122	rosefish.....	363	quinnat.....	55
venus.....	122	rosefishes.....	363	rock.....	280
robalito.....	306				

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
salmon, Ross's arctic.....	60	scamp.....	315	segundo.....	272
Sacramento River.....	55	schoolmaster.....	325	Seneca Lake herring.....	62
sau-qui.....	55	Schuykill cat.....	154	Seneca Lake smelt.....	62
sehago.....	56	scirenga.....	315	sennet.....	255
silver.....	54	scorfaudi.....	370	señorita.....	427, 428, 458
sockeye.....	55	scorpene.....	371	señoritas.....	428
spring.....	55	scorpion.....	371, 487	seran imperial.....	370
sukkegh.....	55	scorpionfishes.....	367	sergeantfish.....	281, 305
tyee.....	55	scourfish.....	261	sergeantfishes.....	281
white.....	279, 297	sculpin.....	371	serra.....	171
of the Colorado.....	114	arctic.....	386	serrana.....	353
salmonete.....	343	black.....	387	serrana hispania.....	353
salmonete amarilla.....	343	brassy.....	386	serrano.....	318
salmon-formed growler.....	207	buffalo.....	381	sesele.....	305
salmon shark.....	19	common.....	386	sesi de lo alto.....	326
salmon trout.....	58	daddy.....	387	shad.....	42, 108, 304, 341
Kamchatka.....	56	eighteen-spined.....	386	Alabama.....	43
of Lake Southerland.....	57	European.....	386	American.....	42
salt-water gar.....	196	great.....	387	broad.....	341
salt-water trout.....	355	lake.....	389	common.....	42
San Bernardino trout.....	59	long-spined.....	386	Connecticut River.....	42
sand conger.....	454	pigny.....	386	gizzard.....	46
sand dab.....	222	red.....	381	hickory.....	42, 46
sand darter.....	287, 288	yellow.....	381, 393	North River.....	64
sand darters.....	289	sculpins, scaled.....	375	of Lake Champlain.....	64
San Diego sole.....	231	scaleless.....	383	Ohio.....	42
sand-diver.....	164	daddy.....	386	Potomac.....	42
sand eel.....	451, 455	scup, common.....	386	young.....	115
sandfish.....	318, 357	scuppaug.....	386	shad lamprey.....	8
sand lance.....	451	sea ape.....	18	shad porgy.....	337
Alaska.....	451	sea bass.....	317, 348	shadwaiver.....	65
sand lances.....	451	black.....	317	shads, gizzard.....	46
sand lancee.....	451	gulf.....	318	shanny, Bosc's.....	462
sand lancees.....	451	of California, white.....	356	nine-lined.....	462
sand perch.....	302, 305	rock.....	318	radiated.....	470
sand pigfish.....	407	sea basses.....	308	shark, basking.....	20
sand pike.....	282	seabats.....	506	bay.....	15
sand roller.....	215	sea bream.....	265	bona.....	20
sand rollers.....	214	sea breams.....	265	bonnethead.....	17
sand shark.....	19	sea cat.....	149	bramble.....	22
sand sharks.....	19	sea catfish.....	148, 149	bullhead.....	10
sand stargazers.....	454	sea catfishes.....	147	California angel.....	22
sand sturgeon.....	35	sea devil.....	32	cat.....	14
sand sucker.....	109, 350	sea devils.....	31, 508	cazon de playa.....	16
sanducha.....	39	sea dog.....	424	cod.....	21
sand whiting.....	350	sea drums.....	352	codfish.....	21
San Pedro fish.....	216	sea eel.....	35	common gray.....	15
sapo.....	486, 487	sea fish, red.....	363	cow.....	11
sardina.....	48, 98, 249	sea fox.....	18	cub.....	15
sardina blanca.....	100	sea goldfish.....	415	dog.....	13
sardina bocana.....	50	seahorse.....	244	dusky.....	16
sardina de España.....	43	common American.....	244	East Indian basking.....	18
sardina de ley.....	43	seahorses.....	240	elephant.....	20
sardina escamuda.....	43	sea lamprey, great.....	8	fish.....	18
sardina machete.....	44	sea mink.....	350	fox.....	18
sardine, California.....	43	sea mullet.....	350	fresh-water.....	16
smooth.....	43	sea owl.....	397	gray.....	15
sardines, scaled.....	43	sea perch.....	307	great blue.....	17
smooth-headed.....	43	sea pike.....	196	hammerhead.....	18
true.....	43	sea poacher.....	395, 397	leopard.....	14
sargassumfish.....	504	sea poachers.....	393	little bullhead.....	22
sargo.....	337, 338	sea possum.....	25	longtail.....	18
sargo rañado.....	332, 338	sea raven.....	269, 393	mackerel.....	19
salmfin.....	129	sea ravens.....	393	Newfoundland.....	14
saucey porgy.....	336	sear.....	257	nurse.....	12, 22
sauger.....	282	sea robin.....	407, 408	oil.....	14
sau-qui salmon.....	55	brown-winged.....	407	salmon.....	19
sauris.....	270	Carolina.....	407	sand.....	19
sauris.....	197	lined.....	407	sharp-nosed.....	14
saury.....	197	red-winged.....	407	shovelhead.....	17
sauter.....	277	small-scaled.....	407	shovelnose.....	18, 19
savaniila.....	39	spotted.....	407	shovel-nosed.....	11
savola.....	262	striped.....	407	sleepers.....	22
sawbelly.....	42	web-fingered.....	407	small blue.....	16
sawfish, common.....	23	sea robins.....	407	soup-fin.....	14
sawfishes.....	23	sea serpent.....	88	spotted.....	14
scabbardfish.....	262	sea smelt.....	350	swell.....	12
scad.....	270	sea snail.....	399, 400	thornback.....	21
big-eyed.....	271	striped.....	400	tiger.....	14
dotted.....	270	sea snails.....	399	whale.....	18
mackerel.....	270	sea trout.....	355	wharf.....	16
scaled sardines.....	43	spotted.....	355	white.....	16
scaled sculpins.....	375	seaweed blenny.....	461	shark pilot.....	279
scaleless sculpins.....	383	sea whitefish.....	357	shark ray.....	22
		sebag salmon.....	56		

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
sharks	10	silverside minnow	119	snake blenny	471
angel	22	silversides	246, 247, 248	spotted	469
basking	20	silver trout	57	snake eels	84
bramble	22	royal	56	snakefish	163, 471
bullhead	10	silver whiting	351	snake mackerel	260
cat	11	silvery gar	196	snake mackerels	261
cow	11	silvery haitail	262	snap mackerel	280
gray	13	silvery minnow	132, 143	snapper	230, 363
great white	20	silvery minnows	123	Bermuda silk	326
hammer-headed	17	silvery rockling	208	blackfin	326
mackerel	19	singingfish	487	dog	325
man-eater	12	single-bearded flyingfish	201	gray	325
nurse	12	sirago	447	Lane	327
sand	19	siscowet	60	mahogany	327
sleepers	21	skate	25	mangrove	313, 325
tectospondylous	21	barndoor	25	Mexican	326
thresher	18	bobtailed	25	Pensacola	326
true	11	California big	26	red	326
whale	18	California common	26	redtail	327
sharksucker	448	clear-nosed	25	silk	326
sharp-finned flyingfish	200	common	25, 26	West Indian	326
sharp-nosed flyingfish	199	little	25	snappers	324
sharp-nosed puffers	500	peck-nosed	25	snapping mackerel	280
sharp-nosed ray	31	prickly	25	snipe eel	83
sharp-nosed shark	14	summer	25	snipe eels	83
sharp-tailed goby	441	winter	25	snipefish	246
shecutts	355	skates	10, 24	snipefishes	245
sheepshead	338, 426	skil	373	snipnose mullet	268
lake	352	skilfishes	373	snook	305
three-banded	358	skimback	103	snowy grouper	311
sheepshead minnow	181	skipjack	42, 197, 251, 259, 279, 280	snub-nosed eels	78
sheepshead porgy	337	skip mackerel	280	soapfish	163, 316
sheepsheads	338	skipper	197	black	316
shellcracker	302	skippick	196	soapfishes	316
shellfish	497	skittle-dog	19, 21	sobaco	492
rock	496	skowitz	54	sobacos	492
sheppy argentine	73	sleepers	435, 436, 440	sockeye salmon	55
shielded darter	283	striped	436	soft-rayed fishes	39
shiner	115, 123, 128, 129, 132	sleepers	435	soldado	235
blunt-nosed	275	sleepers	435	soldierfish	291
goldthread	116	sleepers	22	soldierfishes	234, 235
pike	121	sleepers	21	sole, American	229
pug-nosed	275	slender halfbeak	198	California	222
red-sided	120	shiner	487	gray	228
smelt	132	slippery dick	426, 485	lemon	228
spotted	138	slough bass	267	long-finned	228
swallow minnow	133	small black lamprey	9	Mexican	230
shining bass	304	small blindfish	195	rex	228
shoemaker	274, 279	small blue shark	16	San Diego	231
shore whiting	331	small catfish	154	somka	56
short barracuda	255	small devilfish	32	sorcerers	82
short-finned roach	122	small dolphin	265	soup-fin shark	14
short-headed redhorse	111	small electric ray	27	sourpaw	501
short-nosed gars	37	small green sunfish	300	South American catfishes	156
short-nosed sturgeon	34	small guabin	180	southern chub	277
short-tailed menhaden	44	smallmouth buffalo	102	southern flounder	224
shovelhead shark	17	small-mouthed black bass	298	southern miller's thumb	385
shovel-nosed shark	18, 19	small-scaled sea robin	407	southern mooneye	41
shovelnose shark	35	smear dab	228	southern porgy	336
shovelnose sturgeon	35	smear dabs	228	southern puffer	498
white	35	smelt	132, 143	southern rock bass	303
shovelnose sturgeons	35	American	67	southern stingray	28
Shumard's darter	285	California blue	252	southern striped gurnard	407
sier	257	Cobessiconic	67	southern toothed herring	41
sterring	257	ghost	68	spadefish	36, 358
sierra	257	Kodiak	67	Spaniard	267
sierita	196	little	252	Spanish flag	338, 393
silk snapper	328	pond	68	Spanish flags	303
Bermuda	326	sea	350	Spanish hogfish	425
silver bass	305, 307	Seneca Lake	62	Spanish ladyfish	425
silver carp	103	surf	67	Spanish mackerel	257
silver chub	117, 137	Wilton	67	black-spotted	257
silverfin	129, 132	smelt minnow	143	Monterey	257
silverfish	39, 214, 248, 262	smelt shiner	132	of England	258
hair-finned	275	smelts	66, 67	sparada	496
silver gar	196	ghost	68	spawneater	132
silver hake	214	surf	67	spearfish	103
silver jenny	341	smooth blackfish	424	black	263
silver lamprey	9	smooth cabezon	390	true, of Japan	263
silver-mouthed minnow	142	smooth dogfish	13	spearfishes	450
silver perch	304, 305, 307, 340, 345	smooth-headed sardines	43	pearfish sucker	163
silver salmon	54	smooth hound	13	spearling, ground	387
silverside	247	smooth puffer	497	speck	305
brook	251	smooth sardine	43	speckled bass	297, 298
fresh-water	247	smooth swellfish	497	speckled hen	297, 298
rough	248			speckled hnd	312

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
speckled perch.....	304	stickleback, four-spined.....	239	sucker, May.....	112
speckled porgy.....	337	nine-spined.....	239	Missouri.....	102
speckled trout.....	60	sticklebacks.....	237	mountain.....	104
of Lake Crescent.....	57	nine-spined.....	238	mud.....	107
spet.....	255	stingaree.....	28, 31	northern.....	106
spike.....	282	stingray.....	28	peahp.....	112
spinous dory.....	276	California.....	31	rabbitmouth.....	112
spinous gurnard.....	407	long-tailed.....	31	razorback.....	108
spinous lumpfish.....	398	round.....	29	red.....	106
spinyback blowfish.....	498	southern.....	28	river.....	111
spiny boxfish.....	502	spotted.....	30	Sacramento.....	106
spiny dogfish.....	21	whip.....	28	sailing.....	103
spiny eels.....	75	stingrays.....	28	sand.....	109, 350
split-mouth sucker.....	112	stittise.....	58	spearfish.....	450
splittail.....	116	stockfish.....	214	split-mouth.....	112
spoonbill cat.....	36	stone bass.....	308	spotted.....	107, 109
spot.....	349	stone cat.....	155	striped.....	109
white.....	339	stone cats.....	155	sturgeon.....	106
spottail.....	348	stonelugger.....	107	sweet.....	109
spot-tailed minnow.....	132	stoneroller.....	107, 146	swordfish.....	449
spotted balloonfish.....	502	stonerollers.....	146	Tahoe.....	107
spotted bass.....	297, 348	Storer's chub.....	137	Webbug.....	108
spotted bream.....	299	straw bass.....	297, 305	white.....	105, 110, 111
spotted cabrilla.....	317	strawberry bass.....	302, 305	white-nosed.....	109
spotted cero.....	257	strawberry perch.....	304	winter.....	109
spotted codling.....	213	stair-colored minnow.....	133, 134	suckerel.....	102
spotted cybium.....	257	steaked-cheeked river bass.....	298	suckermouth buffalo.....	102
spotted-finned sunfish.....	304	st eaked lead.....	298	sucker-mouthed minnow.....	142
spotted grouper.....	311	staped anchovy.....	48	suckers.....	102
spotted hake.....	213	staped bass.....	297, 307	carp.....	103
spotted jewfish.....	312	staped dace.....	140	hard-mouthed.....	105
spotted kelpfish.....	458	staped grunt.....	328, 332	mountain.....	104, 105
spotted lycodes.....	474	staped grunts.....	331	razorback.....	108
spotted mackerel.....	257	staped gurnard, northern.....	407	spotted.....	109
spotted minnow.....	145	southern.....	407	suckfish.....	488
spotted moray.....	93	staped marlin.....	263	suckingfish.....	448
common.....	91	staped mud minnow.....	174	Suckley perch.....	304
spotted perch.....	305	staped mullet.....	232	sukkegh salmon.....	55
spotted river bass.....	298	staped mullets.....	232	summer flounder.....	224
spotted rockfish.....	367	striped sea robin.....	407	summer herring.....	42
spotted sea robin.....	407	striped sea snail.....	400	summer skate.....	25
spotted sea trout.....	355	striped sleeper.....	436	Sunapee trout.....	60
spotted shark.....	14	striped sucker.....	109	sunfish.....	20,
spotted shiner.....	138	striped surffish.....	411	115, 274, 275, 298-304, 495,	503
spotted snake blenny.....	469	studsish.....	179	banded.....	304
spotted squeteague.....	355	sturgeon, common.....	84	big-eared.....	300
spotted stargazer.....	454	green.....	84	big-mouthed.....	302
spotted stingray.....	30	lake.....	35	big-nosed.....	299
spotted sucker.....	107, 109	Ohio.....	35	black.....	302
spotted suckers.....	109	Oregon.....	34	black-banded.....	303
spotted sunfish.....	304	red.....	35	blackeye.....	288
spotted-tail minnow.....	129	Sacramento.....	84	black-tailed.....	300
spotted trout.....	305	sand.....	85	bloody.....	300
of Lake Southernland.....	57	short-nosed.....	34	blue.....	298, 301
spotted trunkfish.....	497	shovelnose.....	35	blue-and-green.....	298
spotted weakfish.....	355	white.....	34, 35	blue-mouthed.....	301
sprat.....	44, 51	white shovelnose.....	35	blue-spotted.....	298, 304
springfish.....	385	sturgeons.....	35	brilliant.....	300
spring salmon.....	385	shovelnose.....	35	buffalo.....	298
squaremouth.....	112	sturgeon sucker.....	108	common.....	300, 302
squaretail.....	269	subotaka.....	171	deceptive.....	299
squaretails.....	269	sucker.....	8, 103, 110, 449	eared.....	300
squat.....	22	biglip.....	108	Everglades pigmy.....	305
squawfish.....	113, 114	black.....	108, 107	fresh-water.....	299
Colorado River.....	114	black winter.....	109	gilded.....	301
Sacramento.....	114	blue.....	146	green.....	298
squeteague.....	355	bluehead.....	104	horned-eared.....	299
spotted.....	355	bootleg.....	107	little.....	300
squirrelfish.....	235, 318, 330	brook.....	105	long-eared.....	299, 300
squirrelfishes.....	234, 235, 318	carp.....	103	long-finned.....	304
squirrel hake.....	213	chub.....	108, 109	long-scaled.....	300
squit.....	355	Columbia River.....	108	many-spined.....	304
Stanley's whitefish.....	65	common.....	105	mud.....	303
stargazer.....	454	creek.....	109	ocean.....	508
little.....	385	fine-scaled.....	105	orange-spotted.....	300
spotted.....	454	flannel-mouthed.....	107	pigmy.....	305
stargazers.....	454	gourdseed.....	102	red.....	299
sand.....	454	gray.....	108	red-eyed.....	300, 303
star-headed minnow.....	178	halelp.....	112	red-spotted.....	300
starry flounders.....	226	hog.....	107	river.....	299
steel-backed chub.....	146	humpback.....	108	rock.....	304
steel-colored minnow.....	129	June.....	108	round.....	304
steelhead.....	58	large-scaled.....	110	small green.....	300
Stewart White trout.....	59	long-nosed.....	106	spotted.....	304
stickleback, brook.....	238	Lost River.....	108	spotted-finned.....	304
common eastern.....	238				

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
sunfish, wide-mouthed	302	threadfins	256	trout, golden, of south fork	
sunfish bass	303	threadfish	274	of Kern River	59
sunfishes	297, 299	eight-threaded	256	of Sunapee Lake	60
sunny	302	thread herring	44	of Volcano Creek	59
sun bass	302	three-angled trunkfishes	497	of western tributaries	
sun perch	299, 300, 303, 305	three-banded sheepshead	358	of Kern River	59
sun trout	302	" " " " " "	208	gray	355
surfish, bigmouth	411	" " " " " "	208	Great Lakes	59
black	411	" " " " " "	358	great, of Lake Tahoe	57
striped	411	three-toothed lamprey	10	greenback	57
wall-eyed	410	thresher	18	green	297, 298
white	411	thresher sharks	18	Jordan's	57
surfishes	409	thunderpumper	352	Kamchatka salmon	56
surf smelt	67	thunderpumpers	352	Kamloops	58
surf smelts	67	tiburon	17	Kern River	58
surf whitings	351	tiburon ballenas	18	Lac de Mabre	61
surgeon, blue	362	ticky-ticky	178, 185	Lake Tahoe	57
common	362	tiger shark	14	long-headed, of Crescent	
surgeonfishes	362	tigrone	14	Lake	57
surmullet	343	tilefish	357	Lower California	59
Susan's darter	288	timber crappie	304	Mackinaw	59
swago bass	298	timucu	195	Malma	61
swallow, black	452	tinker mackerel	258	Marston	61
swallowminnow shiner	133	tinmouth	304, 305	McCloud River rainbow	58
swallowers	95	tiñosa	273	mountain	298
black	452	tin perch	204	Nelson's	59
sweet sucker	109	tintoreva	18	nissuee	58
swago	298	trante	262	no-shoe	58
swellbelly	408	tranties	262	Oquassa	61
swellfish	408, 499	tru	163	Pyramid Lake	57
smooth	497	toadfish	457, 498	rainbow	57
swellfishes	498	toadfishes	487	of fish-culturists	58
swellingfish	498	poison	488	red rock	374
swell shark	12	tobaccobox	25, 299, 302	red-spotted	61
" " " " " "	12	tobacco trumpetfish	245	Rio Grande	57
swelltoads	501	toerco	354	river	282
swingletail	18	logue	59	rock	373
switchtail	85	tomcod	210	Roosevelt	59
swordfish	196	California	210	rose	59
common	264	tomcods	210	royal silver	56
real	263	tomtate	331	salmon	58
swordfishes	264	tomtates	331	of Lake Southerland	57
swordfish sucker	449	tonguefish	231	salt-water	355
swordtail	193	tonguefishes	230	San Bernardino	59
symbranchoid eels	76	toothed herring	40	sea	355
		southern	41	spotted	355
		top minnow	177, 178, 185	silver	57
		Holbrook's	185	speckled	60
		top minnows	184	of Lake Crescent	57
		toro	273, 497	spotted	305
		torpedo	27	of Lake Southerland	57
		California	27	Stewart White	59
		great	27	Sunapee	60
		toipedofish	27	Truckee	57
		torsk	214	Verde	114
		toter	107	Volcano Creek	59
		totuava	356	Waha Lake	56
		treefish	369	white	297, 298
		trembler	27	yellowfin	57
		trench mullet	253	Yellowstone	56
		" " " " " "	491		
		" " " " " "	243	hout bass	298
		" " " " " "	215	trout perch	214, 215, 298
		" " " " " "	17	trucha	254
		alekey	212	Truckee trout	57
		aurora	60	true eels	77
		blueback	61	true fishes	33
		of Lake Crescent	57	true sardines	43
		brook	60	true sharks	11
		of western Oregon, etc.	58	true spearfish of Japan	263
		brown	298	trumpetfish	245, 246
		bull	61	tobacco	245
		coast-range	58	unarmed	245
		Colorado River	57	trumpetfishes	244
		Columbia River	56	trunkfish	496
		cut-throat	56	common	497
		Dolly Varden	61	horned	497
		Dublin Pond	60	spotted	497
		Eagle Lake	58	trunkfishes	496
		eastern brook	60	three-angled	497
		emerald	58	tschaviche	54
		Evermann's	59	tschawyttscha	55
		fresh-water	297	tswam	108
		Gila	114	tulpi	64
thread eels	83				
thread filefish	494				
threadfin	256				

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
tullibee.....	64	web-fingered sea robin.....	407	whiting.....	214
Manitoulin.....	64	Webug sucker.....	108	California.....	350
tuna, Allison's.....	260	weichman.....	235	Carolina.....	350
leaping.....	259	welshman.....	297	northern.....	350
tunnies.....	259	western lamprey.....	10	of Lake Winnepesaukee.....	64
great.....	259	West Indian lancelet.....	7	sand.....	350
runny, little.....	258	West Indian pipefish.....	241	shore.....	351
turbot.....	491	West Indian snapper.....	326	silver.....	351
ocean.....	492	West Indies long-finned al- bacore.....	260	surf.....	351
Tuscumbia darter.....	293	whale shark.....	18	wide-mouthed sunfish.....	302
tusk.....	214	whale sharks.....	18	widowfish.....	366
tyee salmon.....	65	wharf shark.....	16	wildow cat.....	152
tyrantfish.....	262	whiffs.....	218	Wilton smelt.....	67
ulke.....	386	Whipple's darter.....	292	windfish.....	115, 117
ulmina.....	98	whipsnake eel.....	88	windowpane.....	217
ulus.....	272	whip stingray.....	28	windowpanes.....	217
unarmed trumpetfish.....	245	whip-tailed rays.....	28	winninsh.....	56
unernak.....	477	whirligig mullet.....	253	winter flounder.....	227
unicornfish.....	495	whitebait.....	41, 247, 248	winter flounders.....	227
unspotted balloonfish.....	502	white bass.....	297, 303, 307	winter skate.....	25
		white-bill.....	43	winter sucker.....	109
vagmar.....	216	whitebone porgy.....	337	black.....	109
vaca.....	321	white cat.....	152, 153	wolf eel.....	472
vacas.....	320	white crappie.....	304	wolfish.....	171, 472
vacuocua.....	345	white croaker.....	353	wood bass.....	208
Vahl's lycodes.....	474	white-eye.....	282	worm eels.....	82
vaqueta de los colores.....	361	whitefish.....	44, 64, 114	wreckfish.....	308
variegated goby.....	446	Bear Lake.....	65	wreckfishes.....	308
variegated minnow.....	181	Bonneville.....	65	wrymouth.....	471
Venus roach.....	122	common.....	65		
Verde trout.....	114	Coulter's.....	65	xuel.....	272
verrugata.....	351	humpback.....	65	xuel de castilla.....	276
verrugata.....	340	Kennicott's.....	65		
verrugato.....	349	Labrador.....	64	yaito.....	259
vijaca.....	419	Lake Erie.....	65	yawmouth perch.....	302
victorfishes.....	258	Lake Superior.....	65	yellow angel.....	361
vicuda.....	255	Lauetta.....	62	yellow-backed rock cod.....	369
vieja.....	432, 434	Manitoba.....	64	yellow bass.....	297, 298, 307
vieja colorada.....	433	Menominee.....	65	yellowbelly.....	302
vieja muger.....	433	Musquaw River.....	65	yellow bream.....	302
viejias.....	431	Nelson's.....	65	yellow caranx.....	273
Virginia lancelet.....	7	Rocky Mountain.....	65	yellow cat.....	153, 155
viuva.....	366	round.....	65	yellow-finned shiner.....	260
viviparous perch.....	409	sea.....	357	yellow-finned grouper.....	311, 313
viviparous perches.....	409	Stanley's.....	65	yellow-finned roncadore.....	350
volador.....	202, 263, 263	Yellowstone.....	65	yellowfin trout.....	57
Pacific.....	263	whitefishes.....	61, 64	yellowfish.....	309, 374
volantin.....	272	white grunt.....	331	yellow goatfish.....	343
Volcano Creek trout.....	59	white hake.....	213	yellow grouper.....	314
voraz.....	328	white lake bass.....	307	yellow grunt.....	330
		whitemouth drummer.....	349	yellow jack.....	272
Wachna cod.....	210	white mullet.....	109, 253	yellow mackerel.....	27, 273
Waha Lake trout.....	56	white mullets.....	253	yellowmouth.....	356
walleye.....	282	white-nosed sucker.....	109	yellow Ned.....	281
wall-eyed herring.....	42	white perch.....	304, 305, 307, 352, 410	yellow perch.....	281, 298, 299, 302
wall-eyed pike.....	282	white salmon.....	279, 297	yellow pike.....	282
wall-eyed pollack.....	209	of the Colorado.....	114	yellow pondporch.....	297
wall-eyed surfhish.....	410	white sea bass of California.....	358	yellow sculpin.....	381, 393
warmouth.....	302	white shark.....	16	yellow-spotted rock cod.....	369
black.....	302	white sharks, great.....	20	Yellowstone trout.....	56
warmouth bream.....	302	white shovelnose sturgeon.....	35	Yellowstone whitefish.....	65
warmouth perch.....	302	white spot.....	339	yellowtail.....	276, 279, 327, 345, 349
waterbelly.....	64	white sturgeon.....	34, 35	yellow-tailed croaker.....	350
wawwhunnekesuog.....	257	white sucker.....	105, 110, 111	yellowtail rockfish.....	364
weakfish, bastard.....	355	white surfhish.....	411	young grunt.....	331
common.....	355	white-tailed remora.....	448	young shad.....	115
spotted.....	355	white trout.....	297, 298	zapatero.....	277
weakfishes.....	353				

LIST OF SCIENTIFIC NAMES

	Page		Page
abacurus, Eleotris	437	Acanthocottus patris	389
aboti, Syngnathus	240	profundorum	392
abotti, Anchovia	50	scorpioides	386
Anchoviella	50	scorpius	386
Hydrophlox	127	sellaris	388
Notropis chalybæus	127	Acanthocybidae	256
Osmerus	67	Acanthocybium	256
mordax	67	petus	256
abbreviata, Chimæra	33	solandri	256
abbreviatus, Bathyalopex	33	Acanthoderma	261
Naucleus	278	temminckii	261
Abeona	409	acanthoderma, Thyrsites	261
aurora	409	acanthognathus, Lophodolus	509
throwbridgi	409	Acantholebias	373
aberrans, Hypoplectrus	321	Acantholepis	68
Hypoplectrus unicolor	321	Acantholiparis	401
Liopropoma	316	opercularis	401
Perca	316	Acanthonotus	75
abildgaardi, Sparisoma	432	nasus	75
Sparus	432	quadricornis	497
abildgardi, Searus	431	Acanthonus	433
Ablennes	197	armatus	433
hians	197	spinifer	433
Aboma	443	Acanthophaelus	191
chiquita	443	melanzonus	191
etheostoma	443	acanthophorus, Serranus	317
lucretiae	443	acanthopoma, Draconetta	450
(Abrams) baleatus, Cyprinus	117	Acanthopterygi	232
smithi, Cyprinus	41	Acanthosoma	508
Abrams bosc	350	carinatum	508
versicolor	115	Acanthostracion	497
(Abrams) balteatus, Cyprinus	118	polygonus	497
Abudefduf	414	Acanthuridae	362
analogus	415	Acanthurus	362
declivifrons	415	alia	363
marginatus	415	bahianus	363
taurus	415	brevis	362
troschelli	415	broussoneti	362
abyssicola, Coregonus	66	cæruleus	362
Prosopium	66	crestonis	363
Raja	26	ctenodon	363
abyssorum, Nematonurus	204	glaucopeareus	363
acadian, Cottus	393	hepatus	362
acadianus, Glyptocephalus	228	hirudo	362
Acantharchus	303	phlebotomus	362
pomotis	303	striatus	363
Acanthemblemaria	465	strigosus	363
spinosa	465	subarmatus	362
Acanthenchelyidae	89	tractus	363
Acanthenchelys	89	triostegus	362
spinicauda	89	zebra	362
Acanthias	21	Acara adpersum	418
americanus	21	bartoni	419
suckleyi	21	cæruleopunctata	423
vulgaris	21	pinima	332
acanthias, Monoceratias	508	pitamba	327
Squalus	21	rectangulare	421
Acanthicus canensis	161	tetramerus	423
Acanthidium	21	acara-pinima, Pristipoma	333
calcea	21	Acanthorhinus	21
pustillum	21	accensum, Plectropoma	321
Acanthion	276	accensus, Hypoplectrus unicolor	321
acanthistius, Bodianus	309	Accipenser macrostomus	35
Enneistius	309	ohiensis	35
Acanthochinus	457	serotinus	35
chaperi	457	acclimdens, Cyclothone	72
Acanthocottus	386	acclivis, Larimus	344
æneus	386	Acedia	231
bubalis	386	nebulosa	231
groenlandicus	387	Acentrolophus	267
labradoricus	388	maculatus	267
laticeps	388	maculosus	267
mitchilli	386	Aceratidae	511
mucosus	387	acervum, Cybium	267
ocellatus	387	Acharina	229
octodecemspinosus	386	achigan, Bodianus	296

List of scientific names—Continued

	Page		Page
Achiridae.....	229	Acomus.....	105
Achirophichthys typus.....	89	(Acomus) generosus, Catostomus.....	104
Achirus.....	229	griseus, Catostomus.....	107
barnharti.....	230	guzmaniensis, Catostomus.....	104
comifer.....	230	lactarius, Catostomus.....	107
fasciatus.....	229, 230	acoupa, Cheilodipterus.....	354
fonsecensis.....	230	Cynoscion.....	354
inscriptus.....	229	Acrocheilus.....	112
klunzingeri.....	229	alutaceus.....	112
lineatus.....	229, 230	acrolepis, Macrurus.....	203
mazatlanus.....	230	Nematonurus.....	203
ornatus.....	231	acronotus, Carcharias.....	16
scutum.....	229	Squalus.....	16
achirus, Achirus.....	229	Acropoecilia.....	190
Pleuronectes.....	229	(Acropoecilia) tridens, poecilia.....	190
Achrophichthys.....	89	Acronurus.....	362
typus.....	89	caeruleatus.....	362
Acipenser.....	33	carneus.....	362
acutirostris.....	34	fuscus.....	362
aleutensis.....	34	nigriculus.....	363
(Antaeus) agassizii.....	34	Acrotidae.....	269
alexandri.....	34	Acrotus.....	269
buffalo.....	35	willoughbyi.....	269
cinnamomi.....	35	Acteis.....	460
oligopeltis.....	34	culebra.....	460
ayresi.....	34	moorei.....	460
bairdi.....	34	Actinochir.....	401
brachyrhynchus.....	34	major.....	401
brevirostris.....	34	Actinopteri.....	33
brevirostrum.....	34	aculeata, Chrysophrys.....	336
carbonarius.....	35	aculeatus, Chelmon.....	359
caryi.....	34	Chnus.....	470
dekayi.....	34	Etmopterus.....	21
fulvescens.....	33, 35	Gasterosteus.....	237
girardi.....	34	Halientichthys.....	507
glabra.....	34	Lophius.....	507
halops.....	34	Prognathodes.....	359
helops.....	34	aculeatus (oculatus), Serranus.....	328
holbrookii.....	34	aculeis, Gasterosteus.....	239
huso.....	33	acuminata, Halecula.....	46
(Huso) anasimos.....	35	Murana.....	85
anthracinus.....	35	acuminatus, Etrumeus.....	46
atelasps.....	35	Grammistes.....	353
copei.....	35	Myrichthys.....	85
kirtlandi.....	37	Pareques.....	353
lamarii.....	35	acuminatus umbrosus, Eques.....	353
megalasps.....	35	Acus.....	243
mertianus.....	35	seguoreus.....	243
paranasimos.....	35	heckeli.....	244
rafinesqui.....	35	martinicensis.....	243
richardsoni.....	35	maxima squamosa viridis.....	36
rosarium.....	35	pellucidus.....	244
kennicottii.....	34	acus, Atopichthys.....	54
laevis.....	35	Gymnotus.....	101, 486
lagenarius.....	36	Sphyræna.....	196
lecontei.....	34	Syngnathus.....	240
lesueurii.....	34	Tylosurus.....	197
hopeltis.....	35	acuta, Loricaria.....	161
macrorhinus.....	34	Myliobatis.....	31
macrostomus.....	35	Ophisoma.....	79
maculosus.....	35	Perca.....	281
medirostris.....	34	acute scabra, Lophius.....	505
microrhynchus.....	34	acuticeps, Oligocottus.....	391
milberti.....	34	Oxyottus.....	391
mitchilli.....	34	acutirostris, Acipenser.....	84
ohlensis.....	35	Corvina.....	346
oxyrhynchus.....	34	Ichthyopus.....	84
oxyrhynchus.....	34	Lutjanus.....	325
platyrhynchus.....	35	Pareuphagus.....	315
putnami.....	34	Pristis.....	23
rhynchæus.....	35	Serranus.....	315
rubicundus.....	35	acutirostris lata, Cerna.....	315
rupertianus.....	35	acutiventris, Alfaro.....	193
ruthenus.....	34	Acutomentum.....	395
schypa.....	34	alutum.....	396
stellatus.....	34	brevispine.....	396
storeri.....	34	clavilatum.....	396
sturio.....	33, 34	eigenmanni.....	396
transmontanus.....	34	entomelas.....	396
vultharius.....	34	hopkinsi.....	396
Acipenseridae.....	33	maedonaldii.....	396
acipenserinus, Agonus.....	394	ovale.....	396
Podotheus.....	394	protergerum.....	396
ackleyi, Raja.....	25	rufum.....	396
		wilsoni.....	396

List of scientific names—Continued

	Page		Page
acutum, Cichlasoma	422	affine, Myctophoma	169
Hemulon	329	Plectropoma	321
acutus, Eucocetus	199	Scopelus	169
Fodiator	199	affinis, Algansea	116
Searus	434	Argyropelecus	74
Synodus	164	Atherinops	232
Thorichthys	422	Atherinopsis	232
adamantinus, Litholepis	38	Auchenopterus	457
Adina	180	Carapus	486
multifasciata	180	Caulolatilus	357
adina, Fundulus	179	Centropomus	306
Adioryx	236	Chelodipterus	296
suborbitalis	236	Chilomycterus	502
adronodacus, Salmo	59	Chimarra	33
adobe, Agosia	141	Chinostomus	121
Apocope	141	Cremnobates	457
Adonis	460	Exocetus	201
pavoninus	460	Exonantes	201
adscensionis, Trachinus	310	Fierasfer	486
adscensionis, Epinephelus	310	Gila	114
adspersa, Parapetenia	418	Heros	422
Rhinoptera	31	Heterandria	185
adspersum, Acara	418	Hypoplectrus unicolor	321
adspersus, Labrus	424	Isopisthus	353
Pantogolabrus	424	Lucania	175
adusta, Corvina	347	Pimelodus	152
adustus, Coeusius	137	Pseudocheirodon	96
Eupomacentrus	414	Psychichthys	33
Gobiesox	489	Stomias	69
Julidio	428	Syngnathus	242
Ophioscion	347	Syrtetes	242
Pomacentrus	414	Thorichthys	422
Pseudojulis	428	Thynnus	258
Æglefinus	210	agassizi, Carelparis	400
linnaei	210, 211	Ipnops	171
æglefinus, Gadus	210, 211	agassizi, Acipenser (Antaceus)	34
Melanogrammus	211	Alepocephalus	51
Ælurichthys	147	Bathysaurus	164
isthmensis	148	Brama	265
longispinus	148	Calhonymus	450
panamensis	148	Carelparis	400
scutatus	148	Chlorophthalmus	165
ænea, Cichla	303	Chologaster	194
æneolus, Notropis	143	Cratinus	317
æneus, Acanthocottus	386	Cylindrosteus	37
Astyanax	96	Dieromita	480
Cottus	386	Forbesella	194
Pimelodus	155	Holconotus	411
Tetragonopterus	96	Hyperprosopon	410
ænigmaticus, Icosteus	269	Ipnops	171
æpyptera, Ammocetes	10	Leptoichthys	53
æquatoris, Talismania	53	Liparis	400
Æquidens	423	Pimephales	145
œruleopunctata	433	Salmo	60
æquidens, Cuius	437	Salvelinus	60
Otolithus	356	Scorpæna	371
Prionodes	319	Toxichthys	410
Serranus	319	Ageneiosus	159
æquoreus, Acus	243	brevifilis	159
Syngnathus	243	caucanus	159
æreus, Sebastodes	367	aggregatus, Cyrtogaster	409
æsculapius, Alepidosaurus	171	agulis, Gadus	209
Plagyodus	171	Agonidae	393
æsupus, Boleosoma	287	Agonostoma globiceps	255
æstivalis, Clupea	42	microps	254
Gobio	138	nasutum	254
Macrhybopsis	138	peroides	254
Pomolobus	42	Agonostomus	254
æstivalis marconis, Hybopsis	138	salvini	254
æstuaris, Paralichthys	224	telfairi	254
æthallion, Citharichthys	218	Agonus	395
æthallus, Carcharias	17	acipenserinus	394
Æthoprora	168	anne	394
effulgens	168	barkani	394
lucida	168	cataphractus	395
Ætobatidae	30	decagonus	394
Ætobatis	30	dodecaædron	394
Ætobatus	30, 31	dodecaædrus	394
californicus	31	vulsus	395
frémynvilli	31	Agosia	140
goodei	31	adobe	141
laticeps	30	chrysogaster	140
ater, Alphestes	312	falcata	141
Epinephelus	312	klamathensis	141
Gymnothorax	91	metallica	140

List of scientific names—Continued

	Page		Page
Agosia novemradiata	141	Albula	39
robusta	140	conorhynchus	39, 40
umatilla	141	erythrocheilus	40
yarrowi	141	forskali	39
Agriosphyræna	255	forsteri	40
agua-bonita, Salmo	59	goreensis	40
Salmo mykiss	59	neogumalca	40
aguaji, Trisotropis	314	parrae	40
aguiha, Peixe	195	plumieri	40
Ahlia	82	rostrata	40
egmontis	82	seminuda	40
Ahynnodontophis	94	vulpes	39
stigmatotus	94	albula, Mugil	253
aguila, Lachnolaimus	425	Albulidae	39
Ailurichthys	147	Albuloides	39
pinnimaculatus	148	albus, Bryttus	302
ailurus, Pimelodus	153	album, Hamulon	320
aimara, Macrodon	99	Moxostoma	110
aiy, Pallasia	394	Alburnellus	123
alabamæ, Alosa	43	altipinnis	127
Claricola	292	amœnus	123
Etheostoma whipplei	292	arge	124
Notropis	126	dilectus	123
alalunga, Bothrocaraopsis	477	jaculus	123
Scomber	260	jemezianus	125
Scombrus	260	matutinus	126
alalunga, Germa	260	micropterus	124
Scomber	260	percobromus	121
alascana, Asterotheca	396	simus	135
alascanus, Ammodytes	451	Alburnops	133
Argyrosomus	62	blennius	133
Leucichthys	62	heterodon	136
Sebastolobus	364	illecebrosus	134
Xenochirus	396	longirostris	135
alata, Fissala	408	nubilis	144
alatus, Arus	148	plumbeolus	128
Lampanvetus	167	saladanus	132
Nyctimaster	167	shumardi	134
Prionotus	408	taurocephalus	145
Alausa	42	Alburnus amabilis	121
Alausella	41	americanus	350
Alaron	190	dilectus	125
alba, Rogenia	41	formosus	129
albacauda, Echeueis	448	lepidulus	125
Albacora	250	lineolatus	135
albacores, Neothunnus	260	megalops	125
Scomber	260	nitidus	123
Albatrossia	204	oligaspis	125
pectoralis	204	rubellus	123
albatrossis, Osmerus	67	rubrifrons	124
Thaleichthys	67	socius	124
albatrossus, Macrourus	206	umbratilis	125
albeolus, Chirope	136	zonatus	123
Luxilus	129	alburnus, Perca	350
Notropis	136	albus, Atherinichthys	251
Notropis megalops	129	Centropomus	307
albescens, Echeueis	449	Coregonus	62, 65
Paraliparis	404	Coregonus clupeaformis	65
Remorina	449	Gymnotus	101
albicaudus, Auchenopterus	457	Eriscion	356
albidactylus, Exocoëtus	202	Lepisosteus	37
albidum, Hamulon	329	Otolithus	356
albidus, Gadus	214	Parasaphirhynchus	35
Labra	307	Ptychostomus	110
Osmerus	163	Alcidae	375
Pimelodus	150, 153	thoburni	375
Ptychostomus	110	Aldrovandia	75
albifimbria, Scorpæna	371	goodii	75
albifrons, Gymnotus	101	gracilis	75
albigutta, Cathetostoma	454	macrochir	75
Kathetostoma	454	pallida	75
Paralichthys	224	rostrata	75
albiguttus, Paralichthys	224	Alectis	274
albimentis, Gymnothorax	93	crinitus	274
Lycodontis	93	Aledon	503
alpinus, Lucifer	70	storeri	503
albirostris, Corythoichthys	243	Alepes	271
Hippichthys	243	melanoptera	271
Prionotus	408	Alepidosaurus asculapius	171
albolaseata, Scorpæna	371	borealis	171
albolineatus, Fundulus	177	(caulopus) poeyi	171
albonimaculatus, Crossias	389	serra	171
Serranus	317	alepidota, Chænopsis	465
albopunctatus, Gobius	440	alepidotum, Gobiosoma	446

List of scientific names—Continued

	Page		Page
alepidotus, Chætodon	265	Alloctotus	391
Derepodichthys	479	embryum	391
Gobius	445, 446	Allogambusia	188
Lucioblennius	465	cana	188
Alepisauridae	171	tridentiger	188
Alepisaurus	171	Allosmerus	67
æsculapius	171	attenuatus	67
altivelis	171	Allosomus	61, 64
azureus	171	Allotus	300
ferox	171	humilis	300
Alepocephalidae	51, 54	Allotriognathus	215
Alepocephalus	51	Allurus	402
agassizii	51	almeida, Belone	195
asperifrons	52	Alopecias	18
bairdii	52	vulpes	18
convexifrons	52	Alopias	18
fundulus	52	macrourus	18
macropterus	53	vulpinus	18
productus	51	Alopidæ	18
rostratus	51	Alosa	42
tenebrosus	52	alabama	43
Aleposomus	53	apicalis	43
copel	53	bishopi	43
aletes, Hudsonius	132	clupea	42
aleutensis, Acipenser	34	cyanonotus	42
Lyconectes	471	lineata	42
aleuticus, Sebastodes	365	menhaden	44
Zalopyr	365	obiensis	42
aleutica, Raja	20	præstabilis	42
aleuticus, Cottus	385	sapidissima	42
Sebastosomus	365	teres	46
Thecopterus	392	alosoides, Amphiodon	41
alexandri, Acipenser (Antaeus)	34	alpense, Leucichthys	63
Rosicola	365	Alpharaia	24
Sebastodes	365	Alphestes	312
alexandri, Orthogoriscus	503	afer	312
Alexurus	437	chloropterus	312
armiger	437	lightfooti	313
alfari, Cichlasoma	419	multiguttatus	313
Parapetenia	419	Alpsmaris	163
Alfaro	193	risso	163
acutiventralis	193	alta, Xurina	137
acutiventris	193	altavela, Pteroplatea	29
culturatus	193	Raja	29
alfredi, Kryptophanaron	234	alternana, Scarus	434
Algansea	115	alternata, Perca mitchilli	307
affinis	116	alticola, Catostomus	105
bicolor	122	Alteus	463
dugesii	122	saltatorius	463
formosa	116	alticus, Salarias	463
lacustris	122	altifrons, Astatheros	420
obesa	116	Heros	420
rubescens	122	altipinna, Belone	197
saliei	116	altipinnus, Alburnellus	127
stigmatura	116	Hydrophlox	127
tarascorum	116	Loricaria	162
tincella	115	Micropogon	349
Algoma	144	altivelis, Alepisaurus	171
amara	144	Corallicola	456
aha, Tentoga	424	Cremnobates	456
ahala, Acanthurus	424	Sebastolobus	364
alcese, Squalus	363	Serranus	310
aliciolus, Trachurus	119	altus, Bubalichthys	102
Alinectes	279	Chormemus	278
alpes, Salmo	402	Hudsonius	137
alithostomus, Diapterus	60	Oligophites	278
Alitardius	342	Præacanthus	323
corallinus	377	Pseudopriacanthus	323
Allector	484	aluta, Sciæna	344
chelona	484	alutaceus, Acrocheilus	112
allector, Dolopenthyx	509	Alutarius amphacanthoides	496
alleganiensis, Moxostoma	111	macracanthus	496
Salmo	60	obliteratus	496
alleni, Rathbunella	452	Alutera	495
alletteratus, Euthynnus	258	cinerea	496
Scomber	258	guntheriana	496
Alinectes	403	monoceros	496
alhsi, Crossias	389	picturata	495
allisoni, Neothunnus	260	punctata	495
Thunnus	260	Aluteres berardi	496
Allochur	402	cuspidanda	495
Alloclinus	456	Aluteres anginosus	496
holderi	456	culturifrons	495
		holbrookii	495

List of scientific names—Continued

	Page		Page
<i>Aluterus venosus</i>	495	<i>Ameiurus nebulosus catulus</i>	154
<i>alutum</i> , <i>A. autmentum</i>	366	<i>platycephalus</i>	154
<i>alutus</i> , <i>Apogon</i>	296	<i>ponderosus</i>	152
<i>Apogonichthys</i>	296	<i>prosthistus</i>	154
<i>Nector</i>	346	<i>vulgaris</i>	153
<i>Sebastichthys</i>	366	<i>americana</i> , <i>Lucioperca</i>	282
<i>Alvarius</i>	282, 294	<i>Manta</i>	32
<i>fonticola</i>	295	<i>Morone</i>	307
<i>lateralis</i>	294	<i>Morhua</i>	210
<i>alveata</i> , <i>Trigonorhina</i>	24	<i>Perca</i>	281, 307
<i>alveatus</i> , <i>Bathytroctes</i>	52	<i>Raja</i>	25
<i>alvifrons</i> , <i>Bathytroctes</i>	52	<i>Scorpaena</i>	363
<i>alvordii</i> , <i>Cottus</i>	385	<i>americanum</i> , <i>Apogon</i>	295
<i>Alvordius</i>	283	<i>americanus</i> , <i>Acanthias</i>	21
<i>aspro</i>	283	<i>Alburnus</i>	350
<i>crassus</i>	283	<i>Ammodytes</i>	451
<i>evermanni</i>	284	<i>Amphiprion</i>	308
<i>evides</i>	284	<i>Apogon</i>	285
<i>guntheri</i>	283	<i>Balistes</i>	493
<i>macrocephalus</i>	284	<i>Blennius</i>	473
<i>maculatus</i>	283	<i>Cyprinus</i>	115, 350
<i>maxinkuckiensis</i>	284	<i>Enchelyopus</i>	213
<i>ouachute</i>	284	<i>Eques</i>	352, 353
<i>peltatus</i>	283	<i>Esox</i>	172, 173
<i>phoxocephalus</i>	284	<i>Esox lucius</i>	173
<i>roanoka</i>	284	<i>Hemirhamphus</i>	393
<i>sellaris</i>	284	<i>Histiophorus</i>	263
<i>spillmani</i>	283	<i>Istiophorus</i>	263
<i>Alysia</i>	168	<i>Labrus</i>	424
<i>loricata</i>	168, 169	<i>Lophius</i>	504
<i>amabilis</i> , <i>Alburnus</i>	124	<i>Macrozoarces</i>	473
<i>Notropis</i>	124	<i>Menticirrhus</i>	330
<i>amara</i> , <i>Albana</i>	144	<i>Petromyzon</i>	8
<i>Dionda</i>	144	<i>Pleuronectes</i>	227
<i>amarillas</i> , <i>Guatavere</i>	309	<i>Polynemus</i>	256
<i>amarus</i> , <i>Hudsonius</i>	132	<i>Polyprion</i>	308
<i>amates</i> , <i>Phallichthys</i>	190	<i>Pseudopleuronectes</i>	227
<i>Pocilia</i>	190	<i>Scorpaena</i>	393
<i>ambassis</i> , <i>Sargus</i>	336	<i>Squalus</i>	19
<i>ambiguus</i> , <i>Lutianus</i>	327	<i>amethystino</i> , <i>Maurolleus</i>	73
<i>Merluccius</i>	214	<i>amethystinus</i> , <i>Salmo</i>	59
<i>Mesopron</i>	327	<i>Ama</i>	38, 295
<i>Ambledon</i>	352	<i>binotata</i>	295
<i>bubalis</i>	102	<i>calva</i>	38
<i>concinus</i>	352	<i>canina</i>	38
<i>grunniens</i>	352	<i>cinerea</i>	38
<i>lineatus</i>	352	<i>guadalupensis</i>	296
<i>neglectus</i>	352	<i>immaculata</i>	40
<i>saturnus</i>	348	<i>lentiginosa</i>	38
<i>Ambloplites</i>	303	<i>marmorata</i>	38
<i>cavifrons</i>	303	<i>occidentalis</i>	38
<i>rupestris</i>	303	<i>ocellcauda</i>	38
<i>amblops</i> , <i>Felichthys</i>	159	<i>ornata</i>	38
<i>Rutilus</i>	133	<i>pliquotii</i>	38
<i>Trachoristes</i>	159	<i>reticulata</i>	38
<i>Amblyopsidae</i>	194	<i>retrosella</i>	295
<i>Amblyopsis</i>	194, 195	<i>subcaerulea</i>	38
<i>mexicanus</i>	448	<i>thompsoni</i>	38
<i>spelæus</i>	195	<i>viridis</i>	38
<i>amblyopsis</i> , <i>Cultus</i>	437	<i>Amiatus</i>	38
<i>Eleotris</i>	437	<i>calva</i>	38
<i>Amblyopus brevis</i>	448	<i>Amnichthys</i>	297
<i>sagitta</i>	448	<i>diapterus</i>	297
<i>Amblypomacentrus</i>	413	<i>Amnidae</i>	38
<i>Amblyraja</i>	24, 25	<i>Amitra</i>	404, 405
<i>amblyrhynchus</i> , <i>Caranx</i>	271	<i>liparina</i>	404, 405
<i>Hemicarax</i>	271	<i>Amitrichthys</i>	404
<i>Amblyseion</i>	344	<i>Amitrus</i>	404
<i>argenteus</i>	344	<i>Amiurus</i>	153
<i>Ameiuridae</i>	151	<i>brachyacanthus</i>	154
<i>Ameiurus</i>	153	<i>brunneus</i>	154
<i>bolli</i>	153	<i>cragini</i>	154
<i>brachyacanthus</i>	154	<i>lophius</i>	153
<i>catus</i>	153	<i>meridionalis</i>	152
<i>dugei</i>	153	<i>niveiventris</i>	153
<i>erebennus</i>	153	<i>obesus</i>	154
<i>lacustris okeechobeensis</i>	152	<i>prosthistus</i>	154
<i>marmoratus</i>	154	<i>Ammocetes</i>	8
<i>melas</i>	154	<i>apoptera</i>	10
<i>mispliensis</i>	154	<i>aureus</i>	10
<i>natalis</i>	153	<i>bicolor</i>	8
<i>natalis analis</i>	153	<i>borealis</i>	9, 10
<i>nebulosus</i>	154	<i>cibarius</i>	10
		<i>concolor</i>	9

List of scientific names—Continued

	Page		Page
<i>Ammocetes unicolor</i>	9	<i>analogus, Abudedefduf</i>	415
<i>Ammocetus</i>	8	<i>Epinephelus</i>	310
<i>Ammocrypta</i>	282, 289	<i>Euchistodus</i>	415
<i>beanii</i>	289	<i>Kyphostus</i>	339
<i>clara</i>	288	<i>Pimelepterus</i>	339
<i>gelida</i>	289	<i>analostana, Cyprinella</i>	129
<i>vivax</i>	288	<i>Eregalia</i>	129
<i>ammocryptus, Tetraodon</i>	499	<i>Anarhichadide</i>	472
<i>Ammodytes</i>	451	<i>Anarhichas</i>	472
<i>alascanus</i>	451	<i>lupus</i>	472
<i>americanus</i>	451	<i>orientalis</i>	472
<i>dubius</i>	451	<i>Anarrhichas</i>	329
<i>personatus</i>	451	<i>Anarrhichas karak</i>	472
<i>septipinnis</i>	451	<i>latifrons</i>	472
<i>tobianus</i>	451	<i>leopaidus</i>	472
<i>vittatus</i>	451	<i>lepturus</i>	472
<i>Ammodytidae</i>	451	<i>maculatus</i>	472
<i>Ammopleuropis</i>	230	<i>minor</i>	472
<i>amoenus, Albunellus</i>	123	<i>strigosus</i>	472
<i>Notropis</i>	123	<i>vomerinus</i>	472
<i>Amore pixuma</i>	436	<i>Anarrhichthyidae</i>	422
<i>amorea, Gobius</i>	436	<i>Anarrhichthys</i>	472
<i>Amorpocephalus</i>	430	<i>felis</i>	472
<i>granulatus</i>	430	<i>ocellatus</i>	472
<i>amphacanthoides, Alutarius</i>	496	<i>anasimos, Acipenser (Huso)</i>	35
<i>Amphigonopterus</i>	409	<i>Acanthias canescens</i>	161
<i>aurora</i>	409	<i>anceps, Cottus</i>	386
<i>Amphilophus</i>	420, 422	<i>Anchisomus</i>	498
<i>freibeli</i>	422	<i>geometricus</i>	499
<i>labiatus</i>	422	(<i>Anchisomus</i>) <i>caudacinctus, Prilonotus</i>	500
<i>lobochilus</i>	422	<i>Anchoa</i>	47, 50
<i>Amphiodon</i>	41	<i>Anchova</i>	47
<i>alosoides</i>	41	<i>Anchovia</i>	50
<i>Amphioides</i>	7	<i>abbotti</i>	50
<i>Amphiovis</i>	7	<i>arenicola</i>	49
<i>amphiovis, Cantherines</i>	494	<i>brevirostrum</i>	51
<i>Monocanthus</i>	494	<i>cheilostoma cayoium</i>	49
<i>Amphiprion americanus</i>	308	<i>clupeoides</i>	50
<i>australis</i>	308	<i>eigenmannia</i>	50
<i>mateuelo</i>	235	<i>elongata</i>	49
<i>Amphistichus</i>	410	<i>filifera</i>	50
<i>argenteus</i>	410, 411	<i>lepidentostole</i>	49
<i>heermanni</i>	411	<i>macrolepidota</i>	50
<i>similis</i>	411	<i>parva</i>	49
<i>Amphotistius</i>	28	<i>platygyrea</i>	49
<i>dipterurus</i>	29	<i>scitula</i>	49
<i>sabinus</i>	29	<i>trinitatus</i>	50
<i>say</i>	28	<i>anchovia, Sardinella</i>	43
<i>ampla, Makaira</i>	263	<i>Sardina</i>	43
<i>amplexocollis, Saitothodus</i>	360	<i>Anchoviella</i>	47
<i>Amplva</i>	50, 51	<i>abbotti</i>	50
<i>balboæ</i>	50	<i>arenicola</i>	49
<i>amplus, Searus</i>	432	<i>argyrophana</i>	48
<i>Tetrapturus</i>	263	<i>astilbe</i>	49
<i>ampullaceus, Ophiognathus</i>	95	<i>branchiomelas</i>	49
<i>Saccopharynx</i>	95	<i>cayoium</i>	49
<i>Anablepidæ</i>	194	<i>cheilostoma</i>	48
<i>Anableps</i>	194	<i>compressa</i>	50
<i>anableps</i>	194	<i>cubana</i>	48
<i>dowei</i>	194	<i>cultrata</i>	48
<i>tetraphthalmus</i>	194	<i>curta</i>	48
<i>anableps, Anableps</i>	194	<i>delicatissima</i>	48
<i>Anacanthini</i>	202	<i>eigenmannia</i>	50
<i>Anacartus guatemalensis</i>	98	<i>elongata</i>	49
<i>anagallinus, Lepomis</i>	300	<i>epsetus</i>	48
<i>Anago</i>	80	<i>exigua</i>	47
<i>macurus</i>	80	<i>filifera</i>	50
<i>nitens</i>	80	<i>guanensis</i>	49
<i>anago, Conger</i>	80	<i>helleri</i>	48
<i>analagutta, Pomacentrus</i>	414	<i>ischana</i>	48
<i>analis, Ameiurus natalis</i>	153	<i>lepidentostole</i>	49
<i>Clinocottus</i>	390	<i>lucida</i>	48
<i>Conger</i>	80	<i>lyolepis</i>	49
<i>Eupomacentrus</i>	413	<i>marichus</i>	47
<i>Hyperprosepon</i>	410	<i>mitchilli</i>	48
<i>Hypoclinichthys</i>	410	<i>mundiola</i>	50
<i>Lutjanus</i>	328	<i>naso</i>	49
<i>Mesoprius</i>	328	<i>panamensis</i>	50
<i>Notacanthus</i>	76	<i>parva</i>	49
<i>Oligocottus</i>	390	<i>per fasciata</i>	47
<i>Orthogoriscus</i>	503	<i>perthecata</i>	47
<i>Pomacentrus</i>	413	<i>platygyrea</i>	49
<i>Seyris</i>	274	<i>rastralis</i>	50
<i>Umbrina</i>	350	<i>roberti</i>	49
		<i>scitula</i>	49

List of scientific names—Continued

	Page		Page
<i>Anchoviella scofieldi</i>	49	<i>Anisotremus scapularis</i>	332
<i>spinifera</i>	50	<i>serrula</i>	333
<i>trinitatus</i>	50	<i>splenatus</i>	332
<i>Ancistrus</i>	161	<i>surinamensis</i>	332
<i>aspidolepis</i>	161	<i>tæniatus</i>	332
<i>chagresi</i>	160	<i>virginicus</i>	332
<i>guacharote</i>	161	<i>anisurum, Movostoma</i>	109
<i>mediana</i>	161	<i>anisurus, Catostomus</i>	109
<i>medians</i>	161	<i>anjerensis, Gobius</i>	442
<i>planiceps</i>	161	<i>anna-carolina, Mugilomorus</i>	39
<i>spinosus</i>	160	<i>anna, Agonus</i>	394
<i>Ancyrodon</i>	354	<i>Cottus</i>	384
<i>jaculidens</i>	354	<i>annectans, Gambusia</i>	187
<i>parvipinnis</i>	353	<i>annectens, Coccothrus</i>	217
<i>ancyrodon, Lonchurus</i>	354	<i>Gambusia</i>	187
<i>Macrodon</i>	354	<i>Prapichthys</i>	187
<i>Ancylopsetta</i>	224	<i>annectens hesperis, Prapichthys</i>	187
<i>dendritica</i>	224	<i>annularis, Centropristis</i>	319
<i>quadrocellata</i>	224	<i>Nauderus</i>	278
<i>andree, Rhinoscopelus</i>	168	<i>Pomoxis</i>	304
<i>Scopelus</i>	168	<i>Sparus</i>	338
<i>androsiensis, Gobiesol</i>	489	<i>annulata, Melanura</i>	174
<i>Angelichthys</i>	361	<i>annulatum, Eloglossum</i>	146, 174
<i>cularis</i>	361	<i>annulatus, Antennarius</i>	506
<i>isabelita</i>	361	<i>Chelichthys</i>	499
<i>townsendi</i>	361	<i>Tetrodon</i>	499
<i>angelus, Squatina</i>	22	<i>anogenus, Hybopsis</i>	136
<i>anginosus, Aluterus</i>	406	<i>Notropis</i>	136
<i>anglorum, Lumpus</i>	398	<i>anolis, Saurus</i>	164
<i>anguliformis, Ophichthys (Sphagebranchus)</i>	84	<i>anomala, Dekaya</i>	357
<i>Sphagebranchus</i>	84, 91	<i>Anomalopidae</i>	234
<i>Anguilla</i>	77	<i>anomalum, Camptostoma</i>	146
<i>anguilla</i>	77	<i>anomalous, Caulolatilus</i>	357
<i>aterima</i>	77	<i>Rutilus</i>	146
<i>australis</i>	77	<i>Anoplagonus</i>	397
<i>blephura</i>	77	<i>inermis</i>	398
<i>chrysypa</i>	77	<i>Anoplarchus</i>	468
<i>cubana</i>	77	<i>cristagalli</i>	468
<i>laticauda</i>	77	<i>cristagalli archolepis</i>	468
<i>lutea</i>	77	<i>insignis</i>	468
<i>novaeorleansis</i>	77	<i>purpurescens</i>	468
<i>novaterre</i>	77	<i>Anoplogaster</i>	233
<i>oceanica</i>	77	<i>cornutus</i>	233
<i>punctatissima</i>	79	<i>Anoploploma</i>	373
<i>rostrata</i>	77	<i>fimbria</i>	373
<i>tenuirostris</i>	77	<i>merlangus</i>	373
<i>texana</i>	77	<i>Anoplopomidae</i>	373
<i>tyrannus</i>	77	<i>anoplos, Uranoscopus</i>	454
<i>wabashensis</i>	77	<i>anoplos, Thyriscus</i>	379
<i>xanthomelas</i>	77	<i>Anopsus</i>	8
<i>anguilla, Anguilla</i>	77	<i>Anosmius</i>	500
<i>Ictalurus</i>	152	<i>Anostomidae</i>	100
<i>Muraena</i>	77	<i>Ansonia cuvieri</i>	264
<i>anguillarlis, Blennius</i>	470, 473	<i>ansonii, Salmo</i>	55
<i>Tribranchus</i>	77	<i>Antacea</i>	10
<i>Anguillidae</i>	77	<i>Antaceus</i>	34
<i>anguilliformis, Pholidichthys</i>	465	<i>Antelochimara</i>	33
<i>Anguillidae</i>	77	<i>chaetirhamphus</i>	33
<i>anguina, Muraena</i>	90	<i>Antennaria avalonis</i>	506
<i>anguineus, Nerophis</i>	243	<i>Antennaridae</i>	504
<i>anguis, Lycodes</i>	475	<i>Antennarius</i>	505
<i>anguliceps, Macrourus</i>	206	<i>annulatus</i>	506
<i>Macrurus</i>	206	<i>astroscopus</i>	506
<i>angulifer, Cichlaurus</i>	416	<i>avalonis</i>	506
<i>Heros</i>	416	<i>corallinus</i>	506
<i>angulosus, Balistes</i>	492	<i>inops</i>	505
<i>Canthidermis</i>	492	<i>leopardinus</i>	505
<i>angusticeps, Belone</i>	196	<i>multicellatus</i>	506
<i>Coregonus</i>	136	<i>nuttalli</i>	506
<i>Sphceroides</i>	498	<i>ocellatus</i>	505
<i>Strongylura</i>	196	<i>pleurophthalmus</i>	505
<i>Tetrodon</i>	498	<i>principis</i>	505
<i>angustidens, Macrostoma</i>	166	<i>radiosus</i>	506
<i>angustifrons, Epinephelus</i>	311	<i>reticularis</i>	505
<i>Paraliparis</i>	405	<i>sanguineus</i>	505
<i>Serranus</i>	311	<i>scaber</i>	505
<i>angustimanus, Merluccius</i>	214	<i>sonnagli</i>	500
<i>anisacanthus, Dolopichthys</i>	509	<i>strigatus</i>	505
<i>Anisarchus</i>	470	<i>tenebrosus</i>	505
<i>medius</i>	470	<i>tenuifilis</i>	505
<i>Anisochæstodon</i>	359	<i>tigris</i>	500
<i>Anisotremus</i>	352	<i>verrucosus</i>	506
<i>bicolor</i>	352	<i>antennatus, Cyclichthys</i>	502
<i>davidsoni</i>	352	<i>Diodon</i>	502
<i>interruptus</i>	352	<i>Anthias</i>	322

List of scientific names—Continued

	Page		Page
<i>Anthias aquilonarias</i>	328	<i>Apocope adobe</i>	141
<i>caballerote</i>	325	<i>carringtonii</i>	140, 141
<i>chernia</i>	311	<i>covesii</i>	141
<i>formosus</i>	330	<i>falcata</i>	141
(<i>Hemianthias</i>) <i>peruanus</i>	322	<i>henshawi</i>	141
<i>jocoi</i>	325	<i>klamathensis</i>	141
<i>quartus rondeleti</i>	326	<i>nevadensis</i>	141
<i>rubirubia</i>	327	<i>nubila</i>	141
<i>saponaceus</i>	316	<i>oscula</i>	140
<i>striatus</i>	311	<i>robusta</i>	141
<i>tenuis</i>	322	<i>umatilla</i>	141
<i>vivanus</i>	322	<i>valifera</i>	141
<i>anthias, Labrus</i>	322	<i>vulnerata</i>	141
<i>anthracinus, A. cypenser (Huso)</i>	35	<i>yarrovi</i>	141
<i>antica, Algansea</i>	122	(<i>Apocope</i>) <i>nevadensis, Rhinichthys</i>	141
<i>anticus, Hesperoleucus</i>	122	<i>velifera, Rhinichthys</i>	141
<i>Antigonia</i>	358	<i>apoda, Perca</i>	325
<i>capros</i>	358	<i>Pleuronectes</i>	229
<i>Antigonidae</i>	358	<i>Apodes</i>	77
<i>antillarum, Conodon</i>	333	<i>apodes, Lutjanus</i>	325
<i>antillarum, Caranx</i>	273	<i>Apodichthys</i>	466
<i>Chilomycterus</i>	502	<i>flavidus</i>	466, 467
<i>Cyclichthys</i>	502	<i>fuorum</i>	467
<i>Fundulus</i>	175	<i>inornatus</i>	467
<i>Sicydium</i>	447	<i>sanguineus</i>	467
<i>Talismania</i>	53	<i>univittatus</i>	467
<i>Antimora</i>	211	<i>violaceus</i>	466
<i>microlepis</i>	211	<i>virescens</i>	467
<i>rhina</i>	211	<i>apodus, Lutjanus</i>	326
<i>rostrata</i>	211	<i>Apogon</i>	38, 295
<i>viola</i>	211	<i>alutus</i>	296
<i>antistius, Chænobryttus</i>	303	<i>americanum</i>	295
<i>antonensis, Pimelodus</i>	153	<i>americanus</i>	295
<i>antrostomus, Idiacanthus</i>	70	<i>atricaudus</i>	295
<i>anuli, Pimphales</i>	146	<i>atrodorsatus</i>	295
<i>Apareiodon</i>	100	<i>binotatus</i>	295
<i>compressus</i>	100	<i>dovii</i>	295
<i>dariensis</i>	100	<i>guadalupensis</i>	296
<i>Apeltes</i>	239	<i>imbarbis</i>	295
<i>quadracus</i>	239	<i>maculatus</i>	295
<i>apeltes, Gasterosteus</i>	239	<i>moluccensis</i>	295
<i>Apeltinae</i>	239	<i>pandionis</i>	296
<i>Aphia</i>	246	<i>pigmentarius</i>	295
<i>meridionalis</i>	246	<i>retrosella</i>	295
<i>Aphanopus</i>	262	<i>rex-mullorum</i>	295
<i>carbo</i>	262	<i>ruber</i>	295
<i>minor</i>	262	<i>sellicauda</i>	295
<i>Aphrododerus</i>	215	<i>apogon, Borophryne</i>	509
<i>cookianus</i>	215	<i>Apogonichthys</i>	296
<i>Aphoristia</i>	230	<i>alutus</i>	296
<i>atricauda</i>	231	<i>melampodus</i>	296
<i>diomedea</i>	231	<i>perdix</i>	296
<i>marginata</i>	231	<i>puncticulatus</i>	296
<i>nebulosa</i>	231	<i>stellatus</i>	296
<i>ornata elongata</i>	231	<i>strombi</i>	296
<i>pigra</i>	230	<i>Apogonidae</i>	295
<i>pusilla</i>	231	<i>Apomotis</i>	298
<i>Aphredoderidae</i>	215	<i>cyaneus</i>	298
<i>Aphredoderus</i>	215	<i>euryurus</i>	299
<i>gibbosus</i>	215	<i>ischyurus</i>	299
<i>sayanus</i>	215	<i>phenax</i>	299
<i>Aphyonius</i>	484	<i>punctatus</i>	299
<i>gelatinosus</i>	484	<i>symmetricus</i>	299
<i>mollis</i>	484	<i>appendiculatus, Centropomus</i>	305
<i>apia, Pirati</i>	313	<i>Exocoetus</i>	201
<i>apiarius, Saranus</i>	308	<i>appendix, Lepomis</i>	299
<i>apiatus, Lepomis</i>	299	<i>Lethenteron</i>	9
<i>apicalis, Aloa</i>	43	<i>Petromyzon</i>	9
<i>Echeneis</i>	449	<i>approximans, Polynemus</i>	256
<i>Apomichthys</i>	230	<i>Pomadasya</i>	334
<i>dumerli</i>	230	<i>Aprior</i>	15
<i>nebulosus</i>	230	<i>ariomus</i>	328
<i>Apistes exul</i>	370	<i>aprior, Gerres</i>	342
<i>Apleion</i>	290, 298	<i>Apriorodon</i>	15
<i>pottii</i>	290	<i>isodon</i>	15
<i>Aplites</i>	298	<i>punctatus</i>	15
<i>Aplodinotus</i>	352	<i>Apristurus</i>	12
<i>grunniens</i>	352	<i>brunneus</i>	12
<i>neglectus</i>	352	<i>profundorum</i>	12
<i>Aplodon</i>	339	<i>Apodon</i>	474
<i>margaritifera</i>	339	<i>corteziana</i>	474
<i>Aplurus</i>	261	<i>cortezianus</i>	474
<i>simplex</i>	261	<i>Aprolepis</i>	444
<i>Apocope</i>	140	<i>barbata</i>	444

List of scientific names—Continued

	Page		Page
Apsedobanchus	354	arctifrons, Calamus	337
Apsicephalus	499	Citharichthys	219
Apsilus dentatus	328	Arctoscopus	357
Apterichthys	84	japonicus	357
selachops	84	Arctozenus	165
Apteronotus	101	borealis	165
passan	101	coruscans	165
rostratus	101	arctum, Siphostoma	242
Apterurus	31	arcturus, Salmo	61
fabroni	31	Salvelinus	61
Apthalmichthys	89	arctus, Syngnathus	242
caribbeus	89	arcuatum, Haemulon	330
javanicus	89	Hyperprosopon	410
Aptocyclus	398	arcuatus, Bathygadus	202
ventricosus	398	Chetodon	360
apua, Bodianus	313	Gadomus	202
Trisotropis venenosus	313	Pomacanthus	360
apus, Platyroctes	53	ardens, Catostomus	105
aquaedulcis, Gymnothorax	93	Hypsilepis	126
aquaedulcis, Muraena	91, 93	Lythrurus	126
Rahula	91	ardeola, Belone	195
aquila, Raja	31	ardesaca, Gila	118
Raja	31	ardesiacus, Lepomis	301
Sciæna	61	arenaceus, Citharichthys	219
aquilarum, Salmo	58	arenata, Umbra	351
aquilonaris, Anthias	328	arenatus, Arus	160
Etelades	328	Pomacanthus	323
aquosus, Pleuronectes	217	Rhynchichthys	140
arabatchi, Salmo	55	Rypticus	316
arabicus, Chanos	40	arenicola, Anchovia	49
araceaga, Pseudoscarus	434	Anchoviella	49
Scarus	433, 434	Carapus	498
aracansanus, Notropis	124	Fierasfer	498
Notropis telecopus	124	Gillellus	455
aræ, Pristigaster	12	arenosus, Gadus	210
aræa, Atherina	246	argalus, Belone	196
aræopus, Catostomus	106	Strongylura	196
Aramaca	218	arge, Alburnellus	124
aramaca, Hemirhombus	218	Chirostoma	250
Pleuronectes	218	Esloparum	250
Rhombus	218	Kuhlia	305
arangoi, Chcerophilus	426	Notropis	124
arara, Haemulon	330	argentata, Ciliata	208
Serranus	311	Motella	208
aratus, Mesoprius	327	argentatus, Astyanax	96
Raizero	327	Platygyrus	128
Arbaciosa	490	argentea, Bathyluopea	281
eos	491	Muraena	77
humeralis	490	Sciæna	334
rhessodon	490	Selene	275
rupestris	491	Sphyræna	255
zebra	491	Steindachneria	203
arborifer, Linophryne	510	argenteolus, Lychnopoles	72, 73
arcansanus, Etheostoma zonale	289	argenteum, Hyperprosopon	410
Nanostoma	289	punctatum, Hyperprosopon	410
arcansanus, Notropis	124	argenteus	9
Notropis telecopus	124	Amblysoion	344
Archaulus	377	Amphistichus	411
biseriatus	377	Brycon	98
archidum, Elattarchus	345	Centronotus	277
Odontoscion	345	Diplodus	338
Archocentrus	420	Eucinostomus	340
centrarchus	420	Fario	55
immaculatus	420	Hyperprosopon	410
nigrofasciatus	420	Kurtus	323
spilurus	420	Larimus	344
spinossissimus	420	Leuciscus	117
archolepis, Anoplarchus cristigalli	468	Micropogon	349
Anoplarchus purpureus	468	Petromyzon	9
Archomenidia	248	Sarchirus	36
sallei	248	Sargus	338
Archopercæ	313	Saurus	277
Archoplites	303	Sparus	337
interruptus	303	Synodus	40
Archosargus	338	Trachinotus	277
aries	338	Trichiurus	262
probatoccephalus	338	Argentina	68
arctica, Liparis	400	carolina	39
arcticum, Benthosema	170	glossodonta	40
arcticus, Chironectes	504	pennanti	73
Galeocerdo	14	pretiosa	67
Gymnogaster	216	sialis	68
Salmo	66	silus	68
Scopelus	170	sphyraena	68
Squalus	14	striata	68
Trachipterus	216	syrtensium	68

List of scientific names—Continued

	Page		Page
Argentinidae.....	68	Arius commersonn.....	149
argentinus, Pimelodus.....	152	dasycephalus.....	150
argentissimus, Gasterosteus.....	238	elaturus.....	149
Plagopterus.....	147	emmeline.....	151
argentiniventris, Lütjanus.....	326	equestris.....	149
Mesopomus.....	326	evermanni.....	151
argentosa, Dionda.....	144	furthi.....	151
Arges.....	159	grandcaissus.....	149
sabalo.....	159	guatemalensis.....	150
Argo.....	265	hypophthalmus.....	151
steindachneri.....	265	insculptus.....	149
argus, Cephalopholis.....	309	kessleri.....	149
Gymnothorax (Limamurana).....	90	laticeps.....	150
Murana.....	90	liropus.....	151
Pleuronectes.....	217	melanopus.....	150
Squalus.....	12	milberti.....	149
argyrea, Scarina.....	262	multiradiatus.....	151
Sciæna.....	341	osculus.....	149
Argyrosus.....	275	planceps.....	149
brevoorti.....	276	platypogon.....	149
cerstedii.....	275	seemani.....	150
pacificus.....	276	spixii.....	150
setipinnis.....	275	steindachneri.....	151
unimaculatus.....	275	taylori.....	151
vomere.....	275	truncatus.....	149
argyrosus, Pogonichthys.....	116	arizone, Pantosteus.....	104
Argyreus.....	139	Arlina.....	287
dulcis.....	139	atrinipinnis.....	286
notabilis.....	141	effulgens.....	287
nubilis.....	140	arlingtonia, Gambusia.....	177
osculus.....	141	armata, Bairdiella.....	346
rubripinnis.....	128	Sciæna.....	273
argyreus, Fario.....	55	armatus, Aspidophorus.....	395
Argyrosus brevoorti.....	276	Centropomus.....	306
filamentosus.....	275	Dactylobatus.....	27
mauricei.....	275	Leptocottus.....	390
mitchilli.....	275	Nector.....	346
setifer.....	275	Nematonurus.....	203
triacanthus.....	275	Serranus.....	313
argyrus, Hybodonathus.....	144	armiger, Aleurus.....	437
argyroleucos, Bodianus.....	345	arnillo, Mesopomus.....	328
argyro-leucos et euguis, Bodianus.....	345	Arnion.....	252
Argyropeleus.....	74	Arnoglossus ventralis.....	220
affinis.....	74	arnoldi, Limia.....	191
durvilli.....	74	Arothron.....	499
hemigymnus.....	74	erethizon.....	500
lychnus.....	74	testudinatus.....	499
olferi.....	74	Artedi.....	8
argyrophana, Anchoviella.....	48	artedi, Coregonus.....	63
argyrophanus, Engraulis.....	48	Leucichthys.....	63
argyrops, Sparus.....	336	Rhamphichthys.....	101
argyrosoma, Damalichthys.....	412	artedi busselli, Leucichthys.....	63
Argyrosomus.....	61	Artediellus.....	377
alascanus.....	62	atlanticus.....	377
ariensis.....	63	carnaticus.....	378
hoys.....	64	clarki.....	377
huronius.....	62	macanthus.....	378
johannæ.....	63	pacificus.....	377
nigripinnis.....	62, 63	unicinatus.....	377
sisco.....	62	artedi, Polynemus.....	256
tulibee.....	64	Artedius.....	377
zenithicus.....	64	asperulus.....	376
argyrosomus, Damalichthys.....	412	fenestralis.....	376
Argyrotæna.....	451	lateralis.....	377
argyrurus, Coryphæna.....	265	notospilotus.....	376
argyrus, Pimelodus.....	152	pugetensis.....	375
aries, Archosargus.....	338	quadriseriatus.....	376
Sargus.....	338	artesan, Nivicola.....	292
Aridia.....	147	Poculichthys.....	292
Ariodes.....	150	Arthrophallus.....	184
ariommus, Aprion.....	328	arubensis, Pœcilia vanderpoli.....	193
Paranotropis.....	125	arundinaceus, Syngnathus.....	240
Photogenus.....	125	ascanni, Silus.....	68
Ariopsis.....	149	Asclichthys.....	392
Ariosoma.....	79	rhoderus.....	392
caudale.....	80	ascendens, Syngnathus.....	241
gilberti.....	80	ascensionis, Holocentrus.....	235
opisthophthalma.....	80	Perca.....	235
Arius.....	150	ascita, Pimelodus.....	158
alatus.....	148	asellus, Cheilichthys.....	500
arenarius.....	150	Asematicthys.....	390
arenatus.....	150	taylori.....	390
arius.....	150	asper, Auchenopterus.....	456
assimilis.....	150	Cottus.....	383
brandti.....	148	Diodon.....	501
caerulescens.....	150		

List of scientific names—Continued

	Page		Page
asper, Exerpes.....	456	Asthenurus.....	208
Hexagrammos.....	373	atripinnis.....	208
Macrurus.....	203	astilbe, Anchoviella.....	49
Pleuronectes.....	227	Stolephorus.....	49
Taractes.....	265	astillero.....	49
aspera, Lumanda.....	227	astori, Petromyzon.....	10
asperifrons, Alepocephalus.....	52	Astrapogon.....	286
asprinna, Cottus.....	384	stellatus.....	286
asprerrinus, Balistes.....	492	Astroblepidae.....	159
Cottus.....	384	Astroblepus.....	159
Mylobatis.....	30	grixi.....	159
Pteromyxus.....	30	purpuratus.....	160
asperus, Serranus.....	310	Astrodermus.....	264
asperulus, Artedius.....	376	guttatus.....	264
Aspicottus.....	381	Astrolutes.....	376
bison.....	381	fenestralis.....	376
taurinus.....	381	Astronesthes.....	71
aspidolepis, Ancistrus.....	161	gemmifer.....	71
Chætoctomus.....	161	niger.....	71
Aspidophoroides.....	397	Astronesthes nigra.....	71
bartoni.....	397	richardsoni.....	71
grœnlandicus.....	397	Astronesthidae.....	71
güntheri.....	397	Astroscoptes.....	454
inermis.....	397	guttatus.....	454
monopterygius.....	397	y-gracum.....	454
olrik.....	397	zephyreus.....	454
tranquebar.....	397	astroscopus, Antennarius.....	506
Aspidophoroididae.....	397	Astyanax.....	96
Aspidophorus.....	395	æneus.....	96
armatus.....	395	argentatus.....	96
decagonus.....	394	brevortii.....	97
europeus.....	395	emperador.....	97
quadricornis.....	393	fasciatus.....	97
spinosissimus.....	184	macrophthalmus.....	97
aspidurus, Urolophus.....	30	mexicanus.....	96
Urotrygon.....	30	nasutus.....	97
Aspisurus.....	362	petenensis.....	97
asprella, Crystallaria.....	288	ruberrimus.....	97
asprellus, Pleurolepis.....	288	scabripinnis.....	97
Radulinus.....	379	Asymmetron.....	7
asprigenis, Poecilichthys.....	291	lucayanum.....	7
aspro, Alvordius.....	283	atæniatus, Chætodon.....	360
assimilis, Arius.....	150	Sarothrodus.....	360
Galeichthys.....	150	atchafalaya, Signalosa.....	46
Astatheros.....	420, 422	vanhyningi, Signalosa.....	47
altifrons.....	420	atelaspi, Acipenser (Huso).....	35
basilaris.....	421	Atelaxia.....	216
citrinellus.....	422	Ateleobranchium.....	207
dorsatus.....	420	pterotum.....	207
erythræum.....	422	aterrima, Anguilla.....	77
evermanni.....	420	athabasca, Leucichthys.....	62
globosum.....	421	Atheresthes.....	221
granadense.....	421	stomias.....	221
hedricki.....	421	Atherina.....	246
heterodontus.....	420	area.....	246
lauræ.....	421	boscu.....	247
lethrinus.....	420	boyeri.....	246
longimanus.....	420	brownii.....	48
maculicauda.....	421	carolina.....	246
margaritifer.....	421	evermanni.....	246
montezuma.....	421	harringtonensis.....	246
oblongus.....	421	hepsetus.....	246, 247
octofasciatus.....	421	humboldtiana.....	250
rectangularis.....	421	laticeps.....	246
robertsoni.....	420	martinica.....	248
rostratum.....	421	menidia.....	247
septemfasciatus.....	421	microps.....	246
sexfasciatus.....	421	microstoma.....	246
tuyrensis.....	421	mordax.....	67
asterias, Blennius.....	461	notata.....	247
Galeus.....	13	sardina.....	247
Mustelus.....	13	stipes.....	246
Spheroides.....	498	tænata.....	246
Sphæroides.....	498	velcana.....	246
Urolophus.....	30	viridescens.....	247
asterius, Blennius.....	461	vomerina.....	250
Asternopteryx.....	468	Atherinella.....	249
gunelliformis.....	468	eriarcha.....	249
Asternotremis.....	215	evermanni.....	248
mesotrema.....	215	panamensis.....	249
Asterotheca.....	396	Atherinichthys.....	250
alascana.....	396	albus.....	251
infraspinata.....	396	brevis.....	250
pentacantha.....	396	gracilis.....	247
		grandoculis.....	251

List of scientific names—Continued

	Page		Page
<i>Atherinichthys guatemalensis</i>	249	<i>atracaudus</i> , <i>Apogon</i>	295
<i>pschylepis</i>	249	<i>Symphurus</i>	231
<i>salae</i>	248	<i>atrilabiatus</i> , <i>Callionymus</i>	450
<i>Atherinidae</i>	246	<i>atrilatus</i> , <i>Zygometes</i>	185
<i>Atherinoides</i>	250	<i>atrilobata</i> , <i>Chromis</i> (<i>Furcaria</i>).....	412
<i>atherinoides</i> , <i>Chriodorus</i>	199	<i>Furcaria</i>	412
<i>Clupea</i>	51	<i>atrimana</i> , <i>Monolepis</i>	220
<i>Notropis</i>	123	<i>atrimanus</i> , <i>Caranx</i>	271
<i>Pterengraulis</i>	51	<i>Hemicaranx</i>	271
<i>Atherinomorus</i>	246	<i>Parasphyraenops</i>	297
<i>Atherinops</i>	252	<i>atrimentis</i> , <i>Menidia peninsulae</i>	247
<i>affinis</i>	252	<i>atripectus</i> , <i>Porogadus</i>	482
<i>insularum</i>	252	<i>atrides</i> , <i>Ditrema</i>	411
<i>cedrosensis</i>	252	<i>Lythrurus</i>	126
<i>guadalupae</i>	252	<i>Phanerodon</i>	411
<i>oregonia</i>	252	<i>atripinis</i> , <i>Arlina</i>	286
<i>regis</i>	252	<i>Asthenurus</i>	208
<i>Atherinopsis</i>	252	<i>Goodea</i>	183
<i>affinis</i>	252	<i>atriventer</i> , <i>Stomias</i>	69
<i>californiensis</i>	252	<i>atrocaudalis</i> , <i>Hybopsis</i>	133
<i>sonora</i>	252	<i>Notropis cayuga</i>	133
<i>tenuis</i>	251	<i>atrocaudus</i> , <i>Apogon</i>	295
<i>Athlennus</i>	197	<i>atro-cyaneus</i> , <i>Pomacentrus</i>	413
<i>atinga</i> , <i>Chilomycterus</i>	502	<i>atrodorsatus</i> , <i>Apogon</i>	295
<i>Diodon</i>	502	<i>atromaculata</i> , <i>Estrella</i>	287
<i>Guamaicu</i>	502	<i>atromaculatus</i> , <i>Cyprinus</i>	117
<i>atkinsi</i> , <i>Gasterosteus</i>	238	<i>Semotilus</i>	117
<i>atlantica</i> , <i>Elacate</i>	281	<i>atromaculatus thoreauanus</i> , <i>Semotilus</i>	117
<i>Emblemaria</i>	464	<i>atronasus</i> , <i>Cyprinus</i>	139, 140
<i>Scorpena</i>	370	<i>Rhinichthys</i>	140
<i>ruscula</i>	370	<i>atropurpureum</i> , <i>Ophidium</i>	469
<i>atlanticum</i> , <i>Oreosoma</i>	232	<i>atropurpureus</i> , <i>Epigeeichthys</i>	469
<i>Oreosoma</i>	232	<i>atrubens</i> , <i>Sebastes</i>	365
<i>atlanticus</i> , <i>Archieillus</i>	377	<i>Zalopyr</i>	365
<i>Benthodesmus</i>	262	<i>atrovirens</i> , <i>Sebastesichthys</i>	365
<i>Bregmaceros</i>	208	<i>Zalopyr</i>	365
<i>Callorhynchus</i>	32	<i>attavilla</i> , <i>Desyatis</i>	29
<i>Dibranchius</i>	508	<i>attenuatus</i> , <i>Cyclichthys</i>	502
<i>Megalops</i>	39	<i>attenuata</i> , <i>Vinciguerra</i>	73
<i>Neoliparis</i>	399	<i>attenuatum</i> , <i>Chirostoma</i>	250
<i>Promethus</i>	261	<i>attenuatus</i> , <i>Allosmerus</i>	67
<i>Rupiscartes</i>	463	<i>Careproctus</i>	403
<i>Salarias</i>	463	<i>Halosaurus</i>	75
<i>Sparus</i>	310	<i>Lycodapus</i>	479
<i>Tarpon</i>	39	<i>Mauroliscus</i>	73
<i>Tetragonurus</i>	269	<i>Osmerus</i>	67
<i>Thynnus</i>	280	<i>Paraliparis</i>	404
<i>atomarium</i> , <i>Sparisoma</i>	431	<i>Rhinoliparis</i>	405
<i>atomarius</i> , <i>Scarus</i>	431	<i>Attulus</i>	350
<i>Atopichthys</i>	54	<i>Atule</i>	270
<i>acus</i>	54	<i>atwoodi</i> , <i>Carcharias</i>	20
<i>cinctus</i>	54	<i>aubeenaubel</i> , <i>Etheostoma</i>	291
<i>cingulus</i>	54	<i>Oligocephalus</i>	291
<i>dentatus</i>	54	<i>aubrieti</i> , <i>Lutjanus</i>	327
<i>ensunculus</i>	54	<i>Auchenipterus</i>	456
<i>falcidens</i>	54	<i>Auchenistius</i>	457
<i>longidens</i>	54	<i>stahli</i>	457
<i>lychnus</i>	54	<i>Auchenopterus</i>	456
<i>obtusius</i>	54	<i>affinis</i>	457
<i>opichthys</i>	54	<i>albicaudus</i>	457
<i>scarius</i>	54	<i>asper</i>	456
<i>Atopoclinidae</i>	465	<i>cingulatus</i>	457
<i>Atopoclinus</i>	465	<i>fajardo</i>	457
<i>ringens</i>	465	<i>fasciatus</i>	457
<i>Atractoscion</i>	356	<i>grandioomis</i>	457
<i>nobilis</i>	356	<i>integripinnis</i>	457
<i>phoxocephalus</i>	356	<i>mexicanus</i>	457
<i>Atractosteus</i>	38	<i>monophthalmus</i>	456, 457
<i>luctus</i>	38	<i>nox</i>	457
<i>tristocchus</i>	38	<i>rubescens</i>	457
<i>tropicus</i>	38	<i>suctorum</i> , <i>Lobotes</i>	323
(<i>Atractosteus</i>) <i>berlandieri</i> , <i>Lepidosteus</i>	38	<i>Auctospina</i>	368
<i>ferox</i> , <i>Lepisosteus</i>	38	<i>auriculata</i>	368
<i>atramentatus</i> , <i>Symphurus</i>	231	<i>dalli</i>	369
<i>atraria</i> , <i>Perca</i>	317	<i>audax</i> , <i>Histiophorus</i>	263
<i>Siboma</i>	119	<i>audens</i> , <i>Menidia</i>	247
<i>Tigoma</i>	119	<i>Aulastome marcgravi</i>	245
<i>atrarius</i> , <i>Fimelodus</i>	154	<i>Auliscops</i>	239
<i>Xenomystax</i>	81	<i>spinescens</i>	239
<i>atratum</i> , <i>Myctophum</i>	169	<i>auliscus</i> , <i>Siphostoma</i>	240
<i>atricauda</i> , <i>Aphoristia</i>	231	<i>Syngnathus</i>	240
<i>Hydrargira</i>	174	<i>Anlophalus</i>	190
<i>atricaudata</i> , <i>Deuteron</i>	96	<i>elongatus</i>	190
<i>atricaudatus</i> , <i>Gephyrocharax</i>	96	<i>Aulorhynchidae</i>	239

List of scientific names—Continued

	Page		Page
<i>Aulorhynchus</i>	239	<i>australis</i> , Esor.....	173
<i>flavidus</i>	239, 240	<i>Remilegia</i>	449
<i>Aulostoma</i>	244	<i>Teuthis</i>	362
<i>cinereum</i>	245	<i>australis</i> , Icelus.....	379
<i>coloratum</i>	245	<i>austrina</i> , <i>Myxostoma</i>	110
<i>Aulostoma macleayi</i>	245	<i>austrinum</i> , <i>Moxostoma</i>	110
<i>Aulostomus</i>	244	<i>Auxis</i>	259
<i>Aulostomidae</i>	244	<i>rochei</i>	259
<i>Aulostomus</i>	244	<i>thazard</i>	259
<i>chinensis</i>	244	<i>avalonis</i> , <i>Antennaria</i>	506
<i>cinereus</i>	245	<i>Averuncus</i>	395
<i>maculatum</i>	245	<i>emmelane</i>	395
<i>aurantiacus</i> , <i>Balistes</i>	284, 495	<i>sterletus</i>	395
<i>Cottogaster</i>	284, 495	<i>averruncus</i> , <i>Kathetostoma</i>	454
<i>auratus</i> , <i>Chelio</i>	426	<i>avitus</i> , <i>Chologaster</i>	194
<i>auriga</i> , <i>Monacanthus</i>	494	<i>avocetta</i> , <i>Nemichthys</i>	83
<i>aurantiacus</i> , <i>Hypohomus</i>	285	<i>Avocetina</i>	83
<i>aurata</i> , <i>Coryphaena</i>	265	<i>elongata</i>	83
<i>Monana</i>	132	<i>gila</i>	83
<i>Salema</i>	339	<i>infans</i>	83
<i>auratum</i> , <i>Pristipoma</i>	333	<i>Awaous</i>	442
<i>auratus</i> , <i>Diapterus</i>	342	<i>nelsoni</i>	442
<i>Gadus</i>	210	<i>axillaris</i> , <i>Pristipoma</i> (<i>Hemulopsis</i>).....	333
<i>Holocentrus</i>	309	<i>axillaris</i> , <i>Boreocottus</i>	387
<i>Mullus</i>	343	<i>Brachydeuterus</i>	333
<i>barbatus</i>	343	<i>Diapterus</i>	342
<i>Notemigonus</i>	115	<i>Gerres</i>	342
<i>Pseus</i>	268	<i>Myoxocephalus</i>	387
<i>Scomber</i>	258	<i>axinophrys</i> , <i>Xystes</i>	395
<i>aurea</i> , <i>Loricaria</i>	162	<i>Axyris</i>	377
<i>aureolum</i> , <i>Moxostoma</i>	110	<i>barringtoni</i>	377
<i>aureolus</i> , <i>Catostomus</i>	109, 110	<i>aya</i> , <i>Chaetodon</i>	360
<i>Diapterus</i>	342	<i>Lutianus</i>	326
<i>Gerres</i>	342	<i>Neomans</i>	453
<i>Salvelinus</i>	60	<i>Aylopon</i>	322
<i>Xenotus</i>	300	<i>martineensis</i>	320
<i>aureo-ruber</i> , <i>Sparus</i>	432	<i>ayresi</i> , <i>Acipenser</i>	34
<i>aureoviridis</i> , <i>Sphyraena</i>	305	<i>Centropomus</i>	318
<i>aureus</i> , <i>Ammocetes</i>	10	<i>Lepidosteus</i>	37
<i>Carangus</i>	272	<i>Parophrys</i>	225
<i>Caranx</i>	270	<i>Ayresia</i>	413
<i>Chaetodon</i>	360	<i>punctipinnis</i>	413
<i>Fundulus</i>	177	<i>ayresi</i> , <i>Lampetra</i>	10
<i>Heros</i>	422	<i>Petromyzon</i>	10
<i>Sparus</i>	302	<i>Sebastodes</i>	367
<i>Thorichthys</i>	422	<i>Sebastomus</i>	367
<i>auriculata</i> , <i>Auctospina</i>	368	<i>Azevia</i>	221
<i>Mobula</i>	31	<i>panamensis</i>	221
<i>auriculatus</i> , <i>Sebastes</i>	368	<i>querna</i>	221
<i>aurifrons</i> , <i>Gnathypops</i>	453	<i>azorica</i> , <i>Coryphaena</i>	265
<i>auriga</i> , <i>Chaetodon</i>	359	<i>Azteca</i>	143
<i>Dules</i>	330	<i>Aztecula</i>	143
<i>aurita</i> , <i>Sardinella</i>	43	<i>lermae</i>	143
<i>Selema</i>	339	<i>mexicana</i>	143
<i>auritus</i> , <i>Brachydeuterus</i>	333	<i>vittata</i>	143
<i>Labrus</i>	299	<i>aztecus</i> , <i>Notropis</i>	143
<i>Lepomis</i>	299	<i>azurea</i> , <i>Hermosilla</i>	340
<i>Macrodon</i>	99	<i>azureus</i> , <i>Alepisaurus</i>	171
<i>aurofrenatum</i> , <i>Sparisoma</i>	432	<i>Carcharias</i>	16
<i>aurofrenatus</i> , <i>Scarus</i>	432	<i>Galeichthys</i>	180
<i>auroguttatus</i> , <i>Zygonectes</i>	177	<i>Azurina</i>	412
<i>aurolineatum</i> , <i>Myetophum</i>	170	<i>hirundo</i>	412
<i>aurolineatum</i> , <i>Bathystoma</i>	331	<i>azurissimus</i> , <i>Microspathodon</i>	416
<i>Hemulon</i>	331		
<i>auro-punctatus</i> , <i>Callyodon</i>	431	<i>babeocki</i> , <i>Rosicola</i>	365
<i>auro-punctatus</i> , <i>Cryptotomus</i>	431	<i>Sebastodes</i>	365
<i>aurora</i> , <i>Abeona</i>	409	<i>bacalau</i> , <i>Gobius</i>	441
<i>Amphigonopterus</i>	409	<i>badius</i> , <i>Euctenogobius</i>	441, 442
<i>Caprophonus</i>	358	<i>Fundulus heteroclitus</i>	176
<i>Catostomus</i>	106	<i>Rhinichthys</i>	140
<i>Eosebastes</i>	366	<i>Bagre</i>	147
<i>Fario</i>	55	<i>bahiensis</i>	148
<i>Sebastichthys</i>	366	<i>eydouxii</i>	148
<i>aurorubens</i> , <i>Centropristis</i>	327	<i>isthmensis</i>	148
<i>Rhomboplites</i>	328	<i>marina</i>	148
<i>aurovittatus</i> , <i>Mesoprius</i>	327	<i>panamensis</i>	148
<i>Ansonia</i>	264	<i>pinnulmaculatus</i>	148
<i>cuvieri</i>	264	<i>scutatus</i>	148
<i>australasicus</i> , <i>Pneumatophorus</i>	253	<i>secunda</i> , etc.....	153
<i>Scomber</i>	253	<i>bagre</i> , <i>Silurus</i>	148
<i>australe</i> , <i>Etheostoma</i>	293	<i>Bagrus proops</i>	148
<i>Zophodon</i>	117	<i>sundiacus</i>	149
<i>australis</i> , <i>Amphiprion</i>	308	<i>(Sciaen)</i> <i>emphysetus</i>	148
<i>Clinocottus</i>	390	<i>bahamensis</i> , <i>Pisces viridis</i>	432
<i>analis</i>	390	<i>Unicornis piscis</i>	495
<i>Echeneis</i>	449		

List of scientific names—Continued

	Page		Page
bahamensis, Vulpes.....	39	Balistes kleini.....	496
bahianus, Acanthurus.....	363	laevis.....	495
Rhombus.....	217	liberensis.....	492
bahiensis, Bagre.....	148	lineopunctatus.....	493
Cypselurus.....	202	linguistula.....	496
Engraulis.....	40	longus.....	493
Exocoetus.....	202	macrops.....	492
Galeichthys.....	148	macropterus.....	449
baileyi, Cyprinodon.....	182	maculatus.....	493
Cyprinodon maculatus.....	182	melanopterus.....	493
bailloni, Caesiomorus.....	276	mento.....	493
Trachinotus.....	276	monoceros.....	495, 496
Baillonus.....	276	monoceros unicolor.....	496
bailloui, Gasterosteus.....	237	moribundus.....	492
Baiodon.....	417	naufagiun.....	491
fasciatus.....	417	nigra.....	493
Baione.....	60	nitidus.....	493
fontinalis.....	60	notatus.....	493
Balostoma.....	223	piceus.....	493
brachialis.....	230	polylepis.....	492
bairdi, Acipenser.....	34	powelli.....	492
Calionymus.....	4,0	punctatus.....	492
Cottus.....	383	radula.....	493
Macourus.....	205	ringens.....	493
bairdiana, Sphyrænop.....	297	rutus.....	493
bairdianus, Sphyrænop.....	297	sandwichensis.....	493, 494
Syngnathus.....	242	sandwichensis.....	494
Syrtetes.....	242	scapus.....	494
Bairdiella.....	345	schoepfi.....	495
armata.....	346	scolopax.....	246
bedoti.....	343	scripta.....	495
chrysura.....	345	scriptus.....	495
ensifera.....	345	sobaco.....	492
icistia.....	345	spilopterygius.....	492
ronchus.....	346	sufflamen.....	492
veracutus.....	346	taeniopterus.....	491
bairdi, Alepocephalus.....	52	triangularis.....	492
Bathymyzon.....	8	unicorn.....	496
Cottus.....	385	veres.....	492
Gasterostomus.....	95	venula.....	491
Mitchillina.....	52	willungben.....	493
Petromyzon (Bathymyzon).....	8	Balsadichthys.....	491
Pomacentrus.....	415	whitell.....	183
Salmo.....	61	balsana, Thyridina.....	183
Stegastes.....	415	balsanus, Istalarius.....	249
Bajacalifornia.....	53	Melanis.....	248
burragei.....	53	balsas, Poecilops.....	248
Bajonado.....	336	balteatus, Cyprinus (Abiamis).....	118
balonado, Calamus.....	336	Eques.....	353
Sparus.....	336	Heros.....	418
balantioptthalmus, Scomber.....	271	Pomacanthus.....	360
balao, Hemiramphus.....	199	Richardsomus.....	118
balboa, Amplova.....	50	Theraps.....	418
Fundulus.....	179	Thynnus.....	260
Oxyzygonectes.....	179	Upeneus.....	343
baldwini, Prionodes.....	319	banana, Butyrinus.....	39, 40
balearica, Muræna.....	79	Chonophorus.....	442
balearicum, Ophisoma.....	80	Gobius.....	442
balearus, Cyprinus (Abiamis).....	117	bancrofti, Torpedo.....	27
balias, Chirus.....	374	Barathrodemus.....	482
Balistes.....	491	manatinus.....	482
americanus.....	493	Barathronus.....	484
angulosus.....	492	bicolor.....	484
asperimus.....	492	barbara, Arolepis.....	444
aurantiacus.....	495	Siphostoma.....	240
barbatus.....	496	Syngnathus.....	240
bellus.....	491	barbata, Brotula.....	479
bridé.....	492	Lonchirus.....	352
broccus.....	494	Loricaria.....	161, 162
buniva.....	491, 493	Pallasina.....	394
capistratus.....	492	Rhamdia.....	158
caprinus.....	491	barbatula, Læmonema.....	213
caprisus.....	491	barbatulum, Læmonema.....	213
carolinensis.....	491	barbatum, Echiostoma.....	70
chinensis.....	494	Læmonema.....	476
electricus.....	493	Ophidion.....	485
ciliaris.....	492	barbatus, Balistes.....	496
ciliatus.....	494	Chautilodus.....	71
curassavicus.....	493	Enchelyopus.....	479
equestris.....	491	barbatus, Mullus.....	343
forcipatus.....	492	Siphagonus.....	394
fuliginosus.....	491	barbatus auratus, Mullus.....	343
guttatus.....	492	barbiger, Macrourus.....	205
heckell.....	493	Macrurus.....	205
ispidus.....	494		

List of scientific names—Continued

	Page		Page
barbiger, Peristedion	406	Bathylagus milleri	89
Peristedium	406	pacificus	89
barbuda, Laja	496	Bathymaster	451
Barbulifer	446	caeruleofasciatus	451
cauthœus	446	hypoplectus	452
papillosus	446	jordanii	452
barbulifer, Rhinoliparis	405	signatus	451
bardus, Minomus	104	bathymaster, Bregmaceros	208
barkani, Agonus	394	Bathymasteridae	451
barnharti, Achirus	230	Bathymyzon	8
baronis-mulleri, Pimelodus	157	bairdii	8
Rhamdia	157	(Bathymyzon) bairdii, Petromyzon	8
barracuda, Esox	255	Bathynectes	480
Sphyræna	255	compressus	480, 481
barracudina, Paralepis	165	gracilis	481
barrati, Etheostoma	294	laticeps	483
Hololepis	294	Bathyonus	480
barreto, Gobioides	448	catena	481
bartholomæi, Caranx	272	pectoralis	482
Elaphotoxon	272	tenia	481
bartoni, Acara	419	Bathyophis	70
Aspidophoroides	397	ferox	70
Chrostoma	250	Bathypasma	401
Cylindrostæus	37	ovigerum	401
Parapetena	419	bathypbila, Cyclothone	72
Bascanichthys	86	bathypbilum, Neostoma	72
bascanium	86	Bathypteroidæ	106
bascanoides	86	Bathypterois	106
cylindricus	86	longifilis	106
panamensis	86	pectoralis	106
peninsulæ	86	quadrifilis	106
scuticaris	86	ventralis	106
bascanium, Bascanichthys	86	Bathysaurus	104
Cœcula	86	agassizii	104
Bascanius	230	ferox	104
tædifer	230	mollis	104
bascanoides, Bascanichthys	86	Bathystoma	331
basilaris, Astatheros	421	aurolineatum	331
Heros	421	rimator	329, 331
Bassogigas	482	striatum	331
gilli	482	Bathystomus	331
stelliferoides	482	Bathytroctes	52
Bassozetus	480	alveatus	52
catena	480, 481	alvifrons	52
compressus	481	homopterus	53
nasus	481	inspector	52
normalis	480, 481	microlepis	52
tenia	481	stomias	52
batabana, Corvula	345	Batis	24
batabanus, Johnius	345	radula	24
bathœctor, Salmo	57	batis, Raja	24, 25
Bathyaonius	395	Batoidei	23
nigripinnis	395, 396	Batrachoides	486
Bathyalopex	33	boulengeri	487
abbreviatus	33	goldmani	487
plumbeus	33	pacifici	487
bathybii, Paraliparis	404	surinamensis	486
bathybius, Cynicoglossus	228	tau	486
Embassichthys	228	variegata	487
Histiobranchus	78	vernulus	487
Synaphobranchus	78	Batrachoididæ	486
Bathylupea	281	Batrachus	487
argentea	281	celatus	487
hoskynii	281	margaritatus	487
Bathylupeidæ	281	pacifici	487
Bathylagus	203	porosissimus	487, 488
arcuatus	202	surinamensis	486
cavernosus	207	tau pardus	487
cottoides	203	Batrictius	486
favosus	203	battaræ, Orthragoriscus	508
longifilis	202	bayamonensis, Gobiionellus	441
macrops	203	Gobius	441
melanobranchus	203	bayanus, Pomadasys	334
Bathygobius	440	Pomadasys	334
fuscus	440	bdellium, Petromyzon	9
soporator	440	Bdellostoma stoutii	8
Bathylaco	164	beadlei, Synchoglanis	151, 152
nigricans	164	beani, Beloncharax	100
Bathylagidæ	68	Caranx	272
Bathylagus	68	Ctenolucius	100
antarcticus	68	Heros	419
benedicti	69	Kirtlandia	248
borealis	69	Melampheus	233
euryops	69	Membras	248

List of scientific names—Continued

	Page		Page
beani, Ophidion.....	485	beringi, Crossias.....	389
Parapetenia.....	419	beringiana, Stelgastrops.....	379
Plectromus.....	233	beringianum, Stelgistrum.....	379
Poecilichthys.....	287	beringianus, Cyclogaster (Neohpans).....	399
Triglops.....	380	Neohpans.....	399
beanni, Ammocrypta.....	589	berlandieri, Lepidosteus (Atractosteus).....	38
Limanda.....	227	Mugil.....	253
Prionotus.....	408	bermude, Branchiostoma.....	7
Serrivomer.....	84	Chromis.....	413
beanorum, Pontinus.....	372	Fundulus.....	175
beardslei, Salmo.....	57	heteroclitus.....	175
Salmo gairdneri.....	57	Furcaria.....	413
beckwithi, Cyprinella.....	130	bermударum, Callionymus.....	450
becuna, Sphyræna.....	255	bermudensis, Carapus.....	486
bedoti, Bairdiella.....	246	Lefroyia.....	481
bedotis, Sciæna (Bairdiella).....	246	Bermudichthys.....	430
beldingii, Cottus.....	384	subfurcatus.....	430
belis, Silurus.....	149	bernardini, Catostomus.....	101
belizanus, Belonesox.....	184	bertheloti, Crnus.....	268
belizianus, Eleotris (Culicus).....	436	Berycida.....	234
bella, Hypoclydonia.....	197	Berycoides.....	234
Bellator.....	408	beryllina, Menidia.....	247
egretta.....	408	beryllina cerea, Menidia.....	247
militaris.....	408	beryllinum, Christoma.....	247
bellicus, Nocomis.....	137	beryllinus, Cryptotomus.....	431
bellus, Balistes.....	491	Gobiosox (Rimicola).....	490
Lythrurus.....	126	Rimicola.....	490
Mimulus.....	126	Beryx.....	234
Belone almeida.....	195	decadactylus.....	234
altipinna.....	197	splendens.....	234
angusticeps.....	196	Betaraia.....	25
ardeola.....	195	betaurus, Cirrhitus.....	358
aigatus.....	196	biaculeatus, Gasterosteus.....	237
caribbea.....	197	Bibronia.....	230
cigonella.....	195	igulata.....	230
ciassa.....	196	bicaudalis, Chapinus.....	497
depressa.....	195	Ostracion.....	496, 497
diplotænia.....	196	bicolor, Algansea.....	122
exilis.....	196	Ammocoetes.....	8
galeata.....	197	Anisotremus.....	332
gerania.....	196	Barathronus.....	484
hians.....	197	Chlopsis.....	82
jonesi.....	197	Exocoetus.....	201
latimana.....	197	Grammocoetus.....	197
maculata.....	197	Pristipoma.....	332
melanochira.....	196	Rondeletia.....	171
microps.....	196	Rypticus.....	316
notata.....	195	Siphateles.....	122
pacifica.....	197	Smeeticus.....	316
raphidoma.....	196	Tigoma.....	120
scutator.....	196	bicornis, Cottus.....	378
stolzmanni.....	196	Icelus.....	378
strongylura.....	195	bifasciatum, Thalassoma.....	429
subtruncata.....	195	bifasciatus, Cichlauius.....	416
tinnu.....	195	Heros.....	416
truncata.....	196	Labrus.....	428, 429
Belonesox.....	184	Mullus.....	343
belizanus.....	184	Upeneus.....	343
Belonidae.....	195	bifasciatus torquatus, Labrus.....	429
Belonichthys beani.....	100	bifrenata, Hybopsis.....	135
Belonopsis.....	83	bifrenatus, Hybopsis.....	135
leuchtenbergii.....	83	bigibbus, Kyphosus.....	339
bellus, Mimulus.....	126	biguttatus, Coelognathus.....	146
bendirei, Cottus.....	384	Labrisomus.....	460
Potomacottus.....	384	Malacoctenus.....	460
benedicti, Bathylagus.....	69	Semotilus.....	137
bengalensis, Holocentrus.....	324	bilinearis, Merluccius.....	214
benotti, Myctophum.....	169	Stomodon.....	214
Scopelus.....	169	bilineata, Lepidopsetta.....	226
Benthocometes.....	482	Platessa.....	226
robustus.....	482	Skiffia.....	183
Benthodesmus.....	282	Umbra pygmaea.....	174
atlanticus.....	282	bilineatum, Pristipoma.....	332
elongatus.....	282	bilineatus, Characodon.....	183
Benthosauridae.....	166	Doryichthys.....	243
Benthosaurus.....	166	billingsiana, Cyprinella.....	131
grallator.....	166	biloba, Corvina.....	349
Benthosema.....	170	bilobus, Blepasas.....	382
arcticum.....	170	Histiocottus.....	382
mulleri.....	170	bimaculata, Percina.....	281
berardi, Aluteres.....	496	bimaculatus, Chaetodon.....	380
bergall, Ctenolabrus.....	424	Clinus.....	460
bergii, Pontinus.....	372	Halicheres.....	426
Scorpaena.....	372	Malacoctenus.....	460
bergii, Macrourus.....	205	Poecilodes.....	187
		Pseudoxiphophorus.....	187

List of scientific names—Continued

	Page		Page
bimaculatus Sayris.....	197	Blennius gunnellus.....	471
Xiphophorus.....	187	hantz.....	462
bimaculatus jonesi, Pseudoxiphophorus.....	188	hermimer.....	469
bimaculatus tæniatus, Pseudoxiphophorus.....	188	lævis.....	461
binghami, Xyrichtys.....	430	lampetraformis.....	470, 471
binoculata, Raja.....	26	lumpenus.....	470
binotata, Amia.....	295	marmoreus.....	461
binotatus, Apogon.....	295	microstomus.....	461
Bipinnula.....	260	multifilis.....	461, 462
bipinnulata, Seriola.....	279	murænoides.....	467
bipinnulatus, Elagatis.....	279	nuchifilus.....	461
birgei, Leucichthys.....	64	oceanicus.....	461
birostratus, Gurnardus.....	406	ocellaris.....	460
Prionotus.....	406	pholis.....	461
birostris, Manta.....	32	pilicornis.....	461
Raja.....	32	polaris.....	478
biserialis, Ophichthys.....	88	polyactocephalus.....	466
Opichthys.....	88	punctatus.....	462, 469
biseriatus, Archaulus.....	377	regius.....	213
bishopi, Aloa.....	43	saliens.....	463
bison, Aspicottus.....	381	serpentinus.....	470, 471
Carpiodes.....	103	stearnsi.....	460
Lepisosteus.....	37	striatus.....	462
bispinosus, Gasterosteus.....	238	tænia.....	468
Gladilunculus.....	238	truncatus.....	461
MelamphaPs.....	233	vinctus.....	461
Melichthys.....	493	blennius, Alburnops.....	133
Mylobatis.....	31	Etheostoma (Rhothoea).....	287
Plectromus.....	233	Hybopsis.....	133
bisselli, Coregonus tullibee.....	63	Ulocentra.....	287
Leucichthys artedi.....	63	Blennophis.....	464
bis-trispinus, Bodianus.....	316	webbii.....	464
bistrispinus, Bodianus.....	316	Bleparichthys.....	274
Promicropterus.....	316	Blepharis.....	274
bisus, Scomber.....	259	major.....	274
bitematus, Eumycterias.....	500	sutor.....	274
bivittata, Elacate.....	281	blepharis, Carangoides.....	274
Iridio.....	423	blephura, Anguilla.....	77
bivittatus, Hybopsis.....	119	Blepsias.....	382
Labrus.....	426	bilobus.....	382
Serranus.....	318	cirrhosus.....	382
blackfordi, Lutianus.....	326	oculofasciatus.....	382
Lutjanus.....	326	trilobus.....	382
Neomænis.....	453	ventricosus.....	380, 381
Yarrella.....	72	Blepsidæ.....	382
blackfordi, Lutianus.....	326	blochii, Bodianus.....	425
Lutjanus.....	326	Caranx.....	271
Blakea.....	458	Orthragoriscus.....	503
blanchardi, Gasterostea.....	239	Piramutana.....	159
Neoclinus.....	458	boa, Esor.....	69
bleekeriana, Chirocentron.....	45	bocourti, Herichthys.....	423
Pellona.....	45	Neetroplus.....	423
Bleekeridæ.....	461	Bodianus.....	309, 425
Blenitrachus.....	462	acanthistius.....	309
Blennicottus.....	391	achigan.....	298
clarki.....	377	albostratus.....	325
globiceps.....	391	apua.....	313
bryosus.....	391	argyroleucos.....	345
recalvus.....	391	argyro-leucos et exiguus.....	345
Blennidæ.....	460	bistrispinus.....	316
blennioides, Etheostoma.....	283, 286	bis-trispinus.....	316
Blenniolus.....	461	blochi.....	425
blenniopeca, Hyostoma.....	286	bodianus.....	309, 425
Blennius.....	460	costatus.....	349
americanus.....	473	diplotænia.....	425
angularis.....	470, 473	fasciatus.....	325
asterias.....	461	guativeri.....	309
asterius.....	461	guttatus.....	309
bosquianus.....	462	jaguar.....	235
brevipinnis.....	461	marginatus.....	313
chisso.....	212	pallidus.....	345
chuss.....	213	pentacanthus.....	225
ciliatus.....	473	pulchellus.....	425
crintus.....	461	rufus.....	425
cristatus.....	461	rupestris.....	303
dolichogaster.....	468	stellatus.....	309
europæus.....	467	stellifer.....	346
fabbrii.....	460	striatus.....	325
favosus.....	461	triourus.....	323
fimbriatus.....	473	vivanet.....	325
fuorum.....	461	bodianus, Bodianus.....	309, 425
galerita.....	460	bœkei, Callionymus.....	450
geminatus.....	461, 462	Doratonotus.....	429
gentilis.....	462	Melanorhinus.....	249
gracilis.....	471	Moringua.....	89

List of scientific names—Continued

	Page		Page
Bogmarus.....	216	borealis, Notorhynchus.....	11
islandicus.....	216	Oligocottus.....	390
bogmarus, Trachypterus.....	216	Opsopœodus.....	115
Bogoslovius.....	204	Paralepis.....	165
clarki.....	204	Petromyzon.....	10
firmisquamis.....	204	Pimelodus.....	152
Boleichthys.....	293	Pœcilichthys.....	291
eos.....	294	Sphyræna.....	255
exilis.....	293, 294	Squalus.....	21, 22
fusiformis.....	294	Boreocottus.....	387
gracilis.....	294	axillarus.....	387
warreni.....	294	Boreogadus.....	209
whipphi.....	292	saida.....	209
boleoides, Cottus.....	385	Boreogaleus.....	14
Radulinus.....	379	boreus, Esox.....	173
Boleosoma.....	287, 288	Borophryne.....	509
ætopus.....	287	apogon.....	509
camurum.....	288	Borborys.....	175
effulgens.....	287	boschna, Pomadasys.....	334
fusiforme.....	294	Pristipoma.....	334
gracile.....	294	bosci, Abramis.....	350
lepidæ.....	290	Gobiosoma.....	446
maculiceps.....	287	Gobius.....	442, 445, 446
maculatum.....	287	Notemigonus.....	115
messum.....	287	Pimelepturus.....	339
nigrum.....	287	sicula, Pimelepturus.....	340
olimstedii.....	287	boscii, Atherina.....	247
brevipinnis.....	287	Gobius.....	442
phlox.....	287	Leuciscus.....	115
stigmaeum.....	287	Seriola.....	280
susane.....	288	bosquianus, Blennius.....	462
tassellatum.....	287	Chasmodes.....	462
tessellatum.....	285, 287	bosquianus, Chasmodes.....	460
vexillare.....	288	bosqui, Pimelepturus.....	340
boleosoma, Gobius.....	439	bostoniensis, Catostomus.....	105
Rhinocephalus.....	439	Muraena.....	77
bolli, Ameiurus.....	153	Bothidae.....	217
bolliani, Hippoglossina.....	223	Bothragonus.....	396
Opsopœa.....	115	swani.....	396
Opsopœodus.....	115	Bothrocara.....	477
Scarus.....	433	mollis.....	477
Bollmannia.....	443	pusilla.....	477
boqueronensis.....	443	remigera.....	477
chlamydes.....	443	Bothrocaropsis.....	477
macropoma.....	443	alalonga.....	477
ocellata.....	443	elongata.....	477
stigmatura.....	443	rictolata.....	477
bombifrons, Heloperca.....	301	Bothrolæmus.....	276
Pomotis.....	301	Bothus.....	217
Bonaci cardinal.....	313	boucardi, Leuciscus.....	132
bonaci, Serranus.....	314	Montana.....	132
Trisotropis.....	314	Pœcilia.....	192
xanthosticta, Mycteroperca.....	314	Pristipoma.....	334
bonapartei, Lampanyctus.....	167	bougainvillei, Sebastes.....	370
Pœcilocephalus.....	87	boulengeri, Batrachoides.....	487
Bonapartia.....	72	Mycteroperca.....	313
pedalota.....	72	Plecotomus.....	160
bonariense, Hæmulon.....	329	Trisotropis.....	313
bonariensis, Seriola.....	280	bouvieri, Salmo.....	56
bonasus, Bubalichthys.....	102	Salmo purpuratus.....	56
Rhinoptera.....	31	bovinum, Electropoma.....	321
bonita, Gambusia.....	188	bovinus, Cyprinodon.....	181
Priapella.....	188	Hypoclinemus unicolor.....	321
Boops.....	417	rubroflaviflatus, Cyprinodon.....	181
boops, Caranx.....	272	bowersi, Labichthys.....	84
Hybopsis.....	134	Mycteroperca.....	314
Macrourus.....	205	Rhynchichthys.....	328
Macrurus.....	205	Bowersia.....	328
Myctophum.....	160	virescens.....	328
Notropis.....	134	bowersianus, Careproctus.....	403
Ostracion.....	503	bowmani, Plagyrrus.....	128
Scomber.....	270	Box.....	417
Trachurus.....	272	boyerj, Atherina.....	246
boqueronensis, Bollmannia.....	443	brachiale, Sparisoma.....	433
borea, Luciopeca.....	282	brachialis, Balistoma.....	230
borealis, Alepidosaurus.....	171	Scarus.....	433
Ammocetes.....	9	brachiatus, Diodon.....	501
Arctozenus.....	165	Brachioptilon.....	32
Bathylagus.....	69	hamiltoni.....	32
Canlopius.....	171	brachiusculus, Grammicolepis.....	232
Fierasfer.....	471	brachyacanthus, Ameiurus.....	154
Icelinus.....	376	Amiurus.....	154
Lampetra.....	10	brachycentrus, Nauderus.....	273
Molvella.....	208	brachycephala, Rhamdia.....	157
Nivicola.....	291		

List of scientific names—Continued

	Page		Page
brachycephalus, Exocoetus.....	200	Brachyrhamphichthys.....	101
Pimelodus.....	157	Brachyrhaphis.....	186
Syngnathus.....	241	cascajalensis.....	187
Uranichthys.....	87	episcopi.....	187
Bramidae.....	265	olomina.....	187
Bramiformes.....	265	parismina.....	187
Bramocharax.....	99	rhabdophora.....	186
bransfordi.....	99	terrabensis.....	186
elongatus.....	99	Brachyrhinus.....	320
Bramopsis.....	410	brachyrhynchus, Acipenser.....	34
mento.....	410	Rhynchichthys.....	235
branchialis, Ammocetus.....	8	brachysoma, Epinephelus.....	311
branchiomelas, Anchoviella.....	49	Brachysomophis.....	89
Stolephorus.....	49	crocodilinus.....	89
Branchiostegidae.....	357	horridus.....	89
Branchiostoma.....	7	brachysomus, Calamus.....	337
bermudae.....	7	Sparus.....	337
californiense.....	7	brachyurus, Oxydontichthys.....	88
caribaeum.....	7	bradfordi, Porocottus.....	388
floridae.....	7	Brama.....	265
lanceolatum.....	7	agassizii.....	265
lubricum.....	7	brevoorti.....	265
pelagicum.....	7	chilensis.....	265
platea.....	7	dussumieri.....	265
virginiae.....	7	marina cauda forcipata.....	265
Branchiostomidae.....	7	oreini.....	265
Branchistegus.....	357	parrae.....	426
Branchyconger.....	78	rai.....	265
brandieriana, Cœcilia.....	84	saussurii.....	265
Branderius.....	84	brama, Cynædus.....	338
brandti, Arius.....	148	brasiliensis, Torpedo.....	27
branicki, Pristipoma.....	334	Vomer.....	275
Rhoniciscus.....	334	brasiliensis corallina, Narcine.....	27
Brannerella.....	464	braytoni, Notropis.....	135
brasiliensis.....	464	Bregmaceros.....	208
sluteri.....	464	atlanticus.....	208
bransfordi, Bramocharax.....	99	bathymaster.....	208
Loricaria.....	162	longipes.....	208
Rhamdia.....	157	maclellandii.....	208
brasilianus, Eugerres.....	342	Bregmaceroida.....	208
Gerres.....	332	breve, Chirostoma.....	250
brasiliense, Pristipoma.....	332	brevibarbe, Lepophidium.....	484
brasiliensis, Capeuna.....	331	Ophidium.....	484
Guabicoara.....	330	brevibarbis, Chalinura.....	204
brasiliensis, Brannerella.....	464	brevicauda, Pomoxys.....	304
Centropristis.....	320	Salmo.....	56
Chupea.....	40	brevicaudata, Brevoortia tyrannus.....	44
Conger.....	78	brevicaudum, Siphostoma.....	241
Erythrinus.....	99	brevicaudus, Diaphanichthys.....	79
Esoc.....	199	Syngnathus.....	241
Gobius.....	448	Breviceps.....	147
Hemiramphus.....	199	filamentosus.....	147
Hippoglossus.....	224	breviceps, Evorthodus.....	438
Hypostomus.....	160	Gasterosteus.....	239
Labrus.....	426	Gobius.....	440
Mugil.....	253	Larimus.....	344
Narcine.....	27	Moxostoma.....	111
Paralichthys.....	224	Pomacentrus.....	413
Plagusia.....	231	Pomotis.....	300
Scorpena.....	371	Porogadus.....	482
Thynnus.....	258	Ptychostomus.....	111
brachychira, Trachurops.....	271	bravifilis, Agenesiosus.....	159
brachychirus, Trachurops.....	271	brevimanus, Eugerres.....	343
Brachydenterus.....	333	Gerres.....	343
auritus.....	333	Tetragonopterus.....	96
axillaris.....	333	brevipes, Lycodes.....	475
corvineformis.....	333	brevipinna, Somniosus.....	21, 22
leuciscus.....	333	brevipinne, Ditrema.....	409
nitidus.....	333	Pristipoma.....	335
Brachygenys.....	330	(Hæmulopsis).....	335
chrysargyreus.....	331	brevipinnis, Bleennius.....	461
Brachyistius.....	409	Microlepidotus.....	335
frenatus.....	409	Thynnus.....	258
Brachymullus.....	343	brevirostre, Peristedion.....	405
Brachyopsis verrucosus.....	394	Peristethus.....	405
xyosternus.....	394	brevirostris, Acipenser.....	34
brachypoda, Gasterosteus pungitius.....	239	Chasmistes.....	108
Pungitius.....	239	Cololabis.....	198
Brachyprotopon.....	228	Engraulis.....	51
brachyptera, Echenis.....	449	Gerres.....	342
Remoropsis.....	449	Hypopomus.....	101
brachypterus, Holocentrus.....	236	Hypoprion.....	17
Lutjanus.....	326	Macrogathus.....	199
Lutjanus.....	326	Rhamphichthys.....	101
Pimelodus.....	157	Rhynchoerarias.....	511
Zygonectes (Micriscus).....	185	Saurus.....	163

List of scientific names—Continued

	Page		Page
brevirostris, Scombrox.....	198	Brycon.....	98
Syngnathus.....	240	argenteus.....	98
brevirostrum, Acipenser.....	34	cephalus.....	98
Anchovia.....	51	dentex.....	98
Hemulon.....	330	falcatus.....	98
Stolephorus.....	50	guatemalensis.....	98
brevis, Acanthurus.....	362	oligolepis.....	99
Amblyopus.....	448	petrosus.....	98
Atherinichthys.....	250	striatulus.....	98
Campostoma.....	146	Bryconamericus.....	97
Centropomus.....	306	cascajalensis.....	97
Dasyatis.....	29	emperador.....	97
Engraulis.....	51	exodon.....	97
Fundulus parvipinnis.....	176	Bryolophus.....	466
Tytilastes.....	448	lysium.....	466
Xiphophorus.....	194	bryoporus, Spratelloides.....	41
helleri.....	194	Bryostemma.....	466
brevispina, Richia.....	293	decoratum.....	466
Richia.....	293	nugator.....	466
brevispine, Acutomentum.....	366	polyactocephalum.....	466
brevispinus, Sebastichthys proriger var.....	366	tarsodes.....	466
brevvoorti, Argyreus.....	276	Brysetæres.....	488
Argyrosus.....	276	pinniger.....	488
Brama.....	265	Bryssophilus.....	488
Euleptorhamphus.....	198	Bryttus.....	298
Poecilurichthys.....	96	albulus.....	302
brevvoorti, Poecilurichthys.....	97	fasciatus.....	304
Brevvoorti.....	44	gloriosus.....	304
pakonus.....	44	humilis.....	300
tyranus.....	44	mineopas.....	299
brevicaudata.....	44	oculatus.....	300
brevvoorti, Astyanax.....	97	punctatus.....	298, 299
Euleptorhamphus.....	198	reticulatus.....	299
briareos, Chilomycteris.....	502	signifer.....	299
Cylichthys.....	502	unicolor.....	299
brices, Chastodon.....	359	Bunaleichthys.....	102
brimleyi, Coccoth.....	127	altus.....	102
Notropis.....	127	bonasus.....	102
bristolensis, Cyclogaster.....	399	bubalinus.....	102
Neoliparis.....	399	niger.....	102
broccus, Balistes.....	494	urus.....	102
brodamus, Cottus.....	395	bubalina, Cyprinella.....	130
Brontes.....	159	bubalinus, Bubaleichthys.....	102
prenadilla.....	213	Leuciscus.....	130
Brosme.....	213	bubalis, Acanthocottus.....	386
brosme.....	214	Ambodon.....	102
brosme, Brosme.....	214	Cottus.....	388
Gadus.....	213, 214	Ichthyobus.....	102
Brosmerus.....	213	Ictobus.....	102
flavescens.....	213, 214	buccanella, Lutjanus.....	326
brosmitana, Lota.....	212	Mesopion.....	326
Brosminus.....	213	buccata, Ericymba.....	141, 142
marginatus.....	479	bucciferus, Labrisomus.....	450
Brosomphycus.....	479	Labrosomus.....	459
marginatus.....	479	bucco, Moxostoma.....	110
ventralis.....	480	Ptychostomus.....	110
verrilli.....	480	Buccone.....	354
Brotula.....	479	prædatoria.....	354
barbata.....	479	bucculentus, Chonophorus.....	442
Brotulidæ.....	479	bucephalus, Macrourus.....	205
broussoneti, Acanthurus.....	362	Macurus.....	205
broussoneti, Gobioides.....	448	buchanani, Hybopsis.....	134
Umbrina.....	350	Notropis.....	134
browni, Lucania.....	181	buetikoferi, Tilapia.....	423
Solea.....	229	buffalo, Acipenser (Antaceus).....	85
brownii, Atherina.....	48	bufo, Lophus.....	487
Hemiramphus.....	199	Scorpena.....	370
Vomer.....	274, 275	bulbiceps, Macrourus.....	205
bruceus, Echinorhinus.....	22	Macurus.....	205
Squalus.....	22	Maynea.....	177
brunnea, Lycogramma.....	476	bullaris, Cyprinus.....	423
Maynea.....	476	bulleri, Paraneotroplus.....	423
brunneus, Amiurus.....	154	Frionodes.....	319
Apristurus.....	12	Serranus.....	319
Catulus.....	12	buniva, Balistes.....	491, 493
Ectenias.....	265	burchamii, Icelinus.....	376
Gobius.....	440	burgall, Ctenolabrus.....	424
Gymnothorax.....	92	burragei, Bajacalifornia.....	53
Hippocampus.....	244	burtonianus, Leuciscus.....	121
Ilyophis.....	78	buseki, Sicydium.....	447
Rivulus.....	180	butleri, Poecilia.....	192
Serranus.....	314	butlerianus, Poecilichthys.....	294

List of scientific names—Continued

	Page		Page
Butyrinus.....	39	californiensis, Diapterus.....	341
banana.....	39, 40	Coryrhamphus.....	243
Bythites.....	480	elongatus, Eucinostomus.....	341
fuscus.....	480	Eucinostomus.....	341
		Medialuna.....	358
caballa, Cybium.....	257	Ophisurus.....	88
Caballerote.....	325	Otolithus.....	356
caballerote, Anthias.....	325	Polynemus.....	256
caballus, Caranx.....	272	Scorpius.....	358
Paratractus.....	272	Syngnathus.....	240
caboverdianus, Ginglymostoma.....	12	Typhlogobius.....	447
cabreræ, Rhamdia.....	157	Xenichthys.....	324
Cabrilla.....	311	Xenistius.....	324
Cæcilia.....	84	callarias, Gadus.....	210
Cæcorhamdia.....	158	Salmo.....	61
urichi.....	158	Callaus.....	348
cærulea, Cheonda.....	120	deliciosa.....	348
Clupea.....	41	Callechelys.....	86
Coryphæna.....	434	guichenoti.....	86
Erogala.....	130	muræna.....	86
Etheostoma.....	291	peninsula.....	86
Meletta.....	43	Callichthyidae.....	160
Sardina.....	43	Callichthys (Holosternum) thoracatus.....	160
cæruleatus, Acronurus.....	362	lævigatus.....	160
cæruleoaurum, Harpe.....	425	littoratus.....	160
cæruleofasciatus, Bathymaster.....	451	Callidulus.....	320
cærulescens, Arius.....	150	flaviventris.....	320
Galectichthys.....	150	Calliodon retractus.....	431
cæruleus, Acanthurus.....	362	Callionymidae.....	450
Carcharias.....	16	Callionymus.....	450
Oligocephalus.....	289, 291	agassizi.....	450
Photogenis.....	130	atrilabiatus.....	450
Scarus.....	434	bairdi.....	450
Squalus.....	17	bermударum.....	450
Cæsiomorus.....	276	boekei.....	450
bailloni.....	276	callurus.....	450
cæsius, Paraconodon.....	331	himantophorus.....	450
Pomadasys.....	331	lyra.....	450
caguite, Sicydium.....	447	pauerradiatus.....	450
calablepta, Dermatocheir.....	97	pelagicus.....	409
Calamus.....	336	sancti-eustatii.....	450
arcifrons.....	337	callipteryx, Campostoma.....	146
bajonado.....	336	callisema, Episema.....	132
brachysomus.....	337	Moniana.....	132
calamus.....	336	callisoma, Herpetoichthys.....	88
kendalli.....	337	callistia, Erogala.....	130
leucosteus.....	337	callistius, Photogenis.....	130
macrops.....	337	calliura, Cyprinella.....	129
medius.....	337	Etheostoma.....	298
megacephalus.....	336	Mycteroperca.....	315
penna.....	337	Callurus.....	298
pennatula.....	336	diaphanus.....	299
plumatula.....	336	floridensis.....	303
proridens.....	336	formosus.....	299
calamus, Calamus.....	336	melanops.....	302, 303
Pagellus.....	336	microps.....	299
calcarata, Scorpæna.....	371	murinus.....	299
calcea, Acanthidium.....	21	punctulatus.....	298
caliente, Xenodermus.....	184	calliurus, Callionymus.....	450
calientis, Goodea.....	184	loglossus.....	438
Hybopsis.....	134	callolepis, Harengula.....	43
Notropis.....	134	Heros.....	422
californica, Alausa.....	43	Thorichthys.....	422
Maynea.....	477	callopterus, Cypselurus.....	202
Morrhua.....	210	Exocoetus.....	202
Squatina.....	22	Calloptilum.....	208
Tetranarce.....	27	mirum.....	208
Torpedo.....	27	callorhini, Therobromus.....	68
californicus, Aetobatus.....	31	Callorhynchus atlanticus.....	32
Cypselurus.....	202	centrina.....	32
Exocoetus.....	202	Calliodon.....	433, 434
Halicheres.....	428	auro-punctatus.....	431
Hippoglossus.....	223	compressus.....	434
Mustelus.....	13	microps.....	435
Myliobatis.....	31	ustus.....	431
Myliobatus.....	31	calliodon, Cyclopterus.....	399
Oxyjulis.....	428	Neoliparis.....	399
Paralichthys.....	223	calobrense, Cichlasoma.....	417
Stereolepis.....	308	calobrensis, Cichlasoma.....	417
californiense, Branchiostoma.....	7	calopteryx, Serranus.....	319
Myctophum.....	169	Calotomus.....	431
californiensis, Atherinopsis.....	252	xenodon.....	431
Chilomycterus.....	502		
Cyprinodon.....	181		

List of scientific names—Continued

	Page		Page
calva, <i>Amia</i>	38	canus, <i>Macrurus</i>	206
<i>Amiatus</i>	38	<i>Maerurus</i>	206
<i>Calycelepidotus</i>	381	<i>Capellaria</i>	505
<i>spinus</i>	381	<i>capellei</i> , <i>Lophotes</i>	215
<i>Camarina</i>	338	<i>capense</i> , <i>Scyllium</i>	11
<i>nigricans</i>	338, 339	<i>capensis</i> , <i>Carchaodon</i>	20
<i>camchaticus</i> , <i>Artediellus</i>	378	<i>capensis</i> , <i>Loricaria</i>	162
<i>camelopardalis</i> , <i>Mycteroperca</i>	315	<i>Capeuna brasiliensis</i>	331
<i>Serranus</i>	315	<i>capuana</i> , <i>Serranus</i>	331
<i>campechanus</i> , <i>Lutjanus</i>	326	<i>capillaris</i> , <i>Zeus</i>	275
<i>Mesoprion</i>	326	<i>capillatus</i> , <i>Chnus</i>	459
<i>campbelli</i> , <i>Moxostoma</i>	109	<i>capistratus</i> , <i>Balistes</i>	492
<i>Salmo</i>	61	<i>Chatodon</i>	359
<i>camperi</i> , <i>Scomberesox</i>	197	<i>Tetrodon</i>	500
<i>camperi</i> , <i>Scomberesox</i>	197	<i>capite plagioplateo</i> , <i>Eleotris</i>	436
<i>campi</i> , <i>Signalosa mexicana</i>	47	<i>striato</i> , <i>Perca marina</i>	330
<i>Campostoma</i>	146	<i>capito</i> , <i>Macrurus</i>	205
<i>anomalum</i>	146	<i>Maerurus</i>	205
<i>brevis</i>	146	<i>Poromitra</i>	223
<i>callipteryx</i>	146	<i>caprinus</i> , <i>Balistes</i>	491
<i>formulosum</i>	147	<i>Otrynter</i>	336
<i>gobionum</i>	146	<i>Stenotomus</i>	336
<i>hippops</i>	146	<i>Capriscus</i>	491
<i>mormyrus</i>	146	<i>minima</i>	496
<i>nasutum</i>	146	<i>porcus</i>	491
<i>ornatum</i>	146	<i>capriscus</i> , <i>Balistes</i>	491
<i>pricei</i>	147	<i>caprodes</i> , <i>Etheostoma</i>	282
<i>Campylodon</i>	75	<i>Percina</i>	283
<i>fabricii</i>	75	<i>Sciaena</i>	282, 283
<i>camtharum</i> , <i>Pristipoma</i>	335	<i>Caprophonus</i>	358
<i>camtschaticus</i> , <i>Crossias</i>	339	<i>aurora</i>	358
<i>camura</i> , <i>Chola</i>	129	<i>capros</i> , <i>Antigonia</i>	358
<i>Erogala</i>	129	<i>captiva</i> , <i>Goodea</i>	184
<i>Vallantia</i>	288	<i>caracasenses</i> , <i>Hemiloricaria</i>	161
<i>camurum</i> , <i>Boleosoma</i>	288	<i>caramuru</i> , <i>Muraenophis</i>	91
<i>Etheostoma</i>	292	<i>Carangidae</i>	239, 269
<i>camurus</i> , <i>Nothonotus</i>	290	<i>Carangichthys</i>	273
<i>Poecilichthys</i>	290	<i>typus</i>	273
<i>cana</i> , <i>Allogambusia</i>	188	<i>Carangoides blepharis</i>	274
<i>Gambusia</i>	188	<i>dorsalis</i>	274
<i>canadensis</i> , <i>Cynoperca</i>	282	<i>gallichthys</i>	274
<i>Salmo</i>	60	<i>carangoides</i> , <i>Seriolophus</i>	280
<i>canadensis</i> , <i>Lucioperca</i>	282	<i>Carangops</i>	271
<i>canadus</i> , <i>Gasterosteus</i>	281	<i>Carangus aureus</i>	272
<i>Rachycentron</i>	281	<i>esculentus</i>	273
<i>canaliculatus</i> , <i>Icelus</i>	378	<i>hippoides</i>	273
<i>canariensis</i> , <i>Chnus</i>	459	<i>carangus</i> , <i>Scomber</i>	273
<i>Pteraplatea</i>	29	<i>Caranx</i>	271, 273
<i>cancrivorus</i> , <i>Pisodonophis</i>	85, 86	<i>amblyrhynchus</i>	271
<i>candalaris</i> , <i>Profundulus</i>	174	<i>antillarum</i>	271
<i>candidissimus</i> , <i>Leptocephalus</i>	79	<i>atrimanus</i>	271
<i>candidus</i> , <i>Careproctus</i>	403	<i>aureus</i>	270, 272
<i>canensis</i> , <i>Acanthicus</i>	161	<i>bartholomaei</i>	272
<i>Leptoancistrus</i>	161	<i>beani</i>	272
<i>canescens</i> , <i>Ancanthias</i>	161	<i>blochii</i>	271
<i>canicula</i> , <i>Squalus</i>	11	<i>caballus</i>	272
<i>canina</i> , <i>Amia</i>	38	<i>chrysurus</i>	345
<i>caninus</i> , <i>Scopelinus</i>	169	<i>cibi</i>	272
<i>caninus</i> , <i>Lachnolaimus</i>	425	<i>daubentonii</i>	273
<i>Pajellus</i>	336	<i>defensor</i>	273
<i>canis</i> , <i>Cynias</i>	13	<i>dumerilii</i>	279
<i>Salmo</i>	54	<i>erithyrus</i>	273
<i>Squalus</i>	13	<i>esculentus</i>	273
<i>canna</i> , <i>Hemulon</i>	329	<i>falcatus</i>	271
<i>Cannorhynchus</i>	245	<i>fallax</i>	272
<i>Cantharus nigro-maculatus</i>	305	<i>fasciatus</i>	272
<i>Canthermes</i>	493	<i>frontalis</i>	273
<i>amphioxys</i>	494	<i>fürthii</i>	271
<i>carolæ</i>	494	<i>gardi</i>	272
<i>nasutus</i>	493, 494	<i>hasselti</i>	270
<i>pullus</i>	494	<i>heterophygus</i>	271
<i>Canthidermis</i>	492	<i>hippoides</i>	273
<i>angulosus</i>	492	<i>hippos</i>	273
<i>maculatus</i>	493	<i>ignobilis</i>	273
<i>sobaco</i>	492	<i>iridinus</i>	271
<i>sufflamen</i>	492	<i>kurra</i>	270
<i>willoughbei</i>	493	<i>latus</i>	272
<i>Canthigaster</i>	500	<i>lepturus</i>	272
<i>punctatissimus</i>	500	<i>leucurus</i>	271
<i>rostratus</i>	500	<i>lugubris</i>	273
<i>Canthigasteride</i>	500	<i>macarellus</i>	270
<i>Canthirhynchus</i>	337	<i>macrophthalmus</i>	271
<i>Canthogaster lobatus</i>	498	<i>margnatus</i>	272
<i>cantrani</i> , <i>Tylosurus</i>	197	<i>medusicola</i>	272
<i>cantrani</i> , <i>Tylosurus</i>	196	<i>melampygus</i>	272

List of scientific names—Continued

	Page		Page
<i>Caranx orthogrammus</i>	274	<i>Carcharius isodon</i>	15
<i>otrynter</i>	274	<i>punctatus</i>	15
<i>panamensis</i>	273	<i>carcharius, Carcharodon</i>	15
<i>pisquetos</i>	272	<i>Carcharodon</i>	20
<i>pisquetus</i>	272	<i>capensis</i>	20
<i>punctatus</i>	270, 273	<i>carcharias</i>	15, 20
<i>richardi</i>	272	<i>rondeleti</i>	20
<i>sanctæ-helenæ</i>	270	<i>verus</i>	20
<i>secundus</i>	272	<i>carcinognathus, Gymnothorax</i>	93
<i>semispinosus</i>	270	<i>cardenal, Bonaci</i>	313
<i>stellatus</i>	273	<i>cardonæ, Coralliozetus</i>	464
<i>symmetricus</i>	270	<i>Carelliparis</i>	400
<i>tenebrosus</i>	273	<i>agassizii</i>	400
<i>trachurus japonicus</i>	270	<i>dennyi</i>	400
<i>vinctus</i>	272	<i>herschelinus</i>	400
<i>Caranxomorus plumiearatus</i>	270	<i>tunicatus</i>	400
<i>Carapidae</i>	486	<i>Caremitra</i>	401
<i>carapinus, Coryphænoides</i>	205	<i>sima</i>	401
<i>Carapo</i>	101	<i>Carencheli</i>	77
<i>carapo, Gymnotus</i>	101, 486	<i>Careproctus</i>	402
<i>Carapus</i>	101, 486	<i>attenuatus</i>	403
<i>affinis</i>	486	<i>bowersianus</i>	403
<i>arenicola</i>	486	<i>candidus</i>	403
<i>bermudensis</i>	486	<i>colletti</i>	402
<i>Carauna</i>	309	<i>ectenes</i>	402, 403
<i>carauna, Serranus</i>	309	<i>furcellus</i>	403
<i>carbo, Aphanopus</i>	262	<i>gelatinosus</i>	402
<i>carbonaria, Percina</i>	283	<i>gilberti</i>	402
<i>Pileoma</i>	283	<i>longifilis</i>	403
<i>carbonarium, Hæmulon</i>	330	<i>longipinnis</i>	403
<i>carbonarius, Acipenser</i>	35	<i>melanurus</i>	402, 403
<i>Gadus</i>	209	<i>mollis</i>	403
<i>carbunculus, Etelis</i>	328	<i>opisthotremus</i>	403
<i>Carcharhinus</i>	15	<i>ostentum</i>	402
<i>cerdale</i>	16	<i>phasma</i>	402
<i>menisorrhæ</i>	16	<i>ranulus</i>	402
<i>natator</i>	17	<i>reinhardi</i>	402
<i>velox</i>	16	<i>simus</i>	401
<i>Carcharias</i>	15	<i>spectrum</i>	402
<i>acronotus</i>	16	<i>carbæum, Branchiostoma</i>	7
<i>æthalorus</i>	17	<i>carbæus, Sargus</i>	338
<i>atwoodi</i>	20	<i>carbææ, Belone</i>	197
<i>azureus</i>	16	<i>Ilsha</i>	45
<i>cæruleus</i>	16	<i>carbæus, Chloroscombrus</i>	276
<i>cerdale</i>	16	<i>Cælorhynchus</i>	207
<i>comersonii</i>	15	<i>Haliæutichthys</i>	507
<i>falciformis</i>	16	<i>Macrurus</i>	207
<i>falcipinnis</i>	16	<i>Tylosurus</i>	197
<i>fronto</i>	17	<i>carbæus, Aphthalmichthys</i>	89
<i>griseus</i>	19	<i>carinatum, Acanthosoma</i>	503
<i>hemiodon</i>	17	<i>Siphostoma</i>	240
<i>henlei</i>	16	<i>carinatus, Diodon</i>	503
<i>isodon</i>	15	<i>Labiichthys</i>	84
<i>lalandi</i>	15	<i>Placopharynx</i>	111
<i>lamia</i>	15, 20	<i>Salmo</i>	56
<i>lamiella</i>	15	<i>Syngnathus</i>	240
<i>limbatus</i>	16, 17	<i>carinicauda, Halosaurichthys</i>	74
<i>longurio</i>	15	<i>carletoni, Leuciscus</i>	116
<i>macloti</i>	17	<i>carminale, Tripterygium</i>	458
<i>menisorrhæ</i>	15	<i>carminalis, Eneapterygius</i>	458
<i>microps</i>	16	<i>carminatus, Cælorhynchus</i>	207
<i>milberti</i>	16, 17	<i>Macrurus</i>	207
<i>natator</i>	17	<i>carminifer, Macrourus</i>	206
<i>nicaraguensis</i>	16	<i>Macrurus</i>	206
<i>obesus</i>	14	<i>carinatus, Pteropodus</i>	369
<i>obscurus</i>	16	<i>Sebastichthys</i>	369
<i>oxyrhynchus</i>	15, 16	<i>carneus, Acronurus</i>	362
<i>platyodon</i>	15	<i>Sicyases</i>	400
<i>platyrhynchus</i>	15	<i>carolæ, Cantherines</i>	404
<i>porosus</i>	16	<i>carolina, Argentina</i>	39
<i>(Prionodon) limbatus</i>	16	<i>Atherina</i>	246
<i>milberti</i>	16	<i>Trigla</i>	406-408
<i>mullerii</i>	16	<i>carolinæ, Potamocottus</i>	383, 385
<i>punctatus</i>	15	<i>carolinensis, Balistes</i>	491
<i>remotus</i>	16	<i>Cestreus</i>	354
<i>sorrhæ</i>	15	<i>Clupea</i>	44
<i>taurus</i>	15, 19	<i>Gobius</i>	440
<i>temminckii</i>	15	<i>Hyperistius</i>	305
<i>tigris</i>	19	<i>Otolithus</i>	355
<i>velox</i>	16	<i>Seriola</i>	279
<i>carcharias, Carcharodon</i>	15, 20	<i>carolinus, Gasterosteus</i>	276, 277
<i>Squalus</i>	15, 20	<i>Labrus</i>	424
<i>(Carcharias) terre-novæ, Squalus</i>	14	<i>Lipophrys</i>	461
<i>Carcharidae</i>	13	<i>Merulinus</i>	407
		<i>Pholis</i>	461

List of scientific names—Continued

	Page		Page
carolinus, <i>Priacanthus</i>	322, 323	Catonotus flabellaris.....	292, 383, 385
<i>Pteraclis</i>	266	kennicotti.....	293
<i>Trachinotus</i>	277	lineolatus.....	293
carpintis, <i>Herichthys</i>	423	Catostomidae.....	102
<i>Neotrophus</i>	423	Catostomus.....	105, 106
carpio, <i>Carpoides</i>	103	(<i>Acomus</i>) <i>generosus</i>	104
<i>Catostomus</i>	103, 109	griseus.....	107
<i>Cyprinodon</i>	180	<i>guzmaniensis</i>	104
<i>Floridichthys</i>	180	<i>lactarius</i>	107
<i>Carpoides</i>	103	<i>alticolus</i>	105
<i>bison</i>	103	<i>anisurus</i>	109
<i>carpio</i>	103	<i>aræopus</i>	106
<i>cutisanserinus</i>	103	<i>ardens</i>	105
<i>cyprinus</i>	103	<i>aureolus</i>	109, 110
<i>darnalis</i>	103	<i>aurora</i>	106
<i>difformis</i>	103	<i>bernardini</i>	106
<i>elongatus</i>	104	<i>bostoniensis</i>	105
<i>grayi</i>	103	<i>bubalus</i>	102
<i>labiosus</i>	103	<i>carpio</i>	103, 109
<i>meridionalis</i>	104	<i>catostomus</i>	106
<i>microstromus</i>	104	<i>chamberlaini</i>	107
<i>niger</i>	102	<i>chloropteron</i>	105
<i>nummifer</i>	103	<i>clarkii</i>	104
<i>seleus</i>	103	<i>commersoni</i>	105
<i>taurus</i>	102	<i>communis</i>	105
<i>thompsoni</i>	103	<i>conchos</i>	107
<i>tumidus</i>	103	<i>congestus</i>	106
<i>urus</i>	102	<i>cypho</i>	103
<i>vacca</i>	103	<i>cyprinus</i>	103
<i>velifer</i>	103	<i>discobolus</i>	107
<i>vitulus</i>	102	<i>duquesnei</i>	110
<i>carringtoni</i> , <i>Apocope</i>	140	<i>elongatus</i>	110
<i>carringtoni</i> , <i>Apocope</i>	141	<i>erythrus</i>	110
<i>caryi</i> , <i>Acipenser</i>	34	<i>fasciatus</i>	109
<i>Embiotoca</i>	412	<i>fasciolaris</i>	109
<i>Hypsurus</i>	412	<i>fecundus</i>	108
<i>cascahiensis</i> , <i>Brachyrhaphis</i>	187	<i>flexuosus</i>	105
<i>Bryconameris</i>	97	<i>forsterianus</i>	105, 106
<i>Gambusia</i>	187	<i>gibbosus</i>	109
<i>cassidy</i> , <i>Embiotoca</i>	411	<i>gla</i>	110
<i>castanea</i> , <i>Sidera</i>	92	<i>gracilis</i>	105
<i>castaneola</i> , <i>Sparus</i>	265	<i>griseus</i>	107
<i>castaneum</i> , <i>Macrostoma</i>	167	<i>hudsonius</i>	106
<i>castaneus</i> , <i>Gymnothorax</i>	92	<i>insignis</i>	105, 106
<i>Ichthyomyzon</i>	9	<i>labiatus</i>	106
<i>Notoscopelus</i>	167	<i>latipinnis</i>	107
<i>castelnaudi</i> , <i>Cylindrosteus</i>	37	<i>lesneuri</i>	111
<i>castor</i> , <i>Pontinus</i>	372	<i>longirostrum</i>	106
<i>Catabelia</i>	166	<i>macrocheilus</i>	106
<i>brachyichthys</i>	166	<i>macrolepidotus</i>	111
<i>Catostyx</i>	480	<i>maculosus</i>	107
<i>rubrostris</i>	480	<i>megastomus</i>	107
<i>sumus</i>	480	<i>melanops</i>	109
<i>catalina</i> , <i>Neothunnus</i>	260	<i>melanotus</i>	137
<i>Catalineta</i>	361	<i>microps</i>	106
<i>Catalufa</i>	323	<i>nanomyzon</i>	106
<i>catalufa</i> , <i>Priacanthus</i>	323	<i>nebuliferus</i>	104
<i>Cataphracti</i>	363	<i>nigricans</i>	107
<i>Cataphractus schoneveldii</i>	395	<i>etowanus</i>	107
<i>cataphractus</i> , <i>Agonus</i>	395	<i>occidentalis</i>	106
<i>Cottus</i>	395	<i>oneida</i>	110
<i>Gasteracanthus</i>	237	<i>pallidus</i>	105
<i>Gasterosteus</i>	237	<i>planiceps</i>	107
<i>cataractæ</i> , <i>Gobio</i>	139	<i>plebeus</i>	104
<i>Rhinichthys</i>	139	<i>pocatello</i>	107
<i>cataractus</i> , <i>Leucosomus</i>	117	<i>reticulatus</i>	105
<i>catena</i> , <i>Bassozetus</i>	480, 481	<i>retropinnis</i>	107
<i>Bathyonus</i>	481	<i>rex</i>	108
<i>catenata</i> , <i>Echidna</i>	94	<i>rhotocetus</i>	107, 108
<i>Pocilla</i>	179	<i>richardsoni</i>	106
<i>catenatum</i> , <i>Xenisma</i>	179	<i>snyderi</i>	106
<i>catenatus</i> , <i>Gymnothorax</i>	94	<i>sonoriensis</i>	106
<i>catenula</i> , <i>Muraenophis</i>	94	<i>sucklii</i>	105
<i>catasbei</i> , <i>Pomotis</i>	302	<i>tahoensis</i>	107
<i>catharina</i> , <i>Pristipoma</i>	333	<i>texasus</i>	108
<i>Cathetostoma albigutta</i>	454	<i>tsilicoensis</i>	106
<i>Cathorops</i>	151	<i>tuberculatus</i>	109
<i>gulesa</i>	151	<i>utawana</i>	105
<i>hypophthalma</i>	151	<i>velifer</i>	103
<i>Cathocheilichthys</i>	341	<i>vittatus</i>	109
<i>Catonotus</i>	292, 293	<i>xanthopus</i>	107
<i>fasciatus</i>	293	<i>catostomus</i> , <i>Catostomus</i>	106
<i>flabellare cumberlandicum</i>	293	<i>Cyprinus</i>	105, 106
<i>lineolatus</i>	293	<i>Phenacobius</i>	142

List of scientific names—Continued

	Page		Page
Catulus.....	11, 17	cayanus, Pristigaster.....	45
brunneus.....	12	cayennensis, Citharichthys.....	218, 219
cephalus.....	11	Otolithus.....	356
saxatilis.....	11	Vomer.....	275
uter.....	12	Cayenne.....	448
xanurus.....	12	guichenoti.....	448
catulus, Ameiurus nebulosus.....	154	caymanensis, Gambusia.....	186
(canicula), Scylliorhinus.....	13	cayorum, Anchovyia cherosstoma.....	49
Galeus.....	13	Anchoviella.....	49
Gobius.....	440	Corythoichthys.....	243
Pimelodus.....	154	Hippichthys.....	243
catulus, Haustor.....	153	Ogilbia.....	479, 480
Serranus.....	311	cayuga, Cottus gracilis.....	386
Silurus.....	153	Eucalia inconstans.....	238
caucanus, Ageneiosus.....	159	Hybopsis.....	133
cauda forcipata, Brama marina.....	265	Notropis.....	133
cauda nigra, Perca marina.....	330	cayuga atrocandialis, Notropis.....	133
caudacinctus, Prilonotus (Anchisomus).....	500	Cebidichthyidae.....	466
caudacuta, Motella.....	209	Cebidichthys.....	466
caudafurcatus, Pimelodus.....	152	cristagalli.....	466
caudale, Ariosoma.....	80	violaceus.....	466
caudalis, Congermurena.....	80	cedroscensis, Athernops insularum.....	252
Iridio.....	427	celatus, Batrachus.....	487
Julis.....	427	Centaurus.....	503
Leptocephalus.....	81	centrale, Cichlasoma.....	419
Leptocephalus (Microconger).....	81	centralis, Parapetema.....	419
Microconger.....	81	Centrarchidae.....	297
Mixonus.....	483	Centrarchus.....	304
Pomacentrus.....	413	hexacanthus.....	304, 305
caudanotatus, Mesoprion.....	326	interruptus.....	303
caudata, Lamna.....	16	macropterus.....	304
Motella.....	209	maculosus.....	303
caudatus, Lepidopus.....	262	obscurus.....	298
Paraliparis.....	404	pentacanthus.....	303
caudicula, Conger.....	79	pomotis.....	303
caudilumbatus, Conger.....	79	tetraacanthus.....	419
Echelus.....	79	centrarchus, Archocentrus.....	420
caudimacula, Hemulon.....	329	Heros.....	420
Sargus.....	338	centrina, Callorhynchus.....	32
audimaculata, Strongylura.....	195	Centriscus.....	245
caudimaculatus, Hemietocetus.....	199	scelopar.....	245
caudofasciata, Lima.....	191	Centristhmus.....	319
Caularchus.....	488	signifer.....	319
maendricus.....	488	Centroblennius.....	470
caulus, Signistes.....	391	Centrochromis.....	415
Caulistius.....	488	rudis.....	415
papillifer.....	488	centrognathos, Zanelus.....	362
Caulolatilus.....	357	centrognathus, Zanelus.....	362
affinis.....	357	Centrolabrus, Zanelus.....	424
anomalus.....	357	exoletus.....	424
chrysops.....	357	Centrolophidae.....	267
cyanops.....	357	Centrolophus.....	267
microps.....	357	lparis.....	267
princeps.....	357	morio.....	267
Caulolepis.....	232	niger.....	267
longidens.....	232	pompius.....	267
Caulophryne.....	511	Centronotus.....	278, 487
jordani.....	511	argenteus.....	277
Caulophrynidae.....	511	conductor.....	278
Caulopus.....	171	cristagalli.....	468
altivelis.....	171	fasciatus.....	467
borealis.....	171	gardneri.....	281
(Caulopus) poeyi, Alepidosaurus.....	171	islandicus.....	471
caulopus, Homesthes.....	463	laetus.....	468
caurina, Clarkina.....	118	spinousus.....	281
caurinus, Cyprinus.....	118	Centropholis.....	266
Cyprinus (Lenciscus).....	118	goodii.....	266
Leuciscus.....	112, 118	petersi.....	266
Mylocheilus.....	118	centropleura, Uranidea semiscabra.....	383
Pteropodus.....	369	Centropomidae.....	305
Sebastes.....	369	Centropomus.....	305
cauthocum, Gobiosoma.....	446	affinis.....	306
cauthocus, Barbulifer.....	446	albus.....	307
cavalla, Cybium.....	257	armatus.....	306
Sierra.....	257	appendiculatus.....	305
cavernosus, Bathygadus.....	207	brevis.....	306
Hymenoccephalus.....	207	cuvieri.....	306
cavifrons, Ambloplites.....	303	ensiferus.....	306
Diagramma.....	335	grandoculatus.....	306
Hemitripterus.....	363	luciopeca undecimalis.....	305
Icelinus.....	375	luteus.....	281
Tarandichthys.....	375	medius.....	306
Caxis.....	325	mexicanus.....	306
caxis, Sparus.....	325	nigrescens.....	306
cayannensis, Lutjanus.....	354	parallelus.....	306

List of scientific names—Continued

	Page		Page
<i>Centropomus pectinatus</i>	306	<i>Ceratias, holbolli</i>	508
<i>pedimacula</i>	306	<i>Ceratichthys</i>	137
<i>robalito</i>	306	<i>physignathus</i>	136
<i>scaber</i>	306	<i>prothemius</i>	136
<i>undecimalis</i>	305, 306	<i>smithi</i>	145
<i>undecim-radiatus</i>	305	<i>squamulentus</i>	136
<i>unionensis</i>	306	<i>vigilax</i>	137, 144, 145
<i>viridis</i>	306	<i>Ceratidae</i>	508
<i>Centropristes</i>	317	<i>Ceratobatus</i>	32
<i>ayresi</i>	318	<i>robertsi</i>	32
<i>fuscus</i>	319	<i>Ceratocottus</i>	381
<i>merus</i>	312	<i>dicerca</i>	381
<i>nigricans</i>	317	<i>lucasi</i>	382
<i>ocyurus</i>	318	<i>Ceratoptera</i>	32
<i>rufus</i>	318	<i>Ceratiscopelus</i>	167
<i>striatus</i>	317	<i>maderensis</i>	167
<i>subligarius</i>	321	<i>Ceratichthys</i>	137, 144
<i>tabacarius</i>	319	<i>cumingii</i>	138
<i>Centropristis</i>	317	<i>cyclotus</i>	137
<i>annularis</i>	319	<i>hyalinus</i>	137
<i>aurorubens</i>	327, 323	<i>hypsinotus</i>	138
<i>brasilensis</i>	320	<i>labrosus</i>	138
<i>displurus</i>	320	<i>leptocephalus</i>	137
<i>macrophthalmus</i>	328	<i>lucens</i>	137
<i>macropoma</i>	318	<i>micropogon</i>	137
<i>subligarius</i>	320	<i>monacus</i>	138
<i>Centropyge</i>	361	<i>physignathus</i>	136
<i>clarionensis</i>	361	<i>prothemius</i>	136
<i>lunata</i>	361	<i>sallen</i>	116
<i>passer</i>	361	<i>smithi</i>	145
<i>Centrosyllium</i>	21	<i>squamulentus</i>	136
<i>fabricii</i>	21	<i>sterletus</i>	139
<i>Centrosymnus</i>	21	<i>stigmaticus</i>	137
<i>coelolepis</i>	21	<i>ventricosus</i>	141
<i>centroura, Raja</i>	28	<i>vigilax</i>	137, 144, 145
<i>centrourus, Pastinachus</i>	28	<i>zanemus</i>	138
<i>Centruropus</i>	28	<i>Ceratius uranoscopus</i>	508
<i>spadiceus</i>	87	<i>Ceratiscopelus</i>	167
<i>cepediana, Megalops</i>	46	<i>maderensis</i>	167
<i>cepedianum, Dorosoma</i>	46	<i>cercostigma, Cyprinella</i>	130
<i>cepedianum exile, Dorosoma</i>	46	<i>Erogala</i>	130
<i>cepedianus, Lophotes</i>	215	<i>Cerdale</i>	473
<i>Priacanthus</i>	323	<i>ionthas</i>	473
<i>Cephacandia</i>	408	<i>cerdale, Carcharias</i>	16
<i>Cephalacanthus</i>	408	<i>Carcharhinus</i>	16
<i>spindrella</i>	408	<i>Scytalina</i>	473
<i>Cephalocassis</i>	149	<i>Cerdalidae</i>	472
<i>cephaloides, Cottus</i>	389	<i>Cerdo</i>	210
<i>Cephalopholis</i>	309, 425	<i>cerea, Memidia beryllina</i>	247
<i>argus</i>	309	<i>Cerna</i>	310
<i>fulvus</i>	309	<i>acutirostris lata</i>	315
<i>fulvus punctatus</i>	309	<i>macrogenis</i>	315
<i>ruber</i>	309	<i>nebulosa</i>	315
<i>stellatus</i>	309	<i>sicana</i>	312
<i>taenios</i>	309	<i>cernium, Polyprion</i>	308
<i>Cephalopis nigripinnis</i>	309	<i>ceruinum, Lepophidium</i>	484
<i>Cephaloptera johni</i>	32	<i>Leptophidium</i>	484
<i>manta</i>	32	<i>cervinus, Ptychostomus</i>	111
<i>vampyrus</i>	32	<i>Scartomyzon</i>	111
<i>Cephalopterus</i>	31	<i>Teretulus</i>	111
<i>giora</i>	32	<i>cervus, Synanceia</i>	381
<i>hypnotomus</i>	32	<i>Cestracion</i>	18
<i>manta</i>	32	<i>francisci</i>	10
<i>olferi</i>	32	<i>pantherinus</i>	10
<i>vampyrus</i>	32	<i>quoyi</i>	10
<i>Cephaloscyllium</i>	11	<i>Cestraciontes</i>	10
<i>uter</i>	12	<i>Cestreus</i>	354
<i>Cephalus</i>	503	<i>carolinensis</i>	354
<i>cephalus, Brycon</i>	98	<i>predatorius</i>	354
<i>Catulus</i>	11	<i>Cestrorhinus</i>	18
<i>Gobiesox</i>	483, 489	<i>cetaceus, Squalus</i>	20
<i>Mugil</i>	252, 253	<i>Cetengraulis</i>	51
<i>Paraliparis</i>	404	<i>edentulus</i>	51
<i>Scylliorhinus</i>	11	<i>garmani</i>	51
<i>Semotilus</i>	117	<i>gilberti</i>	51
<i>Cephola trachyptera</i>	216	<i>mysticetus</i>	51
<i>Cephus</i>	210	<i>Cetomimidae</i>	172
<i>Cepolopis</i>	477	<i>Cetomimus</i>	172
<i>cerapatus, Opsanus</i>	487	<i>gilli</i>	172
<i>cerasinus, Gobiesox</i>	489	<i>storeri</i>	172
<i>Hypsilepis cornutus</i>	128	<i>Cetorhinidae</i>	20
<i>Luxilus</i>	128	<i>Cetorhinus</i>	20
<i>Ceratacanthus</i>	495	<i>marinus</i>	20
<i>schoepli</i>	495	<i>shaviansus</i>	20
<i>Ceratas</i>	508	<i>Cetunculi</i>	171

List of scientific names—Continued

	Page		Page
Chænobryttus.....	302	chagresi, Pimelodus.....	188
antistius.....	303	Thyrina.....	249
gulosus.....	302	Xenocara.....	160
Chænomugil.....	254	chalceum, Pristipoma.....	335
proboscideus.....	254	chalceus, Orthopristis.....	335
Chænophryne.....	510	Chalcinopsis.....	98
longiceps.....	510	chagresensis.....	98
Chænopsetta.....	223	striatulus.....	98
Chænopsidae.....	465	chalcogramma, Theragra.....	209
Chænopsis.....	465	chalcogrammus, Gadus.....	209
alepidota.....	465	chalcogrammus fucensis, Pollachius.....	209
lucius.....	465	chalmus, Epinephelus.....	315
ocellata.....	465	Chalinura.....	204
ocellatus.....	465	brevibarbis.....	204
chaetirhamphus, Antelochumera.....	33	filifera.....	204
Chaetodipterus.....	358	occidentalis.....	204
faber.....	358	serrula.....	204
zonatus.....	358	simula.....	204
Chaetodon.....	359	spinulosa.....	204
alepidotus.....	266	Chalinurus.....	204
arcuatus.....	360	challengeri, Notacanthus.....	76
atæniatus.....	360	Polyacanthotus.....	76
aureus.....	360	chalybeus, Hybopsis.....	127
aunga.....	359	Hydrophlox.....	127
aya.....	360	chalybeus abbotti, Notropis.....	127
bimaculatus.....	360	chalybeus, Chlorophthalmus.....	165
bricei.....	359	Hyphalonedrus.....	165
capistratus.....	359	chamaeleonticeps, Lopholatilus.....	357
chirurgus.....	362	chamberlaini, Catostomus.....	107
ciliaris.....	361	Chasmistes.....	107
collaris.....	359	Cottus.....	385
cornutus.....	362	Hudsonius.....	132
couaga.....	362	Notropis.....	132
cyprinaceus.....	340	chamensis, Ophichthus.....	88
ephippium.....	359	Chanida.....	40
faber.....	358	Channomuræna.....	95
glaucus.....	376	cubensis.....	95
gracilis.....	360	vittata.....	95
humeralis.....	360	Channo-Muræna.....	95
lanceolatus.....	352, 353	Chanos.....	40
lineolatus.....	359	arabicus.....	40
littoricola.....	360	cyprinella.....	40
longirostris.....	359	chanos, Mugil.....	40
lutescens.....	360	chantenay, Raja.....	25
marginatus.....	415	chapale, Chirostoma.....	251
mauriti.....	415	Falcula.....	142
nigrirostris.....	360	Falcularius.....	142
ocellatus.....	360	Chapalichthys.....	184
ornatissimus.....	359	encaustus.....	184
oviformis.....	358	chaperi, Acanthoclinus.....	457
paræ.....	361	Parachnus.....	457
paru.....	360	Chapinus.....	496
plumeri.....	358	bicaudalis.....	497
rhomboides.....	276, 277	chapmani, Psenes.....	269
sargoides.....	415	Tænioconger.....	82
saratius.....	414	Characidium.....	100
sedentarius.....	360	fasciatum.....	100
sohar.....	362	marshi.....	100
sordidus.....	414	Characiniæ.....	95
squamulosus.....	361	Characodon.....	182
striatus.....	359	bilineatus.....	183
triangulum.....	359	eiseni.....	183
tricolor.....	361	encaustus.....	184
tristegus.....	362	ferrugineus.....	183
unicornis.....	362	furcatus.....	183
unicolor.....	360	garmani.....	182
unimaculatus.....	359	geddesi.....	182
chaetodon, Cottus.....	487	lateralis.....	182, 183
Mesogonistius.....	303	luitpoldi.....	183, 184
Pomotis.....	303	multiradiatus.....	183
Chaetodontidae.....	359	variatus.....	182
Chaetodontops.....	359, 360	Charax maxilla superiore longiore.....	100
Chaetostoma.....	160	chartes, Platycephalus.....	221
fischeri.....	161	charybdis, Lepomis.....	303
loborhynchus.....	160	Chasmistes.....	108
Chaetostomus.....	160	brevirostris.....	108
aspidolepis.....	161	chamberlaini.....	107
fischeri.....	161	copel.....	108
heteracanthus.....	161	cujus.....	108
latifrons.....	160	fecundus.....	108
chagresensis, Chalcinopsis.....	161	horus.....	108
chagresi, Ancistrus.....	160	luxatus.....	108
Meridia.....	249	stomias.....	108
Pimelodella.....	158	tahoensis.....	107

List of scientific names—Continued

	Page		Page
Chasmodes.....	462	Chilomycterus californiensis.....	502
bosquianus.....	462	puncticulatus.....	502
jenkinsi.....	463	Chilorhinus.....	83
novemlineatus.....	462	suensonii.....	83
quadri-fasciatus.....	462	Chimæra.....	32
saburrae.....	462	abbreviata.....	33
Chatoëssus.....	46	affinis.....	33
ellipticus.....	46	collei.....	33
eumorphus.....	44	cristata.....	32
mexicanus.....	46, 47	mediterranea.....	32
signifer.....	44	mirabilis.....	33
Chauliodontidae.....	71	monstrosa.....	32
Chauliodus.....	71	plumbea.....	33
barbatus.....	71	Chimæridæ.....	32
macouni.....	71	Chimæroides.....	32
richardsoni.....	71	chinensis, Aulostomus.....	244
sloani.....	71	Bahstes.....	494
Chaunacidae.....	506	Tachysurus.....	150
Chaunax.....	506	chionostigma, Echidna.....	94
nuttingi.....	506	chlostictus, Entomacrodus.....	463
pictus.....	506	Salaras.....	463
Cheilichthys.....	499	chiquita, Aboma.....	443
annulatus.....	499	Gobius.....	443
asellus.....	500	Chirocentrodon.....	45
furthi.....	499	bleekeriana.....	45
furthi.....	499	tæniatus.....	45
pachygaster.....	499	Chirolophidae.....	466
politus.....	499	Chirolophus japonicus.....	466
testudineus.....	499	Chironectes.....	505
trichocephalus.....	499	arcticus.....	504
(Cheilichthys) pachygaster, Tetiodon.....	499	laevigatus.....	504
Cheilo auratus.....	423	mentzel.....	505
Cheilodipterus.....	296	mentzeli.....	508
acoupa.....	354	multiocellatus.....	508
affinis.....	296	pictus.....	504
heptacanthus.....	280	principis.....	505
lineatus.....	296	scaber.....	505
macrodon.....	296	sonntagii.....	508
Chelonemus.....	117	tenebrosus.....	505
Cheilotrema.....	348	tigris.....	508
Cheiragonus.....	393	tumidus.....	504
Cheirodon gorgonæ.....	96	Chiroopsis constellatus.....	373
insignis.....	96	nebulosus.....	373
Chelmo pelta.....	359	Chirostoma.....	250
Chelmon aculeatus.....	359	arge.....	250
chelonus, Allector.....	484	attenuatum.....	250
Cheloniger.....	333	bartoni.....	250
chemnitz, Notacanthus.....	75	beryllinum.....	247
cheneyi, Cottogaster.....	286	breve.....	250
Imostoma.....	286	chapale.....	251
Cheonda.....	117, 118	consocium.....	251
cærulea.....	120	crystallinum.....	251
cooperi.....	118	diaz.....	251
copei.....	119	estor.....	250, 251
egregia.....	118	grandocule.....	251
evermanni.....	118	guatemalensis.....	249
humboldtii.....	119	humboldtianum.....	250
hydrophlox.....	119	jordani.....	250
microdon.....	118	labaræ.....	250
modesta.....	120	lermæ.....	251
montana.....	119	lucius.....	251
Cherna.....	311	mezguntal.....	250
cherna, Anthias.....	311	ocotlane.....	251
chesteri, Phycus.....	213	patzcuaro.....	250
Urophycis.....	213	promelas.....	251
Chiasmodon.....	452	regani.....	250
niger.....	452	seculum.....	251
subniger.....	452	sphyraena.....	251
Chiasmodontidae.....	452	vagrans.....	248
Chiasmodus.....	452	vomarina.....	250
chickasawensis, Luvulus.....	130	zirahuen.....	250
chiuahua, Moniana.....	132	Chirurgus, Chaetodon.....	362
Notropis.....	132	Chirus.....	373
Chilaria.....	435	ballas.....	374
chilensis, Brama.....	265	guttatus.....	373
Sarda.....	259	maculo-seriatus.....	373
chilensis, Exocoetus.....	200	ordinatus.....	374
Pelamys.....	259	pictus.....	374
chiliticus, Hybopsis.....	127	trigrammus.....	374
Hydrophlox.....	127	Chirus, Phytichthys.....	469
Chlomycterus.....	502	Xiphister.....	469
affinis.....	502	chisoyensis, Pœonia.....	192
antillarum.....	502	chisso, Blennius.....	212
atinga.....	502		
briareos.....	502		

List of scientific names—Continued

	Page		Page
Chitonotus.....	375	Chromis epileurorum.....	280
megacephalus.....	375	fenestrata.....	416
pugetensis.....	375	(Furcata) striolata.....	412
chittendeni, Cyclopetta.....	221	fusco-maculatus.....	419
chlamydes, Bollmannia.....	443	chromus, Hæmulon.....	329
chleivastes, Gymnothorax.....	93	Pogonius.....	352
sidera.....	93	Sparus.....	413
Chlopsis.....	82	Chrosomus.....	113
bicolor.....	82	dakotensis.....	113
equatorialis.....	82	eos.....	113
gilberti.....	82	ervthrogaster.....	113
chlora, Chloia.....	135	oreas.....	113
Chlorichthys.....	428	pyrrhogaster.....	113
chloris, Searus.....	432	chrosomus, Hybopsis.....	127
chloristia, Codoma.....	129	Hydrophlox.....	127
Erogala.....	129	chrysargyreum, Hæmulon.....	330, 331
chlorocephalus, Hybopsis.....	127	chrysargyreus, Brachygenys.....	331
Hydrophlox.....	127	chryseus, Rhinoberyx.....	235
chloropteron, Catostomus.....	105	Chrysipetra.....	414
chloropteron, Plectropoma.....	312	Chrysitis, Dionda.....	144
chloropterus, Alphestes.....	312	chrysocephalus, Luxilus.....	128
Chlorophthalmidæ.....	165	Luxilus cornutus.....	128
Chlorophthalmus.....	165	chrysochloris, Pomolobus.....	41, 42
agassizi.....	165	chrysogaster, Agosia.....	140
chalybeus.....	165	chrysoleuca, Corvina.....	346
mento.....	165	chrysoleucas, Leucosomus.....	117
traculentus.....	165	Notemigonus.....	117
Chloroscombrus.....	276	chrysoleucus, Nector.....	346
caribæus.....	276	chrysomelanurus, Sparus.....	311
chrysurus.....	276	chrysomelas, Pteropodus.....	369
orquea.....	276	Sebastichthys.....	369
stirurus.....	276	Chrysoperca.....	307
chlorosoma, Vailantia.....	288	interrupta.....	307
chlorostictus, Sebastichthys.....	367	Chrysophrys aculeata.....	336
Sebastomus.....	367	chrysops, Caulolatilus.....	357
chlorostomus, Trisotropis.....	314	Latlus.....	357
chlururum, Plectropoma.....	321	Lepibema.....	307
Chlorurus.....	433	Ophisurus.....	88
chlururus, Hypoplectrus unicolor.....	321	Pera.....	307
chocoensis, Gephyrocharax.....	96	Sparus.....	336
Chœroichthys.....	243	Stenotomus.....	336
valenciennesi.....	243	chrysopsis, Hydodon.....	41
Chœrojulis arangoi.....	426	chrysoptera, Pera.....	335
grandisquamis.....	426	chrysopterum, Hæmulon.....	329
chœrostoma, Anchoviella.....	48	Sparisoma.....	432
chœrostomus, Engraulis.....	48	chrysopteris, Leuciscus.....	117
chogset, Labrus.....	424	Orthopristis.....	335
chogset fulva, Labrus.....	424	Pera.....	335
Chologaster.....	194	Searus.....	432
agassizi.....	194	chrysos, Scomber.....	272
avitus.....	194	Chrysotodus.....	216
cornutus.....	194	chrysotus, Fundulus.....	177
papilliferus.....	194	Zygonectes.....	177
Chondrostoma pullum.....	146	chrysura, Bairdiella.....	345
Chondropterygi.....	10	chrysurus, Caranx.....	345
Chonophorus.....	442	Chloroscombrus.....	276
banana.....	442	Dipterodon.....	345
buculentus.....	442	Glyphidodon.....	415
contractus.....	442	Glyphisodon.....	415
mexicanus.....	442	Ocyurus.....	327
nelsoni.....	442	Scomber.....	276
chordatus, Saccopharynx.....	95	Sparus.....	327
Choregon.....	66	Stegastes.....	415
Chornemus altus.....	278	chrysus, Eupomacentrus.....	414
palometa.....	278	chrysypa, Anguilla.....	77
saltans.....	277	chucunaque, Rivulus.....	180
Chlorististum.....	315	chucunaque succubi, Rivulus.....	180
rubrum.....	315	chuss, Blennius.....	213
Chriodorus.....	199	Urophycis.....	213
atherinoides.....	199	cibarius, Ammocetes.....	10
Chriolax.....	407	Cibi amarilla.....	272
Chriolepis.....	438	cibi, Caranx.....	272
minutillus.....	438	Cibotia.....	496
Chriomitra.....	257	cicatricosus, Ballistes.....	493
concolor.....	257	Pleuronectes.....	227
Chriope.....	133, 135, 136	cicatrifer, Lycodes.....	475
albeolus.....	136	Cichla ænea.....	303
Chriopeops.....	175	fasciata.....	298
goodei.....	175	floridana.....	297
Christoma beryllinum.....	247	minima.....	298
pennsulæ.....	247	ohiensis.....	298
Chromides.....	412	storeria.....	304
Chromis.....	280, 413	Cichlasoma.....	416
bermudæ.....	413	acutum.....	422
enchrysurus.....	413	alfari.....	419

List of scientific names—Continued

	Page		Page
<i>Cichlasoma calobrense</i>	417	<i>circularis</i> , Raja.....	24
centrale.....	419	Raja.....	24
dorsatum.....	420	<i>circumnatus</i> , <i>Scarus</i>	433
eigenmanni.....	417	<i>cirratus</i> , <i>Ginglymostoma</i>	12
evermanni.....	420	<i>cirratus</i> , <i>Phycus</i>	212
frontale.....	417	Squalus.....	12
gadovii.....	417	<i>Cirrhitidae</i>	358
globosum.....	421	<i>Cirrhitichthys</i>	358
granadense.....	421	<i>Cirrhitidae</i>	358
hedricki.....	421	<i>Cirrhitus</i>	358
laurei.....	421	<i>Cirrhitus</i>	358
lethrinus.....	420	<i>betasurus</i>	358
maculicauda.....	421	<i>maculatus</i>	358
meeki.....	416	<i>rivulatus</i>	358
milleri.....	417	<i>carrhosum</i> , <i>Lepisoma</i>	459
mojarra.....	417	<i>carrhosus</i> , <i>Blepsias</i>	382
multifasciatum.....	418	Hypostomus.....	160
nigrum.....	418	Trachinus.....	382
punctatum.....	417	<i>Cirrostromi</i>	7
robertsoni.....	420	Cisco.....	62, 63
septemfasciatum.....	421	cismontanus, <i>Coregonus williamsoni</i>	65
sexfasciatum.....	421	Prosopium williamsoni.....	65
spinosissimum immaculatum.....	420	<i>Citharichthys</i>	218, 219
steindachneri.....	419	<i>aethalon</i>	218
teapei.....	417	<i>arctifrons</i>	219
tenuis.....	419	<i>arenaceus</i>	219
tuyrensis.....	421	<i>cayennensis</i>	218, 219
umbriferum.....	417	<i>dinoceros</i>	219
zonatum.....	417	<i>fragilis</i>	218
(<i>Cichlasoma</i>) <i>spinossissimum</i> , <i>Heros</i>	420	<i>gilberti</i>	219
<i>Cichla</i>	416	<i>guatemalensis</i>	219
angulifer.....	416	<i>latifrons</i>	218
bifasciatus.....	416	<i>macrops</i>	219
calobrensis.....	417	<i>micro</i>	219
deppii.....	416	<i>microstomus</i>	219
fenestratum.....	416	<i>pætilus</i>	218
godmanni.....	416	<i>panamensis</i>	221
guttulatus.....	416	<i>platophrys</i>	219
intermedius.....	416	<i>sordidus</i>	218
meeki.....	416	<i>spilopterus</i>	219
microphthalmus.....	416	<i>stigmaeus</i>	219
melanurus.....	416	<i>sumichrasti</i>	219
umbrifer.....	417	<i>uhleri</i>	219
<i>Cichlidae</i>	416	<i>unicornis</i>	219
<i>cigonella</i> , <i>Belone</i>	195	<i>ventialis</i>	220
<i>ciliaris</i> , <i>Angelichthys</i>	361	<i>xanthostigmus</i>	218
<i>Balistes</i>	492	<i>Citharus</i>	222
<i>Chætodon</i>	361	<i>Citræodus</i>	359
<i>Zeus</i>	274	<i>Citrinellus</i>	420
<i>Ciliata</i>	208	<i>citrinellus</i> , <i>Astatheros</i>	422
<i>argentata</i>	208	<i>Heros</i>	422
<i>glauca</i>	208	<i>Citula</i>	273, 274
<i>ciliatus</i> , <i>Balistes</i>	494	<i>dorsalis</i>	274
<i>Blennius</i>	473	<i>orthogramma</i>	274
<i>Entosphenus</i>	10	<i>sturensis</i> , <i>Oxyloricaria</i>	162
<i>Epinephelus</i>	365	<i>civilis</i> , <i>Hybognathus</i>	144
<i>Monacanthus</i>	494	<i>clara</i> , <i>Menidia</i>	248
<i>Petromyzon</i>	10	<i>clarias</i> , <i>Pimelodus</i>	159
<i>Sebastosomus</i>	365	<i>Silurus</i>	159, 159
<i>cimbrius</i> , <i>Enchelyopus</i>	209	<i>Claricola</i>	292
<i>Gadus</i>	209	<i>alabamæ</i>	292
<i>cinnamomi</i> , <i>Acipenser</i> (<i>Antacæus</i>).....	35	<i>julius</i>	292
<i>cincta</i> , <i>Parapetenia tetracantha</i>	419	<i>squamiceps</i>	292
<i>cinctus</i> , <i>Atopichthys</i>	54	<i>whippili</i>	292
<i>Dactyloscopus</i>	454	<i>clarionensis</i> , <i>Centropyge</i>	361
<i>Heros tetracanthus</i>	419	<i>Holacanthus</i>	361
<i>Julis</i>	427	<i>clarionis</i> , <i>Xesurus</i>	363
<i>cinerea</i> , <i>Alutera</i>	496	<i>clarius</i> , <i>Pimelodus</i>	159
<i>Amia</i>	38	<i>clarki</i> , <i>Arctidiellus</i>	377
<i>Etheostoma</i>	290	<i>Blennicottus</i>	377
<i>Perca marina gibbosa</i>	329	<i>Bogostovius</i>	204
<i>cinereum</i> , <i>Aulostoma</i>	245	<i>clarkii</i> , <i>Catostomus</i>	104
<i>cinereus</i> , <i>Aulostomus</i>	245	<i>Notolepidomyzon</i>	104
<i>Gerris</i>	341	<i>Salmo</i>	56
<i>Macrourus</i>	205	<i>clarkii declivifrons</i> , <i>Salmo</i>	57
<i>Macrurus</i>	205	<i>clarkii jordanii</i> , <i>Salmo</i>	57
<i>Microspathodon</i>	416	<i>clarkii taboensis</i> , <i>Salmo</i>	57
<i>Mugil</i>	341	<i>Clarkina</i>	118
<i>Notemotus</i>	290	<i>caurina</i>	118
<i>pellatus</i> , <i>Turdus</i>	341	<i>clarus</i> , <i>Vigil</i>	288
<i>cingulatus</i> , <i>Auchenopterus</i>	457	<i>clathrata</i> , <i>Labrax</i>	317
<i>Fundulus</i>	177	<i>Paralabrax</i>	317
<i>Pomacanthus</i>	360	<i>clavata</i> , <i>Raja</i>	25
<i>Zygonectes</i>	177	<i>claviformis</i> , <i>Moostoma</i>	109
<i>cingulus</i> , <i>Atopichthys</i>	54		

List of scientific names—Continued

	Page		Page
claviger Cottus.....	381	Clupea macrophthalma.....	43
Enophrus.....	381	mattowacca.....	42
clavilatum, Acutomentum.....	366	mediocris.....	41, 42
clavilatus, Sebastodes.....	366	megalops.....	42
claytoni, Gobius.....	439	membras.....	41
Rhinogobius.....	439	menhaden.....	44
clepsydra, Murena.....	90	micropus.....	46
Clepticus.....	426	minima.....	41
parre.....	426	mirabilis.....	41
clevelandi, Phoxinus.....	118	neglecta.....	44
Clevelandia.....	445	pallasi.....	41
ios.....	445	parvula.....	41, 42
longipinnis.....	445	pulchardus.....	42
rosea.....	445	pseudoharengus.....	42
Clmidae.....	455, 464	pusilla.....	42
Clinocottus.....	390	sadina.....	46
analis.....	390	sapidissima.....	42
australis.....	390	sternica.....	99
australis.....	390	stolifera.....	44
sabarra.....	390	thrissa.....	44
Clinostomus.....	120	thrissina.....	44
affinis.....	121	tyrannus.....	44
elongatus.....	120	vernalis.....	42
funduloides.....	121	villosa.....	66
hydrophlox.....	119	virescens.....	42
margarta.....	116	vittata.....	41
montanus.....	119	clupeiformis, Coregonus.....	62, 64, 65
pandora.....	120	Salmo.....	64
phlegothontis.....	121	clupeiformis albus, Coregonus.....	65
tenuis.....	119	Clupeidae.....	41
vandoisulus.....	121	clupeiformis, Argyrosomus.....	61
clintoni, Lepidosteus.....	37	Coregonus.....	62
Clinus aculeatus.....	470	Clupeoides.....	40
binaculatus.....	460	clupeoides, Anchovia.....	50
canariensis.....	459	Engraulis.....	50
capillatus.....	459	clupeola, Harengula.....	43
delalandi.....	459	Clypeocottus.....	381
evides.....	458	robustus.....	381
gilli.....	460	Cobitis heteroclitia.....	175
gobio.....	455, 466	killifish.....	176
macrocephalus.....	455	macrolepidota.....	176
maculatus.....	470	majalis.....	176
medius.....	470	tiaroga.....	143
mohri.....	471	coccineus, Lycodes.....	476
nebulosus.....	471	Scarus.....	432
nigripinnis.....	456	Coccogenia.....	127
nuchipinnis.....	459	coccogenis, Coccotis.....	127
ocellatus.....	459	Hypsilepis.....	127
ocellifer.....	455	cocoi, Rhinocopelus.....	169
pectinifer.....	459	Scopelus.....	168, 169
praecisus.....	470	Coccolus.....	127
unimaculatus.....	470	annectens.....	217
zonifer.....	469	Coccotis.....	127
Clloia.....	144	brimleyi.....	127
camura.....	129	coccogenis.....	127
chlora.....	135	macdonaldi.....	127
montiregis.....	131	zonatus.....	128
smithi.....	145	zonistius.....	128
velox.....	145	Cochlognathus.....	146
vivax.....	145	biguttatus.....	146
clippertonense, Ostracion.....	496	ornatus.....	146
clodalis, Hiodon.....	40	coccensis, Cotylopus.....	448
Clodipterus.....	296	cocosa, Echinidna.....	94
Clupanodon.....	44	Codoma.....	143
Clupea.....	41	chloristia.....	129
aestivalis.....	42	eurystoma.....	130
africana.....	44	ornata.....	143
alosa.....	42	topea.....	143
anchovia.....	43	trichroistia.....	130
atherinoides.....	51	vittata.....	143
brasiliensis.....	40	oceca, Murana.....	84
caerulea.....	41	Ocecia.....	84
carolinensis.....	44	brandieriana.....	84
elongata.....	41	Ocellophus.....	87
encrasichelus.....	47	compar.....	87
fasciata.....	42	Oceula basanium.....	86
halec.....	41	oceus, Gastrobranchus.....	8
harengus.....	41	ocolepis, Centroscymnus.....	21
var.....	41	coelestinus, Pseudoscarus.....	435
heterura.....	46	Scarus.....	207
hudsonia.....	132	Ocelorhynchus.....	207
indigena.....	42	caribbeus.....	207
lamprotaenia.....	46	carminatus.....	207
leachi.....	41	laville.....	207
lineolata.....	41	occa.....	207
macrocephala.....	40		

List of scientific names—Continued

	Page		Page
Ceolorhynchus scaphopsis.....	207	Compsura.....	95
(Ceolorhynchus) scaphopsis, Macrurus.....	207	gorgonæ.....	96
cenicola, Pœcilia.....	175	heterura.....	95
cœnosus, Pimelodus.....	153	concatenatus, Ostracion.....	496
Pleuromichthys.....	225	concentricus, Urobatis.....	30
cœrulea, Tautoga.....	424	Conchognathus.....	78
cœruleopunctata, Acara.....	423	grimaldi.....	78
Æquidens.....	423	conchorum, Meridia.....	247
cœruleus, Cyclopterus.....	397	conchos, Catostomus.....	107
cœrulo-aureus, Har, e.....	425	concinus, Ambiodon.....	352
cœrulopunctata, Acara.....	423	Gasterosteus.....	239
cognatus, Cottus.....	385, 336	cœncolor, Ammocetes.....	9
Cogrus.....	87	Chromitra.....	237
maculatus.....	87	Euschistodus.....	414
Coleoptera.....	66	Gymnothorax.....	93
colesi, Scorpena.....	370	Ichthyomyzon.....	9
colias, Scomber.....	258	Lycodes.....	475
colinus, Gadus.....	209	Nebrius.....	94
Colinus.....	145	Nexilarius.....	414
parietalis.....	145	Scomberomorus.....	257
collapsus, Moxostoma.....	110	Thyreoidæa.....	92, 93
collapsus, Ptychostomus.....	110	conductor, Centronotus.....	278
collaris, Chætodon.....	359	conferrum, Oresoma.....	232
Colletia.....	163	confertus, Hyborhynchus.....	145
nocturna.....	168	Pimephales.....	145
rafinœquei.....	402	confinis, Pimelodus.....	154
colleti, Careproctus.....	33	Salmo.....	59
collei, Chimæra.....	33	confluentus, Fundulus.....	177
Hydrolagus.....	33	Salmo.....	55
Colocephali.....	90	conformis, Lavinia.....	120
Cololabis.....	188	Siboma.....	120
brevirostris.....	188	Conger.....	79
salra.....	188	anago.....	80
colombianus, Heterandria.....	189	analis.....	80
Colomesus.....	500	brasiliensis.....	78
psittacus.....	500	caudicula.....	79
colpense, Zestidium.....	347	caudilimlatus.....	79
colonensis, Stellifer.....	347	conger.....	79
colonus, Serranus.....	320	esculentus.....	79
colorado, Liza.....	494	impressus.....	80
Lutjanus.....	326	limbatus.....	78
Lutjanus.....	326	macrops.....	79
Perry.....	425	microstomus.....	80
coloratum, Aulostoma.....	245	mordax.....	88
Colotrigla.....	406	obtusa.....	79
stearnsi.....	406	opisthoptalmus.....	80
Colpichthys.....	252	orbignyanus.....	79
regis.....	252	rubescens.....	79
coluber, Gempylus.....	261	verreauxi.....	79
colubrinus, Stomias.....	69	verus.....	79
Columbia.....	215	vulgaris.....	79
transmontana.....	215	conger, Conger.....	79
columbianus, Pantosteus.....	105	Muraena.....	79
Rutilus.....	122	conger brasiliensis, Muraena seu.....	94
Siphateles.....	122	Congermuraena.....	79
Vomer.....	275	caudalis.....	80
columna, Rhinobatus.....	23	fiava.....	80
comatus, Exocoetus.....	201	guppil.....	80
comes, Roccus.....	355	congestum, Moxostoma.....	110
comifer, Achirus.....	230	congestus, Catostomus.....	110
commersonianus, Lophius.....	505	Congrammus.....	454
commersoni, Ovum.....	499	moorei.....	454
commersoni, Arius.....	149	Congrellus.....	79
Carcharias.....	15	gilberti.....	80
Catostomus.....	105	Congridæ.....	79
Cyprinus.....	105	Congrosoma.....	80
Fistularia.....	245	evermanni.....	80
Squalus.....	15	Congrus.....	79
communis, Catostomus.....	105	curridens.....	79
Dasybatus.....	24	coniceps, Murænesox.....	156
Pogonichthys.....	133	conirostris, Pimelodus.....	156
compar, Cœliophis.....	87	connectens, Lethops.....	447
complanata, Moniana.....	131	Conocara.....	53
compressa, Anchoviella.....	50	macdonaldi.....	53
compressus, A pareiodon.....	100	macroptera.....	53
Bassozetus.....	451	conocephala, Gila.....	112
Bathynectes.....	480, 451	conocephalus, Mylopharodon.....	112
Callyodon.....	454	Conodon.....	333
Engraulis.....	47, 50	antillanus.....	333
Gadus.....	212	nobilis.....	333
Hemigrammus.....	97	serifer.....	333
Nauclerus.....	278	Conorhynchichthys.....	156
Rutilus.....	128	Conorhynchus.....	156
Scarus.....	454	nelsoni.....	156
		Conorhynchus.....	39, 156

List of scientific names—Continued

	Page		Page
conorhynchus Albula	39, 40	Coregonus spilonotus	65
Gymnelis	477	stanleyi	65
consocium, Chirostoma	251	thymaloides	66
conspersa, Gila	120	thymallus	64
conspersus, Gymnothorax	92	tulibee	61
Serranus	311	bisselli	63
constanciae, Pomatomichthys	296	williamsoni	65
constellatus, Chiroptis	373	cismontanus	65
Platophrys	217	(Coregonus) harengus, Salmo	62
Sebastichthys	367	lucidus, Salmo	62
Sebastomus	367	sapidissimus, Salmo	64
consuetus, Salmo	54	tulibee, Salmo	64
continuum, Hæmulon	329	coregonus, Morostoma	110
contractus, Chonophorus	442	Ptychostomus	110
conus, Morostoma	111	coretta, Thunnus	260
Ptychostomus	111	Tynnus	260
convergens, Macrourus	206	cornaceus, Eleutheractis	316
Macrurus	206	Rypticus	316
convexifrons, Alepocephalus	52	Corida	426
Pomotis	300	cornus, Hexanchus	11
cookianus, Aphododerus	215	corneta, Fistularia	245
cooperi, Cheonda	118	cornubiensis, Fimelepterus	268
Metanops	218	Rhombus levis	228
Metaponops	218	cornubicus, Squalus	19
Rafa	26	cornutus, Anoplogaster	233
Salmo	55	Chaetodon	362
copei, Acipenser (Huso)	35	Chologaster	194
Aleposomus	53	Cyclichthys	502
Chasmistes	108	Cyprinus	128
Cheonda	119	Holacanthus	361
Cottus	386	Hoplostethus	263
Paraliparis	405	Luxilus	123, 128
Lepidosteus	37	Pleuronectes	225
Leuciscus	119	Silurus	245, 246
Squalius	119	Zanclus	362
Xenodermichthys	53	cornutus cerasinus, Hypsilepis	128
Copelandellus	294	chrysocephalus, Luxilis	128
copelandi, Cottogaster	285	cyaneus, Hypsilepis	128
Rheocrypta	285	gibbus, Hypsilepis	128
Coplandia	304	isolepis, Luxilis	128
eriarcha	304	Notropis	128
copli, Leuciscus	123	Coro coro	333
Corallicola	456	coro, Coro	333
altivelis	456	Sciæna	333
marmoratum	456	coroides, Umbrina	350
nigripinna	456	coronata, Seriola	280
coralina, Narcine brasiliensis	27	coronatus, Cyclopterus	397
corallinum, Cryptotrema	456	Petrometopon cruentatus	308
corallinus, A. lartedius	377	Serranus	308
Antennarius	506	corporalis, Cyprinus	117
Corallozetus	484	Leucosomus	117
cardone	484	corpore variegato, Truttæ	55
Cordylus	257	cortezianus, Aprodon	474
Coregoni	64	coruscans, Arctozenus	165
Coregonidæ	61	Paralepis	165
coregonoides, Paralepis	165	coruscum, Holocentrum	236
Coregonus	64	coruscus, Hologentrus	236
abyssicola	66	Corvina	348
albus	65	acutirostris	346
angusticeps	136	adusta	347
artedi	62, 63	biloba	349
clupeaformis	62, 64, 65	chrysoleuca	346
albus	65	deliciosa	348
clupeiformis	62	dentex	344
couesi	65	fulgens	345
coulteri	65	(Homoprion) furthi	347
hoyi	62	(Johnius) jacobii	348
kennicottii	65	macrops	345
labradoricus	64	microps	346
lavaretus	64	ophioscion	347
lauretæ	62	oxyptera	320
nelsonii	65	richardsoni	352
neohantoni	64	ronchus	346
nigripinnis	63	stearnsi	348
novæ-angliæ	65	stellifera	346
oregonus	65	subæqualis	345
osmeriformis	62	trispinosa	346
paralepis	165	vermicularis	348
preblei	65	corvinaeforme, Hæmulon	333
prognathus	63	corvinaeformis, Brachydeuterus	333
pusillus	62	Hæmulon	333
quadrilaterialis	65	Corvula	345
richardsoni	64	batabana	345
ruber	163	sanctæ-luciæ	345
signifer	66		

List of scientific names—Continued

	Page		Page
<i>Corvula sialis</i>	345	<i>Cottus bubalis</i>	386
<i>Corynolophus</i>	509	<i>cataphractus</i>	395
<i>reinhardtii</i>	509	<i>cephaloides</i>	389
<i>Coryphæna</i>	264	<i>chaetodon</i>	487
<i>argyrurus</i>	265	<i>chamberlaini</i>	385
<i>aurata</i>	265	<i>claviger</i>	381
<i>azorica</i>	265	<i>cognatus</i>	385, 386
<i>caerulea</i>	434	<i>copel</i>	386
<i>dolfin</i>	264	<i>criniger</i>	390
<i>dorado</i>	264	<i>decastrensis</i>	387
<i>equisetis</i>	265	<i>dicerus</i>	381
<i>fasciolata</i>	264	<i>elegans</i>	381
<i>hippurus</i>	264	<i>evermanni</i>	383
<i>immaculata</i>	264	<i>formosus</i>	386
<i>imperialis</i>	264	<i>franklini</i>	385
<i>lessonii</i>	265	<i>glaber</i>	487
<i>lineata</i>	430	<i>glacialis</i>	387
<i>marcgravi</i>	264	<i>gobio</i>	383
<i>nigrescens</i>	317	<i>gobioides</i>	385
<i>novacula</i>	430	<i>gracilis</i>	383, 385
<i>percliformis</i>	267, 268	<i>cayuga</i>	386
<i>plumieri</i>	267	<i>greeni</i>	384
<i>pomplius</i>	267	<i>greenlandicus</i>	386, 387
<i>psittacus</i>	430, 433	<i>gulosus</i>	383
<i>rupestris</i>	205	<i>hemilepidotus</i>	380, 381
<i>scomberoides</i>	265	<i>hexacornis</i>	388
<i>sueri</i>	264	<i>hispius</i>	383
<i>velifera</i>	268	<i>humilis</i>	387
<i>virgata</i>	265	<i>ictalops</i>	385
<i>viamingii</i>	265	<i>indicus</i>	397
<i>Coryphændæ</i>	264	<i>jack</i>	387
<i>Coryphæniformes</i>	264	<i>klamathensis</i>	384
<i>Coryphænoides</i>	204	<i>lelopomus</i>	385
<i>carpinus</i>	205	<i>macrops</i>	384
<i>filicauda</i>	207	<i>maculatus</i>	386
<i>rupestris</i>	204, 205	<i>marginatus</i>	384
<i>sulcatus</i>	207	<i>marmoratus</i>	387
<i>Coryphænoididæ</i>	202	<i>meridionalis</i>	385
<i>Coryphopterus</i>	439	<i>mertensii</i>	387
<i>glaucofrænum</i>	439	<i>mitchilli</i>	386
<i>Corythoichthys</i>	243	<i>monopterygius</i>	397
<i>albirostris</i>	243	<i>niger</i>	387
<i>cayorum</i>	243	<i>nigricans</i>	386
<i>cosmopolita, Micropteryx</i>	276	<i>octodecem-spinosus</i>	386
<i>Cossyphus</i>	425	<i>onychus</i>	386
<i>puellaris</i>	425	<i>pachypus</i>	386
<i>pulchellus</i>	425	<i>perplexus</i>	384
<i>costalesi, Smaragdus</i>	441	<i>philonips</i>	384
<i>costaricensis, Gobiosox</i>	439	<i>pistilliger</i>	389
<i>costatus, Bodianus</i>	349	<i>platycephalus</i>	387, 388
<i>Cottidæ</i>	383	<i>polaris</i>	388
<i>Cottogaster</i>	285	<i>porosus</i>	387
<i>aurantiacus</i>	234, 285	<i>polyacanthocephalus</i>	387
<i>cheneyi</i>	286	<i>princeps</i>	384
<i>copelandi</i>	285	<i>punctulatus</i>	383
<i>putnami</i>	285	<i>quadricornis</i>	388
<i>uranides</i>	285	<i>rhothea</i>	383
(<i>Cottogaster</i>) <i>uranides, Etheostoma</i>	285	<i>ricei</i>	383, 386
<i>cottoides, Bathygadus</i>	203	<i>richardsoni</i>	385
<i>Cottopsis</i>	383	<i>scorpioides</i>	386
<i>gulosus</i>	383	<i>scorpius</i>	386
<i>parvus</i>	383	<i>semiscaber</i>	384
<i>ricei</i>	383, 386	<i>shasta</i>	383
<i>semiscaber</i>	384	<i>stelleri</i>	381
<i>Cottunculus</i>	392	<i>tænipterus</i>	388
<i>microps</i>	392	<i>tenuis</i>	384
<i>thomsonii</i>	392	<i>thomsonii</i>	392
<i>torvus</i>	392	<i>trachurus</i>	381
<i>Cottus</i>	383	<i>tricuspis</i>	389
<i>acadian</i>	393	<i>tripterygius</i>	393
<i>æneus</i>	386	<i>uncinatus</i>	377
<i>aleuticus</i>	385	<i>ventralis</i>	389
<i>alvodii</i>	385	<i>verrucosus</i>	387
<i>anceps</i>	386	<i>villosus</i>	393
<i>annæ</i>	384	<i>viscosus</i>	385
<i>asper</i>	383	<i>wilsoni</i>	385
<i>asperrima</i>	384	<i>Cotyli nigripinnis</i>	439
<i>aspririmus</i>	384	<i>Cotylopus</i>	447
<i>bairdi</i>	383	<i>acutipinnis</i>	447
<i>bairdii</i>	385	<i>coccensis</i>	448
<i>beldingii</i>	384	<i>punctatus</i>	448
<i>bendirei</i>	384	<i>salvini</i>	448
<i>bicornis</i>	378	<i>couaga, Chaetodon</i>	362
<i>boleoides</i>	385	<i>couchi, Dionda</i>	144
<i>brodamus</i>	395	<i>montana</i>	181

List of scientific names—Continued

	Page		Page
Couchia.....	208	crinitus, Alectis.....	274
minor.....	208	Blennius.....	461
couchiana, Limia.....	193	Zeus.....	274
couchianus, Platypoecilus.....	193	cristagalli, Anoploarchus.....	468
couchii, Serranus.....	308	archolepis, Anoploarchus.....	468
couesi, Coregonus.....	65	Cebedichthys.....	466
Prosopium.....	65	Centronotus.....	468
couesii, Apocope.....	141	cristata, Chimera.....	32
Cryptosaras.....	508	cristatus, Blennius.....	461
Couesius.....	136	Pimelodus.....	158
adustus.....	137	cristiceps, Melamphaes.....	233
dissimilis.....	137	Plectromus.....	233
greeni.....	137	cristagalli, Gobius.....	438
plumbeus.....	136	crista-galli, Gobius.....	438
tribrilateralis.....	116	Cristivomer.....	59
Couesius squamulentus.....	136	namaycush.....	59
coulteri, Coregonus.....	65	siscowet.....	60
Prosopium.....	65	cristulata, Scorpena.....	372
courbina, Pogonathus.....	352	cristulatus, Gymnoclonus.....	466
Pogonias.....	352	Crius.....	268
courtadei, Serranus.....	310	bertheloti.....	268
cragini, Amiurus.....	154	croceus, Leuciscus.....	140
Etheostoma.....	291	Rhynchichthys.....	140
Nivicola.....	291	crocodinus, Brachysomophis.....	89
cramerii, Eosebastes.....	367	Ophichthys.....	89
Hybopsis.....	138	crocata, Plectropoma.....	321
Leuresthes.....	251	crocotus, Hypoplectrus unicolor.....	321
Oregonichthys.....	138	croco, Pristipoma.....	334
Sebastodes.....	367	Rhonciscus.....	334
Umbra.....	174	croicensis, Searus.....	433, 434
crassa, Belone.....	196	croker, Sciæna.....	349
Tigoma.....	120	Cromileptes.....	310
crassicauda, Lavina.....	120	cromis, Labrus.....	352
Siboma.....	120	Pogonias.....	352
crassiceps, Cryptotomus.....	431	Crossias.....	389
Plectromus.....	233	albomaculatus.....	389
Scopelus.....	233	allisi.....	389
crassilabre, Moxostoma.....	111	beringi.....	389
crassilabris, Embryx.....	474	camtschaticus.....	389
Geophagus.....	424	crossotus, Etropus.....	219, 220
(Satanoperca).....	424	crotalinus, Embryx.....	473
Lycodopsis.....	474	Lycodopsis.....	473
Ptychostomus.....	111	Crotalopsis punctifer.....	88
crassus, Alvordius.....	283	crotaphus, Julis.....	426, 427
Esox.....	173	Crotolopsis.....	88
Lepidosteus.....	36	quintifer.....	88
craticula, Zygometes.....	178	crudelis, Eurymyctera.....	91
Cratinus.....	317	cruentatus, Entomacrodus.....	463
agassizii.....	317	Labrus.....	322
Crayracion.....	498	Petrometopon.....	308
Creagrutus.....	98	Priacanthus.....	322
notropoides.....	98	Sparus.....	308
simus.....	98	cruentatus coronatus, Petrometopon.....	308
creaseri, Ruscariops.....	378	cruentifer, Omochelys.....	86
crebripunctata, Pteroplatea.....	29	Pisodonophis.....	86
Cremnobates.....	456	crumenophthalmus, Trachurops.....	271
affinis.....	457	crumenophthalmus, Scomber.....	270, 271
altivelis.....	456	crureum, Xiphidium.....	469
fasciatus.....	456, 457	crureus, Squalus.....	119
integripinnis.....	457	crustosum, Peristedion.....	406
marmoratus.....	456	Peristedion.....	406
nox.....	457	Cryptacanthodes.....	471
cremnobates, Labrosomus.....	456	inornatus.....	471
Starksia.....	456	maculatus.....	471
Crenidens simplex.....	339	Cryptacanthodidae.....	471
Crenilabrus.....	425	Cryptops.....	101
microstoma.....	424	Cryptopterenchelys.....	85
crenulate, Myctophum.....	170	frontalis.....	85
crenularis, Tarteletbeania.....	170	puncticeps.....	85
Creolus.....	320	(Cryptopterus) frontalis, Ophichthys.....	85
creolus, Girardinus.....	159	Cryptopterus.....	85
Serranus.....	320	puncticeps.....	85
Cretotretes.....	259	puncticeps.....	85
crescentale, Gobiosoma.....	446	Cryptosaras.....	508
crescentalis, Gobiosoma.....	446	couesi.....	508
Pomacanthus.....	361	cryptosus, Stromateus.....	267
crescentus, Salmo.....	57	Cryptotomus.....	431
Salmo gairdneri.....	57	auropunctatus.....	431
crestonis, Acanthurus.....	363	beryllinus.....	431
Teuthis.....	363	crassiceps.....	431
cretensis, Labrus.....	431	dentilis.....	431
cuniger, Cottus.....	330	retractus.....	431
cunigerum, Siphostoma.....	241	roseus.....	431
Syngnathus.....	241	ustus.....	431

List of scientific names—Continued

	Page		Page
Cryptotrema.....	456	Curimata.....	100
corallinum.....	456	magdalena.....	100
crysoleucas, Cyprinus.....	114, 115	Curimatella.....	100
Notemigonus.....	115	lepidurus.....	100
crysos, Paratractus.....	272	Curimates, Les.....	100
Scomber.....	272	Curimatidae.....	100
Crystallaria.....	282, 288	Curimatus.....	100
asprella.....	288	magdalena.....	100
Crystallichthys.....	401	spilurus.....	100
cyclopius.....	401	curta, Anchoviella.....	48
cyclostigma.....	401	curtiss-jamesi, Harriotta.....	33
mirabilis.....	401	Polistotrema.....	8
crystallina, Thyrina.....	248	curtus, Stolephorus.....	48
crystallinum, Chirostoma.....	251	Vomer.....	275
Ctenochætus.....	363	curvidens, Congrus.....	78
striatus.....	363	curvilineata, Muranophis.....	91
Ctenodax.....	269	curvus, Tetradon.....	497
williamsi.....	269	cuspidata, Aluterus.....	495
Ctenodon.....	363, 470	cuspidatus, Macrourus.....	206
ctenodon, Acanthurus.....	363	Macurus.....	206
Ctenogobius.....	438	cutsanserinus, Carpiodes.....	103
fasciatus.....	438	cuvieri, Ausonia.....	264
tortuga.....	439	Centropomus.....	306
Ctenolabrus bergall.....	424	Gasterosteus.....	237
uninotatus.....	424	Gnathypops.....	453
Ctenolucius.....	100	Opistognathus.....	453
beani.....	100	Tetragonurus.....	269
insculptus.....	100	cuzamilla, Scarus.....	434
cube, Vomer.....	275	cyanea, Furcaria.....	412
cubana, Anchoviella.....	48	cyanella, Mugilops.....	252
Anguilla.....	77	cyanellus, Apomotis.....	298, 299
Cubanichthys.....	174	Ichthyobus.....	102
cubensis.....	174	Lepomis.....	298
cubanus, Engraulis.....	48	Mugilops.....	252
Epinephelus.....	311	cyaneus, Hypsilepis cornutus.....	128
cubensis, Channomuraena.....	95	Cyanichthys.....	502
Cubanichthys.....	174	cærulesus.....	502
Fundulus.....	174	cyanocephalus, Iridio.....	427
Heterandria.....	188	Labrus.....	426, 427
Hynnis.....	274	Lythrurus.....	126
Limia.....	191	cyanoguttatus, Herlichthys.....	422
Toxus.....	188	cyanolene, Sparisoma.....	432
Vomer septapinnis.....	275	cyanonoton, Aloa.....	42
cubera, Lutjanus.....	325	cyanophrys, Naucrates.....	278
Cubiceps multiradiatus.....	268	Psenes.....	268
cubicus, Ostracion.....	496	cyanops, Caulolatilus.....	357
cubifrons, Malthæ.....	507	Cyanopterus, Cypselurus.....	202
cucuri, Prionodon.....	16	Exocoetus.....	202
Cucuyo.....	493	Leuichthys.....	63
Cugupuguacu.....	312	Lutjanus.....	325
cujus, Chasmistes.....	108	Mesoprion.....	325
culebra, Actis.....	460	cyanostigma, Julis.....	426
Malacoctenus.....	460	Cybidae.....	257
Culius.....	436	Cybium.....	257
sequidens.....	437	acervum.....	257
amblyopsis.....	437	caballa.....	277
fuscus.....	436	cavalla.....	257
perniger.....	437	commersoni.....	257
(Culius) belizianus, Electris.....	436	immaculatum.....	257
cultiferum, Pristipoma.....	334	petus.....	256
cutrata, Anchoviella.....	45	sara.....	256
cutratum, Petalosoma.....	193	solandri.....	256
cutratus, Allaro.....	193	verany.....	256
Stolephorus.....	48	Cycloptus.....	101
cuttrons, Aluterus.....	495	elongatus.....	101
culveri, Trachinotus.....	277	nigrescens.....	101
cumberlandicum, Catonotus fiabellare.....	238	Cyclichthys.....	501
Etheostoma.....	238	antillarum.....	502
cumingii, Ceratichthys.....	138	attenuatus.....	502
Oregonichthys.....	138	priareos.....	502
cummingsi, Hybopsis.....	138	cornutus.....	502
Notropis.....	138	fuliginosus.....	502
cuneata, Mollenesia spheonops.....	198	schœphi.....	502
Poecilia.....	198	spinosus.....	502
cupido, Geophagus.....	424	Cyclogaster.....	399
cupresoides, Pimelodus.....	153	bristolensis.....	399
cupreus, Pimelodus.....	153	megacephalus.....	400
Silurus.....	153	(Neoliparis) beringianus.....	399
Trachinotus.....	277	micraspidophorus.....	399
curacao, Gobius.....	439	Cyclogobius.....	398, 399
Rhinocephalus.....	439	Cyclogobius.....	444
curassavicus, Balistes.....	493	cyclolepis, Dolloa.....	203
curema, Mugil.....	253	Microgobius.....	443
Querimana.....	253	Nematonurus.....	203
cutilus, Salmo.....	61	Cyclonarcæ.....	27

List of scientific names—Continued

	Page		Page
Cyclopium	159	Cynoperca	282
humboldtii	159	canadensis	282
purrense	160	grisea	282
cyclopomatus, Serranus	314	Cynoscion	354
Cyclopoeta	220	acoupa	354
chittendeni	221	jamaicensis	355
fimbriata	220	lelarchus	355
querna	221	macdonaldi	355
Cyclopterichthys	398	microlepidotus	355
glaber	398	nothus	355
Cyclopteridæ	397	obliquatus	355
Cyclopteroides	398	othonopteron	355
gyrinops	398	othonopterus	355
Cyclopterus	397	parvipinnis	356
calyodon	399	phoxocephalus	356
coeruleus	397	regalis	355
corenatus	397	squamipinnis	355
gelatinosus	402	thalassinus	355
lineatus	400	xanthulum	356
liparis	399, 400	cypho, Catostomus	108
major	401	Esox	173
minor	400	Cyphosus	339
lumpus	397	elegans	339
minutus	397	cyprinaceus, Chaetodon	340
musculus	400	Cyprinella	130, 131
nudus	459	analostana	129
orbis	397	beckwithi	130
pavoninus	397	billingsiana	131
pyramidatus	397	bubalina	130
spinosus	397	callura	129
stelleri	398	cercostigma	130
ventricosus	398	forbesi	131
Cycloptus lumpus	397	forlonensis	131
cycloptum, Pimeodus	159	gunnisoni	130
cyclops, Liparis	400	lepida	130
cyclopius, Crystallichthys	401	ludibunda	130
cyclostigma, Crystallichthys	401	lugubris	131
Liparis	401	luxioides	131
Cyclostoml	8	macrostoma	131
Cyclothone	72	notata	130
acclinidens	72	rubripinna	131
bathypbila	72	suavis	131
elongata	72	texana	130
lusca	72	umbrosa	130
microdon	72	venusta	130
signata	72	whipplu	129
cyclotis, Ceratichthys	137	cyprinella, Chanos	40
cylindraceus, Rivulus	179	Megastomatobus	102
cylindricus, Bascanichthys	56	Sclerogynathus	102, 103
Cylindrosteus	37	Cyprinidæ	112
agassizii	37	Cyprinodon	180
bartoni	37	baileyi	182
castelnaudii	37	bovinus	181
megalops	37	rubrofluvialis	181
platostomus	37	californiensis	181
productus	37	carpio	180
rafinesquii	37	dearborni	181
scabriceps	37	elegans	181
zadockii	37	eximus	181
(Cylindrosteus) oculatus, Lepisosteus	37	felicianus	181
Cymatogaster	409, 410	gibbosus	181
aggregatus	409	latifasciatus	181
ellipticus	410	macularius	181
larkinsi	410	baileyi	182
minimus	409	maria	181
pulchellus	410	mydrus	180
rosaceus	410	nevadensis	181
cymatogramma, Pileoma	286	parvus	175
cymatotenja, Etheostoma (Hadropterus)	285	riverendi	181
Hypohomus	285	rubrofluvialis	181
Cynædus brama	338	variegatus	180, 181
Cynias	13	Cyprinodontes	174
canis	13	Cyprinodontidæ	174
lunulatus	13	cyprinoides, Gobius	438
Cynichthys	310	Lophogobius	438
CynicoGLOSSUS	228	Cyprinus (Abramis) balteatus	117, 118
bathybius	228	(Abramis?) smithi	141
Cynocephalus	17	americanus	115, 350
cyznodon, Mesoprion	325	atromaculatus	117
Cynodonichthys	179	atrasatus	139, 140
tenuis	179, 180	bullaris	117
Cynoglossa	228	catostomus	105, 106
Cynoglossidæ	230	caurinus	118
cynoglossus, Glyptocephalus	228	commersoni	105
Pleuronectes	221, 228	cornutus	128

List of scientific names—Continued

	Page		Page
<i>Cyprinus corporalis</i>	117	<i>dariensis</i> , <i>Apareiodon</i>	100
<i>crysoleucas</i>	114, 115	<i>Hemibrycon</i>	97
<i>hemiplus</i>	115	<i>Oxylocara</i>	162
<i>leuciscus</i>	117	<i>Parodon</i>	100
<i>(Leuciscus) caurinus</i>	118	<i>Sternopygus</i>	101
<i>gracilis</i>	136	<i>Dasibatis</i>	28
<i>oregonensis</i>	113, 114	<i>dasyplatys</i> , <i>Psodonophis</i>	86
<i>maxillingua</i>	146	<i>Dasyatidae</i>	28
<i>megalops</i>	128	<i>Dasyatis</i>	28
<i>melanurus</i>	128	<i>attavilla</i>	29
<i>oblongus</i>	109	<i>guttatus</i>	28
<i>sucetta</i>	109	<i>hastatus</i>	28
<i>teres</i>	105	<i>longus</i>	28
<i>vittatus</i>	140	<i>schmardae</i>	28
<i>cyprinus</i> , <i>Carpiodes</i>	103	<i>torrei</i>	28
<i>Catostomus</i>	103	<i>Dasybatus dipterurus</i>	29
<i>Sclerognathus</i>	102	<i>Dasybatus</i>	28
<i>Cypselurus</i>	201	<i>dasycephalus</i> , <i>Arius</i>	150
<i>bahiensis</i>	202	<i>Galeichthys</i>	150
<i>californicus</i>	202	<i>Dasycoctus</i>	392
<i>callopterus</i>	202	<i>setiger</i>	392
<i>cyanopterus</i>	202	<i>daubentonii</i> , <i>Caranx</i>	273
<i>fuorum</i>	202	<i>Davidia</i>	495
<i>fuscatus</i>	201	<i>punctata</i>	495
<i>gibbifrons</i>	202	<i>davidsoni</i> , <i>Anisotremus</i>	332
<i>heterurus</i>	201	<i>Monacanthus</i>	494
<i>holubi</i>	202	<i>Pristipoma</i>	332
<i>lineatus</i>	202	<i>davisi</i> , <i>Rabula</i>	91
<i>lukeni</i>	201	<i>dayi</i> , <i>Phoxargyrea</i>	247
<i>nigricans</i>	201	<i>deani</i> , <i>Eosebastes</i>	367
<i>spilonotus</i>	202	<i>Paraliparis</i>	404
<i>xenopterus</i>	202	<i>Polistotrema</i>	9
<i>cypselurus</i> , <i>Prognurus</i>	403	<i>Sebastodes</i>	367
<i>Cypsilurus</i>	201	<i>dearborni</i> , <i>Cyprinodon</i>	181
<i>(Cypsilurus) proceus</i> , <i>Exocoetus</i>	201	<i>Decactylus</i>	105
<i>Cyttus hololepis</i>	232	<i>decadactylus</i> , <i>Beryx</i>	234
		<i>decagonus</i> , <i>Agonus</i>	394
		<i>Leptagonus</i>	394
<i>Dacentrus</i>	412	<i>decagrammus</i> , <i>Hevagrammos</i>	373
<i>lucens</i>	412	<i>Labrax</i>	373
<i>Dactylagnus</i>	455	<i>Decapterus</i>	270
<i>mundus</i>	455	<i>hypodus</i>	270
<i>Dactyleptus</i>	467	<i>macarellus</i>	270
<i>Dactylobatus</i>	27	<i>punctatus</i>	270
<i>armatus</i>	27	<i>sanctae-helenae</i>	270
<i>dactyloptera</i> , <i>Scorpaena</i>	370	<i>Decaptus</i>	279
<i>Dactylopteridae</i>	408	<i>decastrensis</i> , <i>Cottus</i>	387
<i>Dactylopterus</i>	408	<i>Decentrus lucens</i>	412
<i>pirapada</i>	409	<i>decimalis</i> , <i>Serranus</i>	314
<i>volitans</i>	409	<i>declivifrons</i> , <i>Abudefduf</i>	415
<i>dactylopterus</i> , <i>Helicolenus</i>	370	<i>Euschistodus</i>	415
<i>Dactyloscopidae</i>	454	<i>Salmo</i>	57
<i>Dactyloscopus</i>	454	<i>clarkii</i>	57
<i>cinctus</i>	454	<i>Vomer</i>	275
<i>lunaticus</i>	455	<i>declivis</i> , <i>Seriola</i>	280
<i>pectoralis</i>	454	<i>Decodon</i>	425
<i>poeyi</i>	455	<i>puellaris</i>	425
<i>tridigitatus</i>	454	<i>decoratum</i> , <i>Bryostemma</i>	466
<i>zelotes</i>	455	<i>decoratus</i> , <i>Entomacrodus</i>	463
<i>Dactylostomias</i>	69	<i>Iridio</i>	427
<i>filifer</i>	69, 70	<i>Promicropterus</i>	317
<i>dactylosus</i> , <i>Paraliparis</i>	404	<i>decoris</i> , <i>Doratonotus</i>	429
<i>Dactor</i>	488	<i>decurrens</i> , <i>Pleuronichthys</i>	225
<i>dekayi</i> , <i>Oxyrhina</i>	19	<i>defensor</i> , <i>Caranx</i>	273
<i>Dajaus</i>	254	<i>Dekaya</i>	357
<i>microps</i>	254	<i>anomala</i>	357
<i>monticola</i>	254	<i>dekayi</i> , <i>Acipenser</i>	34
<i>percoides</i>	254	<i>Gasterosteus</i>	269
<i>dakotensis</i> , <i>Chrosomus</i>	113	<i>Isuropsis</i>	19
<i>dalli</i> , <i>Gobius</i>	442	<i>Phycis</i>	213
<i>Lythrypnus</i>	442	<i>Pimelodus</i>	153
<i>Dallia</i>	172	<i>Scomber</i>	258
<i>pectoralis</i>	172	<i>Syngnathus</i>	242
<i>dallii</i> , <i>Auctospina</i>	369	<i>delalandi</i> , <i>Malacotenus</i>	459
<i>Pteropodus</i>	369	<i>delalandii</i> , <i>Clinus</i>	459
<i>Dallidae</i>	172	<i>delicatissima</i> , <i>Anchoviella</i>	48
<i>Damalichthys</i>	412	<i>Umbra</i>	172
<i>argyrosomus</i>	412	<i>delicatissimus</i> , <i>Engraulis</i>	48
<i>vacca</i>	412	<i>deliciosa</i> , <i>Corvina</i>	348
<i>damalis</i> , <i>Carpiodes</i>	103	<i>Hybopsis</i>	134
<i>danae</i> , <i>Xenopthalmichthys</i>	69	<i>Moniana</i>	184
<i>dariensis</i> , <i>Darienichthys</i>	189	<i>Delolepis</i>	471
<i>Gambusia</i>	189	<i>giganteus</i>	472
<i>Darienichthys</i>	189	<i>virgatus</i>	472
<i>darienensis</i>	189	<i>Delothyrus</i>	220

List of scientific names—Continued

	Page		Page
delphinus, Minomus	104	Diaphus	168
Pantosteus	104	theta	168
Deltarala	25	Diapterus	297, 342
Deltistes	108	aureolus	342
lucatus	108	axillaris	342
dendrica, Ramularia	224	californiensis	341
dendritica, Aneylosetta	224	downi	341
dendriticum, Siphostoma	242	evermanni	342
dendriticus, Syngnathus	242	gracilis	341
denegatus, Pomacentrus	415	homonymus	341
dennyi, Carelliparis	400	leroyi	341
Liparis	400	olisthostomus	342
dentatus, Apsilus	328	peruvianus	342
Grammatostomias	70	diapterus, Amnicthys	297
Lucifuga	479	Furemanus	476
Lutjanus	325	Lycoodes	476
Paralichthys	224	diaz, Chirostoma	251
Pleuronectes	224	Zoogoneticus	182
Stygicola	479	Dibranchius	506
Tropididius	328	atlanticus	506
Upeneus	343	dicerus, Ceratocottus	381
dentex, Brycon	98	Cottus	381
Corvina	344	Dicerobatus	31
Engraulis	51	Dicholomycter	499
Menidia	247	Dicrolene	483
Odontogobius	344	flamentosa	483
Osmorus	67	intronigra	483
denticulatus, Girardinus	189	nigra	483
dentius, Cryptotomus	431	pullata	483
Scarus	431	Dicrotus, parvipinnis	261
denudata, Gonostoma	72	diego, Pneumatophorus	258
denudatum, Gonostoma	72	Scomber	258
deppii, Cichlaurus	416	diencæus, Eupomacentrus	413
Heros	416	difformis, Carpiodes	103
depraudus, Esoc	173	digitatus, Lycoodes	475
depressa, Belone	195	digitatus, Holcomycteronus	482
Fistularia	245	dignabilis, Pseudopleuronectes	227
Rhamdia	168	digrammus, Pleuronectes	226
depressus, Lonchilurus	352	diguetti, Neomugil	254
Derepodichthyidae	479	Tenisonger	82
Derepodichthys	479	dilecta, Notosema	225
alepidotus	479	dilectum, Notosema	225
Derichthyidae	77	dilectus, Alburnus	125
Derichthys	77	Notropis	125
serpentinus	77	Dilobomycter	499
dermatinus, Lycodapus	478	Dimalacocentrus	430
Salmo	54	dimidiatus, Gasterosteus	237
Dermatocheir	97	Julis	427
Dermatolepis	313	Leucus	122
inermis	313	Serranus	314
punctatus	313	Syngnathus	240
zancus	313	Trisotropis	314
Dermatostethus	240	Dinectus truncatus	35
punctipinnis	240	Dinemus	234
desmarestia, Raja	25	venustus	234
detersor, Julis	429	dinemus, Minnilus	123
detrusus, Gillichthys	444	dinoceros, Citharichthys	219
Deuterodon, atricaudata	96	Diodon	501, 502
deweegeri, Prionodes	319	antennatus	502
Serranus (Paralabrax)	319	asper	501
Diabasis	329	atinga	502
flavolineatus	330	brachiatus	501
obliquatus	330	carinatus	503
parra	329	echinus	501
steindachneri	330	4-maculatus	501
Diabolichthys	32	fuliginosus	502
elliotti	32	geometricus	502
Diacope	324	lineatus	502
viridis	324	holacanthus	501
diadema, Scarus	433	holocanthus	501
Diagramma cavifrons	335	hystrix	501
melanosplum	332	liturosus	501
Dialarchus	391	maculato-striatus	502
synderi	391	maculatus var	501
Diana	264	maculifer	501
semilunata	264	melanopsis	501
diaphana, Hydrargyra	178	meuleni	502
Sternoptix	74	mola	503
diaphanes, Raja	25	multimaculatus	501
Raja	25	nirolineatus	502
Diaphanichthys	79	novem-maculatus	501
diaphanus, Calliurus	299	pilosus	501
Zygonectes	178	punctatus	501
Diaphasia	486	reticulatus	502

List of scientific names—Continued

	Page		Page
<i>Diodon rivulatus</i>	502	<i>Dolopichthys</i>	509
<i>schœpfi</i>	502	<i>anisacanthus</i>	509
<i>sex-maculatus</i>	501	<i>gracilispinus</i>	509
<i>spinossissimus</i>	501	<i>heteracanthus</i>	510
<i>spinosus</i>	502	<i>luetkeni</i>	510
<i>verrucosus</i>	502	<i>microlophus</i>	509
<i>Diodon plumier, le</i>	502	<i>dombey, Gastrobranchus</i>	8
<i>Diodontidae</i>	501	<i>dommicensis, Gambusia</i>	186
<i>diomedean, Aphoristia</i>	231	<i>Lima</i>	191
<i>diomedianus, Hoplunnis</i>	81	<i>Platycephalus</i>	191
<i>Symphurus</i>	231	<i>Pœcilia</i>	191
<i>Dionda</i>	141	<i>dominus, Protoporus</i>	119
<i>amara</i>	144	<i>Doncella</i>	426
<i>argentosa</i>	144	<i>dorado, Coryphæna</i>	264
<i>chrysitus</i>	144	<i>Doratonotus</i>	429
<i>couchi</i>	144	<i>bœkei</i>	429
<i>episcopa</i>	144	<i>decorus</i>	429
<i>grisea</i>	144	<i>megalepis</i>	429
<i>melanops</i>	144	<i>thalassinus</i>	429
<i>nubila</i>	144	<i>Dorinitator</i>	436
<i>papalis</i>	144	<i>gundlachi</i>	436
<i>plumbea</i>	144	<i>lineatus</i>	436
<i>serena</i>	144	<i>maculatus</i>	436
<i>spadicea</i>	144	<i>microphthalmus</i>	436
<i>texensis</i>	144	<i>dormitator, Platycephalus</i>	435
<i>Dioplites</i>	298	<i>dormitatrix, Eleotris</i>	435
<i>variabilis</i>	298	<i>dormitor, Gobiomorus</i>	435
<i>Diplanchnias</i>	503	<i>Dorosoma</i>	46
<i>ncsus</i>	503	<i>cepedianum</i>	46
<i>Diplectrum</i>	318	<i>exile</i>	46
<i>euryplectrum</i>	318	<i>insociabilis</i>	46
<i>formosus</i>	318	<i>notata</i>	46
<i>macropoma</i>	318	<i>Dorosomidae</i>	46
<i>pacificum</i>	318	<i>dorsale, Hæmulon</i>	330
<i>radiale</i>	318	<i>dorsalis, Carangoides</i>	274
<i>sciurus</i>	318	<i>Citula</i>	274
<i>diplema, Semotilus</i>	117	<i>Halatractus</i>	279
<i>Dipleson</i>	286	<i>Hybopsis</i>	134
<i>fasciatus</i>	293	<i>Hypsypops</i>	416
<i>Diplodus</i>	338	<i>Macurus</i>	207
<i>argenteus</i>	338	<i>Mustelus</i>	13
<i>holbrookii</i>	338	<i>Semotilus</i>	117
<i>sargus</i>	338	<i>Seriola</i>	279
<i>diploproa, Eosebastes</i>	367	<i>Stegastes</i>	416
<i>Sebastichthys</i>	367	<i>Umbrina</i>	350
<i>Diplopondyli</i>	11	<i>Vomer</i>	275
<i>diplotænia, Belone</i>	196	<i>dorsatum, Ciclasoma</i>	420
<i>Bodianus</i>	425	<i>dorsatus, Astatheros</i>	420
<i>Harpe</i>	425	<i>Petromyzon marinus</i>	9
<i>Strongylura</i>	196	<i>dorsomaculata, Girella</i>	339
<i>Dipterodon</i>	295	<i>dorso-punctatus, Pomacentrus</i>	413
<i>dipterurus, Amphotistius</i>	29	<i>Dorsuanus</i>	339
<i>Dasybatus</i>	29	<i>Dorychthys</i>	243
<i>Dipturus</i>	24	<i>lineatus</i>	243
<i>dipus, Milordesmus</i>	473	<i>Doryrhamphus</i>	243
<i>discebolus, Catostomus</i>	107	<i>californiensis</i>	243
<i>Discocephali</i>	448	<i>sierra</i>	243
<i>Discopyge</i>	27	<i>dovu, Apogon</i>	295
<i>ommata</i>	27	<i>Gambusia</i>	186
<i>dispar, Scopelengys</i>	170	<i>Gymnothorax</i>	92
<i>Zygonectes</i>	178	<i>Haplochilus</i>	179
<i>dispius, Iridio</i>	427	<i>Heros</i>	418
<i>PlatyGLOSSUS</i>	427	<i>Murena</i>	92
<i>dispiurus, Centropomus</i>	320	<i>Opisthopterus</i>	45
<i>dissimilis, Couesius</i>	137	<i>Oxyzygonectes</i>	179
<i>Erismystax</i>	138	<i>Paraconodon</i>	332
<i>Leucosomus</i>	137	<i>Parapetema</i>	418
<i>Luxilus</i>	138	<i>Pœcilia</i>	192
<i>distinctum, Sparisoma</i>	432	<i>Fristigaster</i>	45
<i>distictus, Scarus</i>	432	<i>Fristipoma</i>	332
<i>Ditrema atripes</i>	411	<i>dowel, Anableps</i>	194
<i>brevipinnis</i>	409	<i>dow, Lepturus</i>	148
<i>orthonotus</i>	411	<i>Thalassophryne</i>	498
<i>Dixonina</i>	40	<i>dowi, Diapterus</i>	341
<i>nemoptera</i>	40	<i>Exocoetus</i>	201
<i>dodecadæron, Oœcilia</i>	394	<i>Selenaspis</i>	148
<i>dodecadærus, Agonus</i>	394	<i>Dracometta</i>	450
<i>dolfin, Coryphæna</i>	264	<i>acanthopoma</i>	450
<i>dolichocephalus, Gobius</i>	442	<i>Dracomettidae</i>	450
<i>dolichogaster, Blennius</i>	468	<i>Drepanopsetta</i>	222
<i>Rhodymenichthys</i>	468	<i>Dropsarus</i>	208
<i>Doliodon</i>	276	<i>drummond-hayi, Epinephelus</i>	312
<i>Dolloa</i>	203	<i>drummondii, Otolithus</i>	355
<i>cyclolepis</i>	203	<i>dubia, Seriola</i>	280
<i>dolomieu, Micropterus</i>	298	<i>dubium, Exoglossum</i>	146

List of scientific names—Continued

	Page		Page
dubius, <i>Ammodytes</i>	481	<i>Echinorhinus</i>	22
<i>Fierasfer</i>	486	<i>brucus</i>	22
<i>Menephorus</i>	309	<i>obesus</i>	22
<i>Serranus</i>	309	<i>echinus</i> , <i>Diodon</i>	501
ductor, <i>Gasterosteus</i>	278	<i>Echiodon</i>	486
<i>Naucrates</i>	278	<i>drummondi</i>	486
dugesii, <i>Algansea</i>	116	<i>Echiopsis</i>	88
<i>Ameiurus</i>	153	<i>Echiostoma</i>	70
<i>Haustor</i>	153	<i>barbatum</i>	70
dugesii, <i>Fundulus</i>	182	<i>margarita</i>	70
<i>Zoogoneticus</i>	182	<i>margaritum</i>	70
dulcis, <i>Argyreus</i>	139	ectenes, <i>Careproctus</i>	403
<i>Rhinichthys</i>	139	<i>Micropogon</i>	349
Dules.....	320	<i>Ectenias</i> , <i>brunneus</i>	265
<i>auriga</i>	320	<i>edentula</i> , <i>Platirostra</i>	36
<i>flaviventris</i>	320	<i>edentulus</i> , <i>Cetengraulis</i>	51
<i>tæniurus</i>	305	<i>Engraulis</i>	51
dumerilii, <i>Squatina</i>	22	<i>Edriolychnus</i>	508
dumerilii, <i>Apionichthys</i>	230	<i>schmidti</i>	508
<i>Caranx</i>	279	<i>edwardi</i> , <i>Sciæna</i>	353
<i>Polycirrus</i>	351	<i>edwardsi</i> , <i>Psenes</i>	269
<i>Polyclemus</i>	351	<i>Stilbiscus</i>	89
<i>Seriola</i>	279	eeltenkee, <i>Myliobatis</i>	30
duodecim, <i>Engraulis</i>	48	<i>effulgens</i> , <i>Ethiopora</i>	168
duplex, <i>Orthopristis</i>	335	<i>Arlina</i>	287
duquesnii, <i>Catostomus</i>	110	<i>Boleosoma</i>	287
durvilli, <i>Argyropelecus</i>	74	<i>Larimus</i>	344
duryi, <i>Etheostoma</i>	286	<i>eglanteria</i> , <i>Raja</i>	25
dussumieri, <i>Brama</i>	265	<i>Raja</i>	25
<i>Seriola</i>	278	<i>egmontis</i> , <i>Ahlia</i>	82
Dussumeria <i>stolifera</i>	46	<i>Myrophis</i>	82
Dussumeriidae.....	45	<i>egregia</i> , <i>Cheonda</i>	118
dux, <i>Lachnolaimus</i>	425	<i>Tigoma</i>	118
divinensis, <i>Platessa</i>	227	<i>egregius</i> , <i>Exocoetides</i>	454
		<i>egretta</i> , <i>Bellator</i>	408
earlii, <i>Phycis</i>	212	<i>Prionatus</i>	408
Ebisus.....	373	<i>eigenmanni</i> , <i>Acutomentum</i>	366
<i>sagamus</i>	373	<i>Cichlasoma</i>	417
Echelidae.....	82	<i>Evarra</i>	142
<i>Echelus</i> <i>caudilimbatus</i>	79	<i>Gobiesox</i>	490
<i>Echemythes</i>	216	<i>Gobius</i>	439
Echeneidae.....	448	<i>Rhinogobius</i>	439
<i>Echeneis</i>	443	<i>Rumicola</i>	490
<i>albacauda</i>	448	<i>Sebastodes</i>	366
<i>albescens</i>	449	<i>Theraps</i>	417
<i>apicalis</i>	449	<i>Eigenmannia</i>	101
<i>australis</i>	449	<i>virescens</i>	101
<i>brachyptera</i>	449	<i>eigenmannia</i> , <i>Anchovia</i>	50
<i>fusca</i>	449	<i>Anchoviella</i>	50
<i>gualcan</i>	449	<i>eiseni</i> , <i>Characodon</i>	183
<i>holbrookii</i>	449	<i>elaborata</i> , <i>Muraena</i>	93
<i>jacobæa</i>	449	<i>elaboratus</i> , <i>Gymnothorax</i>	93
<i>lineata</i>	449	<i>Elacate</i>	281
<i>lunata</i>	448	<i>atlantica</i>	281
<i>metallica</i>	449	<i>bivittata</i>	281
<i>naucrates</i>	448	<i>falcipectus</i>	281
<i>neucrates</i>	448	<i>malabarica</i>	281
<i>neucrotoides</i>	449	<i>motta</i>	281
<i>neuhoffii</i>	449	<i>pondicerriana</i>	281
<i>osteochir</i>	450	<i>Elacatinus</i>	442
<i>pallida</i>	449	<i>oceanops</i>	442
<i>pervia</i>	449	<i>Elagatis</i>	279
<i>postica</i>	449	<i>bipinnulatus</i>	279
<i>quatuordecimlaminatus</i>	449	<i>Elanura</i>	380
<i>remora</i>	449	<i>fordæa</i>	380
<i>remoroides</i>	449	<i>Elaphocottus</i>	389
<i>scutata</i>	449	<i>Elaphotoxon</i>	271
<i>sexdecim-lamellata</i>	449	<i>bartholomæi</i>	272
<i>sphærenarum</i>	449	<i>ruber</i>	271
<i>squalipeta</i>	449	<i>Elapsophis</i>	87
<i>tetrapturorum</i>	450	<i>Elasmobranchii</i>	10
<i>tropica</i>	449	<i>elassochir</i> , <i>Noturus</i>	156
<i>verticalis</i>	449	<i>Elasmodiscus</i>	402
<i>vittata</i>	449	<i>tremebundus</i>	402
Echidna.....	94	<i>elassodon</i> , <i>Hippoglossoides</i>	222
<i>catenata</i>	94	<i>Elassoma</i>	305
<i>chionostigma</i>	94	<i>evergladei</i>	305
<i>cocosa</i>	94	<i>zonata</i>	305
<i>flavoscripta</i>	94	<i>zonatum</i>	305
<i>fuscomaculata</i>	94	<i>Elassomidae</i>	305
<i>nebulosa</i>	94	<i>Elastoma</i>	328
<i>nocturna</i>	94	<i>elater</i> , <i>Malthæ</i>	507
<i>scabra</i>	94	<i>Zelientes</i>	507
echinatum, <i>Leiodon</i>	21	<i>Elattonistius</i>	41
echinita, <i>Scorpena</i>	372	<i>Elatarchus</i>	345
<i>Echinorhinidae</i>	22	<i>archidum</i>	345

List of scientific names—Continued

	Page		Page
elattura, <i>Netuma</i>	149	elongatus, <i>Luxilus</i>	120
elatturus, <i>Arius</i>	149	Megalops.....	39
elcapitanense, <i>Syngnathus</i>	242	Ophiodon.....	374
elcapitanensis, <i>Syngnathus</i>	242	Osmerus.....	67
electricus, <i>Rhinobatus</i>	23	Pomadasis.....	333
elegans, <i>Cottus</i>	381	Pomotis.....	300
Cyphosus.....	339	Sebastes.....	368
Cyprinodon.....	181	Symphurus.....	231
Gasterosteus.....	238	Elopidæ.....	39
Gibbonsia.....	458	Elopoidea.....	39
Gila.....	114	Elops.....	39
Hamulon.....	329	hawaiensis.....	39
Iridio.....	427	inermis.....	39
Kyphosus.....	339	saurus.....	39
Labco.....	109	elucens, <i>Syngnathus</i>	241
Mesoprion.....	328	emarginatum, <i>Scarus</i>	433
monteyensis, <i>Gibbonsia</i>	458	emarginatus, <i>Lobotes</i>	325
Myxodes.....	458	Serranus.....	315
Nanostoma.....	289	Embassichthys.....	228
Orthoragiscus.....	503	bathyblus.....	228
Pimalepterus.....	339	Embiotoca.....	411
Eleginus.....	209	caryi.....	412
nawaga.....	210	cassidy.....	411
Eleotridæ.....	435	jacksoni.....	411
Eleotris.....	436	lateralis.....	411
abacurus.....	437	lineata.....	411
amblyopsis.....	437	ornata.....	411
capite plagioptero.....	436	perspicabilis.....	411
(<i>Cultus</i>) belizianus.....	436	webbi.....	411
dormitatrix.....	435	Embiotocidæ.....	409
grandisquama.....	436	Emblemaria.....	464
guavina.....	436	atlantica.....	464
gyrinus.....	436	marki.....	464
latifrons.....	436	nivipes.....	464
latifasciatus.....	437	oculocirris.....	464
longiceps.....	435	pandionis.....	464
mauriti.....	268	Emblemaridæ.....	464
mugiloides.....	436	emblematicus, <i>Gobius</i>	443
omocyanus.....	436	Microgobius.....	443
perniger.....	437	Scarus.....	435
pictus.....	437	embryum, <i>Allocottus</i>	391
pisons.....	436	Oligocottus.....	391
seminuda.....	438	Embry.....	473
seminudus.....	438	crassilabris.....	473
sima.....	436	crotalinus.....	473
smaragdus.....	437	parallelus.....	473
somnolentus.....	436	embryx, <i>Gerrus</i>	342
tubularis.....	437	emilia, <i>Opsopcesodus</i>	115
elephas, <i>Squalus</i>	20	Emmeckia.....	428
eleuthera, <i>Rabida</i>	156	venusta.....	428
Eleutheractus.....	316	emmelane, <i>Arius</i>	151
coriaceus.....	316	A verruncus.....	395
eleutherus, <i>Noturus</i>	156	Tachysurus.....	151
elizabeth, <i>Lepidosteus</i>	37	Emmelas.....	364
Ellips.....	151	glaucus.....	354
elliott, <i>Thorichthys</i>	422	emmelas, <i>Lepophidium</i>	484
elliott, <i>Diabolichthys</i>	32	Leptophidium.....	484
ellipsoides, <i>Lebias</i>	181	emoryi, <i>Gila</i>	114
ellipticus, <i>Chatoëssus</i>	46	emperador, <i>Astyanax</i>	97
Cymatogaster.....	410	Bryconamericus.....	97
Platophrys.....	217	Empetrichthyidæ.....	182
Pleuronectes.....	217	Empetrichthys.....	182
Ellops.....	34	merriami.....	182
elongata, <i>Anchovia</i>	49	emphæus, <i>Esobastes</i>	367
Anchoviella.....	49	Sebastes.....	367
Aphoristia ornata var.....	231	Emphyus.....	213
Avocettina.....	53	emphysetus, <i>Bagrus</i> (<i>Sciades</i>).....	148
Bothrocaropsis.....	477	Sciadeichthys.....	148
Clupea.....	41	enceomus, <i>Gobius</i>	439
Cyclothone.....	72	Rhinogobius.....	439
Psocilia.....	190	encaustus, <i>Chapalichthys</i>	184
Umbrina.....	351	Characodon.....	184
Umbrula.....	351	encenadæ, <i>Rhinoptera</i>	31
elongatum, <i>Gonostoma</i>	72	Enchelycephali.....	77
elongatus, <i>Aulophallus</i>	190	Enchelycore.....	90
Bramocharax.....	89	euryrhina.....	90
Carpiodes.....	104	nigricans.....	90
Catostomus.....	102	nigrocaustus.....	90
Clinostomus.....	102	Enchelyopus.....	209, 262, 473
Cycleptus.....	102	americanus.....	213
Eucinostomus.....	102	barbatus.....	479
californiensis.....	341	climbrus.....	209
Hispanicus.....	368	mediterraneus.....	208
Labco.....	109	regalis.....	213
Labichthys.....	83		

List of scientific names—Continued

	Page		Page
enchrysurus, Chromis.....	413	Eosebastes deani.....	367
Helases.....	413	diploproa.....	367
Encrasicholus.....	47	emphaeus.....	367
Encrotes.....	180	introniger.....	366
Engraulidae.....	47	melanostomus.....	366
engraulinus, Photogenis leucops.....	123	savcola.....	367
Engraulis.....	47	semianctus.....	367
argyrophanus.....	48	swifti.....	367
brevis.....	51	Eperlanus.....	67
brevirostris.....	51	Ephippidae.....	358
chaerostomus.....	48	ephippium, Chaetodon.....	359
clupeoides.....	50	Plectropoma.....	321
compressus.....	47-50	Ephippus gigas.....	358
cubanus.....	48	zonatus.....	358
delicatissimus.....	48	Epicopus.....	214
dentex.....	51	epieurum, Chromis.....	280
duodecim.....	48	Epigeichthys.....	469
edentulus.....	51	atropurpureus.....	469
grossidens.....	51	Epigonichthyidae.....	7
janeiro.....	51	Epigonus.....	296
lemniscatus.....	48	macrophthalmus.....	296
louisiana.....	48	occidentalis.....	296
macrolepidotus.....	50	epihexodon, Petromyzon.....	10
mitchilli.....	48	Epinephelus afer.....	312
mordax.....	47	Epinephelidae.....	308
mysticetus.....	51	Epinephelus.....	310
nanus.....	47	adscensionus.....	310
panamensis.....	50	analogus.....	310
perfasciatus.....	47	angustifrons.....	311
poeyi.....	51	brachysoma.....	311
productus.....	50	chalinus.....	315
spumifer.....	50	ciliatus.....	365
tricolor.....	48	cubanus.....	311
engymen, Cetengraulis.....	51	drummond-hayi.....	312
Engyophrys.....	220	flavolumbatus.....	311
sancti-laurentii.....	220	guaza.....	310
Enneacanthus.....	303	guttatus.....	311
gloriosus.....	304	jordani.....	314
margarotis.....	304	labriformis.....	311
obesus.....	304	latefasciatus.....	310
pinniger.....	304	lightfooti.....	313
Enneacentrus.....	309	marginalis.....	310
Enneanectes.....	458	mentzeli.....	310
Ennenpterygius.....	458	morio.....	311, 312
carminalis.....	458	mystacinus.....	310
pusillus.....	458	niphobles.....	312
Enneistius.....	309	niveatus.....	311
acanthistius.....	309	ordinatus.....	311
Enophrys.....	381	rosaceus.....	314
claviger.....	381	ruber.....	309, 315
taurinus.....	381	selhecauda.....	311
ensifera, Bardiella.....	345	striatus.....	311, 318
Sciæna.....	345	Epinnula.....	261
ensiferus, Centropomus.....	306	magistralis.....	261
ensis, Gaidropsarus.....	208	episcopa, Dionda.....	144
Motella.....	208	episcopi, Brachyrhaphis.....	187
Sphyraena.....	255	Gambusia.....	187
entemedor, Narcine.....	27	Episema.....	125
Entomacrodus.....	463	callisema.....	132
chiostictus.....	463	jejuna.....	125
cruentatus.....	463	Epitrachys.....	281
decoratus.....	463	epsetus, Anchoviella.....	48
margaritaceus.....	463	Esox.....	48
nigricans.....	463	Eptatretidae.....	8
entomelas, Acutomentum.....	366	equatorialis, Chlopsis.....	82
Sebastichthys.....	366	Raja.....	26
entomophagus, Leucichthys.....	63	Eques.....	352
Entosphenus.....	9	acuminatus umbrosus.....	353
ciliatus.....	10	americanus.....	352, 353
tridentatus.....	9	balteatus.....	353
Enypnias.....	445	lanceolatus.....	353
seminudus.....	445	lineatus.....	353
Eopsetta.....	222	pulcher.....	353
jordani.....	222	punctatus.....	353
eos, Aibaciosa.....	491	viola.....	353
Boleichthys.....	294	equestris, Arius.....	149
Chrosomus.....	113	Balistes.....	491
Gobiesox.....	491	Equetus.....	352
Othonops.....	447	equirostrum, Seomberesov.....	197
Fronotogrammus.....	322	equisetis, Coryphæna.....	265
Sebastodes.....	368	erate, Lobotes.....	323
Sebastomus.....	368	erebennus, Ameiurus.....	153
Eosebastes.....	366	erebus, Muraena.....	92, 98
aurora.....	366	eremogenes, Salmo.....	58
cramerii.....	367		

List of scientific names—Continued

	Page		Page
erethizon, Arothron	500	Erythrinus brasiliensis	99
Eretmichthys	481	macrodon	99
ocella	481	microcephalus	99
pinnatus	481	trahira	99
erlarcha, Atherinella	249	erythrocheilus, Albula	40
Copelandia	304	erythrogaster, Chrosomus	113
Eurystole	249	lunulus	113
Ericara	53	Serranus	312
salmonae	53, 54	erythrogasterum, Pœcilosoma	291
Ericheia	301	erythropterus, Gobiesox	490
Ericosma	284	Icthelus	303
evides	284	Sicyases	490
Ericeteis	460	erythroptera, Pimelodus	152
kalisherae	460	erythrorhynchus, Salmo	61
Ericymba	141	erythrus, Catostomus	110
buccata	142	escambiae, Zygometes	178
ericymba, Sciæna	346	eschrichtii, Oneirodes	509
Stellifer	346	Escolar	260
erensis, Argirosomus	63	niger	260, 261
Leucichthys	63	violaceus	260
Eriepidae	373	esculentus, Carangus	273
Eriepis	373	Conger	79
zonifer	373	Merluccius	214
Erimonax	138	Eslopsarum	250
monachus	138	arge	250
Erimystax	138	Esloscopus	455
dissimilis	138	zelotes	455
watauga	138	esmarkii, Lycodes	474
Erimyzon	108	Esocidae	172
goodii	109	esopus, Labeo	109
oblongus	109	estor, Gila	120
sucetta	109	Esox	172, 173
erinacea, Raja	25	americanus	172
Raja	25	australis	173
erinaceus, Trichocyclops	501	barracuda	255
Erimemus	137	boreus	173
hyalinus	137	brasiliensis	190
hypsionotus	138	crassus	173
labrosus	138	cypho	173
rubrifrons	138	depiaudus	173
storerianus	137	epsetus	48
Eriscion	355	eslor	173
albus	356	fasciatus	173
macdonaldi	356	flavulus	176
nebulosus	355	immaculatus	173
parvipinnis	355	imperialis	197
reticulatus	355	lineatus	173
stolzmanni	356	longirostris	196
virescens	356	lucius americanus	173
xanthulus	356	lucoides	173
erithrus, s. Caranx	273	lugubris	173
Eritrenia	140	malabaricus	99
Erizo	501	marinus	196
guanabana	501	masquinongy	172
erochrous, Hololepis	294	immaculatus	173
Erogala	129	ohiensis	172
analostana	129	niger	173
caerulea	130	nobilior	172
callistia	130	nobilis	172
camura	129	ohiensis	172
cercostigma	130	ornatus	173
chloristia	129	osseus	36
eurystoma	130	ovinus	181
formosa	129	phaleratus	173
galactura	129	pisculentus	176
nivea	129	pisciculus	176
notata	130	porosus	173
pyrrhomelas	129	ravenel	173
stigmatura	129	reticulatus	173
texana	130	salmonatus	163, 172
trichroistia	130	saurus	197
venusta	130	scomberius	173
whipplei	129	spet	255
xanura	130	sphyrena	255
Erotelis	437	synodus	163
smaragdus	437	timucui	195
valenciennesi	437	tudecomlineatus	173
Errax	228	tristichus	38
zachirus	228	umbrosus	173
Erychthys	433	vermiculatus	173
erythraum, Astatheros	422	vittatus	173
Erythraeus	420	vulpec	39
erythraeus, Heros	422	zonatus	176
erythrichthys	420	estor, Chirostoma	250, 251
erythrinoides, Scarus	432	Esox	172

List of scientific names—Continued

	Page		Page
estor, Gila.....	121	Etrumeus.....	46
Estrella.....	287	acuminatus.....	46
atromaculata.....	287	micropus.....	46
esunculus, A topichthys.....	54	sadina.....	46
Etelides.....	328	Eucalia.....	238
aquilonaris.....	328	inconstans.....	238
Etelis.....	328	cayuga.....	238
carbunculus.....	328	pygmaea.....	238
evivurus.....	328	Euchalarodus.....	227
oculatus.....	328	putnami.....	227
Etheostoma.....	286, 293, 294	Eucinostomus.....	340, 341
artesia.....	292	argenteus.....	340, 341
aubeenaubei.....	291	californiensis.....	341
austiale.....	293	elongatus.....	341
barrati.....	294	elongatus.....	341
blennioides.....	283, 286	gula.....	341
caerulea.....	291	gulula.....	341
calliura.....	298	harengulus.....	341
camurum.....	292	meeki.....	341
caprodes.....	282	productus.....	341
cinerea.....	290	pseudogula.....	341
(Cottogaster) uramides.....	285	pseudo-gula.....	341
cragini.....	292	Eucrotus.....	267
cumberlandicum.....	293	ventralis.....	267
duryi.....	286	Euctenogobius.....	441
fiabellais.....	293	badius.....	441, 442
fiabellata.....	293	latus.....	442
fontinalis.....	293	poeyi.....	442
fricksia.....	291	sagittula.....	441
guntheri.....	283	stigmatus.....	441
(Hadropterus) cymatoxenia.....	285	Eucyclogobius.....	444
manguae.....	285	newberryi.....	444
ouachitæ.....	284	Eudulus.....	320
squamatus.....	285	Eugaleus.....	14
hildebrandti.....	291	Eugeris.....	342
lowæ.....	291	brasiliensis.....	342
julæ.....	292	brevimanus.....	343
lepidogenys.....	292	lineatus.....	342
hinsleyi.....	293	mexicanus.....	343
longimana.....	286	plumieri.....	343
luteovinctum.....	291	Eulamia.....	15
macrocephalum.....	284	nicaraguensis.....	16
maculata.....	290	platyrhynchus.....	15
microptera.....	294	eulepidotus, Sphceroides.....	498
micropterus.....	290	Tetrodon (Sphceroides).....	498
nevisense.....	283	eulepis, Microgobius.....	444
manguae spilatum.....	285	Euleptorhamphus.....	198
nigra.....	287	brevoorti.....	198
(Nothonotus) jordani.....	290	brevoorti.....	198
obeyense.....	292	longirostris.....	198
olmstedii.....	287	velox.....	198
pagel.....	292	Eumecesogrammus.....	470
parvipinne.....	292	præcisus.....	470
pellucidum.....	288	Eumicrotremus.....	398
peltatum.....	283	orbis.....	398
phoxocephalum.....	284	phrynoides.....	398
podostemone.....	286	spinosus.....	398
quappella.....	291	eumorphus, Chatoessus.....	44
rev.....	283	Eumycterias.....	500
(Rhthoea) blennius.....	287	bitæniatus.....	500
rupestre.....	290	Eunichthys.....	410
sagitta.....	293	megalops.....	410
scovelli.....	293	Eupomacentrus.....	413
squamiceps.....	293	adustus.....	414
swannanoa.....	289	analis.....	413
tessellata.....	290	chrysus.....	414
tippecanoe.....	292	dieneæus.....	413
tuscumbia.....	293	flavilatus.....	414
(Ulocentra) histrio.....	286	flaviventer.....	414
variata.....	289	fuscus.....	413
verecundum.....	287	gilli.....	414
whipplei alabamæ.....	292	leucurus.....	414
wrighti.....	285	leucostictus.....	413
zonale arcansum.....	289	nepenthe.....	414
etheostoma, Aboma.....	443	otophorus.....	414
Etheostomidae.....	282	paritatus.....	414
ethon, Syngnathus.....	241	planifrons.....	414
Etmopterus.....	21	rectifrenum.....	414
hillianus.....	21	Eupomotis.....	302
etowanus, Catostomus nigricans.....	107	gibbosus.....	299, 302
Etropus.....	219	heros.....	302
crossotus.....	219, 220	holbrooki.....	302
micros.....	219	holbrooki.....	302
microstomus.....	219	longimanus.....	302
rimosus.....	220	europeus, Trachinus.....	270

List of scientific names—Continued

	Page		Page
europeus, Aspidophorus	395	Evorthodus	438
Blennius	467	breviceps	438
Eurymen	401	Evoxymetopon	262
gyrinus	401	tæniatus	262
Eurymyctera	91	exasperata, Platyrrhina	24
euryopa, Hudsonius	132	exasperatus, Zanterys	21
euryops, Bathylagus	69	excisus, Dorythamphus	243
Icelus	378	Execestes	454
Myxostoma	110	egregius	454
Strongylura	196	Exerpes	456
Tylosurus	196	asper	456
euryorus, Apomotis	299	exigua, Anchoviella	47
Lepomis	299	exiguus, Bodianus argyro-leucos et	345
Eurypharyngide	95	Stolephorus	47
euryplectrum, Diplectrum	318	exile, Dorosoma cepedianum	46
euryrhina, Enchelycore	90	Siphostoma	243
Eurystole	249	exilicauda, Lavinia	113
eriarcha	249	exilis, Belone	196
eyrstole, Stolephorus	48	Boleichthys	294
eurystoma, Erogala	130	Hippoglossoides	222
Eurystomus	107	Lyopsetta	222
eurystomus, Photogenis	130	Noturus	156
Eusearus	431	Rabida	156
Euschistodus	414	Strongylura	196
analogus	415	Syrtetes	243
concolor	414	Eximia	390
declivifrons	414, 415	rubellio	390, 391
Eusebastes	363	eximus, Cyprinodon	181
Euselachi	11	Exocetidae	199
Eusphyra	18	Exocetus	200
Eustomatodus	270	acutus	199
Euthynnus	258	affinis	201
alletteratus	258	albidactylus	202
lineatus	259	appendiculatus	201
yaito	239	bahiensis	202
Eutycheilus	352	bicolor	201
evansi, Hybognathus	143	brachycephalus	200
Evarra	142	californicus	202
eigenmanni	142	callopterus	202
flahuacensis	142	chiliensis	200
Eventognathi	102	comatus	201
Evepigymnus	270	cyanopterus	202
evergladii, Elasmoma	305	(Cypsilurus) procer	201
evermanni, Alvordius	284	dowii	201
Arius	151	evolans	200
Astatheros	420	exiliens	200
Atherina	246	fasciatus	200
Atherinella	248	fucorum	202
Cheonda	118	furcatus	201
Cichlasoma	420	georgianus	200
Congrosoma	80	gibbifrons	202
Cottus	383	gryllus	200
Diapterus	342	heterurus	201
Hadropterus	284	hillianus	200
Hepsetia	246	holubi	202
Leuciscus	113	lamellifer	200
Moharra	342	lineatus	202
Salmo	59	lutkeni	201
Scarus	434	maculipinnis	201
Synodus	104	melanurus	201
Tachysurus	151	mento	200
Thynnus	248	mesogaster	200
Evermannia	445	micropterus	199
longipinnis	445	nigricans	201
panamensis	445	novaboracensis	201
osterura	445	nuttalli	201
Evermannichthys	446	obtusirostris	200
metzelaeri	447	orbignianus	200
spongicola	446, 447	parrae	202
evides, Alvordius	284	quadrirremus	201
Clinus	453	roberti	201
Ericosma	284	rondelii	201
Plectobrachius	468	rubescens	201
evionthas Ophichthys	87	rufipinnis	201
Quadrirremus	87	scylla	201
Eviota	438	speculiger	201
epiphanes	438	spilonotopterus	202
personata	438	spilops	201
evurus, Etelis	328	splendens	200
evolans, Exocetus	200	vermiculatus	202
Prionotus	407	vinciguerrae	200
Trigla	407	volador	200
Evolantia	199	volitans	200, 201
microptera	199	volitens	200
Evoplites	324	xenopterus	202

	Page		Page
Evglossum.....	146	fasciatus, Achnrus.....	229
annulatum.....	146	Astyanax.....	37
dubium.....	146	Auchenopterus.....	457
lesuiianum.....	146	Baodon.....	417
maxilliguala.....	146	Bodianus.....	325
mirabile.....	142	Brytus.....	304
nigrescens.....	146	Carant.....	273
spinicephalum.....	146	Catonotus.....	109
exoletus, Labrus.....	424	Catostomus.....	407, 468
Centrolabrus.....	424	Cetronotus.....	456, 457
Econantes.....	200	Ctenomolales.....	304
affinis.....	201	Ctenogobius.....	273
exsilens.....	200	Diplesion.....	283
rondeletti.....	200	Esoy.....	200
rubescens.....	201	Eocoetus.....	351
rufipinnis.....	201	Geyonemus.....	229
speculiger.....	200	Gymnachurus.....	101
vincigueræ.....	200	Gymnotus.....	362
expansum, Ostracium.....	497	Harpurus.....	188
exsilens, Eyoocetus.....	200	Hemirhamphus.....	344
Eyonantes.....	200	Lamius.....	410, 411
extensus, Fundulus.....	179	Mallophagus.....	490
Lycodapus.....	478	Ovodes.....	468
Zygonectes.....	179	Pholis.....	145
Eytrarius.....	139	Pimophales.....	352
tetianemus.....	139	Pogonias.....	309
eyul, Apistes.....	370	Prionodes.....	279
eydouvi, Bagre.....	148	Scomber.....	369
Galeichthys.....	148	Sebastes.....	490
fabbri, Blennius.....	460	Sicyases.....	242
Faber marinus fere quadratus.....	354	Syngnathus.....	279
faber, Chaetodipterus.....	358	Trachurus.....	276
Chaetodon.....	338	Trachynotus.....	272
fabrieu, Campylodon.....	21	Xurel.....	279
Centroscyllium.....	269	Zonichthys.....	318, 319
Gadus.....	470, 471	fascicularis, Serranus.....	109
Gunnellus.....	400	fasciola, Idiacanthus.....	126
Liparis.....	471	fasciolaris, Catostomus.....	126
Lumpenus.....	31	Lythrurus.....	126
Spinax.....	21	Notropis umbratilis.....	369
fabroni, Apterurus.....	81	Sebastichthys.....	231
fajardo, Auchenopterus.....	141	Symphurus.....	264
falcata, Agosia.....	141	fasciolata, Coryphæna.....	203
Apocope.....	315	favosus, Bathygadus.....	461
Mycteroperca.....	315	Blennius.....	108
phenax, Mycteroperca.....	280	fecundus, Catostomus.....	108
Seriola.....	98	Chasmistes.....	147
falcatus, Brycon.....	271	Felichthys.....	159
Caranx.....	188	amblops.....	181
Glaridichthys.....	271	felicianus, Cyprinodon.....	181
Hemicaranx.....	276	Trifarcius.....	153
Labrus.....	315	felinus, Pimelodus.....	315
Serranus.....	425	Serranus.....	472
Sparus.....	276	fels, Anarrhichthys.....	149
Trachinotus.....	280	Galeichthys.....	14
Zonichthys.....	54	Mustelus.....	154
falcidens, Atopichthys.....	16	Pimelodus.....	149
falciformis, Carcharias.....	281	Silurus.....	576
falcipinnis, Carcharias.....	142	fenestralis, Artedius.....	876
Elacata.....	142	Astrolytes.....	416
Falcula.....	142	fenestrata, Chromis.....	416
chapalæ.....	142	fenestratum, Clethrurus.....	171
Falcularius.....	142	ferox, Alepisaurus.....	70
chapalæ.....	272	Bathypops.....	70
fallax, Caranx.....	300	Idiacanthus.....	58
Pomotus.....	278	Lepisosteus.....	38
fanfarus, Naucrates.....	55	(Atractosteus).....	40
Fario.....	55	Macrodon.....	19
argenteus.....	55	Squalus.....	60
argyreus.....	58	Stomias.....	227
new berrii.....	56	feruginea, Limauda.....	226, 227
stellatus.....	57	Platessa.....	183
tsupplitch.....	55	feruginea, Chaacodon.....	90
.....	24	festue, Plabucus.....	36
.....	298	feuille, Polyodon.....	145
Clupea.....	42	fibulatus, Spinicephalus.....	101, 486
Gambusia.....	169	Fierasfer.....	486
Molnesia.....	192	acutus.....	486
Plagusia.....	231	affinis.....	486
Poecilopsis.....	189	arenicola.....	471
Torrentaria.....	293	borealis.....	486
Trigla.....	100	dubius.....	478
fasciatum, Characium.....	335	fierasfer, Lycodapus.....	483
Pristipoma.....	42	filamentosa, Diacolele.....	
.....	169		
.....	192		
.....	231		
.....	189		
.....	293		
.....	100		
.....	335		
.....	42		
.....	169		
.....	192		
.....	231		
.....	189		
.....	293		
.....	100		
.....	335		
.....	42		
.....	169		
.....	192		
.....	231		
.....	189	</	

List of scientific names—Continued

	Page		Page
filamentosus, Argyriosus	275	flaviventer, Eupomacentrus	414
Breviceps	147	Pomacentrus	414
Hemiramphus	199	flaviventris, Callidulus	320
Icelinus	375, 376	Dules	320
Monacanthus	494	flavoceruleus, Halocentrus	310
Tarandichthys	376	flavolimbatus, Epinephelus	311
filicauda, Coryphaenoides	207	flavolineatus, Hemulon	330
filifer, Dactylostomias	69, 70	flavolineatus, Diabasis	330
filifera, Anchovia	50	Mullus	344
Anchoviella	50	Sargus	338
Chalinura	201	flavo-lineatus, Pimelepterus	339
fimbria, Anoplopoma	373	flavopurpurea, Perca	310
Gadus	373	flavoscripta, Echidna	94
fimbriata, Cyclosetta	220	flavoscriptus, Gymnothorax	91
Loricaria	162	flavovittatus, Upeneus	343
Raja	32	flavulus, Esoc	176
Squatina	22	flavus, Turdus	425
fimbriatus, Blennius	473	Noturus	155
Hemirhombus	220	fleurieu, Ostorhinchus	295
Icelinus	376	flevo-lais, Lepomis	298
Paraliparis	404	flevo-lus, Catostomus	105
Serranus	311	flore, Neoliparis	399
Fimbriatorpedo	27	florealis, Platyglottus	426
firmsguamis, Bogoslovius	204	florida, Branchiostoma	7
Macurus	204	Jordanella	180
fischeri, Chastostoma	161	Siphostoma	241
Chastostomus	161	Syngnathus	241
Solea	230	florida, Cichla	297
Tetragonopterus	97	Huro	297
Fissala	408	floridanus, Phycis	213
alata	408	Urophycis	213
fissuratus, Neoliparis	399	floridensis, Callinurus	303
Fistularia	245	Fundulus	176
comersonii	245	Floridichthys	180
cornuta	245	comersonii	180
depressa	245	comersonii	177
immaculata	245	comersonii	302
neo-boracensis	245	comersonii	132
petimba	245	Leptophilypnus	436
serrata	245	Sargosomus	412
tabacalia	245	Tetodon	499
villosa	245	Fodiator	199
Fistularia	245	acutus	199
fistularis, Flagellaria	245	fodiator, Tylosurus	196
fistulata, Syngnathus	241	foetens, Salmo	163
fistulata, Syngnathus	241	Synodus	163, 172
flabellare, Cumberlandicum, Catonotus	293	folum, Polyodon	36
lineolatum, Catonotus	293	fonsecensis, Achirus	230
flabellaris, Catonotus	293, 383	Solea	230
Etheostoma	293	Fonticola	178
flabellata, Etheostoma	293	fonticola, Alvarius	295
flabellicauda, Rivulus	180	Fundulus	175
Flagellaria	245	Fundulus heteroclitus	175
fistularis	245	fontinalis, Etheostoma	293
flagellum, Goniobatus	30	Salmo	60
Raja	30	Salvelinus	60
Saccopharynx	95	forbesi, Cyprinella	131
Flammeo	236	Forbesella	194
marianus	236	agassizii	194
flammeus, Phoxinus	121	papillifera	194
flava, Congomuraena	80	forcipatus, Balistes	492
Hildebrandia	80	Forcipiger	359
Loricaria	160	flavissimus	359
flavescens, Mesoprion	325	longirostris	359
Morone	231	forficata, Elanura	380
Perca	231	forlonensis, Cyprinella	131
Prionodes	319	Nototropis	131
Searus	432	formosa, Algansea	122
Serranus	319	Erogala	129
Sparisoma	433	Guentheridia	500
flavescens, Brosmerus	213, 214	Heterandria	188
flavicauda, Hyporhamphus	311	Lamia	192
flavidus, Apodichthys	406, 467	Mollenesia	192
Aulorhynchus	240	Moniana	129, 131
Sebastodes	364	Perca	318
Sebastosomus	364	formosulum, Campostoma	147
flaviguttatum, Lythrulon	331	formosum, Hamulon	330
flaviguttatus, Hamulon	331	formosus, Alburnus	129
flavilatus, Eupomacentrus	414	Anthias	330
Pomacentrus	414	Callinurus	299
flavipinnis, Hybognathus	144	Cottus	388
Pristigaster	44	Diplectrum	318
flavivittatus, Syngnathus	241	Holacanthus	361
flavissimus, Forcipiger	359	Siphateles	122
		Tetodon	500

List of scientific names—Continued

	Page		Page
formulosum, Campostoma.....	147	Fundulus bermudæ.....	175
forskåli, Albula.....	39	chrysotus.....	177
Glossodus.....	40	cingulatus.....	177
forsteri, Albula.....	40	confluentus.....	177
Scomberesox.....	197	cubensis.....	174
forsterianus, Catostomus.....	105, 106	dugesi.....	182
fossor, Ichthyomyzon.....	9	extensus.....	179
fosteri, Priapichthys.....	189	floridensis.....	176
4-maculatus, Diodon.....	501	fonticola.....	175
fragilis, Citharichthys.....	218	fuscus.....	174
Macrourus.....	206	grandis.....	176
Macurus.....	206	guatemalensis.....	174
francisci, Cestracion.....	10	heterocheilus.....	175
Gyroleurodus.....	10	badius.....	176
franklini, Cottus.....	385	bermudæ.....	175
Pleuronectes.....	227	fonticola.....	175
fraterculus, Mylocheilus.....	112	grandis.....	176
fremebundum, Hæmulon.....	329	macrolepidotus.....	176
fremuvilli, Aetobatus.....	31	kansæ.....	179
Myliobatis.....	31	labialis.....	174
frenata, Loricaria.....	162	lmi.....	178
frenatus, Brachyistius.....	409	majalis.....	175, 178
Iridio.....	427	meeki.....	178
Odontopyxis.....	395	menona.....	178
Sarritor.....	395	mudfish.....	175
Xantodes.....	375	nisiolus.....	175
Zanolepis.....	375	oavacæ.....	174
fretensis, Hybopsis.....	129	ocellaris.....	176
fricksa, Etheostoma.....	291	pachycephalus.....	174
Oligocephalus.....	291	pallidus.....	176
friedrichstahli, Hetero.....	418	parvipinnis.....	176
Parapetema.....	418	brevius.....	176
frigida, Moniana.....	181	punctatus.....	174
frigidus, Lycodes.....	474	rathbuni.....	175, 179
fræbæi, Amphilophus.....	422	rhizophoræ.....	175
frondosum, Sparisoma.....	433	scartes.....	177
frondosus, Scarus.....	433	sciadicus.....	177
frontale, Cichlasoma.....	417	seminolis.....	178
frontalis, Caranx.....	273	similis.....	177
Cryptopterenchelys.....	85	tenellus.....	177
Gastropsetta.....	225	vinetus.....	176
Leuciscus.....	128	viridescens.....	176
Ophichthys (Cryptopterus).....	85	xenicus.....	180
Prionotus.....	408	zebra.....	176
Theraps.....	417	zebrinus.....	179
fronti unicorni, Lophius.....	507	zonatus.....	178
fronto, Carcharias.....	17	fundulus, Alepocephalus.....	52
Nemichthys.....	83	funehris, Gobiesox.....	489
frost, Gadus.....	210	Gymnothorax.....	92
fucensis, Liparis.....	400	Noturus.....	156
Pollachius chalcogrammus.....	209	Rabida.....	156
Theragra.....	209	Furcaria.....	412
fucorum, Apodichthys.....	467	atrilobata.....	412
Blennius.....	461	bermudæ.....	413
Cypselurus.....	202	cyanea.....	412
Exocoetus.....	202	multilineata.....	412
Xerperes.....	467	puncta.....	412
fulgens, Corvina.....	345	(Furcaria) atrilobata, Chromis.....	412
Myriopristis.....	234	fuicatus, Cypselurus.....	201
Priacanthus.....	323	Ictalurus.....	151, 152
Tetragonopterus.....	96	Exocoetus.....	201
fulgida, Meda.....	147	Phanerodon.....	411
fuliginosus, Balistes.....	491	Pimelodus.....	152
Cyclichthys.....	502	Fucella.....	476
Diodon.....	502	furellus, Careproctus.....	402
Synbranchus.....	76	fureidens, Chaacodon.....	183
fuliginotus, Holconotus.....	411	fuicifer, Paranthias.....	320
fulva, Labrus chogset.....	424	Pimelodus.....	152
fulvescens, Acipenser.....	23, 35	Serianus.....	320
fulvo-rufus, Labrus.....	335	fureiger, Icelus.....	378
fulvus, Cephalopholis.....	12	Furcimanus.....	476
Labrus.....	309	diaptesus.....	476
Physiculus.....	212	furerhinus, Pontinus.....	372
punctatus, Cephalopholis.....	309	fucinea, Pachypops.....	340
ruber, Cephalopholis.....	309	Peica.....	349
fumeus, Notropis.....	125	furiosa, Rabida.....	156
funduloides, Clinostomus.....	121	furius, Noturus.....	156
Zygocetes.....	176	furneri, Micropogon.....	349
Fundulus.....	175, 180	Umbrina.....	349
adina.....	179	futhi, Arius.....	151
albolineatus.....	177	Chelichthys.....	499
antillarum.....	175	furthi, Arius.....	151
aureus.....	177	Caranx.....	271
balboæ.....	179	Corvina (Homiprion).....	317

List of scientific names—Continued

	Page		Page
furthi, Hemicarax.....	271	Gadus punctatus.....	213
Pristipoma.....	332	pulpureus.....	209
Tetrodon.....	499	pygmaeus.....	210
Zestis.....	347	ruber.....	214
furva, Perca.....	317	rupestris.....	210
fusca, Echeis.....	449	sada.....	209
Hydrargyria.....	174	tau.....	486, 137
Sciæna.....	352	tenus.....	213
Tautoga.....	424	tomcod.....	210
fuscescens, Icelinus.....	376	tomcodus fuscus.....	210
fuscomaculata, Echidna.....	94	tomcodus luteus.....	210
fusco-maculatus, Chromis.....	419	mytilus.....	210
fusculus, Centropistes.....	319	planchus.....	210
Prionodes.....	319	toisk.....	214
fuscum, Pygidium.....	159	verlagus.....	210
fuscus, Acionurus.....	362	viens.....	209
Bathygobius.....	440	wachna.....	210
Cyathites.....	480	Gaidropsalæ.....	208
Cylus.....	436	Gaidropsalus.....	208
Eupomacentrus.....	413	ensis.....	208
Fundulus.....	174	mustellus.....	208
Gadus tomcodus.....	210	septentrionalis.....	208
Gobius.....	440	gairdneri, Querimana.....	254
Hemirhombus.....	219	gairdneri, Mugil.....	254
Kyphosus.....	339	gairdneri beaudsi, Salmo.....	37
Pomacentrus.....	413	ciensis, Salmo.....	37
Psenes.....	268	gilberti, Salmo.....	58
Serranus.....	315	shasta, Salmo.....	58
Silurus.....	153	stonei, Salmo.....	58
Syngnathus.....	242	caudneri, Salmo.....	58
Syrtetes.....	242	galactura.....	129
Trachinotus.....	276	galacturus.....	129
Xyster.....	339	Galafite.....	493
fusiiforme, Boleosoma.....	294	Galasaccus.....	175
fusiiformis, Boleichthys.....	294	Galasaccus.....	177
tyllæ, Raja.....	25	galatæ.....	197
gabonensis, Yomei.....	275	galatæ.....	389
gabonensis, Ischnomembras.....	217	galatæ.....	389
Gadella.....	211	Silurus.....	159
gracilis.....	211	Tylosurus.....	197
maraldi.....	211	Galei.....	11
Gadidae.....	209	Galeichthys.....	149
Gadomus.....	202	assiniensis.....	150
acutus.....	202	azuricus.....	150
longifilis.....	202	bobiensis.....	148
gadovi, Cichlosoma.....	417	capitulosens.....	150
Theraps.....	417	dasycephalus.....	150
Gadus.....	210	cydonii.....	148
aglinus.....	210, 211	feliceps.....	149
agilis.....	209	felis.....	148, 149
albidus.....	214	gilberti.....	150
arenosus.....	210	guatemalensis.....	150
auratus.....	210	guentheri.....	149
brosme.....	213	jordani.....	150
callarias.....	210	leizinosus.....	149
carbonarius.....	209	longicephalus.....	150
chalcogrammus.....	209	peruvianus.....	149
cimbrius.....	209	semani.....	150
colinus.....	209	surinamensis.....	150
compressus.....	212	venauchen.....	150
fabrici.....	209	Galeocerdo.....	14
fimbrius.....	373	arcticus.....	14
frost.....	210	tigrinus.....	14
gracilis.....	210, 211	galeoides, Otophidium.....	485
heteroglossus.....	210	Galeorhinus.....	14
lacustis.....	152, 212	zyopterus.....	14
lepidion.....	211	galeita, Blennius.....	460
longipes.....	213	Galeus.....	13, 17
lota.....	212	asterias.....	13
macrocephalus.....	210	catulus.....	13
maculosus.....	212	glauco.....	17
maraldi.....	211	maculatus.....	14
merluccius.....	214	melanostomus.....	12
merlus.....	214	mustelus.....	13
molva.....	212	vulpecula.....	13
mori hui.....	210	galeus, Serranus.....	312
mustela.....	208	Squalus.....	13, 14
nanus.....	210	Galleichthys.....	274
nawaga.....	209, 210	Galleichthys, Crenopoma.....	274
ogac.....	210	Galleichthys, Crenopoma.....	494
perisopus.....	209	Galleichthys.....	274
pollachius.....	209	Gallus, Zeus.....	274, 275
polymorphus.....	210	Galtæ, Squalus.....	118
proximus.....	210		

List of scientific names—Continued

	Page		Page
Gambusia.....	184, 185	Gasterosteus aculeus.....	239
annectans.....	187	apetes.....	139
alingtonia.....	177	argentissimus.....	238
bonita.....	188	atkinsi.....	238
cana.....	188	bailloui.....	237
cascadensis.....	187	biaculeatus.....	237
caymanensis.....	186	bispinosus.....	238
daijensis.....	189	blanchardi.....	239
dominicensis.....	189	breviceps.....	239
dovii.....	186	canadus.....	281
episcopi.....	187	carolinus.....	276, 277
fasciata.....	189	cataphractus.....	237
gracilior.....	186	concinus.....	239
gracilis.....	185	cuvieri.....	237
heckeli.....	185	dekayi.....	239
holbrooki.....	184	dimidiatus.....	237
holbrookii.....	185	ductor.....	278
humilis.....	185	elegans.....	238
infans.....	189	gladiunculus.....	238
latipunctata.....	187	globiceps.....	238
manni.....	186	gymnus.....	237
mcneilli.....	186	inconstans.....	238
melapleura.....	185	inopinatus.....	238
modesta.....	185, 192	insculptus.....	237
myersi.....	185	islandicus.....	238
nicaraguensis.....	184, 186	levis.....	239
nobilis.....	185	leimus.....	237
oligosticta.....	186	loricatus.....	237
panuco.....	186	lotharingus.....	239
parismina.....	187	mainensis.....	239
patuensis.....	185	maurus.....	239
picturata.....	185	microcephalus.....	237
plumbea.....	192	micropus.....	238
punctata.....	184, 185	millipunctatus.....	239
"....."	185	nebulosus.....	239
"....."	185	neustrianus.....	238
regani.....	184, 186	noveboracensis.....	237
rhododaphora.....	186	obolarius.....	237
senilis.....	185	occidentalis.....	238, 239
speciosa.....	185	plebeus.....	237
tenabensis.....	186	ponticus.....	237
tridentiger.....	188	pugeti.....	238
turrubaiensis.....	189	pungitius.....	238, 239
umbrae.....	190	"biachypoda....."	239
vittata.....	186	pygmaeus.....	238
wiavi.....	186	quadricus.....	239
yucatanica.....	186	salitru.....	280
Gambusia.....	175, 179	santa-annae.....	238
Gammaria.....	25	semitriatus.....	237
gardeniana, Hiatula.....	424	semiloricatus.....	237
gardeni, Centronotus.....	181	senatus.....	237
Sternopyx.....	266	spinachia.....	239
gardoneus, Leuciscus.....	115	spinosus.....	237
Notemigonus.....	115	teraculeatus.....	237
garmani, Cetengraulis.....	51	trachinua.....	237
Characodon.....	183	wheatlandi.....	238
Girardinus.....	189	williamsi.....	238
Gobius.....	439	microcephalus.....	237
Lepomis.....	300	Gastrea.....	239
Notropis.....	131	Gastrobranchus.....	8
Paraliparis.....	405	coccus.....	8
Rhinogobius.....	439	dombey.....	8
Stolephorus.....	51	Gastropsetta.....	225
Gaimannia.....	445	frontalis.....	225
henigymna.....	445	Gastrosomus.....	95
paradova.....	445	baudii.....	95
tuba.....	445	pacificus.....	95
spongicola.....	446	gausa, Serranus.....	312
ganoi, Indio.....	427	gauza, Epinephelus.....	310
Julis.....	427	Labrus.....	310
Gariupa.....	312	gavialis, Lepisosteus.....	36
nigrita.....	312	gavi, Merlus.....	214
Gasteracanthus.....	237	geddesi, Characodon.....	182
cataphractus.....	237	Heirichthys.....	423
Gasteropellicidae.....	99	Tomocichla.....	423
Gasteropelcus.....	99	gelatinosum, Melanostigma.....	478
humboldtii.....	169	gelatinosus, Aphyonius.....	484
maculatus.....	99	Caetroctus.....	402
sternicia.....	99	Cyclopterus.....	402
gasteropelcus, Trachinus.....	357	gelida, Ammocrypta.....	289
Gasterosteus.....	238	Macrhybopsis.....	139
Gasterosteidae.....	237	gelidus, Gobio.....	138, 139
Gasterosteinae.....	238	geminatus, Blennius.....	461, 462
Gasterosteus.....	237	Hypleurochilus.....	462
aculeatus.....	237		

List of scientific names—Continued

	Page		Page
<i>gemma</i> , <i>Hypoplectrus</i>	321	<i>Gibbonsia metzi</i>	458
<i>Hypoplectrus unicolor</i>	321	<i>gibbosa cinea</i> , <i>Perca marina</i>	320
<i>genauer</i> , <i>Astiosthes</i>	71	<i>Gila</i>	119
<i>Lampanyctus</i>	167	<i>Momana</i>	131
<i>Leucichthys</i>	63	<i>Perca</i>	302, 320
<i>Gempylidae</i>	260	<i>fluviatilis</i>	302
<i>Gempylus</i>	261	<i>Tigoma</i>	119
<i>coluber</i>	261	<i>gibbosus</i> , <i>Aphredodeus</i>	215
<i>ophidianus</i>	262	<i>Catostomus</i>	100
<i>prometheus</i>	261	<i>Cyprinodon</i>	181
<i>serpens</i>	261, 262	<i>Eupomotis</i>	302
<i>generosus</i> , <i>Catostomus</i> (<i>Acomus</i>).....	104	<i>Pomotis</i>	301
<i>Notolepidomyzon</i>	104	<i>gibbsii</i> , <i>Hemilepidotus</i>	381
<i>genizara</i> , <i>Rabirubia</i>	426	<i>Salmo</i>	57
<i>gentilis</i> , <i>Blennius</i>	462	<i>gibbus</i> , <i>Hypsilepis cornutus</i>	128
<i>Hypsoblennius</i>	462	<i>Liparis</i>	400
<i>Genystiemus</i>	335	<i>Lophius</i>	505
<i>luteus</i>	335	<i>Scarus</i>	433
<i>Genyonemus</i>	349	<i>gigas</i> , <i>Ephippus</i>	358
<i>fasciatus</i>	351	<i>Hippoglossus</i>	221
<i>lineatus</i>	349	<i>Perca</i>	310
<i>Genyorge</i>	324	<i>Sciæna</i>	352
<i>Genypterus omostigma</i>	485	<i>Seriola</i>	270
<i>Genytremus</i>	332	<i>Stereolepis</i>	308
<i>interuptus</i>	332	<i>giganteus</i> , <i>Dololepis</i>	472
<i>geometricus</i> , <i>Anchisomus</i>	499	<i>Gila</i>	114
<i>Diodon</i>	502	<i>affinis</i>	114
<i>lineatus</i> , <i>Diodon</i>	502	<i>ardesiaca</i>	118
<i>Tetrodon</i>	499	<i>conocephala</i>	113
<i>Zeus</i>	275	<i>conspersa</i>	120
<i>Geophagus</i>	424	<i>elegans</i>	114
<i>crassilabris</i>	424	<i>emoryi</i>	114
<i>cupido</i>	424	<i>estoi</i>	121
<i>demon</i>	424	<i>gibbosa</i>	119
<i>magasema</i>	424	<i>gracilis</i>	114
(<i>Satanoperca</i>) <i>crassilabris</i>	424	<i>grahami</i>	114
<i>georgianus</i> , <i>Exocoetus</i>	200	<i>grandis</i>	114
<i>Gephyrocharax</i>	96	<i>gula</i>	120
<i>altricaudata</i>	96	<i>microlepidota</i>	112
<i>altricaudatus</i>	96	<i>mutuæ</i>	114
<i>chocoensis</i>	96	<i>nacrea</i>	114
<i>intermedius</i>	96	<i>nigra</i>	119
<i>gerania</i> , <i>Belone</i>	106	<i>pulchella</i>	119, 120
<i>germanus</i> , <i>Notropis</i>	138	<i>robusta</i>	114
<i>Germo</i>	260	<i>seminuda</i>	114
<i>alalunga</i>	260	<i>gila</i> , <i>Catostomus</i>	106
<i>germo</i>	260	<i>gilberti</i> , <i>Ariosoma</i>	80
<i>germo</i> , <i>Germo</i>	260	<i>Careproctus</i>	402
<i>Scomber</i>	260	<i>Cetengaulis</i>	51
<i>germon</i> , <i>Scomber</i>	260	<i>Chlopsis</i>	82
<i>Gerres</i>	341	<i>Citharichthys</i>	219
<i>apron</i>	342	<i>Congrellus</i>	80
<i>atrolus</i>	342	<i>Galeichthys</i>	150
<i>axillaris</i>	342	<i>Hubbesia</i>	248
<i>brasilianus</i>	342	<i>Hypopsis</i>	134
<i>brevimanus</i>	343	<i>Hypsoblennius</i>	462
<i>brevirostris</i>	342	<i>Ilyonus</i>	445
<i>cinereus</i>	341	<i>Imostoma</i>	286
<i>embryx</i>	342	<i>Isesthes</i>	462
<i>gula</i>	340, 341	<i>Lepidogobius</i>	445
<i>havana</i>	342	<i>Notropis</i>	134
<i>jonesii</i>	341	<i>Noturus</i>	156
<i>lineatus</i>	342	<i>Oncocottus</i>	389
<i>mexicanus</i>	343	<i>hevacornis</i>	389
<i>olisthostoma</i>	342	<i>Podothecus</i>	395
<i>patao</i>	342	<i>Pteropodus</i>	369
<i>peruvianus</i>	342	<i>Rabida</i>	156
<i>plumieri</i>	342, 343	<i>Salmo</i>	58
<i>rhombus</i>	342	<i>gairdneri</i>	58
<i>simillimus</i>	342	<i>Sebastodes</i>	369
<i>squamipinnis</i>	342	<i>Stolephorus</i>	51
<i>vaigiensis</i>	341	<i>Ulocentra</i>	286
<i>zebra</i>	342	<i>Gilbertida</i>	392
<i>Gerridae</i>	340	<i>sigulatus</i>	393
<i>Gerris</i>	341	<i>Gilbertina</i>	392
<i>ghini</i> , <i>Orthogoriscus</i>	503	<i>sigalutes</i>	392, 393
<i>gibba</i> , <i>Hystrio</i>	505	<i>Gillulus</i>	455
<i>gibber</i> , <i>Salmo</i>	54	<i>arenicola</i>	455
<i>gibbiceps</i> , <i>Heros</i>	417	<i>ornatus</i>	455
<i>Theraps</i>	417	<i>semicinctus</i>	455
<i>gibbifrons</i> , <i>Cypselurus</i>	202	<i>gilli</i> , <i>Ayocettina</i>	83
<i>Exocoetus</i>	202	<i>Clinus</i>	460
<i>gibbonsi</i> , <i>Holconotus</i>	412	<i>Hyporhamphus</i>	198
<i>Gibbonsia</i>	458	<i>Labichthys</i>	83
<i>elegans</i>	458	<i>Leuciscus</i>	118
<i>montereyensis</i>	458		

Last of scientific names—Continued

	Page		Page
gilli, Malacopterus.....	460	glaucus, Ciliata.....	208
Pholis.....	468	Lichia.....	276
Pleuronectes.....	228	Prionace.....	17
Synchrus.....	393	glaucotranum, Coryphopterus.....	439
gillia.....	441	Rhinogobius.....	439
gillianus, Julis.....	420	glaucopareus, Acanthurus.....	363
Gillias.....	458	Glaucostegus.....	23
jordani.....	438	glaucostictus, Rhinobatus.....	23
Gillchthys.....	444	glaucostigma, Rhinobatos.....	24
detrusus.....	444	Rhinobatus.....	24
guayamasiae.....	444	glaucum, Stuzostedion.....	282
mirabilis.....	444	glaucus, Chaetodon.....	276
y-cauda.....	444	Emmelas.....	364
gilli, Bassogigas.....	482	Galeus.....	17
Cetomimus.....	172	Sebastes.....	364
Eupomacentrus.....	414	Squalus.....	17
Lepomis.....	303	Trachynotus.....	276
Lipogenys.....	75	globiceps, Agonostoma.....	255
Neobythites.....	481	Blennicottus.....	391
Pomacentrus.....	411	brvostus, Blennicottus.....	391
Sebastodes.....	368	Gasterosteus.....	238
Sebastomus.....	368	Joturus.....	255
Stephanoberyv.....	232	Oligocottus.....	391
Xiphophorus.....	192	globosa, Lyosphara.....	503
Xystroplites.....	301, 302	globosum, Astatheios.....	421
Ginglymodi.....	36	Cichlasoma.....	421
Ginglymostoma.....	12	gloriosus, Bryttus.....	304
cnboverdianus.....	12	Enneacanthus.....	304
cirratum.....	12	Glossanua pandionis.....	296
fulvum.....	12	Glossichthys.....	230
Ginglymostomidae.....	12	Glossodon.....	40
giorna, Cephalopterus.....	32	harengoides.....	40
girardi, Acipenser.....	34	heterurus.....	40
Caranx.....	272	glossodonta, Argentina.....	40
Pseudogulis.....	428	Glossodus.....	39
Girardinichthys.....	182	forskali.....	40
innominatus.....	182	Glossoplites.....	302
limnurgus.....	183	gloveri, Salmo.....	56
Girardinus.....	189	glutinosa, Myxine.....	8
creolus.....	189	Glyphodon chrysurus.....	415
denticulatus.....	189	rudis.....	415
garmani.....	189	savathis.....	415
guppil.....	191	taurus.....	415
metallicus.....	189	troscelli.....	415
pleurospilus.....	190	Glyphisodon.....	414
sonoriensis.....	189	chrysurus.....	415
uninotatus.....	188	moucharra.....	415
versicolor.....	191	rubicundus.....	415
Girella.....	338	Glyptocephalus.....	228
dorsomacula.....	339	acadianus.....	228
nigricans.....	339	cynoglossus.....	228
punctata.....	338	pacificus.....	228
simplicidens.....	339	zachirus.....	228
Girellidae.....	338	Gnathanodon.....	273
Giton.....	101	Gnathobolus.....	45
glaber, Cottus.....	487	Gnathocentrum.....	362
Cyclopterichthys.....	398	Gnathanodon.....	273
Pleuronectes fuscus subrotundus.....	230	speciosus.....	273
glabra, Acipenser.....	34	gnathodus, Pseudoscarius.....	434
Liopsetta.....	227	Scarus.....	434
Platessa.....	227	Gnatholepis.....	442
Pleuronectes satis parva et.....	217	thompsoni.....	443
glacialis, Cottus.....	387	Gnathophis.....	80
Liopsetta.....	227	guppil.....	80
Pleuronectes.....	227	heteroganthus.....	80
Scopelus.....	170	porigae.....	80
Squalus.....	22	Gnathypops.....	453
glacoides, Trachynotus.....	276	aurifrons.....	453
gladifer, Istiophorus.....	263	cuvieri.....	453
Gladiunculus.....	238	macrops.....	453
bispinosus.....	238	maxillosa.....	453
gladiunculus, Gasterosteus.....	238	mystacina.....	453
gladius, Histiophorus.....	263	mystacinus.....	453
Notistium.....	263	rhonalea.....	453
Scomber.....	263	scops.....	453
Tylosurus.....	190	snyderi.....	453
Xiphias.....	264	Gobiesocidae.....	488
Glanfostomi.....	33	Gobiasox.....	488
Glandichthys.....	183	adustus.....	489
falcatus.....	188	androsensis.....	489
torralbasi.....	188	cephalus.....	488, 489
uninotatus.....	188	cerasinus.....	489
Glandodon.....	183	costaricensis.....	489
latidens.....	183, 189	eigenmanni.....	490
		eos.....	491

List of scientific names—Continued

	Page		Page
Gobiosor, erythroptus.....	490	Gobius, claytoni.....	439
funebris.....	489	cristagalli.....	438
gyrinus.....	489	crista-galli.....	438
hæres.....	490	curaçao.....	439
humeralis.....	490	cyprinoides.....	438
macrophthalmus.....	489	dalli.....	442
marmoratus.....	488	dolichocephalus.....	442
muscarum.....	490	eigenmanni.....	439
nigripinnis.....	489	emblematicus.....	443
papillifer.....	488	enceomus.....	439
pinniger.....	488	fuscus.....	440
rhessodon.....	490	garmani.....	439
rhodospilus.....	489	gracilis.....	444
(Rimicola) beryllinus.....	490	gronovi.....	268
rupestris.....	491	gulosus.....	443
sancti-martini.....	490	hemigymnus.....	445
strumosus.....	489	hipoliti.....	440
tudes.....	489	homocyanus.....	440
virgatulus.....	489	kraussi.....	441
vittatus.....	489	lacertus.....	440
yuma.....	490	lanceolatus.....	441
zebra.....	491	lepidus.....	444
Gobiodæ.....	438	lineatus.....	440, 446
Gobio æstivalis.....	138	longicaudus.....	441
cataractæ.....	139	lucretiæ.....	443
gelidus.....	138, 139	lyricus.....	441
plumbeus.....	136	mangicola.....	439
vernalis.....	137	mapo.....	440
gobio, Chinus.....	455, 456	martinicus.....	442
Cottus.....	383	mexicanus.....	442
Gobioclinus.....	456	microdon.....	441
Gobioclinus.....	455	nebulopunctatus.....	440
gobio.....	456	nebulopunctatus.....	440
Gobioides.....	448	newberryi.....	444
barreto.....	448	nicholsi.....	440
broussonnetii.....	448	niger.....	440
gobioides, Cottus.....	385	oblongus.....	448
Hypsicometes.....	451	oceanicus.....	441
Gobioidæ.....	448	ocellaris.....	442
Gobiomorus.....	435	padangensis.....	440
dormitor.....	435	paradoxus.....	445
lateralis.....	436	parvus.....	439
Gobionellus.....	440	pisonis.....	436
bayamonensis.....	441	plumieri.....	447
hastatus.....	440, 441	poeyi.....	442
lyricus.....	441	punctillatus.....	440
microdon.....	441	quadriporus.....	439
oceanicus.....	441	sandvicensis.....	440
sagittula.....	441	semifasciatus.....	441
smaragdus.....	441	seminudus.....	445
strigatus.....	441	shufeldti.....	439
stigmaticus.....	441	smaragdus.....	440, 441
gobionum, Campo-stoma.....	146	soporator.....	440
Gobiosoma.....	445	stigmaturus.....	439
alepidotum.....	446	taasica.....	442
bosci.....	446	thalassinus.....	444
cauthocum.....	446	townsendi.....	444
crescentale.....	446	translucens.....	440
crescentalis.....	446	viridi pallidus.....	446
histrio.....	446	wurdemanni.....	441
horsti.....	446	zebra.....	441
ios.....	445	Goboidea.....	435
longipinne.....	445	godmani, Pimelodus.....	157
longum.....	446	Rhambdia.....	157
molestum.....	446	godmanni, Cichlaurus.....	416
multifasciatum.....	446	Heros.....	416
zosterurum.....	445	Rivulus.....	180
Gobius.....	440	godmani, Batrachoides.....	487
albopunctatus.....	440	gomesi, Ophichthus.....	88
alepidotus.....	445, 446	Ophisurus.....	87, 88
amorea.....	436	gomphodon, Oxyrhina.....	19
anjerensis.....	442	Gonenion.....	280
bacalaus.....	441	serra.....	280
banana.....	442	Goniobatis.....	30
bayamonensis.....	441	macroptera.....	30
boleosoma.....	439	Goniodus.....	22
bosci.....	445, 446	Goniomacrus.....	27
boscu.....	442	Gonioplectrus.....	308
brasiliensis.....	448	hispanus.....	308
breviceps.....	440	Gonocephalus.....	408
brunneus.....	440	macrocephalus.....	408
carolinensis.....	440	Gonocheilodon.....	359
catulus.....	440	Gonoptychus.....	362
chiquita.....	443	mœrens.....	362
		mœreus.....	362

List of scientific names—Continued

	Page		Page
Gonostoma.....	72	gracilis, Scopelus.....	169
denudata.....	72	Tigoma.....	119
denudatum.....	72	Umbrina.....	351
elongatum.....	72	Xiphophorus.....	185
microdon.....	72	gracilispinis, Dolopichthys.....	509
Gonostomidae.....	72	gracilipes, Lamnonema.....	213
Goodea.....	183	graciosus, Pimelodus.....	152
atripinnis.....	183	gradiens, Hypsagonus.....	393
calentis.....	184	grællis, Ophidion.....	485
captiva.....	184	grahami, Gila.....	114
luitpoldi.....	184	Oligocephalus.....	291
toweri.....	184	grallator, Benthosaurus.....	166
whitei.....	183	Gramma.....	322
goodei, Aetobatus.....	31	loreto.....	372
Aldrovandia.....	75	Grammateus.....	336
Centropholis.....	206	medius.....	337
Chriopeops.....	175	grammaticum, Thalassoma.....	429
Erinzyon.....	109	Grammatopleurus.....	373
Halosaurus.....	75	Grammatostomus.....	70
Lucania.....	175	dentatus.....	70
Myliobatis.....	31	Grammichthys.....	229
Paralanchurus.....	351	Grammicolepidæ.....	232
Ptilichthys.....	471	Grammicolepis.....	232
Sebastodes.....	364	brachiusculus.....	232
Spinivomer.....	84	Grammiconotus.....	197
Trachinotus.....	277	bicolor.....	197
Urolophus.....	30	Grammistes acuminatus.....	353
Urotrygon.....	30	hepatus.....	335
Zaclemus.....	351	mauriti.....	332
Goodeidæ.....	182	trivittatus.....	331
goou, Macrurus.....	203	granadense, Astatheros.....	421
Nematonurus.....	203	Cichlasoma.....	421
gorbuscha, Onchorhynchus.....	54	grandicassis, Arius.....	149
Salmo.....	54	grandiceps, Microlepidium.....	211
Gorduchthys.....	89	Paraliparis.....	404
irretitus.....	89	Pseudocottus, Auchenocottus.....	457
gorensis, Albulæ.....	40	Pseudocottus, Auchenocottus.....	371
Hynnus.....	274	Pseudocottus, Auchenocottus.....	129
Vomer.....	275	Pseudocottus, Auchenocottus.....	176
Gorgasia.....	77	Pseudocottus, Auchenocottus.....	176
punctata.....	77	Gila.....	114
gorgonæ, Cheironodon.....	96	Pomadasis.....	334
Compura.....	96	Pomadasy.....	334
gorensis, Vomer.....	275	Ptychocheilus.....	114
gouani, Lepidopus.....	262	grandisquamis, Tetracanthus.....	436
Trachipterus.....	216	grandisquamis, Tetracanthus.....	426
gracile, Boleosoma.....	294	Upeneus.....	344
Myctophum.....	169	grandoculatus, Centropomus.....	306
Peristedion.....	405	grandocule, Chirostoma.....	251
gracilicauda, Macrourus.....	206	grandoculus, Atherinichthys.....	251
Macurus.....	206	granulata, Perca.....	281
gracilicirrus, Umbrina.....	350	Raja.....	25
gracilior, Gambusia.....	186	Raja.....	25
gracilis, Aldrovandia.....	75	granulatus, Amorphocephalus.....	430
Atherinichthys.....	247	granulosa, Pristis.....	23
Bathynectes.....	481	Graodus.....	133
Blennius.....	471	nigrotæniatus.....	132, 133
Boleichthys.....	294	grayi, Carpiodes.....	103
Catostomus.....	105	Lepidosteus.....	37
cayuga, Cottus.....	386	grebnitzki, Neoliparis.....	399
Chaetodon.....	360	Greeleya.....	390
Cottus.....	383, 385	rubellio.....	391
Cyprinus (Leuciscus).....	136	greenei, Cottus.....	384
Diapterus.....	341	Porichthys.....	487
Gadella.....	211	Uranidea.....	384
Gadus.....	210, 211	greeni, Couesius.....	137
Gambusia.....	185	Liparis.....	399
Gila.....	114	Neoliparis.....	399
Gobius.....	444	Polypera.....	399
Hippocampus.....	244	grex, Pneumatophorus.....	258
Hybopsis.....	133	Scomber.....	258
Leptocephalus.....	79	grevi, Istiophorus.....	263
Leptogunnellus.....	470	grimaldi, Conchognathus.....	78
Leuciscus.....	128	grisea, Cynoperca.....	282
Lumpenus.....	470	Dionda.....	144
Lycodes.....	475	Lucioperca.....	282
Mollinesia.....	192	Parapetenia tetracantha.....	419
Mollinesia.....	192	Sciæna.....	352
Moniana.....	131	griseolineatus, Syngnathus.....	240
Perca.....	281	griseus, Carcharias.....	19
Photonectes.....	70	Catostomus.....	107
Pimelodus.....	152	Catostomus (Acomus).....	107
Platygobio.....	136	Heros tetracanthus.....	419
Ptychocheilus.....	113, 114	Hexanchus.....	11
Scomber.....	258	Labrus.....	325

List of scientific names—Continued

	Page		Page
griseus, Lutianus.....	325, 326	Sula, Eudnostomus.....	341
Mesoprion.....	325	Gerres.....	340, 341
Saurus.....	163	Gila.....	120
Squalus.....	11	gulo, Holocentrus.....	308
grixalvi, Astroblepus.....	159	gulonellus, Pogonichthys (Platygnobio).....	136
grenlandica, Microstomus.....	68	gulosa, Cathorops.....	151
Nansenia.....	68	gulosus, Chænobryttus.....	302
grenlandicus, Acanthocottus.....	387	Cottopsis.....	383
Aspidophoroides.....	397	Cottus.....	383
Cottus.....	386, 387	Gobius.....	443
Gunnellus.....	468	Microgobius.....	443
Himantolophus.....	509	Parachromis.....	418
Hippoglossus.....	221	Pomotis.....	302, 303
Salmo.....	66	Tachisurus.....	151
Gronias.....	154	gulula, Eucnostomus.....	341
nigrilabris.....	154	gummigutta, Hypoplectrus unicolor.....	321
gronovii, Gobius.....	268	Plectropoma.....	321
Nomeus.....	268	gundlachi, Dormitator.....	436
Ostracion.....	497	gunelliformis, Asternopteryx.....	468
Solea.....	229	Gunnellus.....	467
Zoarces.....	473	fabricii.....	470, 471
grossidens, Engraulis.....	51	grenlandicus.....	468
Lycengrulus.....	51	murenoides.....	468
Lycodapus.....	479	ornatus.....	468
grummiens, Ambloodon.....	352	ruberrimus.....	468
Aplodinotus.....	352	vulgaris.....	467
Mugil.....	352	gunnellus, Blennius.....	467
gryllus, Exocoetus.....	200	Pholis.....	467
Grystes lineatus.....	373	gunneri, Scomber.....	217
Guabicoara brasiliensisibus.....	330	Squalus.....	20
guacamaia, Pseudoscarus.....	435	gunnisoni, Cyprinella.....	130
Scarus.....	435	guntheri, Alvordius.....	283
Guacamaya.....	435	Aspidophoroides.....	397
guacari, Hypostomus.....	160	Halosaurus.....	74
guachanco, Sphyræna.....	255	Hoplopogrus.....	324
guacharote, Ancistrus.....	161	Lampanyctus.....	167
Hypostomus.....	161	Mugil.....	253
guadalupe, Atherinops insularum.....	252	Sphyræna.....	255
guadalupensis, Ama.....	296	Ulcina.....	397
Apogon.....	296	Xiphophorus.....	193
guacan, Echeineis.....	449	Xiphophorus helleri.....	193
Guamaia atinga.....	502	guntheriana, Alutera.....	496
Guamajacu guara.....	501	guppi, Congromuraena.....	80
guanabana, Erizo.....	501	Girardinus.....	191
Guaperva.....	491	Gnathophis.....	80
lata.....	492	Gurnardus.....	406
guara, Guamajacu.....	501	birostratus.....	406
Guarapuca.....	257	gymnostethus.....	406
gusa, Serranus.....	312	loxias.....	406
guatemalensis, Anacyrtus.....	98	ruscarius.....	406
Arius.....	150	xenisma.....	406
Brycon.....	98	guttata, Limamuræna.....	90
Chirostoma.....	249	Perca.....	311
Citharichthys.....	219	Raia.....	28
Fundulus.....	174	Scorpæna.....	371
Galeichthys.....	150	guttatus, Astrodermus.....	264
Pimelodus.....	157	Astroscompus.....	454
Rhamdia.....	157	Balistes.....	492
Rœboides.....	98	Bodianus.....	309
Thyrina.....	249	Chirus.....	373
Guativera amarillas.....	309	Dasyatis.....	28
guativera, Bodianus.....	309	Epinephelus.....	311
Guatucupa juba.....	332	Johnius.....	313
Guavina.....	435, 436	Labrus.....	426
guavina.....	436	Lutianus.....	326
guavina, Eleotris.....	436	Mesoprion.....	326
Guavina.....	436	Ophisurus.....	87
guayamasæ, Gillichthys.....	444	Perca.....	308
guaza, Epinephelus.....	310	Percopsis.....	214, 215
Labrus.....	310	Pomotis.....	304
guntheri, Galeichthys.....	149	Zeus.....	216
Homalogryles.....	310	Zygonectes.....	178
Mugil.....	253	guttavarium, Plectropoma.....	321
Siremo.....	483	guttavarius, Hypoplectrus unicolor.....	321
Guentheridia.....	500	guthrie, Ophichthys.....	87
formosa.....	500	Ophichthys.....	87
guanensis, Anchoviella.....	49	guttulata, Hypopsetta.....	225
Stolephorus.....	49	guttulatus, Cichlaurus.....	416
Strongylura.....	196	Heros.....	416
Tylosurus.....	196	Pisodonophis.....	85
gulchenoti, Callechelys.....	86	Pleuronichthys.....	225
Cayenna.....	448	guzmanensis, Catostomus (Acomus).....	104
gula, Thyrina.....	248	Gymnacanthus.....	389
guineensis, Ostracion.....	497	galeatus.....	389
Guritinga.....	149	Gymnaciniræ.....	229

List of scientific names—Continued

	Page		Page
Gymnachirus.....	229	gymnurus, Gasterosteus.....	237
fasciatus.....	229	gyrans, Querumana.....	253
nudus.....	229	Gyrinichthys.....	403
Gymneipignathus.....	270	minytrems.....	403
Gymneleotris.....	438	gyrinops, Cyclopteroide.....	398
seminuda.....	438	gyrinus, Eleotris.....	436
Gymnelis.....	477	Eurymen.....	401
conorhynchus.....	477	Gobiesox.....	489
pictus.....	477	Schilbeodes.....	155
stigma.....	477	Silurus.....	155
viridis.....	477	Gyropleurodus.....	10
unimaculatus.....	477	francesci.....	10
Gymnelus pictus.....	477		
Gymnocanthus.....	389	Hadropterus.....	283
galeatus.....	389	evermanni.....	284
pistilliger.....	389	maculatus.....	283
tricuspid.....	389	maxinkuckiensis.....	284
Gymnocephalus.....	267	nigrofasciatus.....	283
messinensis.....	267	roanoka.....	284
ruber.....	309	scierus.....	284
Gymnochinus.....	466	serrula.....	284
cristulatus.....	466	sellaris.....	284
Gymnodontes.....	497	schumardi.....	285
Gymnogaster.....	262	tessellatus.....	289
arcticus.....	216	(Hydropterus) cymatotænia, Etheostoma.....	285
gymnogaster, Sicydium.....	447	mangue, Etheostoma.....	285
Gymnomuræna.....	94	ouachitæ, Etheostoma.....	284
doliata.....	94	squamatus, Etheostoma.....	285
nectura.....	95	hæmaturus, Hybopsis.....	145
Gymnonoti.....	101	Hæmulidæ.....	329
Gymnotus acus.....	486	Hæmulon.....	329
carapo.....	486	acutum.....	329
Gymnopsis.....	94	albidum.....	329
Gymnosarda.....	258	album.....	329
gymnostethus, Gurnardus.....	406	arars.....	330
Prionotus.....	406	arcuatum.....	330
Gymnothorax.....	91	aurolineatum.....	331
afer.....	91	bonariense.....	329
albimentis.....	93	brevirostrum.....	330
aqueduleis.....	93	canna.....	329
brunneus.....	92	carbonarium.....	330
carcinognathus.....	93	caudimacula.....	329
castaneus.....	92	chromis.....	329
catenatus.....	94	chrysargyreum.....	330, 331
chlevastes.....	93	chrysopterus.....	329
concolor.....	93	continuum.....	329
conspersus.....	92	corvinæforme.....	333
dovii.....	92	corvinæformis.....	333
elaboratus.....	93	dorsale.....	330
flavoscriptus.....	91	elegans.....	329
funebis.....	92	flaviguttatus.....	331
jordani.....	92	flavolineatum.....	330
(Limamuræna) argus.....	90	formosum.....	330
millaris.....	92	fremebundum.....	329
mordax.....	91	heterodon.....	330
moringa.....	91	hians.....	330
muræna.....	91	jenguano.....	331
nigrocastaneus.....	90	labridum.....	332
obscuratus.....	93	luteum.....	330
ocellatus.....	91, 93	macrostoma.....	329
nigromarginatus.....	93	macrostomum.....	329
saxicola.....	93	maculosum.....	329
picturatus.....	91	margaritiferum.....	331
polygonus.....	92	mazatlanum.....	331
punctarenæ.....	92	melanurum.....	330
punctatus.....	92	microphthalumum.....	329
rostratus.....	91	mowbrayi.....	329
sanctæ-helenæ.....	92	multilineatum.....	330
scriptus.....	92	notatum.....	329
umbrosus.....	90	obtusum.....	332
verilli.....	93	parra.....	329
versipunctatus.....	92	plumieri.....	330
vicinus.....	92	quadrilineatum.....	331
virescens.....	92	quinquelineatum.....	331
Gymnotidæ.....	101	retrocurrans.....	329
Gymnotorpedo.....	27	rimator.....	331
Gymnotus.....	101	schrunkii.....	330
acus.....	101	sciurus.....	330
albifrons.....	101	scudderi.....	330
albus.....	101	serratum.....	329
carapo.....	101	sextasciatum.....	329
fasciatus.....	101	sextasciatus.....	329
macrourus.....	101	similis.....	330
macurus.....	101	steindachneri.....	330
putaol.....	101	subarcuatum.....	330
gymnura, Trygon.....	28	tæniatum.....	330, 331

List of scientific names—Continued

	Page		Page
<i>Hæmulon undecimale</i>	330	<i>Harpe cœruleo-aureus</i>	425
<i>xanthopteron</i>	330	<i>diplotænia</i>	425
<i>Hæmulopsis</i>	333	<i>pectoralis</i>	425
(<i>Hæmulopsis</i>) <i>axillare</i> , <i>Pristipoma</i>	333	<i>rufa</i>	309
<i>brevipinna</i> , <i>Pristipoma</i>	335	<i>harperi</i> , <i>Sphæroides</i>	498
<i>nitidum</i> , <i>Pristipoma</i>	333	<i>Sphæroides</i>	498
<i>hæres</i> , <i>Gobiosox</i>	490	<i>Harporus</i>	362
<i>Sicyases</i>	490	<i>fasciatus</i>	362
<i>Halatractus</i>	279	<i>harringtonensis</i> , <i>Atherina</i>	246
<i>dorsalis</i>	279	<i>harringtoni</i> , <i>Axyrinus</i>	377
<i>halec</i> , <i>Clupea</i>	41	<i>Harriotta</i>	33
<i>Halecomorphi</i>	38	<i>curtiss-jamesi</i>	33
<i>Halecula</i>	46	<i>raleighana</i>	33
<i>acuminata</i>	46	<i>harroweri</i> , <i>Isopisthus</i>	354
<i>Halias</i>	479	<i>harti</i> , <i>Haplochilus</i>	180
<i>Halichæres</i>	426	<i>Rivulus</i>	180
<i>californicus</i>	428	<i>hasselti</i> , <i>Caranx</i>	270
<i>seilheri</i>	428	<i>Paraserranus</i>	318
<i>Halientæa</i>	507	<i>hasta</i> , <i>Lutjanus</i>	334
<i>spongiosa</i>	508	<i>hastata</i> , <i>Pastinaca</i>	28
<i>stellata</i>	507	<i>hastatus</i> , <i>Dasyatis</i>	28
<i>Halientella</i>	508	<i>Gobionellus</i>	440, 441
<i>lappa</i>	508	<i>hastingsi</i> , <i>Lutjanus</i>	326
<i>Halieutichthys</i>	507	<i>Neomamis</i>	326
<i>aculeatus</i>	507	<i>Haustor</i>	152
<i>caribbæus</i>	507	<i>catus</i>	153
<i>reticulatus</i>	507	<i>dugesi</i>	153
<i>smithi</i>	507	<i>lacustus</i>	152
<i>Haliperca</i>	318	<i>lophius</i>	153
<i>halleri</i> , <i>Urobatus</i>	29, 30	<i>lupus</i>	153
<i>Urolophus</i>	29	<i>okeechobeensis</i>	152
<i>Haloporphyrus viola</i>	211	<i>pricei</i>	153
<i>Haloporphyrus</i>	211	<i>havana</i> , <i>Geires</i>	342
<i>halops</i> , <i>Acipenser</i>	34	<i>Xystæma</i>	342
<i>Halosaurichthys</i>	74	<i>havannensis</i> , <i>Muraena</i>	87
<i>Halosauridæ</i>	74	<i>hawaiiensis</i> , <i>Klops</i>	39
<i>Halosauropsis</i>	75	<i>hawaiiensis</i> , <i>Macrohamphosus</i>	246
<i>Halesaurus</i>	74	<i>haydeni</i> , <i>Ptychostomus</i>	109
<i>attenuatus</i>	75	<i>hayi</i> , <i>Hybognathus</i>	144
<i>goodei</i>	75	<i>hearneri</i> , <i>Salmo</i>	60
<i>guntheri</i>	74	<i>heathi</i> , <i>Schedophilus</i>	268
<i>macrochir</i>	75	<i>heckeli</i> , <i>Acus</i>	244
<i>oweni</i>	74	<i>Balistes</i>	493
<i>oweni</i>	74	<i>Gambusia</i>	185
<i>radiatus</i>	74	<i>Nerophis</i>	244
<i>rostrata</i>	75	<i>hedricki</i> , <i>Astatheros</i>	421
<i>rostratus</i>	75	<i>Cichlasoma</i>	421
<i>hamatus</i> , <i>Icelus</i>	378	<i>heermanni</i> , <i>Amphistichus</i>	411
<i>hamiltoni</i> , <i>Brachioptilon</i>	32	<i>heilneri</i> , <i>Labrisomus</i>	459
<i>Hippoglossoides</i>	222	<i>Heiroptera</i>	25
<i>Manta</i>	32	<i>helenæ</i> , <i>Muraena</i>	90
<i>hammondi</i> , <i>Pimelodus</i>	152	<i>Muraenophis</i>	90
<i>hammondi</i> , <i>Percopsis</i>	215	<i>Helases</i>	413
<i>Semotilus</i>	117	<i>enchrysurus</i>	413
<i>hankinsoni</i> , <i>Parartedius</i>	377	<i>insolatus</i>	413
<i>Haplochilus dovii</i>	179	<i>marginatus</i>	413
<i>floripinnis</i>	177	<i>multilineatus</i>	412
<i>harti</i>	180	<i>Helastes</i>	413
<i>melanops</i>	185	<i>Helicolenus</i>	370
<i>Haplodoci</i>	486	<i>dactylopterus</i>	370
<i>haplognatha</i> , <i>Helioperca</i>	301	<i>maderensis</i>	370
<i>haplognathus</i> , <i>Lepomis</i>	301	<i>Helioperca</i>	301
<i>Haplodontotus</i>	352	<i>bombifrons</i>	301
<i>Haplomi</i>	172	<i>haplognatha</i>	301
<i>bardwicki</i> , <i>Ratabouri</i>	89	<i>incisor</i>	301
<i>harengoides</i> , <i>Glossodon</i>	40	<i>macrochira</i>	301
<i>Harengula</i>	43	<i>marginata</i>	301
<i>callolepis</i>	43	<i>nephela</i>	301
<i>clupeola</i>	43	<i>helleri</i> , <i>Anchoviella</i>	48
<i>humeralis</i>	43	<i>brevis</i> , <i>Xiphophorus</i>	194
<i>latula</i>	43	<i>guntheri</i> , <i>Xiphophorus</i>	193
<i>macrophthalmus</i>	43	<i>Heros</i>	422
<i>maculosa</i>	43	<i>ranchovii</i> , <i>Xiphophorus</i>	193
<i>pensaicolæ</i>	43	<i>strigatus</i> , <i>Xiphophorus</i>	193
<i>sardina</i>	43	<i>Thoricthys</i>	422
<i>thryssina</i>	43	<i>Xiphophorus</i>	193
<i>harengulus</i> , <i>Eucinostomus</i>	341	<i>helolepis</i> , <i>Trachyrhynchus</i>	203
<i>harengus</i> , <i>Clupea</i>	41	<i>Helmichthys</i>	79
<i>Lavinia</i>	113	<i>diaphanus</i>	79
<i>Leucichthys</i>	52	<i>Helmictis</i>	79
<i>Myxus</i>	253	<i>heloepis</i> , <i>Trachyrhynchus</i>	203
<i>Salmo</i> (<i>Coregonus</i>).....	62	<i>Trachyrhynchus</i>	203
<i>harfordi</i> , <i>Ptychocheilus</i>	114	<i>Helops</i>	34
<i>harlani</i> , <i>Lepidosteus</i>	37	<i>helops</i> , <i>Acipenser</i>	34
<i>Harpe</i>	425	<i>helvamaculatus</i> , <i>Sebastes</i>	367
<i>cœruleo-aureum</i>	425	<i>Hemiancistrus</i>	161

List of scientific names—Continued

	Page		Page
Hemianthias.....	322	Hemipterus.....	393
peruanus.....	322	americanus.....	393
vivianus.....	322	cavifrons.....	393
(Hemianthias) peruanus, Anthias.....	322	marinatus.....	382, 393
Hemiarus.....	149	villosus.....	393
Hemibranchiu.....	237	Hemiphysophorus.....	184, 185
Hemibrycon.....	97	hemphiliu, Stathmonotus.....	465
dariensis.....	97	henlei, Carcharias.....	16
(Hemibrycon) polyodon, Tetragonopterus.....	271	Rhinichthys.....	13, 14
Hemicarany.....	271	henshalli, Zygonectes.....	177
amblyrhynchus.....	271	henshawi, Rhinichthys.....	141
atrimanus.....	271	henshawi, Salmo.....	67
falcatus.....	271	hantz, Blennius.....	462
furthii.....	271	Hypsoblennius.....	462
leucurus.....	271	hepatus, Acanthurus.....	362
marginatus.....	271	Grammistes.....	362
rhomboides.....	271	Teulius.....	362
zelotes.....	271	Hepsetia.....	246
Hemichætodon.....	359	evei manni.....	246
Hemielectris.....	437	supes.....	246
latifasciatus.....	437	hepsetus, Athena.....	246, 247
Hemierocetus.....	199	heptacanthus, Cheilodipterus.....	280
caudimaculatus.....	199	heraldi, Tetrodon.....	499
Hemigalea.....	14	Heichthys.....	422
Hemigaleus.....	14	bocourti.....	423
microstoma.....	14	carpinth.....	423
pectoralis.....	14	cyanoguttatus.....	422
Hemigrammus.....	98	geddesi.....	423
compressus.....	97	gervonaceus.....	422
minutus.....	97	trocheli.....	423
unilineatus.....	98	underwoodi.....	423
hemigymna, Garmanina.....	443	hermuer, Blennius.....	469
hemigymnus, Argyropelecus.....	74	Labisonus.....	459
Gobius.....	445	Hermulus.....	426
Hemilepidotus.....	380	Hermosilla.....	340
gibbsii.....	381	azulea.....	340
hemilepidotus.....	381	robusta.....	340
jordanii.....	381	Heteros.....	416
lulesii.....	381	affinis.....	422
rapus.....	381	alifrons.....	420
hemilepidotus, Coltus.....	380, 381	angulifer.....	416
Hemilepidotus.....	381	anus.....	422
Hemiodon.....	161	baleatus.....	418
depressus.....	161	basilaris.....	421
platycephalus.....	161	beam.....	419
hemiodon, Carcharias.....	17	bifasciatus.....	416
Hemiodontidae.....	100	callolepis.....	422
Hemiorolichia.....	161	centricetus.....	420
caiacasensis.....	161	(Cichlasoma) spinosissimus.....	420
Hemiplites.....	304	citrinellus.....	420, 422
sumians.....	304	deppii.....	416
Hemipus.....	114	dovi.....	418
lacustris.....	114, 115	erythreus.....	422
hemipus, Cyprinus.....	115	friedrichstahl.....	418
Hemiramphidae.....	198	geddesi.....	423
Hemiramphus.....	199	gibbiceps.....	417
balao.....	199	godmanni.....	416
basiliensis.....	199	guttulatus.....	416
brownii.....	199	helleri.....	422
fasciatus.....	198	heterodontus.....	420
filamentosus.....	199	intermedius.....	416
longirostris.....	198	istanus.....	418
macrochilus.....	199	labiatum.....	422
picati.....	198	labridens.....	419
pleii.....	199	lentiginosus.....	422
richardi.....	198	lobochilus.....	422
roberti.....	198	longimanus.....	420
rose.....	198	maculipinnis.....	422
saltator.....	199	managuensis.....	418
unifasciatus.....	198	margaritifer.....	421
Hemirhombus.....	218	melanopogon.....	421
aramaca.....	218	melanurus.....	416
fimbriatus.....	220	mento.....	419
fuscus.....	219	microphthalmus.....	416
gruneensis.....	218	montezuma.....	421
ovalis.....	218	motaguensis.....	418
Hemustoma.....	433	multispinosus.....	423
reticulata.....	433	nebulifer.....	417
Hemuthemia.....	121	nicaraguensis.....	418
maculata.....	135	nigricans.....	419
mulieriana.....	121	nigrofasciatus.....	420
vittata.....	121	oblongus.....	421
Hemutipteridae.....	393	octofasciatus.....	421

List of scientific names—Continued

	Page		Page
<i>Heros parma</i>	419	<i>heterurus</i> , <i>Cypselurus</i>	201
<i>pavonaceus</i>	422	<i>Exocoetus</i>	201
<i>rostratus</i>	421	<i>Glossodon</i>	40
<i>salvini</i>	418	<i>hexacanthus</i> , <i>Centrarchus</i>	304, 305
<i>severus</i>	416	<i>Dipterodon</i>	295
<i>sieboldii</i>	423	<i>hexacornis</i> , <i>Cottus</i>	388
<i>spilurus</i>	420	<i>Oncocottus</i>	388
<i>teporatus</i>	422	<i>hexagonatus</i> , <i>Stomias</i>	69
<i>tetracanthus cinctus</i>	419	<i>Hexagrammidae</i>	373
<i>griseus</i>	419	<i>Heaagrammos</i>	373, 374
<i>latus</i>	419	<i>asper</i>	373, 374
<i>torralbasi</i>	419	<i>decagrammus</i>	373
<i>triagramma</i>	418	<i>lagocephalus</i>	374
<i>trunaculatum</i>	420	<i>octogrammus</i>	374
<i>trocheli</i>	423	<i>stelleri</i>	373, 374
<i>urophthalmus</i>	418	<i>supercilius</i>	374
<i>heros</i> , <i>Eupomotis</i>	302	<i>Hevagrammus scaber</i>	374
<i>Pomotis</i>	302	<i>hevagrammus</i> , <i>Labrus</i>	374
<i>Herotilapia</i>	423	<i>Hexanchidae</i>	11
<i>multispinosa</i>	423	<i>Hexanchus</i>	11
<i>Hei petoichthys</i>	87	<i>corinus</i>	11
<i>callisoma</i>	88	<i>griseus</i>	11
<i>ornatissimus</i>	87	<i>Hexanemachthys</i>	149
<i>sulcatus</i>	87	<i>surinamensis</i>	150
<i>herschelinus</i> , <i>Carelliparis</i>	400	<i>heyel</i> , <i>Rivulus</i>	180
<i>Liparis</i>	400	<i>hians</i> , <i>Abelennes</i>	197
<i>hertzbergi</i> , <i>Silurus</i>	143	<i>Belone</i>	197
<i>Hespei anthias</i>	328	<i>Hæmulon</i>	330
<i>herpès</i> , <i>Prapichthys annectens</i>	187	<i>Myctophum</i>	169
<i>Hesperoleucus</i>	121	<i>Sayris</i>	197
<i>antæus</i>	122	<i>Hiatula</i>	424
<i>meani</i>	122	<i>gardemana</i>	424
<i>mitulus</i>	122	<i>hiatula</i> , <i>Labrus</i>	424
<i>navarroensis</i>	121	<i>hieroglyphicus</i> , <i>Zygonectes</i>	178
<i>parvipinnis</i>	122	<i>Hieroptera</i>	24
<i>subditus</i>	122	<i>abredonensis</i>	24
<i>symmetricus</i>	121	<i>Hildebrandia</i>	80
<i>venustus</i>	122	<i>flava</i>	80
<i>heteracantha</i> , <i>Rhamdia</i>	158	<i>hildebrandi</i> , <i>Opsanus</i>	487
<i>heteracanthus</i> , <i>Dolopichthys</i>	510	<i>hildebrandti</i> , <i>Etheostoma</i>	291
<i>Heterandria</i>	188	<i>Oligocephalus</i>	291
<i>affinis</i>	188	<i>Hilgendorfi</i>	404, 405
<i>colombianus</i>	189	<i>hilianus</i> , <i>Eimopterus</i>	21
<i>cubensis</i>	188	<i>Exocoetus</i>	200
<i>formosa</i>	188	<i>Spinax</i>	21
<i>formosus</i>	188	<i>Himantolophidae</i>	509
<i>holbrookii</i>	185, 188	<i>Himantolophus</i>	509
<i>lutzi</i>	189, 190	<i>grenlandicus</i>	509
<i>nobilis</i>	185	<i>reinhardtii</i>	509
<i>occidentalis</i>	189	<i>himantophorus</i> , <i>Callionymus</i>	450
<i>ommata</i>	174	<i>Hiodon</i>	40
<i>patruelis</i>	184, 185	<i>clodalis</i>	40
<i>zonata</i>	191	<i>selenops</i>	41
<i>heterandria</i> , <i>Limia</i>	191	<i>tergisus</i>	40
<i>heteroclitia</i> , <i>Cobitis</i>	175	<i>Hiodontidae</i>	40
<i>heteroclitus badius</i> , <i>Fundulus</i>	176	<i>hippiti</i> , <i>Gobius</i>	440
<i>heteroclitus bermudæ</i> , <i>Fundulus</i>	175	<i>Hippichthys</i>	243
<i>heteroclitus fonticola</i> , <i>Fundulus</i>	175	<i>albistris</i>	243
<i>heteroclitus</i> , <i>Fundulus</i>	176	<i>cayorum</i>	243
<i>heteroclitus grandis</i> , <i>Fundulus</i>	176	<i>heptagonus</i>	243
<i>heteroclitus macrolepidotus</i> , <i>Fundulus</i>	176	<i>spicifer</i>	243
<i>heterodon</i> , <i>Alburnops</i>	136	<i>Hippocampinae</i>	244
<i>Hæmulon</i>	330	<i>Hippocampus</i>	244
<i>Hybopsis</i>	136	<i>brunneus</i>	244
<i>richardsoni</i> , <i>Hybopsis</i>	136	<i>fasciularis</i>	244
<i>Heterodontidae</i>	10	<i>gracilis</i>	244
<i>heterodontus</i> , <i>Astatheros</i>	420	<i>heptagonus</i>	243
<i>Heros</i>	420	<i>hudsonius</i>	244
<i>heteroglossus</i> , <i>Gadus</i>	210	<i>ingens</i>	244
<i>Heterognathi</i>	95	<i>laevicaudatus</i>	244
<i>Heterognathus</i>	250	<i>marginalis</i>	244
<i>heterognathus</i> , <i>Gnathophis</i>	80	<i>punctulatus</i>	244
<i>heterolepis</i> , <i>Hybopsis</i>	134	<i>stylifer</i>	244
<i>Notropis</i>	134	<i>zosteræ</i>	244
<i>Heteroml</i>	75	<i>Hippoglossidae</i>	221
<i>Heterophallina</i>	184, 186	<i>Hippoglossina</i>	222
<i>Heterophallus</i>	184	<i>bolmani</i>	223
<i>rachovii</i>	184	<i>macrops</i>	222, 223
<i>Heterosomata</i>	217	<i>sabanensis</i>	223
<i>Heteroprosopon</i>	225	<i>stomata</i>	223
<i>heteropygus</i> , <i>Caranx</i>	271	<i>Hippoglossoides</i>	222
<i>Heterostichus</i>	458	<i>elassodon</i>	222
<i>rostratus</i>	458	<i>exilis</i>	222
<i>heterura</i> , <i>Clupea</i>	46	<i>hamiltoni</i>	222
<i>Compsura</i>	95	<i>jordani</i>	222

List of scientific names—Continued

	Page		Page
Hippoglossoides platessoides.....	222	holbrookii, Sargus.....	338
robustus.....	222	Holcomycteropus.....	481
hippoglossoides, Pleuronectes.....	221	digitatus.....	481, 482
Reinhardtius.....	221	Holconoti.....	409
Hippoglossus.....	221	Holconotus.....	410, 411
brasilensis.....	224	agassizii.....	411
californicus.....	223	fuliginosus.....	411
gigas.....	221	gibbonsi.....	412
greenlandicus.....	221	megalops.....	410
hippoglossus.....	221	rhodoterus.....	410
maximus.....	221	throwbridgei.....	409
ocellatus.....	218	holderi, Alloeolimus.....	456
ponticus.....	221	Starksia.....	456
hippoglossus, Hippoglossus.....	221	holei, Makara.....	263
Pleuronectes.....	221	Hollardia.....	491
hippoides, Carangus.....	273	hollardi.....	491
hippops, Campostoma.....	146	hollardi, Hollardia.....	491
hippos, Caranx.....	273	hollholli, Ceratias.....	508
Acomber.....	273	Holocanthus.....	498
hippurus, Coryphæna.....	264	leionothos.....	498, 499
hirudo, Acanthurus.....	362	lunatus.....	361
Ichthyomyzon.....	9	melanotos.....	497
hirundinaceus, Squalus.....	17	holocanthus, Diodon.....	501
Hirundo.....	409	Holocentridæ.....	234
hirundo, Azurina.....	412	Holocentrum.....	235
Leiocottus.....	390	coruscum.....	236
Hispaniscus.....	368	longipinne.....	235
elongatus.....	368	marianum.....	236
levis.....	368	osculum.....	236
rubrivinctus.....	368	perlatus.....	236
sinensis.....	368	puncticulatus.....	236
zacentrus.....	368	retrospinis.....	236
hispanis, Serrana.....	353	riparium.....	236
hispanum, Plectropoma.....	308	rostratum.....	236
hispanus, Gonioplectrus.....	308	sancti-pauli.....	236
Serrana.....	353	sicclerum.....	236
hispidus, Balistes.....	494	suborbitalæ.....	236
Cottus.....	393	vexillarium.....	236
Monacanthus.....	494	Holocentrus.....	235
var maculatus, Tetrodon.....	493	auratus.....	309
Histiecthyus.....	215	bengalensis.....	324
Histiobranchus.....	78	brachypterus.....	236
bathybius.....	78	coruscus.....	236
infernalis.....	78	flavoceruleus.....	310
Histioclinus.....	456	gulo.....	308
veliger.....	456	meeki.....	236
Histiocottus.....	382	merou.....	310
bilobus.....	382	osculus.....	236
Histiophorus.....	263	rostratus.....	235
americanus.....	263	rufus.....	236
audax.....	263	sancti-pauli.....	236
gladius.....	263	sanguineus.....	363
indicus.....	263	sogo.....	235
Histrio.....	504	striatus.....	235
gibba.....	505	surinamensis.....	323
jagua.....	505	tigrinus.....	319
pictus.....	504	tortugæ.....	236
histrio, Etheostoma (Ulocentra).....	286	vexillarius.....	236
Gobiosoma.....	446	Holocephali.....	32
Lophius.....	504, 505	holocephalus, Scarus.....	434
Scorpena.....	371	Hololepis.....	294
Ulocentra.....	286	barrati.....	294
hiuleus, Stolephorus.....	48	erocrous.....	294
Holocanthus.....	361	hololepis, Cyttus.....	232
clarionensis.....	361	Zenon.....	232
cornutus.....	361	holomelas, Paraliparis.....	404
formosus.....	361	Holorhinus.....	31
lunatus.....	361	Holosteæ.....	36
passer.....	361	Holostom.....	76
strigatus.....	361	Holotrachys.....	235
tubicen.....	361	lima.....	235
tricolor.....	361	holubi, Cypselurus.....	202
holacanthus, Diodon.....	501	Exocoetus.....	202
Neopacilia.....	190	Homalagrystes.....	310
holbrookii, Acipenser.....	84	guentheri.....	310
Aluterus.....	495	Homalopomus throwbridgei.....	214
Echeneis.....	449	Homesthes.....	463
Gambusia.....	184	caulopus.....	463
Heterandria.....	185, 188	homanus, Squalus.....	20
Ophidion.....	485	homocyanus, Gobius.....	440
Ophidium.....	485	homonymus, Diapterus.....	341
Pomotis.....	302	Homopron.....	346
holbrookii, Diplodu.....	338	lanceolatus.....	246
Eupomotis.....	302	subtruncatus.....	345
Gambusia.....	185	(Homopron) furthii, Corvina.....	347
Pomotis.....	302	homopterus, Bathytroctes.....	53

List of scientific names—Continued

	Page		Page
hopkinsi, Acutomentum.....	366	humilis, Pristipoma.....	334
Hynnis.....	274	Tetragonopterus.....	96
Mycteroperca.....	314	humulus, Pagellus.....	337
Plagiogrammus.....	469	huntia, Molva.....	212
Sebastodes.....	366	Huro.....	297
Hopladelus.....	155	floridana.....	297
Hoplias.....	99	nigricans.....	297
malabaricus.....	99	huronenis, Lepisosteus.....	36
microlepis.....	99	huroniis, Argynosomus.....	62
hopliticus, Paricelinus.....	375	Leucichthys sisco.....	62
hoplomystax, Scarus.....	432	(Huso) anasimios, Acipenser.....	35
Sparisoma.....	432	anthracinus, Acipenser.....	35
Hoplopagrus.....	324	atelaspi, Acipenser.....	35
Hoplopagrus.....	324	copei, Acipenser.....	35
guntheri.....	324	kirtlandi, Acipenser.....	35
Hoplosternum.....	160	lamarii, Acipenser.....	35
punctatum.....	160	megalaspi, Acipenser.....	35
punctatus.....	160	nertinanus, Acipenser.....	35
(Hoplosternum) thoracatus, Callichthys.....	160	paranasimios, Acipenser.....	35
Hoplostethus.....	234	platyrhinus, Acipenser.....	35
cornutus.....	233	rafnesquii, Acipenser.....	35
mediterraneus.....	234	reichi, Acipenser.....	35
Hoplunnis.....	81	richardsoni, Acipenser.....	35
diomedianus.....	81	richardsoni, Acipenser.....	35
schmidti.....	81	hyalina, Suidus.....	165
horathi, Hybopsis.....	134	hyalinus, Ceratichthys.....	137
Lepidosteus.....	345	Erismus.....	137
Notropis.....	134	hyalope, Squalus.....	117
horkeli, Rhinobatos.....	23	Hybognathus.....	143
horrens, Prionotus.....	408	argyritus.....	144
horridus, Brachysomophis.....	89	civilis.....	144
horsti, Gobiosoma.....	446	(Dionda) punctifer.....	144
hoskyni, Bathyclupea.....	281	evansi.....	143
hospes, Mugil.....	254	flavipinnis.....	144
Querimana.....	254	hayi.....	144
houghi, Pimelodus.....	152	nigrotæniata.....	144
hoyi, Argynosomus.....	64	nuchalis.....	143, 144
Coregonus.....	62	osmerinus.....	143
Leucichthys.....	64	placitus.....	143
Pimelodus.....	154	procne.....	133
Uranidea.....	386	regius.....	143
hubbardii, Parophrys.....	226	siderius.....	140
Hubbessia.....	248	stramineus.....	133
gilberti.....	248	volucellus.....	133
huberi, Priapichthys.....	197	(Hybognathus) siderius, Hyborhynchus.....	140
hudsonia, Clupea.....	132	Hybopsis.....	117, 133
hudsonicus, Salmo.....	60	æstivalis marconis.....	138
Hudsonius.....	132	anogenus.....	136
aletes.....	132	atrocaudalis.....	133
altus.....	137	bifrenata.....	135
amarus.....	132	bifrenatus.....	133, 135
chamberlaini.....	132	bivittatus.....	119
euryopa.....	132	blennius.....	133
fluvialis.....	132	boops.....	134
hudsonius.....	132	buchanani.....	134
saladanus.....	132	calentis.....	134
hudsonius, Catostomus.....	106	cayuga.....	133
Hippocampus.....	244	chalybeus.....	127
Hudsonius.....	132	chiliticus.....	127
humboldtii, Cheonda.....	119	chlorocephalus.....	127
Cyclopium.....	159	chrosomus.....	127
Myctophum.....	169	crameri.....	138
Sternopygus.....	101	cummingsi.....	133
Stygogenes.....	159	deliciosa.....	134
Tigoma.....	119	straminea.....	134
humboldtiana, Atherina.....	250	deliciosus.....	134
humboldtianum, Chirostoma.....	250	dorsalis.....	134
humboldtii, Gasteropelecus.....	169	fretensis.....	129
humeralis, Arbacia.....	490	gilberti.....	134
Chaetodon.....	360	gracilis.....	133
Gobiesox.....	490	hamaturus.....	145
Harengula.....	43	heterodon.....	136
Julis.....	426	richardsoni.....	136
Leiostomus.....	349	heterolepis.....	184
Oligocephalus.....	293	horatii.....	134
Paralabrax.....	317	jordanii.....	136
Scarus.....	433	kanawha.....	134
Serranus.....	317	kermi.....	133
humeralmaculatus, Sargus.....	338	kendalli.....	135
humile, Pristipoma.....	334	lacertus.....	129
humilis, Allotis.....	300	longiceps.....	133
Bryttus.....	300	longirostris.....	135
Cottus.....	387	maculata.....	135
Gambusia.....	185	meeki.....	139
Myripristes.....	235	missouriensis.....	133

List of scientific names—Continued

	Page		Page
<i>Hybopsis montanus</i>	139	<i>hyostomus, Nocomis</i>	139
<i>nitida</i>	135	<i>hypacanthus, Psednoblennius</i>	465
<i>niveus</i>	129	<i>Hypargyrus</i>	144
<i>nocomis</i>	135	<i>Hypenetus</i>	343
<i>nux</i>	135	<i>Hypentelium</i>	107
<i>ortenburgeri</i>	135	<i>macroplectum</i>	107
<i>ozarcus</i>	135	<i>macropterum</i>	107
<i>phaenna</i>	132	<i>nigricans</i>	107
<i>phenacobius</i>	134	<i>Hyperchoristus</i>	70
<i>piptolepis</i>	134	<i>tanneri</i>	70
<i>proene</i>	133	<i>Hyperistius</i>	304, 305
<i>rubricroceus</i>	125, 127	<i>carolinensis</i>	305
<i>sabinae</i>	135	<i>Hyperoartia</i>	8
<i>soylla</i>	135	<i>Hyperotreta</i>	8
<i>shumardi</i>	134	<i>Hyperprotopon</i>	410
<i>simus</i>	135	<i>agassizii</i>	410
<i>spectrunculus</i>	134	<i>analis</i>	410
<i>stramineus</i>	134	<i>arcuatum</i>	410
<i>tetranemus</i>	139	<i>argenteum</i>	410
<i>timpanogensis</i>	119	<i>punctatum</i>	410
<i>tuditatus</i>	144, 145	<i>argenteus</i>	410
<i>volucella</i>	133	<i>Hyphalonedrus</i>	165
<i>watauga</i>	138	<i>chalybeus</i>	165
<i>winchelli</i>	133	<i>Hyphessobrycon</i>	97
<i>xenoccephalus</i>	117	<i>panamensis</i>	97
<i>Hyborhynchus</i>	145	<i>Hypleurochilus</i>	461
<i>confertus</i>	145	<i>geminatus</i>	462
<i>(Hybognathus) siderius</i>	140	<i>Hypocaranx</i>	273
<i>nigrellus</i>	145	<i>Hypoclydoma</i>	297
<i>notatus</i>	145	<i>bella</i>	297
<i>perspicuus</i>	145	<i>Hypocritichthys</i>	410
<i>punicus</i>	145	<i>analis</i>	410
<i>superciliosus</i>	145	<i>hypodus, Decapterus</i>	270
<i>tenellus</i>	145	<i>Hypohomus</i>	284
<i>Hydrargira atricauda</i>	174	<i>aurantiacus</i>	285
<i>hma</i>	174	<i>cymatotaenia</i>	285
<i>Hydrargyra</i>	175	<i>nianguae</i>	285
<i>diaphana</i>	178	<i>spilotus</i>	285
<i>fusca</i>	174	<i>Hypolophus</i>	28
<i>hma</i>	174	<i>Hypomesus</i>	67
<i>luciae</i>	178	<i>olidus</i>	68
<i>multifasciata</i>	178	<i>pretiosus</i>	67
<i>nigrofasciata</i>	176	<i>hypoplecta, Rathbunella</i>	452
<i>ornata</i>	176	<i>Hypoplectrus</i>	320
<i>similis</i>	177	<i>aberrans</i>	321
<i>swampina</i>	175	<i>gemma</i>	321
<i>trifasciata</i>	176	<i>lamprurus</i>	321
<i>vernalis</i>	176	<i>maculiferus</i>	321
<i>zebra</i>	179	<i>pinnivaria</i>	321
<i>Hydrolagus</i>	33	<i>unicolor</i>	320, 321
<i>collei</i>	33	<i>aberrans</i>	321
<i>waiti</i>	33	<i>accensus</i>	321
<i>Hydrophlox</i>	126	<i>affinis</i>	321
<i>abbotti</i>	127	<i>bovinus</i>	321
<i>altipinnis</i>	127	<i>chlorurus</i>	321
<i>chalybeus</i>	127	<i>crocotus</i>	321
<i>chiliticus</i>	127	<i>gemma</i>	321
<i>chlorocephalus</i>	127	<i>gummigutta</i>	321
<i>chrosomus</i>	127	<i>guttavarius</i>	321
<i>lutipinnis</i>	127	<i>indigo</i>	321
<i>roseus</i>	127	<i>nigricans</i>	321
<i>rubricroceus</i>	127	<i>pinnivarius</i>	321
<i>stonei</i>	127	<i>puella</i>	321
<i>xenoccephalus</i>	127	<i>vitulinus</i>	321
<i>hydrophlox, Cheonda</i>	119	<i>hypoplectus, Bathymaster</i>	452
<i>Clinostomus</i>	119	<i>Hypopites</i>	324
<i>hygomi, Mycetophum</i>	169	<i>Hypopomus</i>	101
<i>Scopelus</i>	169	<i>brevirostris</i>	101
<i>Hylomyzon</i>	107	<i>occidentalis</i>	101
<i>Hymenoccephalus</i>	207	<i>hypopthalmus, Arius</i>	151
<i>cavernosus</i>	207	<i>Hypoprius</i>	17
<i>italicus</i>	207	<i>brevirostris</i>	17
<i>Hynnis</i>	274	<i>longirostris</i>	17
<i>cubensis</i>	274	<i>signatus</i>	17
<i>goreensis</i>	274	<i>Hypopriodon</i>	17
<i>hopkinsi</i>	274	<i>hypopthalmus, Cathorops</i>	151
<i>Hyodon</i>	40	<i>hypopthalmus, Arius</i>	151
<i>amphodon</i>	41	<i>Hyporhamphus</i>	198
<i>chrysopsis</i>	41	<i>gilli</i>	198
<i>selenops</i>	41	<i>roberti</i>	198
<i>vernalis</i>	40	<i>rose</i>	198
<i>Hyostoma</i>	286	<i>snyderi</i>	199
<i>blennioperca</i>	286	<i>tricuspidatus</i>	198
<i>newmani</i>	286	<i>unifasciatus</i>	198
<i>simotermum</i>	286	<i>Hyporthodus</i>	310
<i>hyostoma, Macrhybopsis</i>	139	<i>flavicauda</i>	311

List of scientific names—Continued

	Page		Page
Hyposerranus.....	310	Icelus hamatus	378
hyposticta, Propterygia.....	24	scutiger.....	378
hypostoma, Mobula.....	32	spatula.....	379
Hypostomus.....	160	spinger.....	378
brasiliensis.....	160	tuncialis.....	379
cirrhosus.....	160	vicinalis.....	378
guacari.....	160	Ichthocolla.....	33
guacharote.....	161	Ichthyapus.....	84
plecostomus.....	160	Ichthyomyron.....	9
robinu.....	160	castaneus.....	9
hypostomus, Cephalopterus.....	32	concolor.....	9
Hypsagonus gradiens.....	393	fossor.....	9
swanii.....	396	hurodo.....	9
hypselosoma, Leptocephalichthys.....	79	Ichthys.....	268
hypselura, Rhamdia.....	157	lockingtoni.....	268
hypselurus, Pimelodus.....	157	icista, Bardiella.....	346
hypselopterus, Leuciscus.....	129	Sciæna.....	346
Hypsiometes.....	451	Icosteidae.....	269
gobioides.....	451	Icosteus.....	269
Hypsifario.....	55	ægmaticus.....	269
Hypsilepis.....	128	ictalops, Cottus.....	385
ardens.....	126	Pegedictus.....	385
coccogenis.....	127	Ictalurus.....	151
cornutus cerasinus.....	128	anguilla.....	152
cyaneus.....	128	furcatus.....	151, 152
gibbus.....	128	kevinskii.....	153
galacturus.....	129	macaskeyi.....	153
iris.....	131	meridionalis.....	152
Hypsinotus.....	358	okeechobeensis.....	152
rubescens.....	358	punctatus.....	152
hypsinotus, Ceratichthys.....	138	robustus.....	152
Erinemus.....	138	simpsoni.....	152
Hypsolepis.....	128	Icthelis.....	299
Hypsophrys.....	418	erythropis.....	303
unimaculatus.....	418	megalotus.....	300
Hypsopsetta.....	225	melanops.....	298
guttulata.....	225	ichtheloides, Lepomis.....	303
hypostoma, Mobula.....	32	Ichthyaelurus.....	151
Hypostomus.....	160	Ichthyapus.....	84
plecostomus.....	160	acutirostris.....	84
hypostomus, Cephalopterus.....	32	Ichthyobus.....	102
Hypsagonus.....	393	cyanellus.....	102
gradiens.....	393	rauchi.....	102
quadricornis.....	393	Ichthyophis.....	94
swanii.....	396	vittatus.....	95
hypselura, Rhamdia.....	157	Ictobus.....	102, 352
Hypsiometes.....	451	bubalis.....	102
gobioides.....	451	urus.....	102
Hypsifario.....	54	Idiacanthidae.....	70
hypsinotus, Erinemus.....	138	Idiacanthus.....	70
Hypsoblenius.....	462	antrostomus.....	70
gentilis.....	462	ferox.....	70
gilberti.....	462	ignobilis, Caranx.....	273
hantz.....	462	Scomber.....	273
ionthas.....	462	Ilietis.....	155
piersoni.....	462	Ilisha.....	44
striatus.....	462	abnormis.....	44
Hypsopsetta.....	225	caribbae.....	45
guttulata.....	225	illecebrosus, Zestidium.....	347
Hypsurus.....	412	illecebrosus, Alburnops.....	134
caryi.....	412	Stelliferus.....	347
Hypsypops.....	415	illustris, Lampogrammus.....	484
dorsalis.....	415, 416	Ilyophidæ.....	78
rubicundus.....	415	Ilyophis.....	78
Hysterocarpus.....	412	brunneus.....	78
traski.....	412	Ilypnus.....	445
hystrix, Diodon.....	501, 502	gilberti.....	445
		imberbe, Ophiodon.....	467
Icelidæ.....	375	Peristedion.....	406
Icelinus.....	376	imberbis, Apogon.....	295
borealis.....	376	Mullus.....	295
burchami.....	376	Sciæna.....	348
cavifrons.....	375	Vulsiculus.....	406
filamentosus.....	375	imiceps, Ophioscion.....	348
fimbriatus.....	376	Sciæna.....	348
fuscescens.....	376	immaculata, Amia.....	40
oculatus.....	376	Coryphæna.....	264
quadriseriatus.....	376	Fistularia.....	245
strabo.....	376	Perca.....	307
tenuis.....	376	immaculatum, Cichlasoma spinosissimum.....	420
Icelus.....	378	Cybium.....	257
australus.....	379	immaculatus, Archocentrus.....	420
bicornis.....	378	Esox.....	173
canaliculatus.....	378	masquinongy.....	173
euryops.....	378	Platyptæcilus maculatus.....	193
furciger.....	378	Salmo.....	60
		Synbranchus.....	76

List of scientific names—Continued

	Page		Page
<i>Imostoma</i>	235	<i>insigne</i> , <i>Pimelodus</i>	155
<i>chenevi</i>	236	<i>insignis</i> , <i>Anoplocheilichthys</i>	468
<i>gilberti</i>	236	<i>catostomus</i>	105, 108
<i>longimane</i>	236	<i>cheirodon</i>	96
<i>meadiæ</i>	236	<i>pimelodus</i>	156
<i>podostemone</i>	236	<i>rabida</i>	156
<i>shumardi</i>	235	<i>insociabilis</i> , <i>Dorosoma</i>	46
<i>imperator</i> , <i>Xiphias</i>	264	<i>insolatus</i> , <i>Helias</i>	413
<i>imperialis</i> , <i>Coryphæina</i>	264	<i>inspector</i> , <i>Bathytroctes</i>	52
<i>Esox</i>	197	<i>insulæ-santæ-cruces</i> , <i>Scarus</i>	424
<i>Lutavis</i>	284	<i>insularum</i> , <i>Atherinops</i>	252
<i>Sebastes</i>	370	<i>cedrosensis</i>	262
<i>Zeus</i>	217	<i>muræna</i>	90
<i>impetiginosus</i> , <i>Serranus</i>	310	<i>integripinnis</i> , <i>Auchenopterus</i>	457
<i>implutus</i> , <i>Tetrodon</i>	500	<i>cremnobates</i>	457
<i>impressus</i> , <i>Conger</i>	80	<i>intermedia</i> , <i>Tigoma</i>	119
<i>inæquulobus</i> , <i>Pogonichthys</i>	116	<i>intermedius</i> , <i>Cichla</i>	416
<i>incilis</i> , <i>Querimana</i>	253	<i>gephyrocharax</i>	96
<i>Incisidens</i>	339	<i>heros</i>	416
<i>simplicidens</i>	339	<i>paralepis</i>	165
<i>incisor</i> , <i>Helopercus</i>	301	<i>pomoxys</i>	304
<i>Kyphosus</i>	339	<i>saurus</i>	164
<i>Pimalepterus</i>	339	<i>sudis</i>	165
<i>Pomotis</i>	301	<i>synodus</i>	164
<i>incisus</i> , <i>Lycodes</i>	475	<i>internasalis</i> , <i>Julis</i>	427
<i>inconstans</i> , <i>Eucalia</i>	238	<i>interrupta</i> , <i>Chrysoperca</i>	307
<i>Gasterosteus</i>	238	<i>morone</i>	307
<i>inconstans cayuga</i> , <i>Eucalia</i>	238	<i>perca mitchilli</i>	307
<i>incrassatus</i> , <i>Leucosomus</i>	117	<i>raja</i>	26
<i>indica</i> , <i>Lutodeira</i>	40	<i>raja</i>	26
<i>indicus</i> , <i>Cottus</i>	397	<i>interruptus</i> , <i>Anisotremus</i>	332
<i>Tetrapturus</i>	263	<i>archophites</i>	303
<i>indigena</i> , <i>Clupea</i>	42	<i>centarchus</i>	303
<i>indigo</i> , <i>Hypoplectrus unicolor</i>	321	<i>genytremus</i>	332
<i>Plectropoma</i>	321	<i>luxilus</i>	128
<i>Inermia</i>	340	<i>intertinctus</i> , <i>Mystriophus</i>	88
<i>vittata</i>	340	<i>ophisurus</i>	88
<i>Inermiidae</i>	340	<i>interstitialis</i> , <i>Serranus</i>	314
<i>inermis</i> , <i>Anoplogonus</i>	307	<i>trisotrops</i>	314
<i>Aspidophoroides</i>	397	<i>intronger</i> , <i>Dicrolene</i>	483
<i>Dermatolepis</i>	313	<i>rosebastes</i>	366
<i>Elops</i>	39	<i>sebastichthys</i>	366
<i>Mesoprion</i>	327	<i>introngra</i> , <i>Dicrolene</i>	483
<i>Rabirubra</i>	327	<i>inurus</i> , <i>Zygonectes</i>	185
<i>Raja inornata</i>	26	<i>Ioa</i>	288
<i>Scorpena</i>	371	<i>vigil</i>	288
<i>Serranus</i>	313	<i>vitrea</i>	288
<i>infans</i> , <i>Avocettina</i>	83	<i>Ioglossus</i>	438
<i>Gambusia</i>	189	<i>callurus</i>	438
<i>Nemichthys</i>	83	<i>Ionthes</i> , <i>Cerdale</i>	473
<i>Poecilopsis</i>	189	<i>Hypsoblennius</i>	462
<i>infernalis</i> , <i>Histiobranchus</i>	78	<i>isesthes</i>	462
<i>Muræna</i>	92	<i>ios</i> , <i>Clevelandia</i>	445
<i>infirmis</i> , <i>Novaculichthys</i>	430	<i>Gobiosoma</i>	445
<i>Xyrichtys</i>	430	<i>Iotichthys</i>	121
<i>infraspinata</i> , <i>Asterotheca</i>	396	<i>phlegethoni</i>	121
<i>infraspinata</i> , <i>Xeneretmus</i>	396	<i>iowa</i> , <i>Etheostoma</i>	291
<i>ingens</i> , <i>Hippocampus</i>	224	<i>ohgocephalus</i>	291
<i>ingolfi</i> , <i>Macrourus</i>	206	<i>Ipnopidæ</i>	170
<i>inustus</i>	430	<i>Ipnope</i>	170
<i>mundicorpus</i>	430	<i>agassizi</i>	171
<i>pavonius</i>	430	<i>murrayi</i>	170
<i>Iniomi</i>	163	<i>irideus</i> , <i>Labrus</i>	304
<i>Innominato</i>	87	<i>salmo</i>	58
<i>innominatus</i> , <i>Girardichthys</i>	182	<i>iridia</i> , <i>Salmo</i>	58
<i>inopinatus</i> , <i>Gasterosteus</i>	238	<i>iridinus</i> , <i>Caranx</i>	271
<i>inops</i> , <i>Antennarius</i>	505	<i>Iridio</i>	426
<i>Inopsetta</i>	226	<i>bivittata</i>	426
<i>ischyra</i>	226	<i>caudalis</i>	427
<i>inornata</i> , <i>Lota</i>	212	<i>cyanoccephalus</i>	427
<i>Raja</i>	26	<i>decoratus</i>	427
<i>inornatus</i> , <i>Apodichthys</i>	467	<i>disipulus</i>	427
<i>Cryptocanthodes</i>	471	<i>elegans</i>	427
<i>Microlepidotus</i>	335	<i>frenatus</i>	427
<i>Oligolepides</i>	277	<i>garnoti</i>	427
<i>Pseudogobius</i>	428	<i>kirschli</i>	427
<i>inscriptum</i> , <i>Nanostoma</i>	289	<i>maculipinna</i>	427
<i>inscriptus</i> , <i>Achirus</i>	229	<i>meyeri</i>	427
<i>Nothonotus</i>	289	<i>microstomus</i>	427
<i>Pomotis</i>	300	<i>nicholsi</i>	428
<i>insculpta</i> , <i>Netuma</i>	149	<i>pictus</i>	427
<i>insculptus</i> , <i>Arius</i>	149	<i>poeys</i>	427
<i>Ctenolucius</i>	100	<i>radiata</i>	426
<i>Gasterosteus</i>	237	<i>sellifer</i>	428
<i>Luciocharax</i>	100	<i>semicinctus</i>	427
<i>insignarius</i> , <i>Pimelodon</i>	155, 156	<i>similis</i>	427

List of scientific names—Continued

	Page		Page
Irillion.....	68	jagua, Histro.....	505
oregonus.....	68	Jaguacanguare.....	415
iris, Hypsilepis.....	131	jaguar, Bodianus.....	235
Leuciscus.....	117	jalapae, Xiphophorus.....	193
iradians, Serranus.....	319	jamaicensis, Cynoscion.....	355
irregularis, Theraps.....	417	Otolithus.....	355
irretitus, Gordichthys.....	89	Raja.....	29
Mugil.....	254	janeiro, Engraulis.....	51
irroratus, Monacanthus.....	494	januaria, Umbrina.....	351
irroratus, Monacanthus.....	494	jaok, Cottus.....	387
Isabelita.....	361	Myoxocephalus.....	387
isabelita, Angelichthys.....	361	japonicus, Arctoscopus.....	357
Isaciella.....	335	Caranx trachurus.....	270
ischana, Anchoviella.....	48	Chirolophus.....	466
ischanus, Notemigonus.....	115	Salmo.....	54
Stolephorus.....	48	Scomber.....	258
Ichnomembras.....	247	Trachurus.....	270
gabonensis.....	247	Trichodon.....	357
gabunensis.....	247	jarrovi, Lepidomeda.....	147
ischanus, Notemigonus.....	115	Minomus.....	104
Ichnomembras.....	247	javinicus, Pseces.....	268
ischyra, Inopsetta.....	226	jcauri, Epiplatys.....	125
ischyrus, Apomotis.....	299	Cyprinus, Parachanna.....	125
Lepiopus.....	299	jerezensis, Alburnellus.....	125
Parophrys.....	223	jenguano, Hamulon.....	331
Iesthes.....	462	jenkinsi, Chasmodes.....	463
gilberti.....	462	Synodus.....	163
ionthas.....	462	Zygocetes.....	178
scrutator.....	462	Jenkinsia.....	45
ishinagi, Megaperca.....	308	lamprotaenia.....	46
islandicus, Bogmarus.....	216	stolifera.....	46
Centronotus.....	471	jessae, Oligocephalus.....	291
Gasterosteus.....	238	Poculichthys.....	291
isodon, Apionodon.....	15	Xyrichthys.....	429
Carachinus.....	15	Xyruia.....	429
Mesoprion.....	326	jocu, Anthias.....	325
isodus, Squalus.....	20	Lutianus.....	325
Isogomphodon.....	15	johanna, Agyrosomus.....	63
maculipinnis.....	16	Leucichthys.....	63
isolepis, Isopsetta.....	226	johni, Cephaloptera.....	32
Lepidopsetta.....	223	Johnius batabanus.....	345
Luxilus cornutus.....	123	guttatus.....	313
Notropis cornutus.....	123	nobilis.....	356
Sternotremis.....	215	regalis.....	355
Isopisthus.....	353	savatilis.....	350
affinis.....	353	jonesi, Belone.....	197
harrowei.....	354	Mollemis.....	188
parvipinnis.....	353	Syngnathus.....	241
remifer.....	353	jonesi, Gerres.....	341
Isoplagiodon.....	15	Pseudoxiphophorus bimaculatus.....	188
Isopsetta.....	226	Jordanella.....	180
isolepis.....	223	floridæ.....	180
Isospondyli.....	39	jordani, Bathymaster.....	452
isthmensis, Alurichthys.....	148	Caulophryne.....	511
Bagre.....	148	Chirostoma.....	250
Pocilopsis.....	190	Eopsetta.....	222
Rivulus.....	189	Epinephelus.....	314
Istiophoridae.....	263	Etheostoma (Nothonotus).....	290
Istiophorus.....	263	Galeichthys.....	150
americanus.....	263	Gillias.....	458
greyi.....	263	Gymnothorax.....	92
maguirei.....	263	Hemilepidotus.....	381
volador.....	263	Hippoglossoides.....	222
wrighti.....	263	Hybopsis.....	136
istiana, Parapetenia.....	418	Lamppanyctus.....	167
istlanus, Heros.....	418	Lutianus.....	324
Istlanus.....	151	Lycodes.....	474
balsanus.....	151	Lycondon.....	92
Isuropsis.....	19	Neonænis.....	324
dekayi.....	19	Nothonotus.....	290
Isurus.....	19	Notropis.....	136
oxyrinchus.....	19	Pantosteus.....	104
punctatus.....	19	Raja.....	26
spallanzani.....	19	Ronquillus.....	452
tigris.....	19	Salmo.....	57
Itanra.....	312	clarkii.....	57
itaiara, Promicrop.....	312	Sebastodes.....	304
Serranus.....	312	Tachisurus.....	150
Serranus.....	312	Trisotropis.....	314
Jaboncello.....	316	Jordania.....	375
jacksoni, Embiotoca.....	411	zonope.....	375
jacobae, Echenes.....	449	Jordanidae.....	375
jacobi, Corvina (Johnius).....	348	josephi, Ophidium.....	495
jacobus, Myripristis.....	235	Joturus.....	254
Jacome, Serranus.....	319	globiceps.....	255
jaculidens, Ancyrodon.....	354	pichardi.....	254
jaculus, Alburnellus.....	123	stipes.....	255

List of scientific names—Continued

	Page		Page
juba, Guatucupa.....	332	klunzingeri, Achirus.....	229
Perca.....	332	Solea.....	229
jugalis, Moniana.....	131	kneri, Pristipoma.....	335
jugulares.....	450	kraussi, Gobius.....	441
julia, Claricola.....	292	Kryptophaneron.....	234
julidlo.....	428	alfredi.....	234
adustus.....	428	Kuhliidae.....	305
notospilus.....	428	Kuhlia arge.....	305
Julis.....	428	kumleri, Uranidea.....	385
caudalis.....	427	Kyphosidae.....	339
cinctus.....	427	Kyphosus.....	339
ciotaphus.....	426, 427	analogus.....	339
cyabostigma.....	426	elegans.....	339
detersor.....	429	meisor.....	339
dimidiatus.....	427	lutescens.....	340
garnoti.....	427	metzelaari.....	339
gillianus.....	429	sectatrix.....	340
humeralis.....	426		
internasalis.....	427	labacae, Chirostoma.....	250
lucasanus.....	429	Labeo elegans.....	109
maculipinna.....	427	labialis, Fundulus.....	174
melanocheir.....	427	Profundulus.....	174
modestus.....	428	labiatum, Heros.....	422
nitida.....	429	labiatus, Amphilophus.....	422
nitidissima.....	429	Catostomus.....	106
opalina.....	426	Labichthys.....	84
patatus.....	426	bowersi.....	84
pictus.....	427	carinatus.....	84
principis.....	426	elongatus.....	83
ruptus.....	427	gilli.....	83
semicinctus.....	427	Labidesthes.....	251
		sicculus.....	251
kalisheræ, Ericthe.....	400	labiosus, Carpiodes.....	103
kamloops, Oncorhynchus.....	58	labradoricus, Acanthorottus.....	388
kanawha, Hybopsis.....	134	Coregonus.....	64
Notropis.....	134	Labrax.....	373
kansae, Fundulus.....	179	albidus.....	307
Plancterus.....	179	clathratus.....	317
karak, Anarrhichas.....	472	decagrammus.....	373
Kathetostoma.....	454	lagocephalus.....	374
albigutta.....	454	monopterygius.....	371
averruncus.....	454	multilineatus.....	307
Katsuwonidae.....	258	nebulifer.....	317
Katsuwonus.....	258	nigricans.....	308
pelamis.....	258	notatus.....	307
vagans.....	258	octogrammus.....	373, 374
kaupi, Physiculus.....	211	osculatus.....	307
kaupli, Synaphobranchus.....	78	superciliosus.....	374
kayko, Salmo keta vel.....	54	Labridae.....	424
keimi, Hybopsis.....	133	labridens, Heros.....	419
Notropis.....	133	Parapetania.....	419
kendalli, Calamus.....	337	labridum, Hæmulon.....	332
Hybopsis.....	135	labriformis, Epinephelus.....	311
Sphagebranchus.....	84, 85	Seranus.....	311
Verma.....	85	Labrisomus.....	459
kennedyi, Trachinotus.....	277	biguttatus.....	460
kennerlyi, Moxostoma.....	109	bucciferus.....	459
Oncorhynchus.....	55	heilneri.....	459
Salmo.....	55	herminier.....	459
kennicotti, Acipenser.....	34	lenticinosus.....	459
Catnotus.....	293	microlepidotus.....	459
Coregonus.....	65	nuchipinnis.....	459
Prosopium.....	65	xanti.....	459
Kenoza.....	173	Labroperca.....	310
kentuckiensis, Luxilis.....	137	Labrosomus bucciferus.....	459
Nocomis.....	137	cremnobates.....	456
kessleri, Arius.....	149	microlepidotus.....	459
Netuma.....	149	labrosus, Ceratichthys.....	138
keta, Oncorhynchus.....	54	Erismenus.....	138
kevinskii, Ictalurus.....	153	Zoarces.....	473
killifish, Cobitis.....	176	Labrus adspesius.....	424
kincaidi, Malacocottus.....	388	americanus.....	424
kirschii, Iridio.....	427	auritus.....	299
kirtlandi, Acipenser (Huso).....	35, 37	bifasciatus.....	429
Kirtlandia.....	248	lorquatus.....	429
laciniata.....	248	bivittatus.....	426
vagrans.....	248	brahieri.....	426
klutech, Oncorhynchus.....	54	carolinus.....	424
Salmo.....	54, 55	chogei.....	424
kitt, Microstomus.....	228	fulva.....	424
Pleuronectes.....	228	cromis.....	352
kiyi, Leucichthys.....	64	cruentatus.....	322
klamathensis, Agosia.....	141	cianocephalus.....	427
Apocope.....	141	evoletus.....	424
Cottus.....	384	falcatus.....	276
kleinli, Balistes.....	496	fulvo-maculatus.....	335

List of scientific names—Continued

	Page		Page
Labrus fulvus	309	Lagochila lacera	112
gauza.....	310	Lagodon	337
griseus.....	325	rhomboides.....	337
hexagrammus.....	374	lahontan, <i>Pantosteus</i>	105
hustula.....	424	lalandi, <i>Carcharias</i>	15
trideus.....	304	Scoliodon.....	15
macropterus.....	304	Seriola.....	279
maximus.....	424	lamellifer, <i>Exocoetus</i>	200
onitis.....	424	Lama.....	20
ornatus.....	429	lama, <i>Carcharias</i>	15
palladius.....	299	lamari, <i>Acipenser</i>	35
pentacanthus.....	424	Lepidosteus.....	37
plumieri.....	330	lamiella, <i>Carcharias</i>	15
psittaculus.....	426	Lamiosis.....	15
pulcher.....	426	Lamna.....	19
radiatus.....	426	caudata.....	16
rufus.....	425	nasus.....	19
salmoides.....	298	punctata.....	19
semiruber.....	425	Lamnidae.....	19
sparoides.....	305	lamotteni, <i>Okkelbergia</i>	10
squeteague.....	355	Petromyzon.....	10
maculatus.....	355	Lampadena	167
striatus.....	317	speculigera.....	167
subfuscus.....	424	Lampanyctus	167
tassellatus.....	424	alatus.....	167
tautoga.....	424	gemmifer.....	167
versicolor.....	336	güntheri.....	167
lacera, Lagochila	112	lacerta.....	167
lacerta, Lampanyctus	167	ritteri.....	167
Synodus.....	163	Lampetra	10
lacterinus, <i>Synodus</i>	164	ayresi.....	10
lactosus, <i>Hybopsis</i>	129	boealis.....	10
Luxilus.....	129	spadicea.....	9
lactus, <i>Gobius</i>	440	wideri.....	9
Scomber.....	258	lampetæformis, Blennius	470, 471
Lachnolæmus	425	Lumpenus.....	471
Lachnolaimus	425	Lampreda.....	10
agula.....	425	Lampridæ.....	216
caninus.....	425	Lampus.....	216
dux.....	425	lauta.....	217
maximus.....	425	legus.....	216
psittacus.....	425	Lamprogrammus	484
suillus.....	425	illustris.....	484
lachrymalis, <i>Ptychostomus</i>	111	lamprotenia, <i>Clupea</i>	46
lacinata, <i>Meridia vagrans</i>	248	Jenkinsia.....	46
lacinata, <i>Kirtlandia</i>	248	lamprurus, <i>Hypoplectrus</i>	321
lacrmosus, <i>Scarus</i>	107	Serranus.....	321
lactarius, <i>Catostomus (Acomus)</i>	497	Lampugus	264
Lactophrys	497	neopolitanus.....	265
trigonus.....	497	punctulatus.....	265
lactus, Algansea	116	sigulus.....	265
Gadus.....	152, 212	lanatus, <i>Merluccius</i>	214
Hausdor.....	152	lanceolatus, <i>Percæ</i>	352
Hemipus.....	115	lanceolatus, <i>Chaetodon</i>	352
okeechobeensis, <i>Ameiurus</i>	152	Equetus.....	333
Pomolobus pseudoharengus.....	42	Gobius.....	341
Læmargus	21	Homopron.....	346
Læmonema	213	Lonchurus.....	352
barbatula.....	213	Stellifer.....	346
barbatulum.....	213	lappa, Halieutella	508
gracillipes.....	213	Lappoperca.....	310
melanurum.....	213	Larimus	344
lætabilis, <i>Moniana</i>	131	aclivus.....	344
lætus, <i>Centronotus</i>	468	argenteus.....	344
lævicaudatus, <i>Hippocampus</i>	244	breviceps.....	344
lævigatus, <i>Callichthys</i>	160	effulgens.....	344
Chronectes.....	505	fasciatus.....	344
Lagocephalus	497	punctatus.....	344
Salmo.....	61	lappa, Halieutella	410
Tetraodon.....	497	lappa, <i>Halieutella</i>	161
Læviraja	34	planiceps.....	161
lævis, <i>Acipenser</i>	25	Lasiognathus	510
Balistes.....	495	saccostoma.....	510
Gasterosteus.....	239	lata, <i>Parapetenia tetraacantha</i>	419
Mustelus.....	13	Xurel.....	272
Raja.....	25	Latebrus	267
Squatina.....	22	oculatus.....	267
variegatus, <i>Orbis</i>	499	lateptotus, <i>Serranus</i>	314
lagenarius, <i>Acipenser</i>	36	lateralis, <i>Alvarius</i>	294
Lagocephalus	497	Artedius.....	377
lævigatus.....	497	Characodon.....	183
pachycephalus.....	497	Embriotes.....	411
lagocephalus, Hexagrammos	374	Gobiomorus.....	436
Labrax.....	374	Mylocheilus.....	112
Salmo.....	54	Philypnus.....	436
Lagochila	112	Richardsonius.....	118

List of scientific names—Continued

	Page		Page
laterals, Scarus	432	leipomus, Cottus	385
Scorpaenichthys	377	lelops, Xeneretmus	396
Teniotoca	411	Xenopyxis	396
Zygonectes	177	Leiotomus	348
latera, Tetradon	500	humeralis	349
lateratum, Myctophum	170	lineatus	349
lateratus, Polyipnus	74	anthurus	340
laterum albo, Antennarius	501	Leurus	287
latham, Trachurus	270	leurus, Gasterosteus	287
laticauda, Anguilla	77	lamar, Acipenser (Huso)	35
Rhamdia	157	Lombus	435
atcaudus, Pimelodus	157	maculatus	435
Sechodon	14	lemmoni, Squalius	119
laticeps, Acanthocottus	388	lemniscatus, Engraulis	48
Etobatus	30	Osmerus	163
Arius	150	Lemniscoma	261
Atherina	246	Lencoraja	24
Bathynectes	483	lentiginosa, Murana	90
Megalocottus	388	lentiginosum, Ostracion	496
Mixonus	483	lentiginosus, Amia	38
latidens, Glaridodon	189	Galeichthys	149
Microstomus	223	Heros	418
Poecilopsis	189	Labrisomus	459
latifasciatus, Cyprinodon	181	Rhinobatus	23
Eleotris	437	Rhinobatus	23
Hemieleotris	437	Tachisurus	149
latifrons, Anarrhichas	472	Theraps	418
Citharichthys	218	leonensis, Oligocephalus	291
Eleotris	436	leonina, Moniana	131
Lycichthys	472	leopardinus, Antennarius	505
Paraliparis	405	Platophrys	218
Steigodonotus	379	Rhomboidichthys	218
Styrium	218	leopardus, Anarrhichas	472
Xenochirus	396	Lepadogaster mæandricus	488
Xenopyxis	396	reticulatus	488
Latilus chrysops	357	testar	489
princeps	357	Lephodolus indicus	509
latimaculatus, Ophisurus	85	Lepibema	307
Latimana, Belone	197	chrysops	307
latinasutus, Macrourus	206	lepida, Boleosoma	291
Macurus	206	Cyprinella	130
laticus, Salmo	64	lepidentostole, Anchovia	49
latipinna, Molinnesia	192	Anchoviella	49
latipinnis, Catostomus	107	Lepidia	211
Zanleopsis	375	Lepidion	211
latipunctata, Gambusia	187	verecundum	211
latirostratus, Macrourus	206	Lepidochætodon	359
Macurus	206	lepidogenys, Etheostoma	202
latura, Loricaria	162	Nivicola	222
Loricaria filamentosa	162	Lepidogobius	444
atus, Caranx	272	gilberti	445
Euctenogobius	442	lepidus	444
Heros tetracanthus	419	Lepidolepis trachyrincus	203
Scomber	276	Lepidolepis norvegicus	205
lauræ, Astatheros	421	Lepidomeda	147
Cichlasoma	421	arrovii	147
laurettæ, Coregonus	62	vittata	147
Leucichthys	62	Lepidomegas	279
lauta, Lampris	217	Lepidopsetta	226
Lavinia	113	bilineata	226
conformis	120	isolepis	226
crassicauda	120	Lepidopus	262
exilicauda	113	caudatus	262
harengus	113	elongatus	262
leachi, Clupea	41	gonani	262
leachianus, Thynnus	258	xantusi	262
Lebias ellipsoides	181	Lepidosteus	36
rhomboidalis	181	(Atractosteus) berlandieri	38
Lebistes	191	ayresi	37
pocilloides	191	clintonii	37
reticulatus	191	copei	37
Lebus	374	crassus	36
lebranchus, Mugil	253	elizabeth	37
lecontei, Acipenser	34	grayi	37
lecurus, Nauclerus	278	harlani	37
leel, Symphurus	231	horathi	37
lefoyi, Diapterus	341	lamani	37
Ulema	341	latirostris	37
Lelroyia	488	leptorhynchus	37
bermudensis	488	lesueurii	37
lelarchus, Cynoscion	355	louisianensis	36
Otolithus	355	manjuari	38
Leobatus, sloani	29	milberti	37
Leiocottus	390	otarius	37
hurundo	390	piquotianus	37
Lelodon	21	semiradiatus	36
leionothos, Holocanthus	499	smithi	37

List of scientific names—Continued

	Page		Page
Lepidosteus thompsoni.....	37	Leptocephalus spallanzani.....	79
troosti.....	37	leptocephalus, Ceratichthys.....	137
lepidulus, Alburnus.....	125	Merlangus.....	209
lepidus, Gobius.....	444	Leptoceardale.....	473
Lepidogobius.....	444	longipinnis.....	473
Oligocephalus.....	291	Leptochlichthys.....	53
Lepiopomus ischyrius.....	299	agassizii.....	53
miniatus.....	300	Leptochinus.....	470
Lepisoma.....	459	maculatus.....	470
currhosum.....	459	Leptocoenge.....	82
Lepisosteidae.....	36	perlongus.....	82
Lepisosteus.....	33	Leptocottus.....	390
albus.....	37	armatus.....	390
(Atractosteus) feiox.....	33	Leptodes.....	71
bison.....	37	Leptoflerasfer.....	486
(Cylindrosteus) oculatus.....	37	macurus.....	486
gavialis.....	36	Leptogunnellus.....	470
huronensis.....	36	gracilis.....	470
lineatus.....	37	Leptolucania.....	174
longirostris.....	36	ommata.....	175
osseus.....	36	Leptophidium cervinum.....	484
oxyrinus.....	37	emmelas.....	484
platostomus.....	37	marmoratum.....	484
semiradiatus.....	36	microlepis.....	485
spatula.....	38	pardale.....	485
Lepodus.....	255	profundorum.....	484
saragus.....	255	prolates.....	484
Lepomis.....	299	stigmatistum.....	485
anagalinus.....	300	Leptophilypnus.....	436
apiatus.....	299	fluvialis.....	436
appendix.....	299	Leptops.....	155
ardesiacus.....	301	Leptopus.....	215
aureus.....	299	peregrinus.....	215
charybdis.....	303	Leptorhaphis.....	189
cyanellus.....	298	Leptorhinophis.....	87
euryotus.....	299	Leptorhynchus.....	83
flexuolarius.....	298	leuchtenbeigi.....	83
garmani.....	300	leptorhynchus, Lepidosteus.....	37
gilii.....	303	Odontopyxis.....	395
haplognathus.....	301	Sarritor.....	395
ichtheloides.....	303	Syngnathus.....	240
lirus.....	302	Leptosomus.....	431
longispinus.....	301	xenodon.....	431
macrochirus.....	301	leptosomus, Luvulus.....	115
miniatus.....	300	Lepturus.....	262
mystacalis.....	300	lepturus, Anarrhichas.....	472
nephelus.....	301	Caranx.....	272
notata.....	298	Macrourus.....	207
occidentalis.....	301	Trichiurus.....	262
opthalmicus.....	299	sermae, Aztecula.....	143
pallida.....	298	Chirostoma.....	251
pallidus.....	299	Notropis.....	143
peltastes.....	300	Skiffa.....	183
purpurascens.....	301	Lernichthys.....	183
salmona.....	298	multiradiatus.....	183
solis.....	300	les Alutiques.....	495
symmetricus.....	299	Curimates.....	100
trifasciata.....	298	Lottes.....	212
Leptophidium.....	484	Musteles.....	208
brevibarbe.....	484	Ovoides.....	499
cervinum.....	484	Spheroides.....	498
emmelas.....	484	Steliferes.....	346
marmoratum.....	484	lessonii, Coryphæna.....	265
microlepis.....	485	Lestidiops.....	166
pardale.....	485	sphyrænops.....	166
profundorum.....	484	lesueuri, Acipenser.....	34
prolates.....	484	Catostomus.....	111
stigmatistum.....	485	Moxostoma.....	111
leptacantha, Rabida.....	155	lesueurianum, Exoglossum.....	146
leptacanthus, Noturus.....	155	lesueuri, Lepidosteus.....	37
Leptagonus.....	394	Letharchus.....	85
decagonus.....	394	pacificus.....	85
Leptarius.....	148	velifer.....	81
dowi.....	148	Lethenteron.....	9
Leptecheneis.....	448	appendix.....	9
Leptoancistrus.....	161	Lethops.....	447
canescens.....	161	connectens.....	447
Leptoblennius.....	470	lethostigma, Paralichthys.....	224
Leptocardi.....	7	lethostigmus, Paralichthys.....	224
Leptocephalichthys.....	79	Lethostole.....	250
Leptocephalidae.....	79	Lethotremus.....	398
Leptocephalus.....	54, 79	muticus.....	398
candidissimus.....	79	vinolentus.....	398
gracilis.....	79	lethrinus, Astatheros.....	420
(Microconger) caudalis.....	81	Cichlasoma.....	420
morrisoni.....	54, 79	letonai, Priapichthys.....	190

List of scientific names—Continued

	Page		Page
leuchtenbergii, Belonopsis	83	leucopsarum, Myctophum (Stenobranchius)	168
Leptorhynchus	53	Nanobranchium	168
Leucichthys	61	leucopus, Photogenis	130
alacanus	62	Rhampoberyx	235
alpenæ	63	leucorhinus, Rhynchoceratius	511
artedi	63	leucorhynchus, Rhinobatos	24
bisselli	63	Rhombatus	24
athabascae	62	leucurus, Eupomacentrus	414
birgei	64	Pomacentrus	414
cyanopterus	63	Leucosomus	116, 117
entomophagus	63	cataractus	117
erianis	63	corporalis	117
gemmifer	63	dissumilis	137
harengus	62	incrassatus	117
hoyi	64	occidentalis	113
johannæ	63	pallidus	117
kiyi	63	rhotheus	117
laurettae	62	leucosteus, Calamus	337
lucidus	62	leucostictus, Eupomacentrus	413
macrogathus	62	Pomacentrus	413
manitoulinus	64	leucotema, Pholidichthys	465
nigripinnus	63	leucurus, Caranx	271
nipigon	64	Hemicaranx	271
ontariensis	62	Leucus dimidiatus	122
osmeriformis	62	olivaceus	122
prognathus	63	Leuresthes	251
pusillus	62	cramerii	251
reighardi	63	tenius	251
sisco	62	Leuroglossus	68
supernus	62	stibius	68
tullibee	63	Leurynnis	474
zenithus	64	paucidens	474
Leucicorpus	483	levis, Hispanicus	368
lusciosus	483	Sebastichthys	368
Leucidius	123	lewini, Zygann	18
pectinifer	123	lewis, Squatina	22
leuciodus, Notropis	124	lewisi, Salar	55, 56
Photogenis	124	Salmo	56
Leuciscus	117, 119	hiberiensis, Balistes	492
argenteus	117	hibertate, Meletta	44
bosci	115	Opisthoneura	44
boucardi	132	Lichia quiebra	277
bubalinus	130	lightfooti, Alphestes	313
burtonianus	121	Epinephelus	313
carletoni	116	ligulata, Seriola	280
caurinus	112, 118	Lila barbuda	496
chrysopterus	117	colorado	494
copii	123	Lile	44
croceus	140	stolifera	44
evermanni	118	lima, Holotrachys	235
frontalis	128	Hydrargira	174
gardoneus	115	Loricaria	162
gilli	118	Myripristis	235
gracilis	128	Rineloricaria	162
hypslepterus	129	Lima-muræna	90
iris	117	melanotis	90
longirostris	120	Limanda	226
lutrensis	131	aspera	227
macrolepidotus	116	beanii	227
nachtriebi	116	ferruginea	227
nasutus	139	proboscidea	227
nitidus	117	limanda, Pleuronectes	222
obesus	128	limandoides, Pleuronectes	222
productus	120	limbatus, Carcharias	16
prolixus	146	Carcharias (Prionodon)	16
pulchelloides	117	Conger	78
pulchellus	117	Oxydactichthys	88
pygmaeus	174	limi, Fundulus	178
rotengulus	117	Umbra	174
siuslawi	118	Zygometes	178
springulus	128	Lima	191
storeri	117	arnoldi	191
tineella	115	caudofasciata	191
vandoisulus	121	couchiana	193
vittatus	128	cubensis	191
zunnensis	114	dominicensis	191
(Leuciscus) caurinus, Cyprinus	118	formosa	193
oregonensis, Cyprinus	114	heterandra	191
leuciscus, Brachydeuterus	333	matamorensis	192
Pristipoma	333	melanonotata	191
leucophæus, Macrourus	205	nigrofasciata	191
Macrurus	205	ornata	191
leucops, Photogenis	123	pavonina	191
engraulinus, Photogenis	123	perugia	191
		pseudoloides	192

List of scientific names—Continued

	Page		Page
<i>Limia venusta</i>	175	<i>Laopsetta obscura</i>	228
<i>versicolor</i>	191	<i>putnami</i>	227
<i>vittata</i>	191	<i>lorus</i> , <i>Chasmistes</i>	108
<i>Limnurgus</i>	182	<i>losterus</i> , <i>Phenacobius teretulus</i>	142
<i>variegatus</i>	182	<i>Luostomus</i>	348
<i>limnurgus</i> , <i>Girardinichthys</i>	183	<i>Liparidæ</i>	399
<i>limosa</i> , <i>Myxine</i>	8	<i>liparina</i> , <i>Amitra</i>	405
<i>Pilodictis</i>	155	<i>lparinus</i> , <i>Paraliparis</i>	405
<i>limosus</i> , <i>Pyrodietis</i>	155	<i>Liparis</i>	399
<i>linea</i> , <i>Mesoprion</i>	325	<i>agassizii</i>	400
<i>Syngnathus</i>	241	<i>arctica</i>	400
<i>lineata</i> , <i>Alosa</i>	42	<i>cyclopus</i>	400
<i>Coryphæna</i>	430	<i>cyclostigma</i>	401
<i>Echeneis</i>	449	<i>denneyi</i>	400
<i>Embiotoca</i>	411	<i>fabricii</i>	400
<i>Sciæna</i>	307	<i>fuçensis</i>	400
<i>Tigoma</i>	119	<i>gibbus</i>	400
<i>lineatus</i> , <i>Achirus</i>	230	<i>herschelinus</i>	400
<i>Ambiodon</i>	382	<i>liparis</i>	400
<i>Cyclopterus</i>	400	<i>megacephalus</i>	400
<i>Cypselurus</i>	202	<i>montagui</i>	399
<i>Diodon geometricus</i>	502	<i>mucosus</i>	399
<i>Dormitator</i>	436	<i>osborni</i>	400
<i>Doryichthys</i>	243	<i>pulchellus</i>	400
<i>Eques</i>	353	<i>ranula</i>	402
<i>Esox</i>	173	<i>reinhardi</i>	402
<i>Eugerres</i>	342	<i>tunicata</i>	400
<i>Euthynnus</i>	259	<i>liparis</i> , <i>Centiophorus</i>	267
<i>Exocoetus</i>	202	<i>Cyclopterus</i>	399, 400
<i>Genyonemus</i>	349	<i>Liparis</i>	400
<i>Gobius</i>	440, 446	<i>major</i> , <i>Cyclopterus</i>	401
<i>Grystes</i>	374	<i>minor</i> , <i>Cyclopterus</i>	400
<i>Leiostomus</i>	349	<i>Lipariscus</i>	401
<i>Lepisosteus</i>	37	<i>nanus</i>	401
<i>Micropogon</i>	349	<i>Liparopidae</i>	398
<i>Mugil</i>	253	<i>Liparops</i>	398
<i>Pleuronectes</i>	230	<i>stelleri</i>	398
<i>Smaris</i>	342	<i>Lipogenvidea</i>	75
<i>Sternopygus</i>	101	<i>Lipogenys</i>	75
<i>Trichodon</i>	357	<i>gilli</i>	75
<i>Zygonectes</i>	177	<i>Lipomyzon</i>	108
<i>lineis transversis</i> , <i>Passer</i>	230	<i>Lipophrys</i>	461
<i>lineolata</i> , <i>Clupea</i>	41	<i>carolinus</i>	461
<i>Pelamys</i>	259	<i>liropus</i> , <i>Arius</i>	151
<i>Pœcilia</i>	192	<i>Tachysurus</i>	151
<i>Sarda</i>	259	<i>lirus</i> , <i>Lepomis</i>	302
<i>lineolatum</i> , <i>Catnotus flabellare</i>	293	<i>lythuis</i>	126
<i>lineolatus</i> , <i>Alburnus</i>	135	<i>Nototopis</i>	126
<i>Catnotus</i>	293	<i>lisa</i> , <i>Memidia</i>	240
<i>Metrogaster</i>	409	<i>Xenatherina</i>	249
<i>Pseudoscarus</i>	434	<i>lister</i> , <i>Ostracion</i>	497
<i>Tetrodon</i>	497	<i>lita</i> , <i>Muraena</i>	93
<i>Zygonectes</i>	178	<i>Litholepis</i>	38
<i>lineopinnis</i> , <i>Muraena</i>	92	<i>littoralis</i> , <i>Odontaspis</i>	19
<i>lineopunctatus</i> , <i>Balistes</i>	493	<i>Squalus</i>	19
<i>Xanthichthys</i>	493	<i>Umbrina</i>	351
<i>linguata</i> , <i>Balistes</i>	496	<i>Umbrula</i>	351
<i>Pleuronectes</i>	222	<i>littorcola</i> , <i>Chaetodon</i>	360
<i>lingel</i> , <i>Æglefinus</i>	211	<i>liturosus</i> , <i>Diodon</i>	501
<i>Trachurus</i>	270	<i>lividus</i> , <i>Petromyzon</i>	10
<i>linophora</i>	359	<i>Silurus</i>	153
<i>linophryne</i>	510	<i>livree</i> , <i>Pimelode</i>	156
<i>arborifer</i>	510	<i>liza</i> , <i>Mugil</i>	253
<i>lucifer</i>	510	<i>lobatus</i> , <i>Canthogaster</i>	498
<i>macrodon</i>	510	<i>Sphæroides</i>	495
<i>polypogon</i>	510	<i>lobochilus</i> , <i>Amphilophus</i>	422
<i>linsleyi</i> , <i>Etheostoma</i>	293	<i>Heros</i>	422
<i>Liloglossina</i>	223	<i>Lobotes</i>	323
<i>tetropthalma</i>	223	<i>auctorum</i>	323
<i>tetropthalmus</i>	223	<i>emarginatus</i>	325
<i>lolepis</i> , <i>Lionurus</i>	207	<i>erata</i>	323
<i>Macrusus</i> (<i>Lionurus</i>).....	207	<i>pacificus</i>	324
<i>Xystreurus</i>	223	<i>somnolentus</i>	323
<i>liomonacanthus</i>	493	<i>surinamensis</i>	323
<i>Lioniscus</i>	34	<i>Lobotidae</i>	323
<i>Lionurus</i>	207	<i>lockingtoni</i> , <i>Ichthys</i>	208
<i>holepis</i>	207	<i>Lonchirus</i>	352
(<i>Lionurus</i>) <i>holepis</i> , <i>Macrusus</i>	207	<i>depressus</i>	352
<i>lopeltis</i> , <i>Acipenser</i>	35	<i>lanceolatus</i>	352
<i>lioperca</i>	313	<i>Lorchomisthus</i>	453
<i>liopropoma</i>	316	<i>micrognathus</i>	453
<i>aberrans</i>	316	<i>lonchurum</i> , <i>Opisthognathus</i>	453
<i>longlepis</i>	316	<i>Lonchurus</i>	352
<i>rubra</i>	315	<i>ancylodon</i>	354
<i>liopsetta</i>	227	<i>lonchurus</i> , <i>Opisthognathus</i>	453
<i>glacialis</i>	227	<i>longa</i> , <i>Trygon</i>	28

List of scientific names—Continued

	Page		Page
longicauda, <i>Muraena</i>	91	Lophopsetta.....	217
Rabula.....	91	maculata.....	217
longicaudus, <i>Gobius</i>	441	Lophotes.....	215
longicephalus, <i>Galeichthys</i>	150	capellei.....	215
Tachisurus.....	150	cepedianus.....	215
longiceps, <i>Chenophryne</i>	510	Lophotidae.....	215
Elcotris.....	435	lordii, <i>Salmo</i>	61
Hybopsis.....	133	loreti, <i>Gramma</i>	322
Porogadus.....	452	Loricaria.....	161
longidens, <i>Caulolepis</i>	233	acuta.....	161
longifilis, <i>Gadomus</i>	202	altipinnis.....	162
longifilus, <i>Bathygadus</i>	202	aurea.....	162
Careproctus.....	403	barbata.....	161
longilepis, <i>Liopropoma</i>	316	bransfordi.....	162
longimana, <i>Etheostoma</i>	286	capetensis.....	162
longimane, <i>Imostoma</i>	286	filamentosa, <i>latiura</i>	162
longimanus, <i>Astatheros</i>	420	fimbriata.....	162
Eupomotis.....	302	flava.....	160
Heros.....	420	frenata.....	162
Squalus.....	15	latiura.....	162
Xystroplites.....	302	lima.....	162
longipes, <i>Bregmaceros</i>	208	panamensis.....	162
Gadus.....	213	plecostomus.....	160
Physiculus.....	212	rostrata.....	161, 162
longipinna, <i>Gobiosoma</i>	445	strigilata.....	162
Holocentrum.....	235	tuyrensis.....	162
longipinnis, <i>Careproctus</i>	403	uracantha.....	162
Clevelandia.....	445	variegata.....	161
Evermannia.....	445	Loricarichthys.....	161
Leptocerdale.....	473	Loricaridae.....	160
Stromateus.....	269	loricata, <i>Alysia</i>	169
longirostris, <i>Alburnops</i>	135	loricatus, <i>Gasterosteus</i>	237
Chaetodon.....	359	Macrogathus.....	36
Esox.....	196	Phalangites.....	394
Euleptorhamphus.....	198	lorito, <i>Sparisoma</i>	432
Forcipiger.....	359	Loro.....	434, 435
Hemirhamphus.....	198	Lota.....	212
Hybopsis.....	135	brosimiana.....	212
Hypoprion.....	17	incornata.....	212
Lepisosteus.....	36	maculosa.....	212
Leuciscus.....	120	Lotella.....	211
Lumpenella.....	471	maxillaris.....	211
Lumpenus.....	471	lotharingus, <i>Gasterosteus</i>	239
Saurus.....	163	loubina, <i>Perca</i>	305
longirostrum, <i>Catostomus</i>	106	louisiana, <i>Engraulis</i>	48
longispatha, <i>Peristidium</i>	405	louisiana, <i>Notropis</i>	124
longispathum, <i>Peristedion</i>	405	Syngnathus.....	242
longispinis, <i>Aelurichthys</i>	148	Synctes.....	242
Lepomis.....	301	louisianensis, <i>Lepidosteus</i>	36
Pontinus.....	372	lowei, <i>Polymia</i>	234
longulus, <i>Pomotis</i>	299	lowii, <i>Omosudis</i>	171
longum, <i>Gobiosoma</i>	446	loxias, <i>Gurnardus</i>	406
longurio, <i>Carcharias</i>	15	Prionotus.....	406
Scollodon.....	15	Lucania.....	175, 182
longus, <i>Balistes</i>	493	affinis.....	175
Dasyatis.....	28	browni.....	181
Ophisurus.....	85	goodei.....	175
lophar, <i>Perca</i>	280	parva.....	175
Lopharis.....	280	lucasana, <i>Sphyraena</i>	255
mediterraneus.....	280	lucasanum, <i>Thalassoma</i>	429
Lophidae.....	504	lucasanus, <i>Julis</i>	429
Lophiomus.....	504	lucasi, <i>Ceratocottus</i>	382
setigerus.....	504	lucayanum, <i>Asymmetron</i>	7
Lophius.....	504	lucens, <i>Ceratichthys</i>	137
aculeatus.....	507	Dacentrus.....	412
americanus.....	504	lucetia, <i>Vinciguerra</i>	73
bufo.....	487	lucetius, <i>Mauloleucus</i>	73
gibbus.....	505	luciae, <i>Hydrargyra</i>	178
histrio ocellatus.....	505	Zygoneustes.....	178
piscatorius.....	504	lucida, <i>Aethroprora</i>	168
radiatus.....	507	Anchoviella.....	48
setigerus.....	504	lucius, <i>Leucichthys</i>	62
spectrum.....	505	Luxilus.....	125
tumidus.....	504	Salmo (<i>Coregonus</i>).....	62
vespertilio.....	507	Stolephorus.....	48
viviparus.....	504	Lucifer.....	70
lophius, <i>Amiurus</i>	153	Lucifer, <i>Linophryne</i>	510
Hauator.....	153	Lucifuga.....	479
Lophobranchii.....	240	dentatus.....	479
Lophodolus.....	509	subterranea.....	479
acanthognathus.....	509	subterraneus.....	479
Lophogobius.....	438	Lucioblennius.....	465
cyprinoides.....	438	alepidotus.....	465
Lopholatilus chamaleonticeps.....	357	lucius.....	465
Lopholatilus.....	357	lucioceps, <i>Saurus</i>	169

List of scientific names—Continued

	Page		Page
lucioceps, <i>Synodus</i>	163	Lutianus analis.....	326
Luciocharax.....	100	apodus.....	325, 326
insculptus.....	100	argentiventris.....	326
striatus.....	100	aya.....	326
Lucioperca americana.....	282	blackfordi.....	326
borea.....	282	brachypterus.....	326
canadensis.....	282	buccanella.....	326
grisea.....	282	campechanus.....	326
pepinus.....	282	colorado.....	326
lucioperca undecimalis, <i>Centropomus</i>	305	cyanopterus.....	325
luciopercaneus, <i>Mentiperca</i>	319	griseus.....	325, 326
Serranus.....	319	guttatus.....	326
Lucius.....	172	hastingsi.....	326
Lucius americanus, <i>Esox</i>	173	joeci.....	325
Lucius, <i>Atractosteus</i>	38	jordani.....	324
Chenopsis.....	465	luteus.....	335
Chirostoma.....	251	lutianus.....	324
Lucioblennius.....	465	lutjanoides.....	326
Ptychocheilus.....	114	mahogoni.....	327
lucoides, <i>Esox</i>	173	megalephthalmus.....	327
lucetiae, <i>Aboma</i>	443	novemfasciatus.....	325
Gobius.....	443	synagris.....	326, 327
ludibunda, <i>Cyprinella</i>	130	viridis.....	324
luetkeni, <i>Dolopichthys</i>	510	vivanus.....	326
lugubris, <i>Caranx</i>	273	lutianus, <i>Lutianus</i>	324
Cyprinella.....	131	lutipinnis, <i>Hydrophlox</i>	127
Esox.....	173	Prisigaster.....	45
Malacotenus.....	460	lutjanoides, <i>Lutianus</i>	326
Melamphats.....	233	Ocyurus.....	326
Myxodes.....	460	Lutianus.....	324
Plectromus.....	233	acutirostris.....	325
Xurel.....	273	aubrieti.....	327
lutpoldi, <i>Characodon</i>	184	blackfordi.....	326
Goodea.....	184	brachypterus.....	326
lumbicus, <i>Muraena</i>	76	cayanensis.....	354
Myrophis.....	83	colorado.....	326
luminosum, <i>Myctophum</i>	170	cupera.....	325
Lumpenella.....	471	dentatus.....	325
longirostris.....	471	lunulatus.....	311
Lumpenidae.....	470	novemfasciatus.....	325
Lumpenus.....	470	prieto.....	325
fabrici.....	471	purpureus.....	326
gracilis.....	470	stearnsi.....	325
lampetraeformis.....	471	surinamensis.....	332
longirostris.....	471	torridus.....	326
mackayi.....	471	triangulum.....	348
nubilis.....	471	tridens.....	318
Lumpus.....	397	trilobatus.....	317
anglorum.....	398	verres.....	427
vulgaris.....	398	lutkeni, <i>Cypselurus</i>	201
lumpus, <i>Cyclopterus</i>	397	Exocoetus.....	201
luna, <i>Pomotis</i>	301	Lutodeira.....	40
Zeus.....	216	lutensis, <i>Leuciscus</i>	131
lunaris, <i>Orthogoriscus</i>	503	Moniana.....	131
unata, <i>Centropyge</i>	361	lutzi, <i>Heterandria</i>	190
Echeneis.....	448	Luvardae.....	264
Solea.....	217	Luvarifomes.....	264
lunaticus, <i>Dactyloscopus</i>	455	Luvarus.....	264
lunatus, <i>Holocanthus</i>	361	imperialis.....	264
Platyphrys.....	217	luxatus, <i>Chasmistes</i>	108
Pleuronectes.....	217	Deltistes.....	106
Rhinichthys.....	140	Luxilus.....	113
lunulatus, <i>Cynias</i>	13	luxiloides, <i>Cyprinella</i>	131
Lutjanus.....	310	Luxilus.....	125, 128
Mustelus.....	13	albeolus.....	129
lupus, <i>Anarhichas</i>	472	cerasinus.....	128
Haustor.....	153	chickasavensis.....	130
Pinelodus.....	153	chrysocephalus.....	128
Lusca, <i>Cyclothone</i>	72	cornutus.....	128
luscusos, <i>Leucicorus</i>	453	chrysocephalus.....	128
lutea, <i>Anguilla</i>	77	isolepis.....	128
luteovinctum, <i>Etheostoma</i>	291	dissimilis.....	138
luteovinctus, <i>Oligocephalus</i>	291	elongatus.....	120
lutescens, <i>Chaetodon</i>	360	erythrogaster.....	113
Kyphosus.....	340	interruptus.....	128
Pimblepterus.....	340	kentuckiensis.....	137
luteum, <i>Haemulon</i>	330	lactosus.....	129
luteus, <i>Centropomus</i>	261	leptosomus.....	115
Gadus tomcodus.....	210	lucius.....	125
Genyatemus.....	335	roseus.....	127
Lutianus.....	335	seco.....	115
Noturus.....	155	selene.....	132
Rhinichthys.....	139	zonistius.....	128
Lutjanidae.....	324	lycaodon, <i>Salmo</i>	55
Lutjanus.....	324	Lycenchelys.....	475
ambiguus.....	327	paxillus.....	476

List of scientific names—Continued

	Page		Page
<i>Lycenchelys porifer</i>	476	<i>Lyosphaera</i>	503
<i>verruilli</i>	475	<i>globosa</i>	503
<i>Lycengraulis</i>	51	<i>lyricus, Gobionellus</i>	441
<i>grossidens</i>	51	<i>Gobius</i>	441
<i>poeys</i>	51	<i>lysamus, Bryolophus</i>	463
<i>Lynchonpoles</i>	72	<i>lythrochloris, Xenotis</i>	300
<i>argenteolus</i>	73	<i>Lythrulon</i>	331
<i>lychnus, Argyropelecus</i>	74	<i>flavigutatum</i>	331
<i>Lycas</i>	475	<i>opalescens</i>	331
<i>nebulosus</i>	475	<i>Lythrurus</i>	125
<i>seminudus</i>	475	<i>ardens</i>	125
<i>Lycichthys</i>	472	<i>atropes</i>	125
<i>latrifrons</i>	472	<i>bellus</i>	125
<i>minor</i>	472	<i>cyanocephalus</i>	125
<i>Lycocara</i>	478	<i>fasciolaris</i>	125
<i>parri</i>	478	<i>lurus</i>	125
<i>Lycodalepis</i>	476	<i>maurinus</i>	125
<i>mucosus</i>	476	<i>notropis</i>	125
<i>polaris</i>	476	<i>punctulatus</i>	125
<i>Lycodapodidae</i>	478	<i>roseipinnis</i>	125
<i>Lycodapus</i>	478	<i>umbratilis</i>	125
<i>attenuatus</i>	479	<i>Lythrypnus</i>	442
<i>dermatinus</i>	478	<i>dalli</i>	442
<i>extensus</i>	478		
<i>fiestasfer</i>	478	<i>Macabi</i>	40
<i>grossidens</i>	479	<i>macarellus, Caranx</i>	270
<i>lycodon</i>	478	<i>Decapterus</i>	270
<i>mandibularis</i>	479	<i>macaskeyi, Ictalurus</i>	153
<i>parviceps</i>	478	<i>maclellandii, Bregmaceros</i>	208
<i>Lycodes</i>	474	<i>macdonaldi, Acutomentum</i>	366
<i>anguis</i>	475	<i>Cocobis</i>	127
<i>brevipes</i>	475	<i>Conocara</i>	53
<i>cicatrifer</i>	475	<i>Cynoscion</i>	356
<i>coccineus</i>	476	<i>Eriscion</i>	356
<i>concolor</i>	475	<i>Nannobranchium</i>	168
<i>diapterus</i>	476	<i>Notropis</i>	127
<i>digitatus</i>	475	<i>Penopus</i>	483
<i>esmarkii</i>	474	<i>Salmo</i>	57
<i>frigidus</i>	474	<i>mykiss</i>	57
<i>gracilis</i>	475	<i>Zygoneetes</i>	177
<i>incisus</i>	475	<i>Macdonaldia</i>	76
<i>jordani</i>	474	<i>macellus, Prionistius</i>	380
<i>(Lycodopsis) pacificus</i>	474	<i>macer, Polyprosopus</i>	20
<i>mucosus</i>	476	<i>mackayi, Lumpenus</i>	471
<i>nebulosus</i>	475	<i>Siphostoma</i>	241
<i>palearis</i>	475	<i>Syngnathus</i>	241
<i>pavilloides</i>	476	<i>mackenzii, Salmo</i>	61
<i>paxillus</i>	476	<i>Stenodus</i>	61
<i>perspicillum</i>	474	<i>macloti, Carcharias</i>	17
<i>porifer</i>	476	<i>maclura, Raja</i>	29
<i>reticulatus</i>	474	<i>macouni, Chauliodus</i>	71
<i>serpens</i>	475	<i>Pterygocottus</i>	380
<i>terra-novae</i>	474	<i>macracanthum, Pristipoma</i>	334
<i>turneri</i>	476	<i>macracanthus, Alutarius</i>	496
<i>vahli</i>	474	<i>Pomadasy</i>	334
<i>verruilli</i>	475	<i>Macrhybopsis</i>	138
<i>zoarehus</i>	475	<i>festivus</i>	138
<i>lycodon, Lycodapus</i>	478	<i>gelida</i>	139
<i>Lycodontis albimentis</i>	93	<i>hyostoma</i>	139
<i>Lycodontus</i>	476	<i>marcomis</i>	138
<i>mirabilis</i>	476	<i>meeki</i>	139
<i>Lycodopsis</i>	474	<i>montana</i>	139
<i>crassilabris</i>	474	<i>sterletus</i>	139
<i>crotalinus</i>	473	<i>macrocephala, Olupea</i>	40
<i>pacificus</i>	474	<i>Murana</i>	77
<i>scaurus</i>	474	<i>macrocephalum, Etheostoma</i>	284
<i>Lycogramma</i>	476	<i>macrocephalus, Alvordius</i>	284
<i>brunnea</i>	476	<i>Chnus</i>	455
<i>Lycodontis</i>	91	<i>Gadus</i>	210
<i>jordani</i>	92	<i>Gonocephalus</i>	409
<i>Lyconectes</i>	471	<i>Mnierpes</i>	455
<i>aleutensis</i>	471	<i>Sebastes</i>	372
<i>Lycinema</i>	476	<i>Semotilus</i>	117
<i>barbatum</i>	476	<i>macrocerus, Monacanthus</i>	494
<i>lynx, Pimelodus</i>	153	<i>macrochelus, Catostomus</i>	106
<i>Lyocetus</i>	510	<i>macrochir, Aldrovandia</i>	75
<i>murrayi</i>	611	<i>Halosaurus</i>	75
<i>lyolepis, Anchoviella</i>	49	<i>Sebastes</i>	364
<i>Stolephorus</i>	49	<i>macrochura, Helioperca</i>	301
<i>Lyoliparis</i>	400	<i>macrochirus, Hemirhamphus</i>	199
<i>pulchellus</i>	400	<i>Lepomis</i>	301
<i>Lyomeri</i>	95	<i>Macrodon</i>	99, 354
<i>Lyopomi</i>	74	<i>aimara</i>	39
<i>Lyopsetta</i>	222	<i>ancy lodon</i>	354
<i>jordani exilis</i>	222	<i>auritus</i>	99

List of scientific names—Continued

	Page		Page
Macrodon feror.....	99	Macrostoma.....	166
microlepis.....	99	angustidens.....	166
mordax.....	354	castaneum.....	167
patana.....	99	margaritiferrum.....	166
tereira.....	99	quercinum.....	166
teres.....	99	macrostoma, Cyprinella.....	131
trahira.....	99	Hæmulon.....	329
macrodon, Cheilodipterus.....	296	macrostomum, Hæmulon.....	329
Erythrinus.....	99	Macrostomus.....	166
Linophryne.....	510	macrostomus, Accipenser.....	35
Macrodonophis.....	88	Macrouride.....	202
mordax ophurus.....	38	Macrourus.....	204, 205
macrodon, Squalus.....	19	albatrossus.....	206
macrogenis, Cerna.....	315	anguliceps.....	206
macrognathum, Opistognathus.....	452	bairdi.....	205
Macrognathus.....	245	barbiger.....	205
brevirostris.....	199	bergii.....	205
loricatus.....	36	boops.....	205
macrognathus, Leucichthys.....	62	bucephalus.....	205
macrolepidota, Anchovia.....	50	bulbiceps.....	205
Cobitis.....	176	canus.....	206
macrolepidotum, Moxostoma.....	111	capito.....	205
macrolepidotus, Catostomus.....	111	carminifer.....	206
Engraulis.....	50	cinereus.....	205
Fundulus heteroclitus.....	176	convergens.....	206
Leuciscus.....	116	cuspidatus.....	206
Notropis.....	125	fragilis.....	206
Pogonichthys.....	116	gracilicauda.....	206
Macrolepis.....	295	ingolfi.....	206
macrolepis, Phenagoniates.....	95	latinasutus.....	206
Pontinus.....	372	latirostratus.....	206
Reboides.....	95	lepturus.....	207
macronema, Pimelodus.....	159	leucophaeus.....	205
macronemus, Polynemus.....	256	(Malacocephalus) pectoralis.....	204
macrophthalmus, Astyanax.....	97	orbitalis.....	206
Caranx.....	271	pectoralis.....	204
Centropomus.....	328	steligidolepis.....	205
Epigonus.....	296	tenuicauda.....	206
Gobiesox.....	489	trichurus.....	206
Hypentelium.....	107	(Macrourus) steligidolepis, Macurus.....	205
macropoma, Bollmannia.....	443	macrourus, Alopias.....	18
Centropomus.....	318	Gymnotus.....	101
Diplectrum.....	318	Macrozoarces.....	473
Macrops.....	328	americanus.....	473
macrops, Balistes.....	492	macrurus, Ophiosoma.....	80
Bathygadus.....	203	Macurus.....	205
Calamus.....	337	acrolepis.....	203
Citharichthys.....	219	anguliceps.....	206
Conger.....	79	asper.....	203
Corvina.....	345	barbiger.....	205
Cottus.....	384	boops.....	205
Gnathypops.....	453	bucephalus.....	205
Hippoglossina.....	222, 223	bulbiceps.....	205
Opisthognathus.....	453	canus.....	206
Opisthopterus.....	45	capito.....	205
Pristigaster.....	45	carbburus.....	207
Vacuqua.....	345	carminatus.....	207
macroptera, Conocara.....	53	carminifer.....	206
Goniobatis.....	30	cinereus.....	205
macropterus, Hypentelium.....	107	(Celorhynchus) scaphosia.....	207
macropterus, Alepocephalus.....	53	convergens.....	206
Balistes.....	493	cuspidatus.....	206
Centrarchus.....	304	dorsalis.....	207
Labrus.....	304	firmsquamis.....	204
Thynnus.....	260	fragilis.....	206
macrophthalmus, Astyanax.....	97	goodii.....	203
Centropomus.....	328	gracilicauda.....	206
Dormitator.....	436	ingolfi.....	206
Gobiesox.....	489	latinasutus.....	206
Harengula.....	43	latirostratus.....	206
Priscanthus.....	322	leucophaeus.....	205
Pristipomoides.....	328	(Lionurus) holopis.....	207
Scomber.....	258	longifilis.....	205
Tetragonopterus.....	97	(Macrourus) steligidolepis.....	205
macropterna, Clupea.....	43	(Nematonurus) magnus.....	204
macroptus, Malacoctenus.....	460	occa.....	207
Myxodes.....	460	orbitalis.....	206
Macrobrachphosidae.....	245	suborbitalis.....	203
Macrobrachphosus.....	245	tenuicauda.....	206
macurus, Leptodermis.....	486	trichurus.....	206
Monacanthus.....	493	villosus.....	207
Macrobrachphosus hawaiiensis.....	246	macurus, Anago.....	80
scolopax.....	246	Gymnotus.....	101
macrorhinus, Accipenser.....	34	Oxydenticthys.....	88
macrospila, Paramutana.....	159	Oxydenticthys.....	87

List of scientific names—Continued

	Page		Page
macularius baileyi, Cyprinodon	181	maculosus, Serpens marinus	87
macularius, Cyprinodon	181	Serranus	311
maculata, Belone	197	maderensis, Ceratoscopelus	167
Etheostoma	290	Pseudomuraena	81
Hemitremia	135	Scopelus	167
Hybopsis	135	maderaspatesensis, Pristigasterichthys	310
Lophopselta	217	maderensis, Helicolenus	370
Morone	302	maendricus, Caularchus	488
Nerophis	243	Lepadogaster	488
nigra, Muræna	91	Mænde	340
nigra et viridis, Muræna	92	magasema, Geophagus	424
Perca	310	magdalena, Curmatia	100
Sciæna	436	Curmatia	100
maculiceps, Boleosoma	287	Otolithus	356
maculatofasciatus, Paralabrax	317	Paralichthys	223
Serranus	317	magistralis, Epinnula	261
maculato striatus, Diodon	502	magnoculus, Ophichthus	87, 88
maculatum, Aulostomus	245	Scytalophis	204
Boleosoma	287	magnus, Macrurus (Nematonurus)	327
maculatus, Alfordius	283	mahogoni, Lutianus	239
Anarrhichas	472	Mesopron	239
Apogon	295	maensis, Gasterosteus	176
Aulostomus	245	Pungius	176
Balistes	493	majalis, Colitis	175, 176
Canthodermus	493	Fundulus	401
Cirrhitus	470	major, Actinocor	274
Clinus	87	Blephans	401
Cogrus	386	Cyclopterus lipans	274
Cottus	471	Galichthys	114
Cryptacanthodes	501	Ptychocheilus	263
Diodon	436	Makaira	263
Dormitator	14	ampla	263
Galeus	99	hole	263
Gasteropeleus	283	marlina	263
Hadropterus	193	mazara	263
immaculatus, Platypocilus	355	mitsukurii	263
Labrus squeteague	435	nigricans	504
Lembus	470	makus, Ranzania	281
Leptoclinus	295	malabanco, Elacate	99
Monopron	343	malabancus, Esox	99
Mullus	290	Hoplias	99
Nothonotus	11	Synodus	99
Notorynchus	151, 152, 158	Malacanthide	353
Pimelodus	193	Malacanthus	353
Platypocilus	217	plumieri	353
Pleuronectes	36	Malacocephalus occidentalis	204
Proceros	268	Malacocottus	388
Psenes	316, 317	kincaidii	388
Rhypticus	316	zonurus	388
Rypticus	257	Malacotenus	459
Scomber	257, 258	biguttatus	460
Scomberomorus	311	bimaculatus	460
Serranus	498	culbrebræ	460
Sphæroides	498	delalandi	460
Tetrodon hispidus var.	343	gilli	460
Upeneus	182	lugubris	460
Zoogoneticus	421	macropus	460
maculicauda, Astatheros	421	moorei	460
Cichlosoma	331	ocellatus	459
Orthostoechus	501	puertoricensis	460
maculifer, Diodon	217	varius	459
Platophrys	321	versicolor	460
maculiferus, Hypoplectrus	217	Malacosteidae	71
Pleuronectes	427	Malacosteus	71
maculipinna, Iridio	427	niger	71
Julis	201	malarmat, Peristedion	405
maculipinnis, Exocoetus	422	maliger, Pteropodus	369
Heros	16	Sebastichthys	369
Isogomphodon	230	malispinosus, Monomeiopis	480
Monochir	92	malleus, Zygena	18
Thyrosoides	360	Mallotus	66
maculo-cinctus, Sirothrodus	43	villosus	66
maculosa, Harengula	212	(Mallotus) pacificus, Salmo	66
Lota	87	malma, Salmo	61
Muraena	488	Salvelinus	61
Thalassophryne	373	spectabilis, Salvelinus	61
maculo-seriatus, Chirus	329	Malthæa	506
maculosum, Hamulon	267	truncata	507
maculosus, Acentrolophus	35	Malthæ	506
Acipenser	107	cubifrons	507
Catostomus	303	elater	507
Centrarchus	212	Malthæa nasuta	507
Gadus	268	notata	507
Nomeus	390	Malthopsis	507
Oligocottus	223	spinosa	507
Paralichthys	145	spinulosa	507
Primephales		managuense, Parapetenia	418

List of scientific names—Continued

	Page		Page
managuensis, <i>Heros</i>	418	marina, <i>Pastinaca</i>	29
<i>Pimelodus</i>	157	<i>Perca</i>	363
<i>Rhamdia</i>	157	<i>rubra</i> , <i>Perca</i>	226
manatia, <i>Raia</i>	32	<i>sectatrix</i> , <i>Perca</i>	340
manatinus, <i>Barathrodemus</i>	482	<i>Strongylura</i>	196
Mancalias.....	508	<i>Vulpecula</i>	18
<i>shufeldti</i>	508	marinus, <i>Diabolus</i>	32
<i>uranoscopus</i>	508	<i>dorsatus</i> , <i>Petromyzon</i>	9
mandibularis, <i>Lycodapus</i>	479	<i>Esox</i>	196
manglicola, <i>Gobius</i>	439	<i>foie quadratus</i> , <i>Faber</i>	358
<i>Rhinogobius</i>	439	<i>Gasterosteus</i>	239
mango, <i>Polynemus</i>	256	<i>maculosus</i> , <i>Serpens</i>	87
manitou, <i>Perca</i>	253	<i>Mugilops</i>	252
manitoulinus, <i>Leucichthys</i>	64	<i>Petromyzon</i>	8
manuari, <i>Lepidosteus</i>	38	<i>Sebastes</i>	363
manni, <i>Gambusia</i>	186	<i>Silurus</i>	148
manni, <i>Zygonectes</i>	174	marki, <i>Emblemaria</i>	465
Manta.....	32	mailina, <i>Makaira</i>	263
<i>americana</i>	32	marmorata, <i>Amia</i>	38
<i>birostris</i>	32	<i>Pteroplatea</i>	29
<i>hamiltoni</i>	32	<i>Ulea</i>	393
manta, <i>Cephaloptera</i>	32	<i>Unibranchapertura</i>	76
<i>Cephalopterus</i>	32	marmoratum, <i>Corallicola</i>	456
Mapo.....	440	<i>Lepophidium</i>	481
mapo, <i>Gobius</i>	440	<i>Leptophidium</i>	484
maraldi, <i>Gadella</i>	211	marmoratus, <i>Ameiurus</i>	154
<i>Gadus</i>	211	<i>Cottus</i>	387
marcgravi, <i>Aulastome</i>	245	<i>Cremnobates</i>	456
<i>Coryphæna</i>	264	<i>Gobiesox</i>	488
<i>Rhinobatus</i>	23	<i>Hemitripterus</i>	382, 393
<i>Salmo</i>	100	<i>Pimelodus</i>	154
marconis, <i>Hybopsis astivalis</i>	138	<i>Rhinichthys</i>	139
<i>Macrhybopsis</i>	138	<i>Rivulus</i>	179
Margariscus.....	116	<i>Scorpaenichthys</i>	382
<i>margarita</i>	116	<i>Sphæroides</i>	498
<i>nachtiebi</i>	116	<i>Synbranchus</i>	76
margarita, <i>Clinostomus</i>	116	<i>Tetodon</i>	498
<i>Echostoma</i>	70	<i>Thius</i>	163
<i>Margariscus</i>	116	marmorea, <i>Rabula</i>	91
margaritaceus, <i>Entomacrodus</i>	463	marmoreus, <i>Blennius</i>	461
<i>Salarias</i>	463	<i>Muraenophis</i>	91
margaritatum, <i>Myctophum</i>	169	<i>Syngnathus</i>	241
margaritatus, <i>Butiachus</i>	487	marshi, <i>Characidium</i>	100
<i>Porichthys</i>	487	marstoni, <i>Salmo</i>	61
<i>Tropidichthys</i>	500	<i>Salvelinus</i>	61
margaritifer, <i>Astatheros</i>	421	martæ, <i>Cyprinodon</i>	181
<i>Heros</i>	421	matui, <i>Pristigaster</i>	45
<i>Serranus</i>	311	matutina, <i>Atherina</i>	248
margaritifera, <i>Hæmulon</i>	331	<i>Spicara</i>	340
<i>Apodion</i>	339	martinicensis, <i>Acus</i>	243
<i>Macrostoma</i>	166	<i>Aylopon</i>	320
margaritiferus, <i>Notoscopelus</i>	166	<i>Menticirrhus</i>	351
margaritum, <i>Echostoma</i>	70	<i>Nerophis</i>	243
margatoti, <i>Enneacanthus</i>	304	<i>Novaculichthys</i>	430
marginalis, <i>Epinephelus</i>	310	<i>Ocyanthias</i>	323
<i>Hippocampus</i>	244	<i>Umbrina</i>	351
marginata, <i>Aphoristia</i>	231	<i>Vomer</i>	275
<i>Helioperca</i>	301	<i>Xylichthys</i>	430
<i>Rissola</i>	485	martinicus, <i>Gobius</i>	442
<i>Uranidea</i>	384	<i>Membras</i>	248
marginatum, <i>Ophidium</i>	485	<i>Smaris</i>	340
marginatus, <i>Abudefduf</i>	413	<i>Upeneus</i>	343
<i>Bodianus</i>	313	mas, <i>Salmo</i>	56
<i>Brosimius</i>	479	Mascalongus.....	172
<i>Brosimophycis</i>	479	maschalespilos, <i>Scarus</i>	433
<i>Caranx</i>	272	<i>Sparisoma</i>	433
<i>Chaetodon</i>	415	mascoæ, <i>Moxostoma</i>	110
<i>Cottus</i>	384	<i>masoni</i> , <i>Salmo</i>	58
<i>Helias</i>	413	masquinongy, <i>Esox</i>	172
<i>Hemicaranx</i>	271	<i>immaculatus</i> , <i>Esox</i>	173
<i>Neobythites</i>	481	<i>ohiensis</i> , <i>Esox</i>	172
<i>Notius</i>	156	<i>massachusettsensis</i> , <i>Monacanthu</i>	494
<i>Phycis</i>	212, 213	<i>massiliensis</i> , <i>Polyprion</i>	308
<i>Pimelodus</i>	152	<i>Scorpaena</i>	308
<i>Pomotis</i>	301	Masticura.....	28
<i>Serranus</i>	311	matamorenensis, <i>Limia</i>	192
<i>Symphurus</i>	231	matejuelo, <i>Amphiprion</i>	235
<i>Xuel</i>	272	mathematicus, <i>Tetodon</i>	497
marianum, <i>Holocentrum</i>	236	mattoiwacca, <i>Clupea</i>	42
marianus, <i>Flammeo</i>	236	matutinus, <i>Alburnellus</i>	126
marina, <i>Bagre</i>	148	<i>Lythrurus</i>	126
<i>capite striato</i> , <i>Perca</i>	330	<i>mauricei</i> , <i>Argyrius</i>	275
<i>cauda forcipata</i> , <i>Brama</i>	265	<i>mauriti</i> , <i>Chaetodon</i>	415
<i>cauda nigra</i> , <i>Perca</i>	330	<i>Eleotris</i>	268
<i>gibbosa cinerea</i> , <i>Perca</i>	329	<i>Grammistes</i>	332
<i>Mugilops</i>	252	Maurolicæ.....	73

List of scientific names—Continued

	Page		Page
Maurolicus	73	megacephalus, Cyclogaster	400
amrthystino punctatus	73	Liparis	400
attenuatus	73	Megaderia	94
lucetinus	73	megalaspius, Acipenser (Huso)	35
oculatus	73	Doratonotus	428
pennanti	73	Megalobrycon	98
tripunctulatus	73	Megalocottus	387
maxillaris, Lotella	211	laticeps	388
Murenoides	468	platycephalus	388
Maxillingua	146	megalonotus, Fristis	23
maxillingua, Cyprinus	146	Megalonema punctatum	159
Exoglossum	146	robustum	159
maxillosa, Gnathypops	453	Megaperca	308
maxillosus, Opisthognathus	453	Megalopids	39
Rhinichthys	139	Megalops atlanticus	39
maximus, Cetorhinus	20	cephediana	46
Hippoglossus	221	elongatus	39
Labrus	425	notata	44
Lachnolaimus	425	ogline	44
Squalus	20	megalops alboolus, Notropis	129
maxinkuckiensis, Alvordius	284	megalops, Alburnus	125
Hadropterus	284	Clupea	42
mayeri, Mayerina	81	Cylindrosteus	37
Mayerina	81	Cyprinus	128
mayeri	81	Eumichthys	410
Maynea	476	Holconotus	410
brunnea	476	Micropogon	349
bulbiceps	477	Opsopogon	115
californica	477	Paranotopis	125
patagonica	476	Pimelodus	152
pusilla	477	Trypharodon	115
mazara, Makaira	263	megalocephthalmus, Lutianus	327
Tetrapturus	263	Neomans	327
mazatlana, Seriola	279	megalois, Ichelis	300
Solea	230	Xenotis	300
mazatlanum, Hamulon	331	Megaperca	308
mazatlanus, Achirus	230	ishinagi	308
Zonichthys	279	Megaphalus	488
mcnieli, Gambusia	186	megastoma, Opisthognathus	452
meadia, Imostoma	286	Megastomatobius	102
Ulocentra	286	cyprinella	102
meany, Ruscarius	378	megastomus, Catostomus	107
mearnsi, Hesperoleucus	122	Meladerma	281
Notropis	122	nigerrima	281
Meda	147	Melamphaps beani	233
fulgida	147	bispinosus	233
media, Palometa	267	cristiceps	233
Medialuna	358	lugubris	233
californiensis	358	nycterinus	233
mediana, Ancistrus	161	Melamphaidae	232
medians, Ancistrus	161	melampodus, Apogonichthys	296
Medidae	147	melampterus, Salmo	55
mediocris, Clupea	41, 42	melampygus, Caranx	272
Pomolobus	42	Xurel	272
mediorum, Antennarius puncto	505	Melanichthys	338
Medipellona	45	Melaniris	248
medirostris, Acipenser	34	balsanus	248, 249
mediterranea, Chimera	32	sardina	249
Sauridia	163	Melanobranchus	208
mediterraneus, Enchelyopus	208	melanobranchus, Bathygadus	208
Hoplostethus	234	melanocephalus, Plargyrus	145
Lopharis	280	Melanocetidae	510
Scomber	259	Melanocetus	510
Sternopyx	74	johnsoni	510
medius, Anisarchus	470	(Liocetus) murrayi	511
Calamus	337	niger	510
Centropomus	306	polyactis	510
Clinus	470	melanocheir, Julius	429
Grammateus	37	Stellifer	347
Stromateus	267	Zestis	347
medius, Myoxocephalus	337	melanocheira, Belone	196
medusicola, Caranx	272	melanogaster, Pleuronectes	224
Xurel	272	Pocilia	191
medusophagus, Schedophilus	268	Melanogrammus	210
meeki, Cichlasoma	416	aelefinus	211
Cichla	416	melanopogon, Heros	421
Eucinostomus	341	melanopoma, Polynemus	256
Fundulus	178	melanops, Callurus	302, 303
Holocentrus	236	Catostomus	109
Hybopsis	139	Dionda	144
Machyropsis	139	Haplochilus	185
Microgobius	444	Ichelis	298
Thyrina	249	Minytrema	109
Ulaema	341	Sebastes	364
megacephalus, Calamus	338	Sebastodes	365
Chitonotus	375	Sebastosomus	365

List of scientific names—Continued

	Page		Page
Mesoprion cyanopterus.....	325	microdon, Cheonda.....	118
cynodon.....	325	Cyclothone.....	72
elegans.....	323	Gobionellus.....	441
flavescens.....	325	Gobius.....	441
griseus.....	325	Gonostoma.....	72
guttatus.....	326	Osmerus.....	66
inermis.....	327	Pseudotriakis.....	12
isodon.....	326	Richardsonius.....	118
linea.....	325	Microdontophis.....	87
litura.....	325	altipinnis.....	87
mahogoni.....	327	Microelectris.....	437
oajano.....	327	mindii.....	437
pacificus.....	325	panamensis.....	437
parvus.....	325	Microgadus.....	210
pomacanthus.....	324	proximus.....	210
profundus.....	326	torced.....	210
retrospinus.....	324	micrognathus, Lonchopisthus.....	453
ricardi.....	327	Opisthognathus.....	453
rosaceus.....	328	Microgobius.....	443
sobra.....	326	cyclolepis.....	443
uninotatus.....	327	emblematicus.....	443
vivanus.....	326	eulepis.....	444
vorax.....	328	gulosus.....	443
Mesops.....	424	meeki.....	444
Mesopus.....	67	miraflorensis.....	444
mesotrema, Asternotremia.....	215	signatus.....	443
messieri, Sirembo.....	480	thalassinus.....	444
messinensis, Gymnocephalus.....	267	microlepidota, Gila.....	112
metallica, Agosia.....	140	Microlepidium.....	211
Echeneis.....	449	grandiceps.....	211
metallicus, Girardinus.....	189	Microlepidotus.....	335
Notropis.....	124	brevipinnis.....	335
Metaponops cooperi.....	218	inornatus.....	335
Metonops.....	218	microlepidotus, Cynoscion.....	355
cooperi.....	218	Labrisomus.....	459
metopias, Triglops.....	380	Orthodon.....	112
metopoclampus, Scopelus.....	168	Otolithus.....	355
Metrogaster.....	409	microlepis, Antimora.....	211
lineolatus.....	409	Epicyrus.....	98
Metzelaaria.....	370	Hoplias.....	99
tridecimspinosa.....	370	Lepophidium.....	485
metzelaari, Evermannichthys.....	447	Leptophidium.....	485
Kyphosus.....	339	Macrodon.....	99
metzi, Gibbonsia.....	458	Tristrotis.....	314
meuleni, Diodon.....	502	microlepis, Bathytroctes.....	52
mexicana, Aztecula.....	143	microleptes, Ariphaton.....	309
campi, Signalosa.....	47	microlophus, Dolopichthys.....	509
Pamphers.....	323	Pomotis.....	302
Pocilia.....	192	Micromesurus.....	31
Signalosa.....	47	Micrometrus.....	409
mexicanum, Myctophum.....	168	minimus.....	409
Nannobranchium.....	168	mironemus, Peristedion.....	406
mexicanus, Amblyopsis.....	448	Microperca.....	282, 294
Astyanax.....	96	procliaris.....	294
Auchenopterus.....	457	proliaris.....	294
Centropomus.....	306	punctulata.....	294
Chato'ssus.....	47	microperca, Etheostoma.....	294
Chonophorus.....	442	microphthalum, Hemulon.....	329
Eugerres.....	343	Micropogon.....	349
Gerres.....	343	altipinnis.....	349
Gobius.....	442	argenteus.....	349
Mugil.....	253	ectenes.....	349
Pemphers.....	323	furneri.....	349
Saurus.....	163	lineatus.....	349
Tetragonopterus.....	96	megalops.....	349
meyeri, Iridio.....	427	trifilis.....	349
mezquital, Chirostoma.....	250	undulatus.....	349
miacanthus, Ardiellus.....	378	micropogon, Ceratichthys.....	137
miarchus, Anchoviella.....	47	Nocomis.....	137
Stolephorus.....	47	Micropogonias.....	349
micraspidophorus, Cyclogaster (Neoliparis).....	399	microps, Agonostoma.....	254
Neoliparis.....	399	Atherina.....	246
(Micristius) brachypterus, Zygonectes.....	185	Belone.....	196
Micristodus.....	18	Callurus.....	299
punctatus.....	18	Callyodon.....	435
microcephalus, Erythrinus.....	99	Carcharias.....	16
Gasterosteus.....	237	Caulolatilus.....	357
williamsoni.....	237	Catostomus.....	106
Pleuronectes.....	228	Corvina.....	346
Somniosus.....	22	Cottunculus.....	392
Squalus.....	21, 22	Dajaus.....	254
Microconger.....	81	Nebria.....	354
caudalis.....	81	Otolithus.....	356
(Microconger) caudalis, Leptocephalus.....	81	Pagellus.....	336, 337
Microdesmus.....	472	Pseudoscarus.....	435
dipus.....	472	Stellifer.....	346
retropinnis.....	472	Strongylura.....	196

List of scientific names—Continued

	Page		Page
microptera, Evolantia.....	199	miniatus, Sclerotis.....	300
Rhamdia.....	157	Sebastichthys.....	365
Micropterus.....	297, 298	Zoogoneticus.....	182
dolomieu.....	298	Minniellus.....	133
pseudoplites.....	298	minima, Capriscus.....	496
pseudotites.....	298	Cichla.....	298
salmoides.....	297	Clupea.....	41
micropterus, Etheostoma.....	290	Perca.....	287
Exocoetus.....	199	minimus, Cymatogaster.....	409
Pimelodus.....	157	Micrometrus.....	409
Micropteryx.....	276, 279	miniofrenatus, Scaurus.....	432
cosmopolita.....	276	Minnilus.....	123
micropteryx, Notropis.....	124	bellus.....	126
Alburnellus.....	124	dinemus.....	123
Platysomus.....	275	microstomus.....	133, 134
microphthalmum, Hæmulon.....	329	notatus.....	145
microphthalmus, Cichlaurus.....	416	nigripinnis.....	125
Dormitator.....	436	punctulatus.....	126
Heros.....	416	scepticus.....	123
Tetrabanchus.....	76	xanurus.....	130
Tetragonopterus.....	96	Minomus.....	105
micropus, Clupea.....	46	bardus.....	104
Etrumeus.....	46	delphinus.....	104
Gasterosteus.....	238	jarrovi.....	104
microrhynchus, Acipenser.....	34	platyrhynchus.....	104
microrrhynchus, Pseudoscarius.....	435	minor, Anarrhichus.....	472
micros, Citharichthys.....	219	Aphanopus.....	262
Etopus.....	219	Courchia.....	208
Microspathodon.....	415	Cyclopterus, paris.....	400
azurissimus.....	416	Lyceichthys.....	472
cinereus.....	416	Umia.....	255
microstigmus, Myrophis.....	83	minutulus, Chirolepis.....	438
Microstoma.....	228	minutus, Cyclopterus.....	397
microstoma, Crenilabrus.....	424	Hemiramphus.....	97
Atherina.....	246	Minytrema.....	109
Scartella.....	461	melanops.....	109
Uranidea.....	385	minytremus, Gymnichthys.....	403
Microstomidae.....	68	mirabile, Exoglossum.....	142
Microstomus.....	228	mirabilis, Chimæra.....	33
greenlandica.....	68	mirabilis, Clupea.....	41
kitt.....	228	Crystallichthys.....	401
latidens.....	228	Gillichthys.....	444
pacificus.....	228	Lycodonus.....	476
microstomus, Blennius.....	461	Phenacobius.....	142
Carpiodes.....	104	miraflorensis, Microgobius.....	444
Citharichthys.....	219	mirum, Calloptilum.....	208
Conger.....	80	missillensis, Ameiurus.....	154
Etopus.....	219	mississippiensis, Morone.....	307
Iridio.....	427	Pristis.....	23
Minnilus.....	133, 134	missouriensis, Hybopsis.....	133
Pleuronectes.....	228	mittchilli, Acanthocottus.....	386
microtrachys, Raja.....	25	Acipenser.....	34
mictrura, Pteroplatea.....	29	alterusta, Perca.....	307
Raja.....	29	Anchoviella.....	48
micurum, Syacium.....	218	Argyrosus.....	275
milberti, Acipenser.....	34	Cottus.....	386
Arius.....	149	Engraulis.....	48
Carcharias.....	16	interrupta, Perca.....	307
(Prionodon).....	16	Perca.....	307
Lepidosteus.....	37	Stolephorus.....	48
milbertianus, Syngnathus.....	242	Mitchillina.....	52
miles, Porogadus.....	482	bairdi.....	52
pectoralis, Prionotus.....	407	mitrulus, Hesperoleucus.....	122
milesi, Pimphales.....	145	mitsukuru, Makara.....	263
milharis, Gymnothorax.....	92	Tetrapturus.....	263
Thyrsoidea.....	92	miura, Rabida.....	156
militaris, Bellator.....	408	miurus, Noturus.....	156
Prionotus.....	408	Ophichthys.....	89
Silurus.....	159	Scytalichthys.....	89
miltschitch, Salmo.....	54	Mixonus.....	483
milleri, Bathylagus.....	69	caudalis.....	483
Cichlasoma.....	417	laticeps.....	483
Theraps.....	417	mixtus, Gadus tomcodus.....	210
millipunctatus, Gasterosteus.....	239	mizolepis, Plectromus.....	233
milneri, Nocomis.....	136	Scopelus.....	233
Pagellus.....	337	Mnerpes.....	455
milneriana, Hemitrema.....	121	macrocephalus.....	455
milnerianus, Phoxinus.....	121	Mobula.....	31
minace, Gila.....	114	auriculata.....	31
mindii, Stallifer.....	347	hypostoma.....	32
Syngnathus.....	242	olferi.....	32
mineopas, Bryttus.....	299	thypostoma.....	32
miniaturum, Peristedion.....	405	mobular, Raja.....	31
minatus, Lepidopomus.....	300	Mobulidae.....	31
Lepomis.....	300	modesta, Cheonda.....	120
Rosticola.....	365	Gambusia.....	185, 192

List of scientific names—Continued

	Page		Page
modestus, Julis	428	Moniana	131
Xyrichthys	430	aurata	132
Mœbia	481	boucardi	132
promelas	481	callisema	132
merens, Gonopterus	362	chihuahuæ	132
Moharra	342	complanata	131
evermanni	342	couchi	131
rhombea	342	deliciosa	134
mohavensis, Siphateles	123	formosa	129, 131
mohri, Clinus	471	frigida	131
mojarra, Cichlasoma	417	gibbosa	131
Theraps	417	gracilis	131
Mola	503	jugalis	131
mola	503	lætabilis	131
rotunda	503	leonina	131
mola, Diodon	503	lutrensis	131
Mola	503	nitida	135
Tetrodon	503	preserpina	132
Molacanthus	503	pulchella	131
pallasi	503	robusta	132
Molari	94	rutia	131
molestum, Gobiosoma	446	santamarie	131
Molida	503	tristis	132
Molinesia fasciata	192	Monoceratias	508
Mollienæ	191	acanthias	508
formosa	192	monoceros, Alutera	496
gracilis	192	Balistes	495, 496
jonesi	188	unicolor, Balistes	496
latipinna	192	Monochir maculipinnis	230
petenensis	192	reticulatus	229
sphenops	192	Monolene	220
cuneata	193	atrimana	220
vanderpolli	193	sessilicauda	220
spilura	192	Monomeropus	480
velifera	192	malispinosus	480
Mollesia gracilis	192	Monomitopus	480
latipinna	191	torvus	480
petenensis	192	Monomitra	404
velifera	192	Monopion	295
mollis, Aphyonius	484	maculatus	295
Bathysaurus	164	pigmentarius	295
Bothrocar	477	Monopterhinus	11
Caenoproctus	403	monopterygius, Aspidophoroides	397
Pleuronectes	229	Cottus	397
molluccensis, Apogon	295	Labrax	374
Molva	212	Pleurogrammus	374
hunta	212	monophtthalmus, Auchenopterus	456, 457
molva	212	Monosira	344
vulgaris	212	stahl	344
molva, Gadus	212	monstrosa, Chimæra	32
Molva	212	montagu, Leparis	399
Molvelia	208	montana, Cheonda	119
borealis	208	Macrhybopsis	139
Monacanthidae	493	montanus, Clinostomus	119
Monacanthum proboscideum	495	Hybopsis	139
Monacanthus	494	Thymallus	66
amphioxys	494	Montereya	391
aurea	494	recalva	391
ciliatus	494	montereyensis, Gibbonsia elegans	458
davidsoni	494	Raja	26
filamentosus	494	montezuma, Astatheros	421
gallinula	494	Heros	421
hispidus	494	montezumæ, Xiphophorus	194
irroratus	494	monticola, Dajaus	254
macrocerus	494	Mugil	254
macrurus	493	montregis, Chiola	131
massachusettsensis	494	moorei, Acteis	460
occidentalis	494	Congrammus	454
oppositus	495	Malacoctenus	460
pardalis	493	mordax abbotti, Osmerus	67
paralanus	494	mordax, Atherina	66
piraaca	494	Conger	88
pullus	494	Engraulis	47
punctatus	494	Gymnothorax	91
ruppelli	494	Macrodon	354
setifer	494	Muraena	91
signifer	494	ophiurus, Macrodonophis	88
spilonotus	495	Osmerus	67
striatus	494	Sagenichthys	354
tuckeri	495	spectrum, Osmerus	67
varius	494	morbibundus, Balistes	492
monacanthus, Plectropoma	313	moringa, Gymnothorax	91
monacus, Ceratichthys	138	Muraena	91
Ermonax	138	Moringua	89
mona, Stephanoberyx	232	boekei	89
monensis, Squalus	19	lateralis	89

List of scientific names—Continued

	Page		Page
Moringua linearis.....	89	Mugil berlandieri.....	253
lumbrioides.....	89	brasiliensis.....	253
moringua, Muræna.....	91	cephalus.....	252, 253
Moringuidæ.....	89	chanos.....	40
morio, Centrolophus.....	267	cinereus.....	341
Epinephelus.....	312	curema.....	253
Serranus.....	312	gamardianus.....	254
mormyrus, Campostoma.....	146	guntheri.....	253
Morone.....	307	guntheri.....	253
americana.....	307	grunniens.....	352
flavescens.....	281	hospes.....	254
interrupta.....	307	incilis.....	254
maculata.....	302	irretitus.....	254
mississippiensis.....	307	lebranchus.....	253
pallida.....	307	lineatus.....	253
rufa.....	307	liza.....	253
Moronidæ.....	307	mexicanus.....	253
Morrhua.....	210	monticola.....	254
americana.....	210	obliquus.....	349
californica.....	210	petrosus.....	253
punctatus.....	211	plumieri.....	253
morrhua, Gadus.....	210	proboscideus.....	254
Serranus.....	310	rammelsbergii.....	253
morrisi, Leptocephalus.....	79	salmonius.....	40
Moseleya.....	203	setosus.....	254
moseri, Verasper.....	223	tang.....	253
motaguensis, Heros.....	418	thoburni.....	253
Parapetenis.....	418	trichodon.....	254
Pimelodus.....	157	Mugilidæ.....	252
Rhamdia.....	157	mugiloides, Eleotris.....	436
Motella.....	208	Mugilomorus.....	39
argentata.....	208	anna-carolina.....	39
caudacuta.....	209	Mugilops.....	252
ensis.....	208	cyaneila.....	252
reinhardtii.....	208	cyaneilus.....	252
septentrionalis.....	208	marina.....	252
motta, Elacate.....	281	marinus.....	252
moucharra, Glyphisodon.....	415	mülleri, Benthosema.....	170
mowbrayi, Hæmulon.....	329	Leporinus.....	88
Rhinogobius.....	439	Pempheris.....	323
Moxostoma.....	109	Rhamphichthys.....	101
album.....	110	Salmo.....	170
alleganiensis.....	111	mülleri, Carcharias (Prionodon).....	16
anisurum.....	109	Mulhyphenus.....	343
aureolum.....	110	Mulhæ.....	343
austrium.....	110	Mulloidæ.....	344
breviceps.....	111	rathbuni.....	344
bucco.....	110	Mullus.....	343
campbelli.....	109	auratus.....	343
claviformis.....	109	barbatus.....	343
collapsus.....	110	auratus.....	343
congestum.....	110	bifasciatus.....	343
conus.....	111	flavolineatus.....	344
coregonus.....	110	imberbis.....	295
crassilabre.....	111	maculatus.....	343
kennerlyi.....	109	multifasciata, Adina.....	180
lesueurii.....	111	Hydrargyra.....	178
macrolepidotum.....	111	Parapetenia.....	418
mascoia.....	110	multifasciatum, Cichlasoma.....	418
papulosum.....	109	Gobiosoma.....	446
parvidens.....	110	multifasciatus, Pronotogrammus.....	322
pdiense.....	110	multifilius, Blennius.....	461, 462
poecilurum.....	111	multiguttatum, Plectropoma.....	313
robustum.....	111	multiguttatus, Alphestes.....	313
rupiscartes.....	111	multilineata, Furcra.....	412
tenuis.....	109	Pocilia.....	192
thalassium.....	110	multilineatum, Hæmulon.....	330
trisinatum.....	105	multilineatus, Hellases.....	412
valenciennesi.....	108	Labrax.....	307
victoria.....	108	multimaenellus, Diodon.....	501
mucosum, Xiphidion.....	468	multiocellatus, Antennarius.....	508
mucosus, Acanthocottus.....	387	Chironectes.....	508
Liparis.....	398	multipunctata, Skiffia.....	183
Lycodalepis.....	476	multipunctatus, Xenoderm.....	183
Lycodes.....	476	multiradiatus, Arnis.....	151
Neoliparis.....	399	Characodon.....	183
Xiphister.....	468	Cubiceps.....	183
mucronata, Perca.....	308	Lernionchthys.....	183
mucronatum, Ophidium.....	467	multispinosa, Herotlapia.....	423
mucronatus, Neoconger.....	81	multispinosus, Heros.....	423
Odontomatus.....	45	multiocellata, Muræna.....	92
midfish, Fundulus.....	175	multofasciata, Scæna.....	349
muelleri, Lepidomergas.....	279	mundaola, Anchoviella.....	50
Scopelus.....	170	mundaolus, Stolephorus.....	50
Mugil.....	252, 253	mundiceps, Xyrichtys.....	430
albula.....	253	mundicorpus, Inistius.....	430

List of scientific names—Continued

	Page		Page
mundus, Dactylagnus.....	455	Muraenophis punctata.....	92
Oligoplites.....	278	vicina.....	92
Urotrygon.....	30	Muraenopsis.....	87
Muraena.....	77, 90	triserialis.....	88
acuminata.....	85	murrinus, Calhurus.....	299
afra.....	91	murrayi, Ipnope.....	170
anguilla.....	77	Lyocetus.....	511
anguina.....	90	Melanocetus (Liocetus).....	511
aquea-dulcis.....	91, 93	muscarum, Gobiesox.....	490
argentea.....	77	musculus, Cyclopterus.....	400
argus.....	90	muskoka, Notropis.....	134
balearica.....	79	mustacrus, Serranus.....	310
bostonensis.....	77	Mustela.....	208
clepsydra.....	90	mustela, Gadus.....	208
ceca.....	84	Mustelus, Les.....	208
conger.....	79	mustellus, Gaidropsarus.....	208
down.....	92	Mustellus stellatus.....	13
elaborata.....	93	Mustelus.....	13, 14, 17
erebus.....	92, 93	asterias.....	13
hawaiiensis.....	87	californicus.....	13
helena.....	90	dorsalis.....	13
infernalis.....	92	felis.....	14
insularum.....	90	lævis.....	13
lentiginosa.....	90	lunulatus.....	13
lineopinnus.....	92	mento.....	13
lita.....	93	mustelus.....	13
longicauda.....	91	plebejus.....	13
lumbicus.....	76	mustelus, Galeus.....	13
macrocephala.....	77	Mustelus.....	13
maculata nigra.....	91	Squalus.....	13, 14
et viridis.....	92	muticus, Lethotremus.....	398
maculosa.....	87	Mycteroperca.....	314
melanotis.....	90	bonaci xanthosticta.....	314
meleagris.....	93	boulengeri.....	313
mordax.....	91	bowersi.....	314
moringa.....	91	calhura.....	315
moringua.....	91	camelopardalis.....	315
multocellata.....	92	falcata.....	315
myialeucosticta.....	90	phenax.....	315
myrialeucostictus.....	92	hopkinsi.....	314
nigra.....	79	olfax.....	315
nigricans.....	90	pardalis.....	315
ophis.....	87, 94	phenax.....	315
panamensis.....	91	rosacea.....	314
pantherina.....	93	tigris.....	315
pfeifferi.....	93	venadurum.....	314
picta.....	93	xenarcha.....	314
pinnata.....	78	Myctophidae.....	166
pinta.....	90	Myctophum.....	169
pintita.....	92	affine.....	169
polyuranodon.....	91	astratum.....	170
punctata.....	91	auroaternatum.....	169
ratibouri.....	89	benoiti.....	169
retifera.....	90	boops.....	169
rostrata.....	77	californiense.....	169
sanctæ-helenæ.....	92	crenulare.....	170
savanna.....	78	gracile.....	169
serpentina.....	77	hians.....	169
seu conger brasiliensis.....	93	humboldtii.....	169
siderea.....	93	hygomi.....	169
sordida.....	93	laternatum.....	170
supremo etc.....	79	luminosum.....	170
unicolor etc.....	90	margaritatum.....	169
variegata.....	93	mexicanum.....	168
muræna, Callechelys.....	86	nannochir.....	167
Lycodes.....	475	nitidulum.....	169
Muraenesocidae.....	78	nocturnum.....	168
Muraenesox.....	78	oculeum.....	169
coniceps.....	79	punctatum.....	169
savanna.....	78	procellarium.....	170
tricuspidata.....	78	protoculus.....	168
Muraenidae.....	90	rafinisqueti.....	168
Muraenoblenna.....	8, 94	regale.....	168
olivacea.....	8	remiger.....	169
Muraenoides.....	467	(Stenobranchius) leucopsarum.....	168
maxillaris.....	468	tennetulum.....	170
suef.....	467	townsendi.....	167
muraenoides, Biennius.....	467	mydrus, Cyprinodon.....	180
Gunnellus.....	468	myersi, Gambusia.....	185
Muraenophis.....	90	Mykiss.....	56
caramuru.....	91	mykiss, Salmo.....	56
catenula.....	94	agua-bomta, Salmo.....	59
curvilineata.....	91	Myliobatis.....	31
marmoreus.....	91	acuta.....	31
ocellatus.....	87	asperrimus.....	30
pantherina.....	93	bispinosus.....	31

List of scientific names—Continued

	Page		Page
<i>Myliobatis californicus</i>	31	<i>Myxostoma euryops</i>	110
<i>eeltenkee</i>	30	<i>pocellura</i>	111
<i>freminvilli</i>	31	<i>Myxus harengus</i>	253
<i>goodei</i>	31	<i>Myzopsetta</i>	226
<i>marginatus</i>	31		
<i>vespertilio</i>	31	<i>nachtheibi, Leuciscus</i>	116
<i>Mylocheilus</i>	112	<i>Margariscus</i>	116
<i>caurinus</i>	118	<i>nacaea, Gila</i>	114
<i>fraterculus</i>	112	<i>palnal, Sparactodon</i>	280
<i>lateralis</i>	112	<i>namaycush, Cristivomei</i>	59
<i>Myloleucus</i>	119, 120	<i>Salmo</i>	59
<i>parovanus</i>	121	<i>Nannobranchium</i>	167
<i>pulverentulus</i>	119, 120	<i>leucopsarum</i>	168
<i>thalassinus</i>	122	<i>macdonaldi</i>	168
<i>Mylopharodon</i>	112	<i>mexicanum</i>	168
<i>conocephalus</i>	112, 113	<i>nannochu</i>	167
<i>robustus</i>	112, 113	<i>nigrum</i>	167
<i>Mylorhina</i>	31	<i>regale</i>	168
<i>myops, Salmo</i>	163	<i>Nannocampus</i>	244
<i>Saurus</i>	163	<i>nanus</i>	244
<i>Trachinocephalus</i>	163	<i>subosseus</i>	244
<i>Myoxocephalus</i>	387	<i>nannochir, Myctophum</i>	167
<i>axillaris</i>	387	<i>Nannobachium</i>	167
<i>jaok</i>	387	<i>Nannostomus</i>	289
<i>mednus</i>	387	<i>nanomyzon, Catostomus</i>	106
<i>niger</i>	387	<i>Nanostoma</i>	289
<i>parvulus</i>	387	<i>arcansanum</i>	289
<i>polyacanthocephalus</i>	387	<i>elegans</i>	289
<i>stelleri</i>	387	<i>inscriptum</i>	289
<i>ve, rucosus</i>	387	<i>rupestre</i>	289
<i>myrialeucosticta, Muraena</i>	90	<i>swannanon</i>	289
<i>myrialeucostictus, Muraena</i>	92	<i>thalassium</i>	289
<i>myriaster, Rhomboidichthys</i>	217	<i>vinctipes</i>	289
<i>Myrichthys</i>	83	<i>zonale</i>	289
<i>acuminatus</i>	83	<i>Nansenia</i>	68
<i>oculatus</i>	83	<i>gicnlandica</i>	68
<i>pantostigmus</i>	83	<i>nanus, Engaulis</i>	47
<i>tigrinus</i>	83	<i>Gadus</i>	210
<i>xystus</i>	85	<i>Lipniscus</i>	401
<i>Myriolepis</i>	373	<i>Nannocampus</i>	244
<i>zonifer</i>	373	<i>Narcacion</i>	27
<i>Myriopristis</i>	235	<i>Narcaciontes</i>	27
<i>fulgens</i>	234	<i>Narcetes</i>	52
<i>occidentalis</i>	235	<i>erimelas</i>	52
<i>trachypoma</i>	234	<i>pluriserialis</i>	52
<i>Myripristis</i>	235	<i>stomias</i>	52
<i>humilis</i>	235	<i>Narcine</i>	27
<i>jacobus</i>	235	<i>brasiliensis</i>	27
<i>japonicus</i>	234	<i>corallina</i>	27
<i>lima</i>	235	<i>entemedor</i>	27
<i>occidentalis</i>	235	<i>indica</i>	27
<i>pocilopus</i>	235	<i>nigra</i>	27
<i>Myrophis</i>	82	<i>umbrosa</i>	27
<i>egmontis</i>	82	<i>naresi, Salmo</i>	61
<i>lumbicus</i>	83	<i>Salvelinus</i>	61
<i>microstigmus</i>	83	<i>narinari, Raja</i>	30, 31
<i>punctatus</i>	82, 83	<i>Stoasodon</i>	30
<i>vafer</i>	83	<i>naso, Anchoviella</i>	49
<i>mystacalis, Lepomis</i>	300	<i>Stolephorus</i>	49
<i>mystacina, Gnathypops</i>	453	<i>nasus, Acanthonotus</i>	75
<i>mystacinus, Epinephelus</i>	310	<i>Bassozetes</i>	481
<i>Gnathypops</i>	453	<i>Diplanchias</i>	503
<i>Serranus</i>	310	<i>Lamna</i>	19
<i>mystes, Scorpæna</i>	371	<i>Menticirrhus</i>	351
<i>mystoetus, Cetengraulis</i>	51	<i>Notacanthus</i>	75
<i>Engraulis</i>	51	<i>Squalus</i>	19
<i>mystinus, Sebastichthys</i>	364	<i>Umbra</i>	351
<i>Sebastosomus</i>	365	<i>nasuta, Malthca</i>	507
<i>Mystrophis</i>	88	<i>Netuma</i>	149
<i>interstinotus</i>	88	<i>nasutum, Agonostoma</i>	254
<i>Mytilophagus</i>	410	<i>Campostoma</i>	146
<i>fasciatus</i>	410	<i>nasutus, Astyanax</i>	97
<i>Myxine</i>	8, 94	<i>Cantherines</i>	491, 494
<i>glutinosa</i>	8	<i>Leuciscus</i>	139
<i>limosa</i>	8	<i>Ogcocephalus</i>	507
<i>Myxinidae</i>	8	<i>Trachynotus</i>	276
<i>Myxodagnus</i>	455	<i>natalis, Ameiurus</i>	153
<i>opercularis</i>	455	<i>analis, Ameiurus</i>	153
<i>Myxodes elegans</i>	458	<i>Pimelodus</i>	153
<i>lugubris</i>	480	<i>natator, Carcharhinus</i>	17
<i>macropus</i>	480	<i>Carcharias</i>	17
<i>varius</i>	459	<i>Nauclerus</i>	278
<i>versicolor</i>	480	<i>abbreviatus</i>	278
<i>Myxostoma</i>	109	<i>annularis</i>	278
<i>austriana</i>	110	<i>brachycentrus</i>	278

List of scientific names—Continued

	Page		Page
<i>Myxostoma compressus</i>	278	<i>Nemabrama</i>	234
<i>lecurus</i>	278	<i>Nematistidae</i>	280
<i>triacanthus</i>	278	<i>Nematistius</i>	280
<i>Naucrates</i>	279, 278	<i>pectoralis</i>	280
<i>cyanophrys</i>	278	<i>Nematognathus</i>	147
<i>ductor</i>	278	<i>Nematonurus</i>	203
<i>fanfarus</i>	278	<i>abyssorum</i>	203
<i>moveboracensis</i>	278	<i>acrolepis</i>	203
<i>polysarcus</i>	278	<i>armatus</i>	203
<i>serratus</i>	278	<i>cyclolepis</i>	203
<i>naucrates, Echeis</i>	448	<i>goodii</i>	203
<i>naufragium, Balistes</i>	451	<i>suborbitalis</i>	203
<i>Nautichthys</i>	382	<i>(Nematonurus) magnus, Macrurus</i>	204
<i>oculofasciatus</i>	382	<i>Nematonus</i>	482
<i>pribilovius</i>	382	<i>pectoralis</i>	482
<i>Nautiscus</i>	382	<i>nematophthalma, Merinthus</i>	372
<i>pribilovius</i>	382	<i>nematophthalmus, Sebastes</i>	372
<i>Nautopædium</i>	487	<i>nematopus, Neetroplus</i>	423
<i>porosissimum</i>	487	<i>Physiculus</i>	212
<i>nautopædium, Porichthys</i>	487	<i>Nemichthyidae</i>	83
<i>navarroensis, Hesperoleucus</i>	121	<i>Nemichthys</i>	83
<i>nawaga, Eleginus</i>	210	<i>avocetta</i>	83
<i>Gadus</i>	239, 210	<i>fronto</i>	83
<i>nazuz, Nototropis</i>	126	<i>infans</i>	83
<i>Orcella</i>	126	<i>scolopacea</i>	83
<i>Nealotus</i>	261	<i>scolopaceus</i>	83
<i>tripes</i>	261	<i>Nemocampsis</i>	297, 298
<i>neapolitanus, Lampugus</i>	265	<i>nemoptera, Divonina</i>	40
<i>nebrascensis, Nocomis</i>	137	<i>neo-boracensis, Fistularia</i>	245
<i>Nebria</i>	354	<i>Neobythites</i>	481
<i>nucrops</i>	354	<i>gilli</i>	481
<i>occidentalis</i>	354	<i>marginatus</i>	481
<i>zestus</i>	354	<i>ocellatus</i>	481
<i>Nebria concolor</i>	94	<i>robustus</i>	481
<i>nebularis, Platophrys</i>	217	<i>stelliferoides</i>	482
<i>nebulifer, Hieros</i>	417	<i>Neoclinus</i>	458
<i>Labra</i>	317	<i>blanchardi</i>	458
<i>Paralabra</i>	317	<i>satricus</i>	458
<i>Theraps</i>	417	<i>Neoconger</i>	81
<i>nebuliferus, Catostomus</i>	104	<i>nucronatus</i>	81
<i>nebulopunctatus, Gobius</i>	440	<i>peilongus</i>	82
<i>nebulosa, Acedia</i>	231	<i>vermiformis</i>	81
<i>Aphoristia</i>	231	<i>neogaea, Phille</i>	121
<i>Cerna</i>	315	<i>neogaeus, Phovinus</i>	121
<i>Perca (Percina)</i>	283	<i>neoguinaica, Albula</i>	40
<i>Saurida</i>	164	<i>neohantoni, Coregonus</i>	64
<i>nebulosus, Ameiurus</i>	154	<i>Neoliparis</i>	399
<i>Apionichthys</i>	230	<i>atlanticus</i>	399
<i>catulus, Ameiurus</i>	154	<i>beringianus</i>	399
<i>Chiropsis</i>	273, 374	<i>bristolensis</i>	399
<i>Clinus</i>	471	<i>callyodon</i>	399
<i>Eriscion</i>	355	<i>fissuratus</i>	399
<i>Gasterosteus</i>	239	<i>floræ</i>	399
<i>Lycias</i>	475	<i>grebnitzkii</i>	399
<i>Lycodes</i>	475	<i>greeni</i>	399
<i>Otolithus</i>	354, 255	<i>micraspidophorus</i>	399
<i>Pimelodus</i>	154, 155	<i>mucosus</i>	399
<i>Pteropodus</i>	369	<i>rutteri</i>	399
<i>Sebastes</i>	369	<i>(Neonore) hemirhamphus, Cyclopterus</i>	399
<i>Silurus</i>	155	<i>Neomænis</i>	324
<i>Urobatis</i>	30	<i>blackfordi</i>	453
<i>Urolophus</i>	30	<i>hastingsi</i>	326
<i>Zeus</i>	232	<i>jordani</i>	324
<i>Nectoliparis</i>	402	<i>megalphthalmus</i>	327
<i>pelagicus</i>	402	<i>Neomugil</i>	254
<i>Nector</i>	346	<i>digueti</i>	254
<i>alutus</i>	346	<i>Neomuræna</i>	91
<i>armatus</i>	346	<i>nigromarginata</i>	91, 93
<i>chrysoleucus</i>	346	<i>Neopocilia</i>	190
<i>verre-crucis</i>	346	<i>holacanthus</i>	190
<i>necturna, Gymnomuræna</i>	95	<i>Neostoma</i>	72
<i>necturus, Uropterygius</i>	94, 95	<i>bathypilum</i>	72
<i>Neetroplus</i>	423	<i>Neothunnus</i>	260
<i>bocourti</i>	423	<i>albacores</i>	260
<i>carpintis</i>	423	<i>allisoni</i>	260
<i>nematopus</i>	423	<i>catalinae</i>	260
<i>sieboldii</i>	423	<i>nepenthe, Eupomacentrus</i>	414
<i>nefastus, Pomotis</i>	300	<i>nephela, Heliopecten</i>	301
<i>neglecta, Clupea</i>	44	<i>nephelus, Lepomis</i>	301
<i>neglectus, Amblyodon</i>	352	<i>Pleuromichthys</i>	225
<i>Aploclinotus</i>	352	<i>Sphaeroides</i>	498
<i>nelsoni, Awaous</i>	442	<i>Tetrodon</i>	498
<i>Chonophorus</i>	442	<i>nerka, Onchorhynchus</i>	55
<i>Conorhynchus</i>	156	<i>Salmo</i>	55
<i>Coregonus</i>	65	<i>Nerophis anguineus</i>	243
<i>Salmo</i>	59		

List of scientific names—Continued

	Page		Page
<i>Nerophis heckeli</i>	244	<i>nigricans, Cottus</i>	386
<i>maculata</i>	243	<i>Cypselurus</i>	201
<i>martinicensis</i>	243	<i>Enchelycore</i>	90
<i>netinianus, Acipenser (Huso)</i>	35	<i>Entomacrodus</i>	463
<i>Nettastoma procerum</i>	82	<i>etowanus, Catostomus</i>	107
<i>Nettastomidae</i>	82	<i>Exocoetus</i>	201
<i>Netuma</i>	149, 150	<i>Girella</i>	339
<i>elattura</i>	149	<i>Heros</i>	419
<i>insculpta</i>	149	<i>Huro</i>	297
<i>kessleri</i>	149	<i>Hypentelium</i>	107
<i>nasuta</i>	149	<i>Hypoplectrus unicolor</i>	321
<i>oscula</i>	149	<i>Labrax</i>	308
<i>planiceps</i>	149	<i>Makara</i>	263
<i>platypogon</i>	149	<i>Muræna</i>	90
<i>thalassina</i>	149	<i>Parapetenia</i>	419
<i>neurates, Echeineis</i>	448	<i>Petromyzon</i>	8
<i>neuratroides, Echeineis</i>	449	<i>Pimelodus</i>	152
<i>neusirianus, Gasterosteus</i>	238	<i>Plectropoma</i>	321
<i>nevadensis, Apocope</i>	141	<i>Pseudocottus</i>	363
<i>Cyprinodon</i>	181	<i>Serranus</i>	310
<i>Rhinichthys (Apocope)</i>	141	<i>nigralabrus, Gronias</i>	154
<i>nevisense, Etheostoma</i>	283	<i>nigripinna, Corallicola</i>	456
<i>newberryi, Fario</i>	58	<i>nigripinnis, Argyrosomus</i>	62, 63
<i>Salmo</i>	58	<i>Bathylagus</i>	395, 396
<i>newberryi, Eucyclogobius</i>	444	<i>Cephalopops</i>	309
<i>Gobius</i>	444	<i>Clinus</i>	456
<i>newmani, Hyostoma</i>	286	<i>Coregonus</i>	63
<i>Nexilarius</i>	414	<i>Cotybus</i>	489
<i>concolor</i>	414	<i>Gobiesox</i>	489
<i>nianguae, Etheostoma (Hadropterus)</i>	285	<i>Leucichthys</i>	63
<i>Hypohomus</i>	285	<i>Minnilus</i>	125
<i>spilotum, Etheostoma</i>	285	<i>Promicropterus</i>	317
<i>nicaraguensis, Carcharias</i>	16	<i>Rhypticus</i>	317
<i>Eulamia</i>	16	<i>Siremba</i>	480
<i>Gambusia</i>	184, 186	<i>nigrostrius, Chætodon</i>	360
<i>Heros</i>	418	<i>Sarothiodus</i>	360
<i>Pimelodus</i>	157	<i>ngula, Garupa</i>	312
<i>Rhamdia</i>	157	<i>ngitum, Cichlasoma</i>	418
<i>Theraps</i>	418	<i>ngitus, Seiranus</i>	312
<i>nicholsi, Irigio</i>	428	<i>Theraps</i>	418
<i>Platyglottus</i>	423	<i>nigrocaneus, Enchelycore</i>	90
<i>nicholsi, Gobius</i>	440	<i>Gymnothorax</i>	90
<i>Rhinogobiops</i>	440	<i>nigrocinctus, Sebastes</i>	370
<i>Nicholsina</i>	431	<i>Sebastichthys</i>	370
<i>nieuhofii, Echeineis</i>	449	<i>Sebastodes</i>	364
<i>niger, Astronesthes</i>	71	<i>nigrofasciata, Hydralgva</i>	176
<i>Bubalichthys</i>	102	<i>Limia</i>	191
<i>Carpodes</i>	102	<i>nigrofasciatus, Archocentrus</i>	420
<i>Centrolophus</i>	267	<i>Hadropterus</i>	283
<i>Chiasmodon</i>	452	<i>Heros</i>	420
<i>Cottus</i>	387	<i>nigrolineatus, Diodon</i>	502
<i>Escolar</i>	261	<i>nigro-maculatus, Cantharus</i>	305
<i>Esox</i>	173	<i>nigromaculatus, Rhypticus</i>	316
<i>Gobius</i>	440	<i>nigromarginatus, Neomuræna</i>	91, 93
<i>Lamprogrammus</i>	484	<i>nigromarginatus, Gymnothorax ocellatus</i>	93
<i>Malacoctetus</i>	71	<i>nigrotæniata, Hybognathus</i>	144
<i>Melanocetus</i>	510	<i>nigrotæniatus, Graodus</i>	132, 133
<i>Myoxocephalus</i>	387	<i>nigrum, Boleosoma</i>	287
<i>Sparus</i>	265	<i>Petromyzon</i>	9
<i>Tautora</i>	424	<i>nimbana, Vinciguerra</i>	73
<i>Thyrstes</i>	261	<i>numbarius, Zalargis</i>	73
<i>nigerrima, Meladerma</i>	281	<i>niphobles, Epinephelus</i>	312
<i>nigra, Astronesthes</i>	71	<i>Sparisoma</i>	432
<i>Balistes</i>	493	<i>niphonius, Præacanthus</i>	323
<i>Dicrolene</i>	483	<i>nipigon, Leucichthys</i>	64
<i>Etheostoma</i>	287	<i>nsorius, Fundulus</i>	175
<i>Gila</i>	119	<i>nitens, Anago</i>	80
<i>Muræna</i>	79	<i>Ophisoma</i>	80
<i>maculata</i>	91, 92	<i>Trichurus</i>	262
<i>Narcine</i>	27	<i>nitida, Hybopsis</i>	135
<i>Perca</i>	267	<i>Julis</i>	429
<i>marina cauda</i>	330	<i>Moniana</i>	135
<i>nigrellus, Hyborhynchus</i>	145	<i>Pomotis</i>	300
<i>nigrescens, Centropomus</i>	306	<i>nitidissima, Julis</i>	429
<i>Coryphæna</i>	317	<i>nitidulum, Myctophum</i>	169
<i>Cycleptus</i>	102	<i>nitidum, Pristipoma (Hæmulops)</i>	333
<i>Dorsanurus</i>	339	<i>Thalassoma</i>	429
<i>Exoglossum</i>	146	<i>nitidus, Alourmus</i>	123
<i>Salmo</i>	60	<i>Balistes</i>	493
<i>Symphurus</i>	230	<i>Brachydeuterus</i>	333
<i>Tigoma</i>	120	<i>Leuciscus</i>	117
<i>nigricans, Bathylaco</i>	164	<i>Pomoxis</i>	304
<i>Camarina</i>	338, 339	<i>Salmo</i>	60
<i>Catostomus</i>	107	<i>Tetragonopterus</i>	96
<i>Centropistes</i>	317	<i>nivea, Erogala</i>	129

List of scientific names—Continued

	Page		Page
niveatus, <i>Epinephelus</i>	311	notatus, <i>Pomotis</i>	302
<i>Pomacentrus</i>	416	<i>Porichthys</i>	487
<i>Serranus</i>	310, 311	<i>Semotilus</i>	177
<i>Stegastes</i>	416	<i>Zygonectes</i>	177
niveiventris, <i>Amurus</i>	153	notemigonoides, <i>Notropis</i>	125
niveus, <i>Hybopsis</i>	129	Notemigonus.....	114
Nivicola.....	291	<i>auratus</i>	114, 115
<i>artensis</i>	292	<i>bosci</i>	115
<i>borealis</i>	291	<i>crystoleucas</i>	115
<i>cragini</i>	292	<i>gardoneus</i>	115
<i>lepidogenys</i>	292	<i>ischanus</i>	115
<i>oboyense</i>	292	nothochir, <i>Ophichthys</i>	87
<i>paget</i>	292	<i>Quassiremus</i>	87
<i>punctulata</i>	292	Notonotus.....	282, 288, 290
<i>tippecanoe</i>	292	<i>camurus</i>	290
<i>virgata</i>	292	<i>cinereus</i>	290
nivipes, <i>Emblemaria</i>	464	<i>inscriptus</i>	289
nobiliana, <i>Torpedo</i>	27	<i>jordani</i>	290
nobilior, <i>Esox</i>	172	<i>maculatus</i>	290
nobilis, <i>Atractoscion</i>	356	<i>rufilineatus</i>	290
<i>Conodon</i>	333	<i>tesselatus</i>	290
<i>Esox</i>	172	<i>thalassinus</i>	289
<i>Gambusia</i>	185	<i>vulneratus</i>	290
<i>Heterandria</i>	185	(<i>Notonotus</i>) <i>jordani</i> , <i>L. leosoma</i>	290
<i>Johnius</i>	356	nothus, <i>Cynoscion</i>	355
<i>Perca</i>	333	<i>Otolithus</i>	355
<i>Polyrhina</i>	234	Notidani.....	11
Nocomis.....	137	Notidanus.....	11
<i>bellicus</i>	137	Notistium.....	263
<i>hyostomus</i>	139	<i>gladius</i>	263
<i>kentuckiensis</i>	137	Notogammus.....	469
<i>micropogon</i>	137	<i>rothrocki</i>	469
<i>milneri</i>	136	Notolepidomyzon.....	104
<i>nebrascensis</i>	137	<i>clarki</i>	104
<i>rubifrons</i>	138	<i>generosus</i>	104
nocomis, <i>Hybopsis</i>	135	<i>plebeus</i>	104
<i>Notropis</i>	135	<i>santa-anae</i>	104
nocturna, <i>Colletia</i>	168	Notorhynchus <i>borealis</i>	11
<i>Echidna</i>	94	Notorynchus.....	11
<i>Rabida</i>	156	<i>maculatus</i>	11
nocturnum, <i>Myctophum</i>	168	<i>platycephalus</i>	11
nocturnus, <i>Noturus</i>	156	Notoscopeus.....	166
<i>Poecilophis</i>	94	<i>castaneus</i>	167
nodulosus, <i>Xenodermichthys</i>	53	<i>margaritiferus</i>	166
Nomalus.....	354	<i>quercinus</i>	166
Nomeida.....	268	Notosema.....	225
Nomeus.....	268	<i>dilecta</i>	225
<i>gronovii</i>	268	<i>dilectum</i>	225
<i>maculosus</i>	268	notospilotus, <i>Arctidius</i>	376
<i>oxyurus</i>	268	<i>Parastolytes</i>	376
normalis, <i>Bassozetus</i>	481	notospilus, <i>Julidio</i>	428
noivegia, <i>Perca</i>	363	<i>Pseudojulis</i>	428
norvegicus, <i>Lepidolepis</i>	205	Nototropis <i>forlonensis</i>	131
norwegianus, <i>Squalus</i>	22	<i>lirus</i>	126
notabilis, <i>Argyreus</i>	141	<i>nazas</i>	126
Notacanthidae.....	75	<i>stilbius</i>	125
Notacanthus.....	75	Notropis.....	123
<i>analis</i>	76	<i>eneolus</i>	143
<i>challengeri</i>	76	<i>alabamae</i>	126
<i>chemnitzii</i>	75	<i>albeolus</i>	136
<i>nasus</i>	75	<i>amabilis</i>	124
<i>phasganorus</i>	76	<i>amcenus</i>	123
<i>rissoanus</i>	76	<i>anogenus</i>	136
<i>rostratus</i>	76	<i>arcasianus</i>	124
<i>spinosus</i>	76	<i>arge</i>	124
Notaius.....	149	<i>athermoides</i>	123
notata, <i>Atherina</i>	247	<i>aztecus</i>	143
<i>Belone</i>	195	<i>boops</i>	134
<i>Cypimella</i>	130	<i>braytoni</i>	135
<i>Dorosoma</i>	46	<i>brimleyi</i>	127
<i>Erogala</i>	130	<i>buchanani</i>	134
<i>Lepomis</i>	298	<i>calientis</i>	134
<i>Megalops</i>	44	<i>cayuga</i>	133
<i>Menidia</i>	247	<i>atrocaudalis</i>	133
<i>Perca</i>	281	<i>chalybeus abbotti</i>	127
<i>Malthodes</i>	507	<i>chamberlaini</i>	132
<i>Strongylura</i>	195	<i>chihuahua</i>	132
notatum, <i>Hamulon</i>	329	<i>cornutus isolepis</i>	128
<i>Pristipoma</i>	332	<i>cummingsi</i>	133
notatus, <i>Balistes</i>	493	<i>dilectus</i>	125
<i>Hyborhynchus</i>	145	<i>fumeus</i>	125
<i>Labrax</i>	307	<i>garmani</i>	131
<i>Mimulus</i>	145	<i>germanus</i>	136
<i>Passer lineis transversis</i>	230	<i>gilberti</i>	134
<i>Pimelodus</i>	152	<i>heterodon richardsoni</i>	136

List of scientific names—Continued

	Page		Page
<i>Notropis heterolepis</i>	134	<i>noxi</i> , <i>Auchenopterus</i>	457
<i>horatii</i>	134	<i>Cremnobates</i>	457
<i>jordani</i>	136	<i>noyesi</i> , <i>Scarus</i>	434
<i>kanawha</i>	134	<i>nubila</i> , <i>Apocope</i>	140
<i>keimi</i>	133	<i>Dionda</i>	144
<i>kendalli</i>	135	<i>nubilus</i> , <i>Alburnops</i>	144
<i>lermae</i>	143	<i>Argyreus</i>	140
<i>leuciodus</i>	124	<i>Lumpenus</i>	471
<i>louisianae</i>	124	<i>nuchalis</i> , <i>Hybognathus</i>	143
<i>lythurus</i>	126	<i>Scarus</i>	434
<i>macdonaldi</i>	127	<i>nuchifilus</i> , <i>Blenni</i> us.....	461
<i>macrolepidotus</i>	125	<i>nuchipinnis</i> , <i>Clinus</i>	459
<i>mearnsi</i>	122	<i>Labrisomus</i>	459
<i>megalops albeolus</i>	129	<i>nudus</i> , <i>Cyclopterus</i>	489
<i>metallicus</i>	124	<i>Gymnachirus</i>	229
<i>micropteryx</i>	124	<i>nugator</i> , <i>Brvostemma</i>	466
<i>muskoka</i>	134	<i>nummifer</i> , <i>Carpiodes</i>	103
<i>nocomis</i>	135	<i>Salmo</i>	61
<i>notatus</i>	124	<i>nuttalli</i> , <i>Exocoetus</i>	201
<i>notemigonoides</i>	125	<i>nuttingi</i> , <i>Chaunax</i>	506
<i>nuv</i>	135	<i>Scorpaena</i>	371
<i>orca</i>	126	<i>nuttingi</i> , <i>Antennarius</i>	506
<i>ortenbergeri</i>	135	<i>nux</i> , <i>Hybopsis</i>	185
<i>ozarcanus</i>	135	<i>Notropis</i>	135
<i>phenacobius</i>	134	<i>nycterinus</i> , <i>Melamphas</i>	233
<i>photogenis</i>	123	<i>Plectromus</i>	233
<i>pilsbryi</i>	124	<i>Nyctimaster</i>	167
<i>reticulatus</i>	133	<i>alatus</i>	167
<i>robustus</i>	132	<i>townsendi</i>	167
<i>roseipinnis</i>	126	<i>Nyctophus</i>	169
<i>rubrifrons</i>	124	<i>rafnesquei</i>	169, 169
<i>sabinae</i>	135		
<i>santamariae</i>	131	<i>oaxaca</i> , <i>Fundulus</i>	174
<i>scepticus</i>	123	<i>Profundulus</i>	174
<i>scopiferus</i>	132	<i>Rhamdia</i>	157
<i>socius</i>	124	<i>obesa</i> , <i>Algansea</i>	122
<i>stilbius</i>	125	<i>obesus</i> , <i>Amiurus</i>	154
<i>stonei</i>	127	<i>Carcharias</i>	14
<i>telescopus</i>	124	<i>Echinorhinus</i>	22
<i>arcuatus</i>	124	<i>Enneacanthus</i>	304
<i>topeka</i>	143	<i>Leuciscus</i>	128
<i>umbra</i> , <i>fasciolaris</i>	126	<i>Pomotis</i>	304
<i>umbrier</i>	131	<i>Siphoteles</i>	122
<i>universitatis</i>	124	<i>Trienodon</i>	14
<i>welaka</i>	124	<i>obeyense</i> , <i>Etheostoma</i>	292
<i>notropoides</i> , <i>Creagrutus</i>	98	<i>Nivicola</i>	292
<i>nottil</i> , <i>Zygonectes</i>	178	<i>obliquatus</i> , <i>Cynoscion</i>	355
<i>Noturus</i>	155	<i>Diabasis</i>	330
<i>elassochir</i>	156	<i>Otolithus</i>	355
<i>eleutherus</i>	156	<i>obliquus</i> , <i>Mugil</i>	349
<i>exilis</i>	156	<i>obliteratus</i> , <i>Alutarius</i>	496
<i>flavus</i>	155	<i>oblonga</i> , <i>Platessa</i>	224
<i>funebis</i>	156	<i>oblongior</i> , <i>Pimelepterus</i>	340
<i>furiosus</i>	155, 156	<i>oblongus</i> , <i>Astatheros</i>	421
<i>gilberti</i>	156	<i>Cyprinus</i>	109
<i>leptacanthus</i>	155	<i>Erimyzon</i>	109
<i>luteus</i>	155	<i>Gobius</i>	448
<i>margunatus</i>	156	<i>Heros</i>	421
<i>miurus</i>	156	<i>Paralichthys</i>	224
<i>nocturnus</i>	156	<i>Pleuronectes</i>	224
<i>occidentalis</i>	155	<i>obolarius</i> , <i>Gasterosteus</i>	237
<i>platycephalus</i>	155	<i>obscura</i> , <i>Liopsetta</i>	228
<i>sialis</i>	155	<i>Sternoptix</i>	73
<i>novae-boracensis</i> , <i>Vomer</i>	275	<i>Sternoptyx</i>	73
<i>Novacula</i>	430	<i>obscuratus</i> , <i>Gymnothorax</i>	93
<i>kallasoma</i>	430	<i>Pomacentrus</i>	414
<i>novacula</i> , <i>Coryphaena</i>	430	<i>obscurus</i> , <i>Carcharias</i>	16
<i>Xyrichtys</i>	430	<i>Centrarchus</i>	298
<i>Novaculichthys</i>	430	<i>Pleuronectes</i>	228
<i>infirmus</i>	430	<i>Pomotis</i>	301
<i>martinicensis</i>	430	<i>Squalus</i>	16
<i>rosipes</i>	430	<i>obtusa</i> , <i>Ophisoma</i>	79
<i>ventralis</i>	430	<i>Raja</i>	26
<i>novae-angliae</i> , <i>Coregonus</i>	65	<i>obtusirostris</i> , <i>Exocoetus</i>	200
<i>novaeorleansis</i> , <i>Anguilla</i>	77	<i>obtusum</i> , <i>Hemulon</i>	332
<i>novae-terrae</i> , <i>Anguilla</i>	77	<i>obtusus</i> , <i>Rhinichthys</i>	140
<i>novaeboracensis</i> , <i>Exocoetus</i>	201	<i>Scarus</i>	434
<i>Gasterosteus</i>	237	<i>Squalus</i>	16
<i>Naucrates</i>	278	<i>obvelatus</i> , <i>Prionodon</i>	16
<i>novemfasciatus</i> , <i>Lutjanus</i>	325	<i>Occa</i>	394
<i>Lutjanus</i>	325	<i>verrucosa</i>	394
<i>novemlineatus</i> , <i>Chasmodes</i>	462	<i>occa</i> , <i>Ceolorhynchus</i>	207
<i>Pholis</i>	462	<i>Macrurus</i>	207
<i>novem-maculatus</i> , <i>Diodon</i>	501	<i>Pristis</i>	23
<i>novemradiata</i> , <i>Agosia</i>	141	<i>Occella</i>	394
		<i>dodecaedron</i>	394

List of scientific names—Continued

	Page		Page
occidentalis, <i>Amia</i>	38	oculocirris, <i>Emblemaria</i>	464
Catostomus.....	106	oculofasciatus, <i>Blepsias</i>	382
Chalmura.....	204	<i>Nautichthys</i>	382
Epignus.....	296	oculo-radiato, <i>Turdus</i>	426
Gasterosteus.....	238, 239	Ocyanthus.....	320
Heterandria.....	189	<i>martineensis</i>	320
Hypopomus.....	101	Ocyurus.....	327
Lepomis.....	301	<i>chrysurus</i>	327
Leucosomus.....	113	<i>lutjanoides</i>	327
Malacocephalus.....	204	<i>riggersmoei</i>	327
Monacanthus.....	494	ocyrus, <i>Centropristes</i>	318
Myripristis.....	235	<i>Pimelepterus</i>	340
Myripristis.....	235	<i>Sectator</i>	340
Nebris.....	354	<i>Serranus</i>	318
Noturus.....	155	Odontaspidae.....	19
Poecilopsus.....	189	Odontaspis.....	19
Roeboides.....	98	<i>littoralis</i>	19
Tetranarce.....	27	Odontognathus.....	45
Torpedo.....	27	<i>mucronata</i>	45
<i>Xenotis</i>	301	<i>panamensis</i>	45
occipitalis, <i>Scorpena</i>	371	(<i>Odontognathus</i>) <i>panamensis</i> , <i>Pristigaster</i>	45
oceanica, <i>Anguilla</i>	79	Odontopyxis.....	396
oceanicus, <i>Blennius</i>	461	<i>frenatus</i>	395
<i>Gobionellus</i>	441	<i>leptorhynchus</i>	395
<i>Gobius</i>	441	<i>trispinosus</i>	396
<i>Rhinoscopelus</i>	169	Odontoscion.....	334
Oceanomyzon.....	8	<i>archidum</i>	345
<i>wilsoni</i>	8	<i>dentex</i>	334
oceanops, <i>Elacatinus</i>	442	<i>xanthops</i>	334
ocella, <i>Eretmichthys</i>	481	oerstedii, <i>Argyreus</i>	275
<i>Rhinichthys</i>	139	<i>Selene</i>	275
<i>Venefica</i>	82	<i>Tetragonopterus</i>	96
ocellaris, <i>Fundulus</i>	176	ogac, <i>Gadus</i>	210
<i>Gobius</i>	442	Ogcocephalidae.....	506
<i>Platessa</i>	224	Ogcocephalus.....	506
ocellata, <i>Bollmannia</i>	443	<i>nasutus</i>	507
<i>Chenopsis</i>	465	<i>radiatus</i>	507
<i>Perca</i>	348	<i>vespertilio</i>	507
<i>Raja</i>	25	Ogilbia.....	479
<i>Scianops</i>	348	<i>cayorum</i>	480
<i>Zenopsis</i>	232	<i>ventralis</i>	480
ocellatus, <i>Acanthocottus</i>	337	<i>verillii</i>	480
<i>Anarrhichthys</i>	472	oglina, <i>Megalops</i>	44
<i>Antennarius</i>	505	oglinum, <i>Opisthonema</i>	44
<i>Chenopsis</i>	465	ohiensis, <i>Accipenser</i>	35
<i>Chatodon</i>	360	<i>Alosa</i>	42
<i>Clinus</i>	459	<i>Cichla</i>	293
<i>Gymnothorax</i>	91, 93	<i>Esox</i>	172
<i>Hippoglossus</i>	218	<i>masquinongy</i>	172
<i>Lophius histrio</i>	505	oianco, <i>Mesopomus</i>	327
<i>Malacotenus</i>	459	okeechobeensis, <i>Ameiurus lacustris</i>	152
<i>Muraenopsis</i>	87	<i>Hauator</i>	152
<i>Neobythites</i>	481	<i>Ictalurus</i>	152
<i>ngromarginatus</i> , <i>Gymnothorax</i>	93	Okkelbergia.....	10
<i>Opichthys</i>	87	<i>lamotteni</i>	10
<i>Platophrys</i>	217	olfax, <i>Mycteroperca</i>	314
<i>Pleuronichthys</i>	225	<i>Serranus</i>	314
<i>Rhombus</i>	217	olfersi, <i>Argyroleucus</i>	74
<i>saxicola</i> , <i>Gymnothorax</i>	93	<i>Cephalopterus</i>	32
<i>Zeus</i>	232	<i>Mobula</i>	32
ocellicauda, <i>Amia</i>	38	<i>Sternoptyx</i>	74
ocellifer, <i>Clinus</i>	458	oldus, <i>Hypomesus</i>	68
ocellane, <i>Chirostoma</i>	251	<i>Salmo</i> (<i>Osmerus</i>).....	68
octodecemspinosus, <i>Acanthocottus</i>	386	oligaspis, <i>Alburnus</i>	125
octodecem-spinosus, <i>Cottus</i>	386	Oligocephalus.....	290
octofasciatus, <i>Astatheros</i>	421	<i>aubeanaubei</i>	291
<i>Heros</i>	421	<i>caeruleus</i>	291
octofilis, <i>Trichidion</i>	256	<i>fricksia</i>	291
Octogrammus.....	373	<i>grahami</i>	291
<i>pallasi</i>	374	<i>hildebrandti</i>	291
octogrammus, <i>Hexagrammos</i>	374	<i>humeralis</i>	293
<i>Labrax</i>	374	<i>iowa</i>	291
octonemus, <i>Polyneumus</i>	256	<i>jessiae</i>	291
oculata, <i>Squatina</i>	22	<i>leonensis</i>	291
oculatus, <i>Bryttus</i>	300	<i>lepidus</i>	291
<i>Etelis</i>	328	<i>luteovinctus</i>	291
<i>Icelinus</i>	376	<i>pulchellus</i>	291
<i>Latebrus</i>	297	Oligocottus.....	390
<i>Lepisosteus</i> (<i>Cylindrosteus</i>).....	37	<i>acuticeps</i>	391
<i>Mauroluchus</i>	73	<i>analis</i>	390
<i>Myrichthys</i>	85	<i>borealis</i>	390
<i>Pseudontophis</i>	85	<i>embryum</i>	391
<i>Serranus</i>	328	<i>globiceps</i>	391
oculeatus, <i>Orthogoriscus</i>	503	<i>maculosus</i>	390
oculeum, <i>Myctophum</i>	169	<i>synderi</i>	391

List of scientific names—Continued

	Page		Page
oligodon, Osmerus.....	68	Ophichthus chamensis.....	88
Polymerus.....	256	evionthas.....	87
oligolepis, Brycon.....	99	gomesi.....	88
oligopeltis, Acipenser (Antaceus).....	34	guttifer.....	87
Oligoplites.....	277	magnoculus.....	88
altus.....	278	ocellatus.....	87
inornatus.....	277	ophis.....	87
mundus.....	278	parilis.....	88
palometa.....	278	retropinnis.....	87
refulgens.....	278	rugifer.....	88
saliens.....	277	triserialis.....	88
saurus.....	277	zophochir.....	88
Oligopodes.....	266	Ophichthyidae.....	84
Oligopodus.....	266	Ophichthys.....	87
Oligopus.....	266	biserialis.....	88
oligosticta, Gambusia.....	186	(Cryptopterus) frontalis.....	85
olisthostoma, Gerres.....	342	guttifer.....	87
olisthostomus, Diapterus.....	342	miurus.....	59
olivacea, Poecilia.....	177	nothochir.....	87
olivaceus, Leucis.....	122	pauciporus.....	88
Siphateles.....	122	pisavarius.....	85
olivaris, Opladelus.....	155	retropinnis.....	87
Silurus.....	155	schneideri.....	88
olmstedii, Boleosoma.....	287	(Sphagebranchus) anguiformis.....	84
brevipinnis, Boleosoma.....	287	xysturus.....	85
Etheostoma.....	287	zopochir.....	88
olomina, Brachyrhaphis.....	187	Ophichthus evionthas.....	87
Priapichthys.....	187	ophidianus, Gempylus.....	262
olrki, Aspidophoroides.....	397	Ophidiæ.....	484
Ulcina.....	397	Ophidion.....	485
omiseomaycus, Percopsis.....	215	beani.....	485
Salmo.....	215	grælii.....	485
ommata, Discopyge.....	27	holbrookii.....	485
Heterandria.....	175	imberbe.....	467
Leptolucania.....	175	Ophidium.....	485
Opistognathus.....	453	atropurpureum.....	469
ommatitis, Leptocottus.....	380	brevibarbe.....	484
ommatitis, Ophichthys.....	453	holbrookii.....	485
ommatitis, Platichthys.....	224	josephi.....	485
Omocheilus.....	86	marginatum.....	485
cruentifer.....	86	mucronatum.....	467
omocyanus, Eleotris.....	430	parrisi.....	478
omostigma, Genypterus.....	455	pellucidum.....	79
Otophidium.....	485	stigma.....	477
Omosudis.....	171	taylori.....	485
Omosudis.....	171	unemak.....	477
lowii.....	171	viridis.....	477
Oncoccephalus.....	576	Ophioblennius.....	404
Oncocottus.....	388	steindachneri.....	404
gilberti.....	388	webbi.....	404
hexacornis.....	388	Ophiodon.....	374
gilberti.....	389	elongatus.....	374
quadricornis.....	388	Ophiodontidae.....	374
Oncorhynchus.....	54	Ophiodontidae.....	374
gorbuscha.....	54	Ophionathus.....	95
kamloops.....	58	ampullaceus.....	95
kenneryi.....	55	ophioneus, Sphagebranchus.....	84
keta.....	54	Ophioscion.....	347
kisutch.....	54	adustus.....	347
nerka.....	55	imiceps.....	348
tschawyttscha.....	55	punctatissimus.....	347
oneida, Catostomus.....	110	sclerus.....	348
Oneirodes.....	509	stimulus.....	348
eschrichtii.....	509	strabo.....	347
onitis, Labrus.....	424	typicus.....	347
Tautoga.....	424	ophioscion, Corvina.....	347
Onos.....	208	ophis, Muraena.....	87
rufus.....	208	Ophichthus.....	87
ontariensis, Leucichthys.....	62	Ophisoma.....	79
Thymallus.....	66	acuta.....	79
Triglopsis.....	389	balearicum.....	80
Onus.....	214	macrurum.....	80
riali.....	214	nitens.....	80
onychus, Cottus.....	386	obtusa.....	79
Oonidus.....	499	prorigerum.....	80
opah, Zeus.....	217	Ophisomus.....	467
opalescens, Lythrulon.....	331	Ophisternon.....	76
opalina, Julis.....	426	Ophisuraphis.....	84
opercularis, Acantholiparis.....	401	Ophisurus.....	85
Myxodagnus.....	455	californiensis.....	88
Polydactylus.....	256	chrysops.....	88
Stolephorus.....	51	crocodilinus.....	89
Trichidion.....	256	gomesi.....	88
Ophichthys regius.....	87	guttatus.....	87
Ophichthys.....	87	interstinctus.....	88
biserialis.....	88	latimaculatus.....	85

List of scientific names—Continued

	Page		Page
Ophisurus longus	85	oregonia, Atherinops	252
ophis	87	Oregonichthys	138
parilis	88	crameri	138
rostellatus	88	cumingi	138
sugillatus	88	oregonius, Coregonus	66
xysturus	85	irillion	66
ophryas, Paralichthys	224	Oreosoma	232
Prionotus	407	atlanticum	232
opthalmicus, Lepomis	299	conferrum	232
Opistharthri	11	orientalis, Anarhichas	472
Opisthistius	339	Pelamys	259
Opisthognathidæ	452	Salmo	55
Opisthognathus	452	ornata, Ama	38
cuvieri	453	Aphoristia	231
lonchurum	453	Codoma	143
lonchurus	453	Embiotoca	411
macrognaethum	452	Hydrargyra	176
macrops	453	Lima	191
maxillosus	453	Raja	26
megastoma	452	ornatum, Camposoma	146
micrognathus	453	ornatus, Achirus	231
ommata	453	Cochlognathus	146
ommatum	453	Esox	173
punctatum	453	Gillellus	455
punctatus	453	Gunnellus	468
rhomaleus	453	Labrus	429
scaphiurus	452	Pholis	468
Opisthonema	44	Tetrodon	500
libertate	44	Ornichthys	407
oglinum	44	orqueta, Chlonoscombrus	276
Opisthopterus	45	orsini, Ozodura	503
dovii	45	ortenburgeri, Hybopsis	135
macrops	45	Notropis	135
opisthophthalmus, Ariosoma	80	Orthocolus	66
opisthotremus, Careproctus	403	Orthodon	112
opistophthalmus, Conger	80	microlepidotus	112
Opladelus	155	Orthogoriscus	503
olivaris	155	orthogramma, Citula	274
Oplopoma	374	Orthonopias	391
pantherina	374	triakis	392
oppositus, Monacanthus	495	orthonotus, Ditrema	411
Opsanus	487	Orthoprists	335
cerapalmus	487	chalcæus	335
hildebrandi	487	chrysopterus	335
jardus	487	duplex	335
tau	487	poeyi	335
Opsoprea	115	reddingi	335
bollmani	115	Orthopsetta	218
Opsopocodus	115	Orthostoechus	331
bollmani	115	maculicauda	331
borealis	115	Orthragoriscus alexandri	503
emilæ	115	analis	503
megalops	115	battaræ	503
osculus	115	blochn	503
oquassa, Salmo	61	elegans	503
Salvelinus	61	ghini	503
cræbe, L. Diodon	502	lunaris	503
Orbidus	498	oculeatus	503
orbignianus, Exocoetus	200	redi	503
orbignyanus, Conger	79	retzii	503
Orbis lævis variegatus	499	rondeletii	503
orbis, Cyclopterus	398	solaris	503
Eumicrotremus	398	Orthragus	503
orbitalis, Macrourus	203	Oryzias	259, 260
Macrurus	208	subulatus	260
orbitalis, Pagellus	336	osbeck, Trachinus	310
orbitalis, Pagellus	128	Osbeckia	495
orca, Notropis	128	scripta	495
Orcella	128	osborni, Leparis	400
nasas	128	Typhlichthys	195
orca	128	oscatans, Sciaenæ	347
oreni, Brama	265	oscatans, Zestus	347
Orcula	120	oscula, Apocope	141
creutli, Phoxinus	120	Netuma	149
Tigoma	120	Sciaenæ	352
Oreynus	260	osculatæ, Labray	307
subulatus	260	osculum, Holocentrum	236
ordinatus, Chirus	374	osculus, Argyreus	141
Epinephelus	311	Arus	149
oreas, Chrosomus	113	Holocentrus	236
oregonensis, Cyprinus (Leuciscus)	114	Opsopocodus	115
Ptychocheilus	114	Osmeridæ	66
Rutilus	122	osmeriformis, Coregonus	62
Siphateles	122	Leucichthys	62

List of scientific names—Continued

	Page		Page
<i>osmerinus</i> , <i>Hybognathus</i>	143	<i>Otrynter</i>	336
<i>Osmerus</i>	67	<i>caprinus</i>	336
<i>abbotti</i>	67	<i>otrynter</i> , <i>Caranx</i>	274
<i>albatrossis</i>	67	<i>otsego</i> , <i>Salmo</i>	64
<i>albidus</i>	163	<i>ouachita</i> , <i>Alvordius</i>	284
<i>attenuatus</i>	67	<i>Etheostoma</i> (<i>Hadropterus</i>).....	284
<i>dentex</i>	67	<i>ouananiche</i> , <i>Salmo</i>	56
<i>elongatus</i>	67	<i>Salmo salar</i>	56
<i>lemniscatus</i>	163	<i>ouatalibi</i> , <i>Serranus</i>	309
<i>microdon</i>	66	<i>ovale</i> , <i>Acutomentum</i>	366
<i>mordax</i>	67	<i>Syacium</i>	213
<i>abbotti</i>	67	<i>ovalis</i> , <i>Hemirhombus</i>	213
<i>spectrum</i>	67	<i>Sebastodes</i>	366
<i>oligodon</i>	68	<i>ovicephalus</i> , <i>Sparus</i>	338
<i>sargenti</i>	67	<i>oviformis</i> , <i>Chaetodon</i>	358
<i>spectrum</i>	67	<i>ovigerum</i> , <i>Bathypasma</i>	401
<i>thaleichthys</i>	67	<i>ovinus</i> , <i>Esox</i>	181
<i>viridescens</i>	67	<i>ovis</i> , <i>Sargus</i>	338
<i>Osphyolax</i>	243	<i>Ovoides</i>	499, 500
<i>pellucidus</i>	244	<i>Ovum</i>	499
<i>osseus</i> , <i>Esox</i>	36	<i>Oxybeles</i>	486
<i>Lepisosteus</i>	36	<i>ocybrachium</i> , <i>Sparisoma</i>	432
<i>Ostarlophysi</i>	95	<i>oxybrachius</i> , <i>Scarus</i>	432
<i>ostentum</i> , <i>Careproctus</i>	402	<i>Oxychaetodon</i>	359
<i>osteoichthys</i> , <i>Echeneis</i>	450	<i>Oxycottus</i>	391
<i>Rhombochirus</i>	450	<i>acuticeps</i>	391
<i>osteosticta</i> , <i>Trygon</i>	29	<i>Oxygeneum</i>	113
<i>Ostichthys</i>	234	<i>pulverulentum</i>	113
<i>trachypomus</i>	234	<i>Oxyjulis</i>	428
<i>Ostorhinchus</i>	295	<i>californicus</i>	428
<i>Ostracidae</i>	496	<i>Oxylabrax</i>	305
<i>Ostracion</i>	496	<i>Oxylebidae</i>	374
<i>bicaudalis</i>	497	<i>Oxylebius</i>	374
<i>boops</i>	503	<i>pictus</i>	374
<i>clippertonense</i>	496	<i>Oxyloricaria</i>	162
<i>concatenatus</i>	496	<i>citirensis</i>	162
<i>gronovii</i>	497	<i>dariensis</i>	162
<i>guineensis</i>	497	<i>panamensis</i>	162
<i>lentiginosum</i>	496	<i>Oryodontichthys</i>	87
<i>lister</i>	497	<i>brachyurus</i>	88
<i>punctatum</i>	496	<i>limbatus</i>	88
<i>punctatus</i>	496	<i>macrurus</i>	88
<i>quadricornis</i>	497	<i>oxyptera</i> , <i>Corvina</i>	320
<i>sexocornutus</i>	497	<i>Oxyrhina</i>	19
<i>trigonus</i>	497	<i>dakayi</i>	19
<i>triquetus</i>	496	<i>gomphodon</i>	19
<i>undulatus</i>	497	<i>oxyrhynchus</i> , <i>Acapenser</i>	15, 16
<i>yalei</i>	497	<i>Carcharias</i>	15, 16
<i>Ostracum</i>	496	<i>Tetrodon</i>	500
<i>expansum</i>	497	<i>oxyrinchus</i> , <i>Isurus</i>	19
<i>tricornis</i>	497	<i>Oxyurus</i>	79
<i>Ostracodermi</i>	496	<i>oxyurus</i> , <i>Lepisosteus</i>	37
<i>otarius</i> , <i>Lepidosteus</i>	37	<i>Nomeus</i>	268
<i>Othonops</i>	447	<i>Oxyzygoneetes</i>	179
<i>eos</i>	46	<i>balboae</i>	179
<i>othonops</i> , <i>Perkinsia</i>	355	<i>dovli</i>	179
<i>othonopterus</i> , <i>Cynoscion</i>	353	<i>Ozadura</i>	503
<i>Otolithidae</i>	356	<i>ozarcanus</i> , <i>Hybopsis</i>	135
<i>Otolithus albus</i>	356	<i>Notropis</i>	135
<i>californiensis</i>	355	<i>Ozodura orsim</i>	503
<i>carolinensis</i>	356	<i>pachycephalus</i> , <i>Fundulus</i>	174
<i>cayennensis</i>	355	<i>Lagocephalus</i>	497
<i>drummondi</i>	355	<i>Tetrodon</i>	497
<i>jamaicensis</i>	355	<i>pachygaster</i> , <i>Chellichthys</i>	499
<i>leirarchus</i>	356	<i>Tetrodon</i> (<i>Chellichthys</i>).....	499
<i>magdalenae</i>	355	<i>Pachygnatha</i>	492
<i>microlepidotus</i>	356	<i>Pachygnathus</i>	492
<i>microps</i>	355	<i>Pachylabrus</i>	411
<i>nebulosus</i>	355	<i>variegatus</i>	411
<i>nothus</i>	355	<i>pachylepis</i> , <i>Atherinichthys</i>	249
<i>obliquatus</i>	355	<i>Thyrinops</i>	249
<i>reticulatus</i>	355	<i>Pachynathus</i>	292
<i>rhomboidalis</i>	354	<i>Pachypops</i>	349
<i>squamipinnis</i>	355	<i>furcata</i>	349
<i>stolzmanni</i>	356	<i>pachypops</i> , <i>Cottus</i>	386
<i>thalassinus</i>	355	<i>Pachyurus</i>	189
<i>toe-roe</i>	354	<i>pacificus</i> , <i>Belone</i>	197
<i>virescens</i>	356	<i>pacifici</i> , <i>Batrachoides</i>	487
<i>Otophidium</i>	485	<i>Batrachus</i>	487
<i>galeoides</i>	485	<i>Conodon</i>	331
<i>omostigma</i>	485	<i>Paraconodon</i>	331
<i>scrippsii</i>	485	<i>pacificum</i> , <i>Diplectrum</i>	318
<i>taylori</i>	485	<i>pacificus</i> , <i>Argyreus</i>	276
<i>welshii</i>	485	<i>Arteidiellus</i>	377
<i>otophorus</i> , <i>Eupomacentrus</i>	414	<i>Bathylagus</i>	69
<i>Pomacentrus</i>	414		

List of scientific names—Continued

	Page		Page
<i>pacificus</i> , <i>Gastrostomus</i>	95	<i>panamensis</i> , <i>Caranx</i>	273
<i>Glyptocephalus</i>	228	<i>Citharichthys</i>	221
<i>Larimus</i>	344	<i>Engraulis</i>	50
<i>Letharchus</i>	85	<i>Evermannia</i>	445
<i>Lobotes</i>	323, 324	<i>Hypheosbrycon</i>	97
<i>Lycodes</i>	474	<i>Loricaria</i>	162
(<i>Lycodopsis</i>).....	474	<i>Menticirrhus</i>	351
<i>Lycodopsis</i>	474	<i>Microeleotris</i>	437
<i>Mesoprion</i>	325	<i>Muraena</i>	91
<i>Microstomus</i>	228	<i>Odontognathus</i>	45
<i>Psenes</i>	269	<i>Oxyloricaria</i>	162
<i>Salmo</i> (<i>Mallotus</i>).....	66	<i>Panamichthys</i>	187
<i>Thaleichthys</i>	66, 67	<i>Parapsettus</i>	359
<i>Thynnus</i>	260	<i>Petrometopon</i>	308
<i>Tylosurus</i>	197	<i>Piabuena</i>	99
<i>Verrugato</i>	324	<i>Priapichthys</i>	187
<i>padangensis</i> , <i>Gobius</i>	440	<i>Pustigaster</i> (<i>Odontognathus</i>).....	45
<i>pæstulus</i> , <i>Citharichthys</i>	218	<i>Rabula</i>	91
<i>pagei</i> , <i>Etheostoma</i>	292	<i>Rhencus</i>	334
<i>Nivicola</i>	292	<i>Serranus</i>	308
<i>Pagellus calamus</i>	336	<i>Solea</i>	230
<i>humilis</i>	337	<i>Tetragonopterus</i>	96
<i>microps</i>	336, 337	<i>Umbina</i>	351
<i>mulneri</i>	337	<i>Panamichthys</i>	187
<i>orbicularis</i>	336	<i>panamensis</i>	187
<i>penna</i>	334, 337	<i>pandionis</i> , <i>Apogon</i>	296
<i>Pagrus</i>	337	<i>Emblematis</i>	464
<i>pagrus</i>	337	<i>Xystramia</i>	296
<i>vulgaris</i>	337	<i>pandora</i> , <i>Clinostomus</i>	120
<i>pagrus</i> , <i>Pagrus</i>	337	<i>panduratus</i> , <i>Leiobatus</i>	23
<i>Sparus</i>	337	<i>pannosa</i> , <i>Scorpena</i>	371
<i>Pagellus caninus</i>	336	<i>pantherina</i> , <i>Muraenophis</i>	93
<i>Palamita</i>	259	<i>Oplopoma</i>	374
<i>palearis</i> , <i>Lycodes</i>	475	<i>pantherinus</i> , <i>Cestracion</i>	10
<i>Palmurichthys</i>	267	<i>Ichthyophis</i>	94
<i>perciiformis</i>	268	<i>Pseudaroides</i>	159
<i>Palinurus</i>	267	<i>Pantosteus</i>	104
<i>palladius</i> , <i>Labrus</i>	279, 302	<i>arizone</i>	104
<i>pallasi</i> , <i>Molacanthus</i>	503	<i>columbianus</i>	105
<i>Octogrammus</i>	373, 374	<i>delphinus</i>	104
<i>Pallasia</i>	503	<i>jordani</i>	104
<i>Pallasia</i>	503	<i>lahontan</i>	105
<i>pallasi</i>	503	<i>platyrhynchus</i>	104
<i>pallasi</i> , <i>Clupea</i>	41	<i>santa-anæ</i>	104
<i>Pleuronectes</i>	226	<i>virescens</i>	105
<i>Pallasina</i>	394	<i>pantostigmus</i> , <i>Myrichthys</i>	85
<i>aix</i>	394	<i>panuco</i> , <i>Gambusia</i>	186
<i>barbata</i>	394	<i>papalis</i> , <i>Dionda</i>	144
<i>pallida</i> , <i>Aldrovandia</i>	75	<i>papillifer</i> , <i>Caulistius</i>	488
<i>Echeneis</i>	440	<i>Gobiesox</i>	488
<i>Lepomis</i>	298	<i>papillifera</i> , <i>Forbesella</i>	164
<i>Morone</i>	307	<i>papilliferus</i> , <i>Chologaster</i>	194
<i>pallidus</i> , <i>Bodianus</i>	345	<i>papilio</i> , <i>Melletes</i>	380
<i>Catostomus</i>	105	<i>papillosum</i> , <i>Moxostoma</i>	109
<i>Fundulus</i>	176	<i>Sycaum</i>	218
<i>Labrus</i>	301	<i>papillosus</i> , <i>Barbulifer</i>	446
<i>Lepomis</i>	299	<i>Pleuronectes</i>	218
<i>Leucosomus</i>	117	<i>Ptychostomus</i>	109
<i>Pimelodus</i>	152	<i>Parachromis</i>	418
<i>Platygnathus</i>	136	<i>gulosus</i>	418
<i>Pomotis</i>	301, 302	<i>Paraclinus</i>	457
<i>Salmo</i>	59	<i>chaperi</i>	457
<i>Xystroplites</i>	302	<i>Paraconodon</i>	331
<i>palmipes</i> , <i>Trigla</i>	407	<i>cesius</i>	331
<i>paloma</i> , <i>Trachinotus</i>	277	<i>dovi</i>	332
<i>Palometa</i>	266	<i>pacifici</i>	331
<i>media</i>	267	<i>Paradiodon</i>	501
<i>palometa</i>	267	<i>paradoxa</i> , <i>Garmannia</i>	445
<i>simillima</i>	266	<i>paradoxus</i> , <i>Gobius</i>	445
<i>snyderi</i>	267	<i>Psychrolutes</i>	392
<i>palometa</i> , <i>Chorinemus</i>	278	<i>Stylophthalmus</i>	71
<i>Oligoplites</i>	278	<i>Parafundulus</i>	182
<i>Palometa</i>	267	<i>Paragambusia</i>	184
<i>Stromateus</i>	266, 267	<i>Paragonus</i>	394
<i>Trachinotus</i>	276	<i>Parahemiodon</i>	161
<i>palustris</i> , <i>Poecilichthys</i>	294	<i>typus</i>	161
<i>Pammelas</i>	267	<i>Parajulis</i>	426
<i>pammelas</i> , <i>Melanostigma</i>	478	<i>Paralabrax</i>	317
<i>pampanus</i> , <i>Trachinotus</i>	276	<i>clathratus</i>	317
<i>panamense</i> , <i>Pristipoma</i>	333, 334	<i>humeralis</i>	317
<i>panamensis</i> , <i>Elmichthys</i>	148	<i>maculato-fasciatus</i>	317
<i>Anchoviella</i>	50	<i>nebulifer</i>	317
<i>Antherinella</i>	249	(<i>Paralabrax</i>) <i>dewegeri</i> , <i>Serranus</i>	319
<i>Azevia</i>	221	<i>Paralepis</i>	165
<i>Bagre</i>	148	<i>barracudina</i>	165
<i>Bascanichthys</i>	86	<i>borealis</i>	165

List of scientific names—Continued

	Page		Page
<i>Paralepis coregonoides</i>	165	<i>Parartedius</i>	377
<i>coruscans</i>	165	<i>bankinsoni</i>	377
<i>intermedius</i>	165	<i>Parascaphirhynchus</i>	35
<i>paralepis</i> , <i>Coregonus</i>	165	<i>albus</i>	35
<i>Paralichthys</i>	222	<i>Parascorpæna</i>	370
<i>Paralichthys</i>	223	<i>Paraserranus</i>	318
<i>estuarius</i>	224	<i>parasiticus</i> , <i>Simenichelys</i>	78
<i>albigutta</i>	224	<i>Parasphyrenops</i>	267
<i>albiguttus</i>	224	<i>atrimanus</i>	267
<i>brasilensis</i>	224	<i>Parastrolytes</i>	376
<i>californicus</i>	223	<i>notospilotus</i>	376
<i>dentatus</i>	224	<i>Paratractus</i>	272
<i>lethostigmus</i>	223	<i>caballus</i>	272
<i>maculosus</i>	223	<i>crysos</i>	272
<i>magdalene</i>	223	<i>Parauchenipterus</i>	159
<i>oblongus</i>	224	<i>pardale</i> , <i>Lepophidium</i>	485
<i>ommatum</i>	224	<i>Leptophidium</i>	485
<i>ophyrus</i>	224	<i>pardalis</i> , <i>Monacanthus</i>	493
<i>sinaloe</i>	224	<i>Mycteroperca</i>	315
<i>squamulentus</i>	224	<i>Xystroperca</i>	315
<i>woolmani</i>	224	<i>pardus</i> , <i>Batrachus tau</i>	487
<i>Paraliparis</i>	404	<i>Opsanus</i>	487
<i>albescens</i>	404	<i>Parephippus</i>	358
<i>angustifrons</i>	405	<i>Parepinephelus</i>	315
<i>attenuatus</i>	404	<i>acutirostris</i>	315
<i>bathybi</i>	404	<i>Pareques</i>	353
<i>caudatus</i>	404	<i>acuminatus</i>	353
<i>cephalus</i>	404	<i>pulcher</i>	353
<i>copei</i>	405	<i>punctatus</i>	353
<i>dactylosus</i>	404	<i>umbrosus</i>	353
<i>deani</i>	404	<i>viola</i>	353
<i>fimbriatus</i>	404	<i>Parexocetus</i>	200
<i>garmani</i>	405	<i>mesogaster</i>	200
<i>grandiceps</i>	404	<i>pargus</i> , <i>Mesoprion</i>	325
<i>holomelas</i>	404	<i>Paricelinus</i>	375
<i>latifrons</i>	405	<i>hopliticus</i>	375
<i>liparinus</i>	405	<i>thoburni</i>	375
<i>membranaceus</i>	404	<i>parietalis</i> , <i>Colchus</i>	145
<i>mento</i>	404	<i>parilis</i> , <i>Ophichthus</i>	88
<i>rosaceus</i>	404	<i>Ophichthus</i>	88
<i>ulochir</i>	405	<i>parismina</i> , <i>Brachyrhaphis</i>	187
<i>parallelus</i> , <i>Centropomus</i>	306	<i>Gambusia</i>	187
<i>Embryx</i>	474	<i>parket</i> , <i>Salmo</i>	61
<i>Paralongchurus</i>	351	<i>parma</i> , <i>Heros</i>	419
<i>goodei</i>	351	<i>Parapetenia</i>	419
<i>petersi</i>	351	<i>parmatus</i> , <i>Setarches</i>	372
<i>Paramia</i>	206	<i>parmifera</i> , <i>Raja</i>	26
<i>paranasimos</i> , <i>Acijensei</i> (Huso).....	35	<i>Parodon dariensis</i>	100
<i>Paraneotriplus</i>	423	<i>piracicaba</i>	100
<i>bulleri</i>	423	<i>Parophrys</i>	225
<i>Paranotropsis</i>	125	<i>ayresi</i>	225
<i>ariommus</i>	125	<i>hubbardi</i>	226
<i>jejunos</i>	125	<i>ischyrus</i>	226
<i>megalops</i>	125	<i>vetulus</i>	225, 226
<i>scabriceps</i>	125	<i>parovanus</i> , <i>Myloleucus</i>	121
<i>Paranthias</i>	320	<i>parra</i> , <i>Diabasis</i>	329
<i>turcifer</i>	320	<i>Hæmulon</i>	329
<i>Parapetenia</i>	418	<i>parra</i> , <i>Albula</i>	40
<i>adspersa</i>	418	<i>Brama</i>	426
<i>alfari</i>	419	<i>Chætodon</i>	361
<i>bartoni</i>	419	<i>Clepticus</i>	426
<i>beam</i>	419	<i>Exocetus</i>	202
<i>centralis</i>	419	<i>parraianus</i> , <i>Monacanthus</i>	494
<i>dovii</i>	418	<i>parrii</i> , <i>Lycocara</i>	478
<i>friedrichstahl</i>	418	<i>Ophidium</i>	478
<i>islana</i>	418	<i>parryi</i> , <i>Rhamdia</i>	158
<i>labridens</i>	419	<i>partitus</i> , <i>Eupomacentrus</i>	414
<i>managuense</i>	418	<i>Pomacentrus</i>	414
<i>mento</i>	419	<i>paru</i> , <i>Chætodon</i>	360
<i>motaguensis</i>	418	<i>Pepilus</i>	266
<i>multifasciata</i>	418	<i>Pomacanthus</i>	360
<i>nigricans</i>	419	<i>Stromateus</i>	266
<i>parma</i>	419	<i>Parupeneus</i>	343
<i>salvini</i>	418	<i>parva</i> , <i>Anchovia</i>	49
<i>steindachneri</i>	419	<i>Anchoviella</i>	49
<i>tenuis</i>	419	<i>Echenis</i>	449
<i>tetracantha</i>	419	<i>Lucania</i>	175
<i>cinta</i>	419	<i>Pleuronectes satis</i>	217
<i>grisea</i>	419	<i>parviceps</i> , <i>Lycodapus</i>	478
<i>lata</i>	419	<i>parvidens</i> , <i>Moxostoma</i>	110
<i>toralbasii</i>	419	<i>parvipinna</i> , <i>Etheostoma</i>	262
<i>tr. maculata</i>	420	<i>parvipinnus</i> , <i>Ancylodon</i>	353
<i>urophthalma</i>	418	<i>brevis</i> , <i>Fundulus</i>	176
<i>Parapsettus</i>	359	<i>Cynoscion</i>	356
<i>panamensis</i>	359	<i>Dicorotus</i>	261

List of scientific names—Continued

	Page		Page
parvipinnis, Erisicion	356	pelamides, Scomber	238
Fundulus	176	pelamidis, Rhynchichthys	235
Hesperoleucus	122	pelamis, Katsuwonus	238
Isopisthus	353	Scomber	238
Ptax	261	pelamitus, Scomber	239
parvula, Clupea	41, 42	Pelamys	239
parvulus, Myoxocephalus	387	chilensis	239
parvus, Cottopsis	383	lineolata	239
Cyprinodon	175	orientalis	239
Gobius	439	pelegrinus, Squalus	20
Rhinogobius	439	Pellona	44
Sebastes ruber	368	bleekeriana	45
Upeneus	343	orbignyana	44
passan, Apterionotus	101	pellucida, Salmoperca	214, 215
Passer lineis transversis notatus	230	pellucidum, Etheostoma	288
passer, Centropyge	361	Ophidium	79
Holacanthus	361	pellucidus, Acus	244
Pastinaca	28	Osfhyolax	243, 244
hastata	28	Pleurolepis	268
marina	29	Psenes	269
olivacea	28	Thyrus	220
pastinaca, Raja	28	Vigil	268
Pastinachus	28	Pelodichthys	155
centrourus	28	Pelona	217
schmarda	28	heckeli	217
torrei	28	pelta, Chelmo	359
patagonica, Maynea	476	peltastes, Lepomis	300
patana, Macrodon	99	peltatum, Etheostoma	283
patao, Gerres	342	peltatus, Alvordius	283
patatus, Julis	426	Turdus cinereus	341
patris, Acanthocottus	389	Pempheridae	323
patronus, Brevoortia	44	Pempheris	323
patruelis, Gambusia	185	mexicana	323
Heterandria	184	mexicanus	323
pattersoni, Troglolani	155	mulleri	323
patzcuaro, Chirostoma	250	poeyi	323
paucidens, Leurynnis	474	schomburgkii	323
Salmo	55	touea	323
pauciporus, Ophichthys	88	peninsula, Bascanichthys	85
pauciradiatus, Callionymus	450	Callecheilus	86
Pseudoxiphophorus	188	Chirostoma	247
paucispinis, Sebastes	364	Menidia	247
Sebastodes	364	peninsula atrimentis, Menidia	247
pavo, Xyrichthys	430	penna, Calamus	337
pavonaceus, Herichthys	422	Pagellus	337
Heros	422	pennanti, Argentina	73
pavonina, Lima	191	Maurolicus	73
pavoninus, Adonis	460	Squalus	19
Cyclopterus	397	pennatula, Calamus	336
Inustus	430	Penopus	482
Xyrichthys	430	macdonaldi	482, 483
pavilloides, Lycodes	476	pensacola, Harengula	43
paxillus, Lycenchelys	476	penshinensis, Salmo	56
Lycodes	476	pentacantha, Asterolthea	396
peckianus, Syngnathus	242	pentacanthus, Bodianus	235
peckii, Syngnathus	242	Centrarchus	303
pectinatus, Centropomus	306	Labrus	424
Pristis	23	Xenochirus	396
pectinifer, Chnus	459	Pentanemus	256
Leucidius	123	quinquarius	256
pectoralis, Albafrossia	204	pepinus, Luciopeca	282
Bathyonus	482	pepo, Searus	433
Bathypterois	166	Peprilus	266
Dactyloscopus	454	pari	266
Dallia	172	snyderi	267
Harpe	425	peracutatus, Pleuronectes	226
Hemigaleus	14	Perca	281
Macrourus	204	aberrans	316
(Malacocephalus)	204	acuta	281
Nematistius	280	albumus	350
Nematonus	482	americana	281, 307
Prionotus	407	apoda	325
miles	407	ascensionis	235
Pedalion	503	atraria	317
pedallota, Bonapartia	72	chrysops	307
pedallotus, Zaphotias	72	chrysoptera	335
pedicellaris, Sciaenops	483	chrysopterus	335
Pediculati	504	flavescens	281
pedimacula, Centropomus	306	flavopurpurea	310
Pegedictis	383, 385	fluviatilis	281
Ictalops	383, 385	gibbosa	302
Peixe agulha	195	formosa	318, 330
pelagicus, Callionymus	400	furcata	349
Nectoliparis	402	furva	317
Scomber	264	gibbosa	302, 329
Syngnathus	240-242	gigas	310

List of scientific names—Continued

	Page		Page
<i>Perca gracilis</i>	281	<i>Peristedion brevirostre</i>	405
<i>granulata</i>	281	<i>crustosum</i>	406
<i>guttata</i>	308, 311	<i>gracile</i>	405
<i>guttatus</i>	308	<i>longispathum</i>	405
<i>immaculata</i>	307	<i>malarmat</i>	405
<i>juba</i>	332	<i>miconemus</i>	406
<i>lanceolata</i>	352	<i>miniatum</i>	405
<i>lophar</i>	280	<i>platycephalum</i>	406
<i>lobbina</i>	305	<i>Peristedium barbiger</i>	406
<i>maculata</i>	310	<i>crustosum</i>	406
<i>marina</i>	363	<i>longispatha</i>	405
<i>capite striato</i>	330	<i>platycephalum</i>	406
<i>cauda nigra</i>	330	<i>Peristethus</i>	405
<i>gibbosa cinerea</i>	329	<i>brevirostre</i>	405
<i>rubra</i>	236	<i>peristethus, Podotheus</i>	394
<i>sectatrix</i>	340	<i>Perkinsia</i>	46
<i>melanura</i>	330	<i>othonops</i>	46
<i>minima</i>	287	<i>periatum, Holocentrum</i>	236
<i>mittelli</i>	307	<i>perlongus, Leptoconger</i>	82
<i>alternata</i>	307	<i>Neoconger</i>	82
<i>interrupta</i>	307	<i>pernigae, Culius</i>	437
<i>mucronata</i>	308	<i>Eleo-lis</i>	437
<i>nigra</i>	267	<i>Peropus</i>	382
<i>nobilis</i>	333	<i>perplexus, Coltic</i>	384
<i>norwegica</i>	363	<i>perico, Pseudo-</i>	435
<i>notata</i>	281	<i>Scarus</i>	435
<i>ocellata</i>	348	<i>Petro colado</i>	425
<i>(Perca) nebulosa</i>	283	<i>peti oteti, Pristis</i>	23
<i>philadelphica</i>	318	<i>personata, Eviota</i>	438
<i>punctata</i>	309, 345	<i>personatus, Ammodyte</i>	451
<i>pusilla</i>	295	<i>perspicabilis, Embiotoca</i>	411
<i>rhomboidalis</i>	338	<i>perspicillans, Tetradon</i>	500
<i>robusta</i>	311	<i>peristethus, Peristethus</i>	474
<i>rufa</i>	236	<i>peristethus, Peristethus</i>	145
<i>salmones</i>	282	<i>peristethus, Peristethus</i>	47
<i>saltatrix</i>	280	<i>peristethus, Peristethus</i>	47
<i>saxatilis</i>	307	<i>peristethus, Peristethus</i>	322
<i>sectatrix</i>	340	<i>Hemianthias</i>	322
<i>septentrionalis</i>	307	<i>perugae, Lima</i>	191
<i>serrato-granulata</i>	281	<i>Platycephalus</i>	191
<i>stellio</i>	310	<i>peruvianus, Diapetus</i>	342
<i>striata</i>	331	<i>Galeichthys</i>	149
<i>tota</i>	310	<i>Genes</i>	342
<i>trifurca</i>	318	<i>Pescador</i>	505
<i>undulata</i>	349	<i>Petalosoma</i>	193
<i>unicolor</i>	321	<i>cultratum</i>	193
<i>unimaculata</i>	338	<i>Petalurichthys</i>	193
<i>varia</i>	317	<i>petenensis, Astyanax</i>	97
<i>variabilis</i>	365	<i>Meletta</i>	47
<i>venenosa</i>	313	<i>Mollinesia</i>	192
<i>ventre luteo</i>	302	<i>Pimelodus</i>	158
<i>vitrea</i>	282	<i>Pocilia</i>	192
<i>percellens, Raja</i>	23	<i>Rhamdia</i>	158
<i>Rhinobatos</i>	23	<i>Signalosa</i>	47
<i>Percesoces</i>	246	<i>Tetragonopterus</i>	97
<i>Percidae</i>	281, 282	<i>Petenia</i>	423
<i>perciiformis, Coryphaena</i>	267, 268	<i>splendida</i>	423
<i>Palinurichthys</i>	268	<i>petersi, Centropholis</i>	266
<i>Perca</i>	282	<i>Paralichthys</i>	351
<i>bimaculata</i>	283	<i>petimba, Fistularia</i>	245
<i>caprodes</i>	283	<i>Petrometopon</i>	308
<i>carbonaria</i>	283	<i>cruentatus</i>	308
<i>manitou</i>	283	<i>coronatus</i>	308
<i>nebulosa</i>	282	<i>panamensis</i>	308
<i>rex</i>	283	<i>Petromyzon</i>	8
<i>zebra</i>	283	<i>americanus</i>	8
<i>(Perca) nebulosa, Perca</i>	283	<i>appendix</i>	9
<i>percobromus, Alburnellus</i>	124	<i>argenteus</i>	9
<i>Percoides, Agonostoma</i>	254	<i>(Artedi)</i>	8
<i>Dajaus</i>	254	<i>astori</i>	10
<i>Percomorphi</i>	246	<i>ayreui</i>	10
<i>Percopsidae</i>	214	<i>(Bathymyzon) bailloni</i>	8
<i>Percopsis</i>	214	<i>bdellum</i>	9
<i>guttatus</i>	214, 215	<i>borealis</i>	10
<i>hammondi</i>	215	<i>culatus</i>	10
<i>omismomaycus</i>	215	<i>epithexodon</i>	10
<i>peragrinus, Leptopus</i>	215	<i>fluvialis</i>	10
<i>perezi, Platypodon</i>	16	<i>lamoteni</i>	10
<i>perfasciata, Anchoviella</i>	47	<i>lividus</i>	10
<i>perfasciatus, Engraulis</i>	47	<i>marinus</i>	8
<i>periscopus, Gadus</i>	209	<i>dorsatus</i>	9
<i>Perissias</i>	220	<i>nigricans</i>	8
<i>taniotus</i>	220	<i>nigum</i>	9
<i>Peristedudae</i>	405	<i>plumbeus</i>	10
<i>Peristedion</i>	405	<i>tridentatus</i>	9, 10
<i>barbiger</i>	406	<i>Petromyzonidae</i>	8

List of scientific names—Continued

	Page		Page
Petronasom.....	433	Photogenis, scabriceps.....	125
petiosus, Brycon.....	98	spilopterus.....	129
Mugil.....	253	stigmaturus.....	129
Serranus.....	313	telescopus.....	124
petus, Acanthocybium.....	256	photogenis, Notropis.....	123, 128
Cybium.....	256	Squalus.....	128
pfeifferi, Muraena.....	93	Photonectes.....	70
Phille.....	121	gracilis.....	70
neogaea.....	121	Phoxaegrea.....	247
phænna, Hybopsis.....	132	dayi.....	247
Phænodon.....	71	Phoxinus clevelandi.....	118
ringens.....	71	flammeus.....	121
Phæthomichthys.....	264	milnerianus.....	121
tuberculatus.....	264	neogeus.....	121
phaeton, Pistigaster.....	45	occuti.....	120
Serranus.....	309	phoxocephalum, Cynoscion.....	356
phalæna, Umbriana.....	350	Etheostoma.....	284
Phalangistes.....	395	phoxocephalus, Alvordius.....	284
loricatus.....	394	Atractoscion.....	356
phaleratus, Esox.....	173	phrynoides, Eumicrotornus.....	398
Phallichthys.....	190	Phtheinchthys.....	449
amatus.....	190	tropicus.....	449
pittieri.....	190	Phycis.....	212, 213
Phanerodon.....	411	chesteri.....	213
atripes.....	411	curatus.....	212
finlayi.....	411	dekayi.....	213
P.....	424	earli.....	212
P.....	76	flouidanus.....	213
phasma, Caelepiocetus.....	402	marginatus.....	212, 213
Phenacobius.....	142	rostratus.....	213
centostomus.....	142	tinca.....	212
mhiabilis.....	142	yarrelli.....	213
scopifer.....	142	phycis, Lota.....	211
teretulus.....	142	Phycocetes.....	478
liosternus.....	142	latitans.....	478
uranops.....	112	suspectus.....	478
phenacobius, Hybopsis.....	134	Physiculus.....	211
Notropis.....	134	dalwigkii.....	211
Phenagoniutes.....	95	fulvus.....	212
macrolepis.....	95	kaupii.....	211
wilsoni.....	95	longipes.....	212
phenax, Apomotis.....	209	nematopus.....	212
Myctelopoeca.....	315	rastralliger.....	212
falcat.....	315	physignathus, Ceatichthys.....	136
philadelphica, Pecca.....	318	Platygobio.....	136
philadelphicus, Triloburus.....	318	Physostomi.....	39
philomps, Cottus.....	384	Phytichthys.....	469
Philypnus.....	435	chuus.....	469
laterilis.....	436	versicolor.....	469
philebotomus, Acanthurus.....	362	Piabucina.....	99
philegethontis, Clinostomus.....	121	erythrinoides.....	99
lotichthys.....	121	festæ.....	99
philegethontis, Clinostomus.....	121	panamensis.....	99
phlox, Boleosoma.....	287	picarti, Hemirhamphus.....	198
Ulocentra.....	287	piceus, Balistes.....	493
Phobator.....	389	Melichthys.....	493
phoebe, Prionodes.....	319	pichardi, Joturus.....	254
Serranus.....	319	Picorellus.....	172
Pholidæ.....	466	picta, Muraena.....	93
Pholidichthys.....	465	Scorpena.....	370
anguliformis.....	465	Sidera.....	93
leucotænia.....	465	picturata, Alutera.....	495
Pholis.....	461, 467	Gambusia.....	185
carolinus.....	461	picturatus, Gymnothorax.....	91
fasciatus.....	468	Syngnathus.....	241
gilli.....	468	pictus, Chaunax.....	506
gunnellus.....	467	Chuonectes.....	504
novemlineatus.....	462	Chirus.....	374
ornatus.....	468	Eleuthis.....	437
quadrifasciatus.....	462	Gymnelus.....	477
subbifurcatus.....	470	Histrio.....	504
pholis, Biennius.....	461	Iridio.....	427
Photocorynus.....	509	Julis.....	427
spinceps.....	509	Oxylebius.....	374
Photogenis.....	123	Sciades.....	148
arioninus.....	125	Torpedo.....	27
cæruleus.....	130	Urocentrus.....	467
callistus.....	130	plicuda, Sphyræna sphyræna var.....	255
eurystomus.....	130	plicudilla, Sphyræna.....	255
grandipinnis.....	129	pidiense, Moxostoma.....	110
leucodus.....	124	pidiensis, Ptychostomus.....	110
leucops.....	123	piersoni, Hypsoblenius.....	462
engraulinus.....	130	piger, Symphurus.....	230
leucopus.....	130	pigmentarius, Apogon.....	295
piptolepis.....	134	Monoprius.....	295
pyrrhomelas.....	129	pigra, Aporistia.....	230

List of scientific names—Continued

	Page		Page
<i>pilchardus</i> , Clupea.....	43	<i>Pimelodus</i> , marmoratus.....	154
<i>puleatus</i> , Prionotus.....	407	<i>megalops</i>	152
<i>Pileoma</i>	282	<i>micropterus</i>	157
<i>carbonaria</i>	283	<i>motaguensis</i>	157
<i>cyematogramma</i>	286	<i>natalis</i>	153
<i>semifasciata</i>	282	<i>nebulosus</i>	154, 155
<i>semifasciatum</i>	283	<i>nicaraguensis</i>	157
<i>zebra</i>	283	<i>nigricans</i>	152
<i>pilicornis</i> , Blennius.....	461	<i>notatus</i>	152
<i>Pilodictis</i>	155	<i>pallidus</i>	152
<i>lunosa</i>	155	<i>petenensis</i>	155
<i>pilosa</i> , Solea.....	230	<i>platycephalus</i>	154
<i>pilosus</i> , Diodon.....	501	<i>polycaulus</i>	158
<i>Trichodiodon</i>	501	<i>pullus</i>	154
<i>pilsbryi</i> , Notropis.....	124	<i>punctulatus</i>	155
<i>Pimelepterus</i>	339	<i>quelon</i>	156
<i>analogus</i>	339	<i>rigidus</i>	159
<i>bosci</i>	339	<i>rogersi</i>	158
<i>sicula</i>	340	<i>salvini</i>	157
<i>bosquii</i>	340	<i>spixi</i>	150
<i>cornubiensis</i>	268	<i>viscosus</i>	155
<i>elegans</i>	339	<i>vulgaris</i>	153
<i>flavo-lineatus</i>	339	<i>vulpeculus</i>	154
<i>incisor</i>	339	<i>vulpes</i>	152
<i>lutescens</i>	340	<i>wagneri</i>	157
<i>oblongior</i>	340	<i>Pimelometopon</i>	426
<i>ocyrus</i>	340	<i>pulcher</i>	426
<i>sandvicensis</i>	339	<i>Pimephales</i>	145
<i>Pimelode insignis</i>	155	<i>agassizii</i>	145
<i>livree</i>	156	<i>anuli</i>	146
<i>Pimelodella</i>	158	<i>confertus</i>	145
<i>chagresi</i>	158	<i>fasciatus</i>	145
<i>Pimelodidae</i>	156	<i>maculosus</i>	145
<i>Pimelodon</i>	155	<i>milesi</i>	145
<i>insignarius</i>	155, 156	<i>promelas</i>	145
<i>Pimelodus</i>	153	<i>pingeli</i> , Triglops.....	379
<i>aeneus</i>	155	<i>pinguis</i> , Pleuronectes.....	221
<i>affinis</i>	152	<i>pinima</i> , Acara.....	332
<i>ailurus</i>	153	<i>pinnata</i> , Murena.....	78
<i>albidus</i>	150, 153	<i>pinnatus</i> , Eretmichthys.....	481
<i>antoniensis</i>	153	<i>Synaphobranchus</i>	78
<i>argentinus</i>	152	<i>piniger</i> , Brysseteres.....	483
<i>argyrus</i>	152	<i>Enneacanthus</i>	304
<i>ascita</i>	158	<i>Gobiesox</i>	488
<i>atrarius</i>	154	<i>Rosicola</i>	365
<i>baronis-mulleri</i>	157	<i>Sebastosomus</i>	365
<i>borealis</i>	152	<i>pinnimaculatus</i> , Ailurichthys.....	148
<i>brachycephalus</i>	157	<i>Bagre</i>	148
<i>brachypterus</i>	157	<i>pinnivaria</i> , Hypoplectrus.....	321
<i>catulus</i>	154	<i>pinnivarius</i> , Hypoplectrus unicolor.....	321
<i>caudafurcatus</i>	152	<i>pinnulata</i> , Seriola.....	279
<i>chagresi</i>	158	<i>pinta</i> , Murena.....	90
<i>clarias</i>	159	<i>pintita</i> , Murena.....	92
<i>coenosus</i>	153	<i>piptolepis</i> , Hybopsis.....	134
<i>confinis</i>	154	<i>Photogenus</i>	134
<i>conirostris</i>	156	<i>Piquitinga</i>	48
<i>eristatus</i>	158	<i>piquotianus</i> , Lepidosteus.....	37
<i>cupresoides</i>	153	<i>piquotil</i> , Amia.....	38
<i>cupreus</i>	153	<i>piraaca</i> , Monacanthus.....	494
<i>cyclopum</i>	159	<i>piracicabae</i> , Parodon.....	100
<i>dekayi</i>	153	<i>Piramutana blochii</i>	159
<i>erythroptera</i>	152	<i>macrospila</i>	159
<i>felinus</i>	153	<i>pirapeda</i> , Dactylopterus.....	409
<i>felis</i>	154	<i>Piraliapia</i>	313
<i>furcatus</i>	152	<i>pirense</i> , Cycloptum.....	160
<i>furcifer</i>	152	<i>pirensis</i> , Astroblepus.....	160
<i>godmani</i>	157	<i>pisavarus</i> , Ophichthys.....	85
<i>gracilis</i>	152	<i>pisicatorius</i> , Lophius.....	504
<i>graciosus</i>	152	<i>piscatrix</i> , Pseudorhamdia.....	159
<i>guatemalensis</i>	157	<i>Pisces</i>	33
<i>hammondi</i>	152	<i>paciculus</i> , Esox.....	176
<i>houghi</i>	152	<i>Piscis viridis bahamensis</i>	432
<i>hoyi</i>	154	<i>piscis bahamensis</i> , Unicornis.....	495
<i>hypselurus</i>	157	<i>pusculentus</i> , Esox.....	176
<i>insigne</i>	155	<i>Pisodonophis</i>	86
<i>insignis</i>	156	<i>cancrivorus</i>	86
<i>lateralis</i>	152	<i>daspilotus</i>	86
<i>laticaudus</i>	157	<i>guttilatus</i>	85
<i>leucoptera</i>	152	<i>Pisodontophis</i>	86
<i>lupus</i>	153	<i>guttatus</i> , Ophis.....	85
<i>lynx</i>	153	<i>pisonis</i> , Eleotris.....	436
<i>macronema</i>	159	<i>Gobius</i>	436
<i>maculatus</i>	151, 152, 153, 159	<i>Pisodonophis</i>	86
<i>managuensis</i>	157	<i>cruentifer</i>	86
<i>margnatus</i>	152	<i>oculatus</i>	85
		<i>pisquetus</i> , Caranx.....	272

List of scientific names—Continued

	Page		Page
pitilliger, Cottus	389	platyrhynchus, Acipenser	35
Gymnocanthus	389	Scaphirhynchops	35
pitamba, Acara	327	platostomus, Cylinndrosteus	37
Pithecomyzon	108	Lepisosteus	37
pittieri, Phallichthys	190	Platotichthys	221
Poecilia	190	chartes	221
pututosus, Rhypticus	316	platyargyrea, Anchovia	49
piranga, Serranus	310	Anchoviella	49
piruma, Amore	436	platycephalum, Peristedion	406
placitus, Hybognathus	143	Peristedium	406
Placopharynx	111	Platycephalus dormitator	435
carinatus	111	platycephalus, Ameiurus	164
Plagiogrammus	469	Cottus	387, 388
hopkinsi	469	Megalocottus	388
plagioplateo, Eleotris capite	436	Noturus	155
Plagiostomata	10	Notorynchus	11
Plagusia	230	Pimelodus	154
lactea	230	Platyaster	44
plagusia, Pleuronectes	231	Platygllossus dispilus	427
Symphurus	231	fiorealis	426
Plagopterus	147	nicholsi	428
argentissimus	147	poeyi	427
Plagusia	230, 231	Platygnathus	136
brasilensis	231	gracilis	136
fasciata	231	pallidus	136
lactea	230	physignathus	136
tessellata	231	(Platygnathus) gulonellus, Pogonichthys	136
plagusia, Pleuronectes	230, 231	Platygnathus	328
Symphurus	231	platyodon, Carcharias	15
Plagyodontem	171	Squalus	15
Plagyodus	171	Platypodon	15
æsculapius	171	perezi	16
plana, Platessa	227	Platypterus	193
planci, Tympanomium	503	couchianus	193
Plancterus	179	dominicensis	191
kansæ	179	maculatus	193
zebra	179	immaculatus	193
planiceps, Ancistrus	161	mentalis	192
Arius	149	perugæ	191
Catostomus	107	quitzcoensis	182
Lasiancistrus	161	tropicus	192
Netuma	149	variatus	193
planiceps, Netuma	149	playtypogon, Arius	149
planifrons, Eupomacentrus	414	Netuma	149
Pomacentrus	414	Platyrrhina evasperata	24
Planirostra	36	triseriata	24
edentula	36	Platyrrhinidae	24
planus, Pleuronectes	227	Platyrrhinoides	24
Plargyrus argentatus	128	triseriatus	24
bowmani	128	platyrhinus, Acipenser (Huso)	35
melanocephalus	145	platyrhynchus, Carcharias	15
typicus	128	Eulamia	15
plargyrus, Rutilus	128	Minomus	104
plate, Branchiostoma	7	Pantosteus	104
Platessa	226	Platysomatichthys	221
blimeata	226	stomas	221
dentata	222	Platysomus	274
divinensis	222	micropteryx	275
ferruginea	226, 227	sphinx	275
glabra	227	Platysqualus	17
oblonga	223, 224	tudes	17
ocellaris	224	Platytristes	53
plana	227	apus	53
pusilla	227	plebeus, Catostomus	104
rostrata	227	Gasterosteus	237
platessa, Pleuronectes	226	Notolepidomyzon	104
platessoides, Hippoglossoides	222	plebeus, Mustelus	13
Pleuronectes	222	Plecotomus	160
Platichthys	226	boulengeri	160
rugosus	226	dorso dipterygio	160
stellatus	226	seminudus	160
unibrosus	226	plecostomus, Hypostomus	160
Platrostra	36	Loricaria	160
edentula	36	Plectobranchius	468
Platophrys	217	evides	468, 469
constellatus	217	Plectognathus	491
ellipticus	217	plectrodon, Porichthys	488
leopardinus	218	Plectromus	233
lunatus	217	beani	233
maculifer	217	bispinosus	233
nebularis	217	crassiceps	233
ocellatus	217	crusticeps	233
spinosus	217	lugubris	233
tæbiopterus	220	mizolepis	233
platophrys, Citharichthys	219	nycterinus	233
Platopterus	24	suborbitalis	233

List of scientific names—Continued

	Page		Page
<i>Plectropoma accensum</i>	321	<i>Pleuronichthys guttulatus</i>	225
affine.....	321	nepheles.....	225
bovinum.....	321	ocellatus.....	225
chloropterum.....	312	ritteni.....	225
chlorurum.....	321	verticalis.....	225
crocota.....	321	pleurophthalmus, Antennarius.....	505
ephippium.....	321	pleurospilus, Girardinus.....	190
gummigutta.....	321	Poecilistes.....	190
guttavarium.....	321	pleurostictus, Triglops.....	379
hispanum.....	308	Pleurothvis.....	74
indigo.....	321	Phosteostoma.....	45
melanorhina.....	321	lutipinnis.....	45
monacanthus.....	313	plumatula, Calamus.....	336
multiguttatum.....	313	plumbea, Chimæra.....	33
nigricans.....	321	Dionda.....	144
puella.....	320, 321	Gambusia.....	192
vitulinum.....	321	plumbeolus, Alburnops.....	128
<i>Plectrypops</i>	236	plumbeus, Bathyalopex.....	33
retrospinis.....	236	Couesius.....	136
pleianus, Pseudoscarus.....	435	Gobio.....	136
Scarus.....	435	Petromyzon.....	10
plei, Hemiramphus.....	199	Pseudoscarus.....	435
pleuriticus, Salmo.....	57	rubrilateralis, Couesius.....	116
Pleuraeromylon.....	13	plumierianus, Caranxomorus.....	270
Pleurogrammus.....	374	plurmer, Le Diodon.....	501
monopterygius.....	374	plurmer, Chærodon.....	358
<i>Pleurolepis</i>	288	Coryphæna.....	356
asprellus.....	288	Lingæres.....	343
pellucidus.....	288	Gerres.....	342, 343
<i>Pleuronectes</i>	226	Gobius.....	447
achius.....	229	Hæmulon.....	330
americanus.....	227	Læbrus.....	330
apoda.....	229	Malacanthus.....	356
aquosus.....	217	Mugil.....	253
aramaca.....	218	Polydactylus.....	256
argus.....	217	Scæna.....	333
asper.....	227	Scomber.....	271
cicatricosus.....	227	Scomberomorus.....	257
cornutus.....	225	Scorpena.....	370
cynoglossus.....	221, 228	Sivdium.....	447
dentatus.....	224	plumierianus, Caranxomorus.....	270
diagrammus.....	226	plumieri, Polydactylus.....	256
ellipticus.....	217	pluriserialis, Naretes.....	52
franklini.....	227	plutoma, Raja.....	26
fuscus subrotundus glaber.....	230	pluvialis, Tetradon.....	499
gilli.....	228	pluvius, Salmo.....	61
glabra.....	217	<i>Pneumatophorus</i>	257
glacialis.....	227	australasicus.....	258
hippoglossoides.....	221	diogo.....	258
hippoglossus.....	221	grey.....	258
kitt.....	228	pneumatophorus, Scomber.....	257, 258
lumanda.....	222, 226	pocantello, Catostomus.....	107
lmandoides.....	222	Podager.....	341
lineatus.....	230	podas, Bothus.....	217
linguata.....	222	podoleptus.....	215
lunatus.....	217	podostemone, Etheostoma.....	286
maculatus.....	217	Imostoma.....	286
maculiferus.....	217	Podothecus.....	394
melanogaster.....	224	acipenserinus.....	394
microcephalus.....	228	gilberti.....	395
microstomus.....	228	peristethus.....	394
mollis.....	229	<i>Pœcilia</i>	190
oblongus.....	224	(Acropœcilia) tridens.....	190
obscurus.....	228	amates.....	190
ocellatus.....	217	boucardi.....	192
oculis etc.....	229	butleri.....	192
pallasii.....	226	catenata.....	179
papillosus.....	218	chisoyensis.....	192
peracuatius.....	226	cervicola.....	175
pinguis.....	221	cuneata.....	193
plagus.....	231	dommoceris.....	191
plagusia.....	230, 231	dovi.....	192
plagus.....	227	elongata.....	130
platessa.....	226	limantouri.....	192
platessoides.....	226	lneyata.....	192
quadridens.....	228	melanogaster.....	191
quadrifuberculatus.....	228	melapleura.....	178, 185
quenseli.....	228	mexicana.....	192
satis parva.....	217	multilineata.....	192
saxicola.....	228	olivacea.....	177
stellatus.....	228	petenensis.....	192
surinamensis.....	217	pittieri.....	190
<i>Pleuronectidæ</i>	225	presidionis.....	189
<i>Pleuronichthys</i>	225	reticulata.....	191
cœvus.....	225	reticulatus.....	191
decurrens.....	225	salvatoris.....	192

List of scientific names—Continued

	Page		Page
<i>Poecilia schneideri</i>	190	<i>Pogonias</i>	352
<i>sphenops</i>	192	<i>chromis</i>	352
<i>spilonota</i>	192	<i>courbina</i>	352
<i>spilurus</i>	192	<i>fasciatus</i>	352
<i>surinamensis</i>	190	<i>Pogonichthys</i>	116
<i>tenuis</i>	192	<i>argyreus</i>	116
<i>thermalis</i>	192	<i>communis</i>	136
<i>tridens</i>	190	<i>inaequilobus</i>	116
<i>vanderpolli</i>	193	<i>macrolepidotus</i>	116
<i>arubensis</i>	193	<i>(Platygobio) gulonellus</i>	136
<i>vittata</i>	191	<i>symmetricus</i>	121
<i>vivipara</i>	190	<i>Pogonius chromis</i>	352
<i>Poecilichthys</i>	232, 289	<i>polaris, Blennius</i>	476
<i>artesia</i>	292	<i>Cottus</i>	388
<i>asprigenis</i>	291	<i>Lycodalepis</i>	476
<i>beani</i>	287	<i>Merlangus</i>	209
<i>borealis</i>	291	<i>Porocottus</i>	388
<i>butlerianus</i>	294	<i>Polipturus</i>	257
<i>camurus</i>	290	<i>Pohstotrema</i>	8
<i>jessiae</i>	291	<i>curtiss-jamesii</i>	8
<i>mesesus</i>	287	<i>deani</i>	8
<i>palustris</i>	294	<i>stoutii</i>	8
<i>punctulatus</i>	292	<i>politus, Cheilichthys</i>	499
<i>quiescens</i>	294	<i>Seriphus</i>	353
<i>rufilineatus *</i>	290	<i>Tetraodon</i>	499
<i>sagitta</i>	293	<i>Pollachius</i>	209
<i>sanguifluus</i>	290	<i>chalcogrammus fucensis</i>	209
<i>saxatilis</i>	287	<i>virens</i>	209
<i>spectabilis</i>	291	<i>pollachius, Gadus</i>	209
<i>swami</i>	291	<i>pollicaris, Uranidea</i>	356
<i>variatus</i>	289	<i>pollux, Pontinus</i>	372
<i>versicolor</i>	291	<i>polyacanthocephalus, Cottus</i>	387
<i>virgatus</i>	292	<i>Myoxocephalus</i>	387
<i>vitreus</i>	288	<i>Polyacanthonotus</i>	76
<i>vulneratus</i>	290	<i>challengeri</i>	76
<i>zonalis</i>	289	<i>rostratus</i>	76
<i>Poecilidae</i>	184	<i>polyactis, Melanocetus</i>	510
<i>Poecilopsis</i>	189	<i>polyactiocephalum, Bryostemma</i>	466
<i>balsus</i>	189	<i>polyactiocephalus, Blennius</i>	466
<i>fasciata</i>	189	<i>Polycanthus</i>	239
<i>infans</i>	189	<i>polycaula, Rhamdia</i>	158
<i>isthmensis</i>	190	<i>polycaulus, Pimelodus</i>	158
<i>latidens</i>	189	<i>Polycirrhus</i>	351
<i>occidentalis</i>	189	<i>dumerli</i>	351
<i>presidionis</i>	189	<i>rathbuni</i>	351
<i>turubarensis</i>	189	<i>Polyclemus</i>	351
<i>Poecilistes</i>	189	<i>dumerli</i>	351
<i>pietropilus</i>	190	<i>Polydactylus</i>	256
<i>Poecilopterus</i>	87	<i>opercularis</i>	256
<i>bonapartei</i>	87	<i>plumieri</i>	256
<i>Poecilodes</i>	187	<i>plumieri</i>	256
<i>bimaculatus</i>	187	<i>polygonius, Acanthostracion</i>	497
<i>poeciloides, Lebistes</i>	191	<i>Gymnothorax</i>	92
<i>Lamia</i>	192	<i>polylepis, Balistes</i>	492
<i>Poecilophis</i>	94	<i>Scorpaena</i>	370
<i>nocturnus</i>	94	<i>Polyipnus</i>	74
<i>Poecilopsis isthmensis</i>	190	<i>laternatus</i>	74
<i>poecilopterus, Jubs</i>	426	<i>spinosus</i>	74
<i>poecilopus, Myrpristis</i>	235	<i>polylepis, Verrunculus</i>	492
<i>Rhamphoberyx</i>	235	<i>Polymixia</i>	234
<i>Poecilosoma</i>	289	<i>lowei</i>	234
<i>erythrogastrum</i>	291	<i>nobilis</i>	234
<i>transversum</i>	291	<i>Polymixidae</i>	234
<i>poecilura, Myxostoma</i>	111	<i>polymorphus, Gadus</i>	210
<i>Poecilurichthys</i>	96	<i>Polynemidae</i>	256
<i>breviortii</i>	96, 97	<i>Polynemus</i>	256
<i>pulcher</i>	97	<i>americanus</i>	256
<i>tenuis</i>	97	<i>approximans</i>	256
<i>unilineatus</i>	98	<i>artedii</i>	256
<i>poecilurum, Moxostoma</i>	111	<i>californiensis</i>	256
<i>Poecilus</i>	190	<i>macronemus</i>	256
<i>poeyi, Alepidosaurus (Caulopus)</i>	171	<i>mango</i>	256
<i>Dactyloscopus</i>	455	<i>melanopoma</i>	256
<i>Engraulis</i>	51	<i>octonemus</i>	256
<i>Euctenogobius</i>	442	<i>oligodon</i>	256
<i>Gobius</i>	442	<i>paradiseus</i>	256
<i>Iridio</i>	427	<i>quinquarius</i>	256
<i>Lycengraulis</i>	51	<i>sexradiatus</i>	409
<i>Orthopristis</i>	335	<i>virginicus</i>	256
<i>Pempheris</i>	323	<i>Polyodon</i>	36
<i>Platyglottus</i>	427	<i>feuille</i>	36
<i>Siphostoma</i>	241	<i>folium</i>	36
<i>Syngnathus</i>	241	<i>spatula</i>	36
<i>Synodus</i>	164	<i>polyodon, Tetragonopterus (Hemibrycon)</i>	97
<i>Pogonathus</i>	352	<i>Polyodontidae</i>	36
<i>courbina</i>	352		

List of scientific names—Continued

	Page		Page
Polypersa.....	399	Pomotis convexifrons.....	300
greeni.....	399	elongatus.....	300
polypogon, Linophryne.....	510	fallax.....	300
Polypriion.....	308	gibbosus.....	301
americanus.....	308	gulosus.....	302, 303
cernium.....	308	guttatus.....	304
massiliensis.....	308	heros.....	302
Polyprosopus.....	20	holbroeckii.....	302
macer.....	20	incisor.....	301
Polypterichthys.....	244	inscriptus.....	300
valentini.....	244	longulus.....	299
polysarcus, Naucrates.....	278	luna.....	301
polyspilos, Rhombus.....	223	microlophus.....	302
Polyuranodon.....	91	nefastus.....	300
polyuranodon, Muraena.....	91	nitida.....	300
Pomacanthodes.....	360	notatus.....	302
zonipectus.....	360, 361	obesus.....	305
Pomacanthus.....	360	obscurus.....	301
arcuatus.....	360	pallidus.....	301
baileatus.....	360	popeii.....	300
cingulatus.....	360	ravenelli.....	302
crescentalis.....	361	rubri-cauda.....	299
paru.....	360	sanguinolentus.....	300
quinquecinctus.....	360	solis.....	300
Pomacanthus, Mesoprion.....	324	speciosus.....	301, 302
Pomacentridae.....	412	vulgaris.....	302
Pomacentrus.....	413	pomotis, Acantharehus.....	303
adustus.....	414	Centrarchus.....	303
analigutta.....	414	Pomoxis.....	304
analis.....	413	annularis.....	304
atro-cyaneus.....	413	nitidus.....	304
bairdii.....	415	sparoides.....	305
breviceps.....	413	Pomoxys berricauda.....	304
caudalis.....	413	intermedius.....	304
denegatus.....	415	protacanthus.....	304
dorso-punicans.....	413	pompilius, Centrolophus.....	267
flavilatus.....	414	Pompilus.....	267
flaviventer.....	414	pompilus, Coryphæna.....	267
fuscus.....	413	Thynnus.....	267
gilli.....	414	ponderosus, Ameiurus.....	152
leucurus.....	414	pondicerriana, Elacate.....	281
leucostictus.....	413	ponticus, Gasterosteus.....	237
lividus.....	413	Hippoglossus.....	221
niveatus.....	416	Pontinus.....	372
obscuratus.....	414	beanorum.....	372
otophorus.....	414	bergii.....	372
partitus.....	414	castor.....	372
planifrons.....	414	fucirhinus.....	372
quadrigutta.....	416	longispinus.....	372
rectifrenum.....	414	macrolepis.....	372
variabilis.....	413	pollux.....	372
xanthurus.....	413	rathbuni.....	372
Pomadassus bayanus.....	334	sierra.....	372
elongatus.....	333	popeii, Pomotis.....	300
grandis.....	334	popoche, Xystrosus.....	142
Pomadasy.....	334	porcus, Capriscus.....	401
approximans.....	334	Scorpena.....	370
bayanus.....	334	Porichthys.....	487
boschnæ.....	334	greenel.....	487
cæsius.....	331	margaritatus.....	487
grandis.....	334	nautopodium.....	487
macracanthus.....	334	notatus.....	487
productus.....	334	plectrodon.....	488
sinuosus.....	335	porifer, Lycenchelys.....	476
starr.....	334	Lycodes.....	476
templei.....	334	Porobronchus.....	486
Pomatonichthys.....	208	linearis.....	486
constanciae.....	208	Porochinus.....	470
Pomatomidae.....	280	rotbrocki.....	470
Pomatopus.....	280	Porocottus.....	388
saltatrix.....	280	bradfordi.....	388
skib.....	280	polaris.....	388
telescopium.....	206	quadratus.....	388
Pomatopriion.....	415	quadrifilis.....	388
Pomatopsetta.....	222	sellaris.....	388
Pomolobus.....	41, 42	Porogadus.....	482
æstivalis.....	42	atripectus.....	482
chrysocloris.....	41	breviceps.....	482
medicaris.....	42	longiceps.....	482
pseudoharengus.....	42	miles.....	482
lacustris.....	42	pomelas.....	481
Pomotis.....	299	Poromitra.....	233
bombifrons.....	301	capito.....	233
breviceps.....	300	Poronotus.....	267
catesbei.....	302	simillimus.....	266
chætodon.....	303	triacanthus.....	267

List of scientific names—Continued

	Page		Page
porosissimum, Nautopædium	488	Prionodes fuscus	319
porosissimus, Batrachus	487, 488	phoebe	319
porosus, Carcharias	16	tabacaria	319
Cottus	387	tigrinus	319
Esox	173	Prionodon	17
Scoliodon	15	cucuri	16
Squalus	15	obelatus	16
postica, Echeoneis	449	(Prionodon) limbatus, Carcharias	16
Potamocottus	383	milberti, Carcharias	16
bendirei	384	Prionotus	497
caroline	383, 385	alatus	408
punctulatus	383	albipinnatus	408
zopherus	385	beanni	408
pottsi, Aplesion	290	birostratus	408
Rafinesquellus	290	evolans	407
powelli, Balistes	492	frontalis	408
precisus, Clinus	470	gymnostethus	408
Eumesogrammus	470	horrens	408
predatoria, Buccone	354	loxias	408
predatorius, Cestreus	354	miles pectoralis	407
prestabilis, Alosa	42	militaris	408
prestigiator, Serranus	319	ophryas	407
prayensis, Pseudupeneus	343	pectoralis	407
preblei, Coregonus	65	pileatus	407
Prosopium	65	punctatus	407
prenadilla, Brontes	159	quiescens	408
presidionis, Pœcilia	189	rosus	407
Pœciliopsis	189	rubio	407
pretiosa, Argentina	67	ruscarius	408
pretiosus, Hypomesus	67	sarritor	407
Ruvettus	261	scutulus	407
Trachichthys	234	stearnsi	408
Priacanthichthys	310	stephanophrys	407
maderaspatisensis	310	strigatus	407
Priacanthidae	322	tribulus	408
Priacanthus	322	xenisma	408
altus	323	Prionurus punctatus	363
arenatus	322	Pistida	23
carolinus	322	Pristidurus	12
catalufa	322	Pristigaster	45
cepedianus	323	cayanus	45
cruentatus	322	dovi	45
fulgens	322	flavipinnis	44
macrophthalmus	322	lutipinnis	45
niphonius	322	macrops	45
serrula	323	marti	45
Priapella	188	(Odontognathus) panamensis	45
bonita	188	phaeton	45
Priapichthys	187	tartoon	45
annectans	187	Pristipoma	334
hesperis	187	acara-pimma	333
fosteri	189	auratum	333
huberi	187	bicolor	332
letonai	190	bilineatum	332
olomina	187	boschma	334
panamensis	187	boucardi	334
pribilovius, Nautichthys	382	branicki	334
Nautiscus	382	brasiliense	332
pricei, Campostoma	147	brevipinne	335
Haustor	153	camtharum	335
Villarius	152, 153	catharine	333
Pricus	10	chalcum	335
prieto, Lutjanus	325	croco	334
Prionotus	500	cultiferum	334
(Anchisomus) caudacinctus	500	davidsoni	332
Prinospina	364	dovi	332
princeps, Caulolatilus	357	fasciatum	335
Cottus	384	furthi	332
Latilus	357	(Hæmulopsis) axillare	333
princeps, Antennarius	505	brevipinne	335
Chironectes	505	nitidum	333
Julis	426	humile	334
Prionodon	180	humilis	334
Prionodontophis	91	kneri	335
Prionace	17	leuciscus	333
glauca	17	macracanthum	334
Prionatus egretta	408	melanopterum	332
Prionistius	380	notatum	332
macellus	380	panamense	333
Prionodes	319	productum	334
æquidens	319	ramosum	334
baldwini	319	rodo	333
bulleri	319	scapulare	332
deweegeri	319	splendatum	332
fasciatus	319	serrula	333
flavescens	319	trilineatum	332

List of scientific names—Continued

	Page		Page
Pristipomes, les.....	334	Promicrops.....	312
Pristipomus.....	334	itaiara.....	312
Pristipomoides.....	328	quinquefasciata.....	312
macrophthalmus.....	328	Promicropus.....	316
typus.....	328	bistrispinus.....	316
Pristis.....	23	decoratus.....	317
acutirostris.....	23	nigripinnis.....	317
granulosa.....	23	Pronotogrammus.....	322
megalodon.....	23	eos.....	322
microdon.....	23	multifasciatus.....	322
mississippiensis.....	23	proops, Bagrus.....	148
occa.....	23	Sciadeichthys.....	148
pectinatus.....	23	Propterygia.....	24
zephyreus.....	23	hyposticta.....	24
pristis, Squalus.....	23	prorates, Lepophidium.....	484
Pristiurus.....	12, 13	Leptophidium.....	484
arae.....	12	proridens, Calamus.....	336
melastomas.....	12	proriger, Gnathopis.....	80
xanurus.....	12	Sebastichthys.....	366
Pristocantharus.....	335	var brevispinis, Sebastichthys.....	366
Pristopomus.....	334	Squalus.....	120
Proarthri.....	10	prorigerum, Acutomentum.....	366
probatocephalus, Archosargus.....	338	Ophisoma.....	80
Sparus.....	338	proserpina, Moniana.....	132
proboscidea, Limanda.....	227	Prosopium.....	65
proboscideum, Monacanthum.....	495	abyssicola.....	66
proboscideus, Chanomugil.....	254	conesi.....	65
Mugil.....	254	couleri.....	65
procellarium, Myctophum.....	170	kennicotti.....	65
procera, Venefica.....	82	preblei.....	65
Proceros.....	36	quadrilaterale.....	65
maculatus.....	36	spilonotus.....	65
vittatus.....	36	stanleyi.....	65
procerum, Nettastoma.....	82	williamsoni.....	65
Prochilus.....	436	cismontanus.....	65
proene, Eacoetus (Cypsilurus).....	201	Prospinus.....	312
Hybognathus.....	133	prosthemus, Ceratichthys.....	136
Hybopsis.....	133	prosthismus, Ameiurus.....	154
Proctostegus.....	261	protacanthus, Pomovys.....	304
proctostegus.....	264	proteus, Salmio.....	54
proctostegus, Proctostegus.....	264	protoculus, Nyctophum.....	168
producta, Rhinobatus.....	24	Protoporus.....	119
Ulama.....	341	dominus.....	119
productum, Pristipoma.....	334	proxima, Seriola.....	279
productus, Alepocephalus.....	51	proximus, Gadus.....	210
Cylindrosteus.....	37	Microgadus.....	210
Engraulis.....	50	prunosus, Gadus tomcodus.....	210
Eucinostomus.....	341	Pseudoblechnus.....	465
Leuciscus.....	120	hypacanthus.....	465
Merlangus.....	214	Psempylus prometheus.....	261
Merluccius.....	214	Psenes.....	268
Pomadasy.....	334	auratus.....	268
Rhinobatos.....	24	chapmani.....	269
proellaris, Microperca.....	294	cyanophrys.....	268
profundorum, Acanthocottus.....	292	edwardsi.....	269
Aristurus.....	12	fuscus.....	268
Lepophidium.....	484	javanicus.....	268
Leptophidium.....	484	maculatus.....	268
Scylliorhinus.....	12	pacificus.....	269
Xenognathus.....	52	pellucidus.....	269
Zesticelus.....	392	regulus.....	268
Profundulus.....	174	Psettichthys.....	222
candalaria.....	174	melanostictus.....	222
labialis.....	174	sordidus.....	213
oaxaca.....	174	umbrosus.....	226
punctatus.....	174	Pseudageniosus.....	159
profundus, Mesoprius.....	326	pseudaplates, Micropterus.....	208
Prognathodes.....	359	Pseudaroides.....	158
aculeatus.....	359	pantherinus.....	159
prognathus, Coregonus.....	63	Pseudarius.....	150
Leucichthys.....	63	Pseudocheirodon.....	96
Prognurus.....	403	affinis.....	96
cypselurus.....	403	Pseudochromide.....	322
prolixus, Leuciscus.....	146	pseudogula, Eucinostomus.....	341
promelas, Chirostoma.....	251	pseudo-gula, Eucinostomus.....	341
Mecbia.....	481	pseudoharengus, Clupea.....	42
Pimephales.....	145	lacustris, Pomolobus.....	42
Porogadus.....	481	Pomolobus.....	42
Promethus.....	261	Pseudohemiodon.....	161
atlanticus.....	261	pseudohispanica, Sardinia.....	43
promethus, Gampylus.....	261	Pseudophilus.....	428
Promethichthys.....	261	adustus.....	428
Psempylus.....	261	girardi.....	428
Promethichthys.....	261	inornatus.....	428
promethus.....	261	melanotis.....	428

List of scientific names—Continued

	Page		Page
<i>Pseudojulis notospilus</i>	428	<i>Pteroplatea micrura</i>	29
<i>venustus</i>	428	<i>rava</i>	29
<i>Pseudoloricaria</i>	161	<i>vallanti</i>	29
<i>Pseudomonacanthus</i>	493	<i>Pteropodus</i>	369
<i>Pseudomoringua</i>	89	<i>caratus</i>	369
<i>Pseudomurena</i>	91	<i>caurinus</i>	369
<i>maderensis</i>	91	<i>chrysomelas</i>	369
<i>Pseudopleuronectes</i>	227	<i>dallii</i>	369
<i>americanus</i>	227	<i>gilberti</i>	369
<i>dignabilis</i>	227	<i>maliger</i>	369
<i>pseudoplates, Micropterus</i>	298	<i>nebulosus</i>	369
<i>Pseudopriacanthus</i>	323	<i>rastralliger</i>	369
<i>altus</i>	323	<i>vevillaris</i>	369
<i>serrula</i>	323	<i>Pteropsandæ</i>	451
<i>pseudoptiles, Micropterus</i>	298	<i>Pteropus</i>	369
<i>Pseudorhamdia</i>	158	<i>pterotum, Ateleobranchium</i>	207
<i>piscatrix</i>	159	<i>pteryura, Strongylura</i>	196
<i>Pseudorhombus</i>	223	<i>pteryurus, Tylosurus</i>	196
<i>vorax</i>	224	<i>Pterygocottus</i>	380
<i>Pseudoscarus</i>	435	<i>macomi</i>	380
<i>aracanga</i>	434	<i>Pterygostedion imberbe</i>	406
<i>coelestinus</i>	435	<i>Ptilichthyidae</i>	471
<i>gnathodes</i>	434	<i>Ptilichthys</i>	471
<i>guacamaya</i>	435	<i>goodii</i>	471
<i>lineolatus</i>	434	<i>Ptychocheilus</i>	113
<i>microps</i>	435	<i>gracilis</i>	113, 114
<i>microrrhinos</i>	435	<i>grandis</i>	114
<i>perrioi</i>	435	<i>harfordi</i>	114
<i>plelanus</i>	435	<i>lucius</i>	114
<i>plumbeus</i>	435	<i>major</i>	114
<i>simpler</i>	435	<i>oregonensis</i>	114
<i>teniopterus</i>	433	<i>umpqua</i>	114
<i>Pseudoscopus</i>	452	<i>vorax</i>	114
<i>scriptus</i>	452	<i>Ptycholepis</i>	40
<i>Pseudosebastes</i>	370	<i>Ptychostomus</i>	109
<i>Pseudotriakidae</i>	12	<i>albidus</i>	110
<i>Pseudotriakis</i>	12	<i>albus</i>	110
<i>microdon</i>	12	<i>breviceps</i>	111
<i>Pseudoxiphophorus</i>	187	<i>bucco</i>	110
<i>bimaculatus</i>	187	<i>cervinus</i>	111
<i>jonesi</i>	188	<i>collapsus</i>	110
<i>teniatus</i>	188	<i>conus</i>	111
<i>pauciradiatus</i>	188	<i>coregonus</i>	110
<i>reticulatus</i>	187	<i>crassilabris</i>	111
<i>Pseudupeneus</i>	343	<i>haydeni</i>	109
<i>præyensis</i>	343	<i>lachrymalis</i>	111
<i>Psilonotus</i>	500	<i>papillosus</i>	109
<i>psittacinus, Serranus</i>	319	<i>pidensis</i>	110
<i>psittaculus, Labrus</i>	426	<i>robustus</i>	111
<i>psittacus, Colomesus</i>	500	<i>thalassinus</i>	110
<i>Coryphæna</i>	430, 433	<i>velatus</i>	109
<i>Lachnolaimus</i>	425	<i>Fudiano verde</i>	426
<i>Tetrodon</i>	500	<i>vermelho</i>	425
<i>Xyrichtys</i>	430	<i>puella, Hypoplectrus unicolor</i>	321
<i>Psychichthys</i>	33	<i>Plectropoma</i>	320, 321
<i>affinis</i>	33	<i>puellaris, Cossyphus</i>	425
<i>Psychrolutes</i>	392	<i>Decodon</i>	425
<i>paradoxus</i>	392	<i>puertoricensis, Malacoctenus</i>	460
<i>zebra</i>	392	<i>pugetensis, Arctidius</i>	375
<i>Psychromaster</i>	293	<i>Chitonotus</i>	375
<i>tuscumbia</i>	293	<i>pugeti, Gasterosteus</i>	238
<i>Ptix</i>	261	<i>pulchella, Gila</i>	119, 120
<i>parvipinnis</i>	261	<i>Leuciscus</i>	120
<i>Pteraclidae</i>	266	<i>Moniana</i>	131
<i>Pteraclidus</i>	266	<i>Tigoma</i>	120
<i>Pteracis</i>	266	<i>pulchelloides, Leuciscus</i>	117
<i>carolinus</i>	266	<i>pulchellus, Bodianus</i>	425
<i>trichipterus</i>	266	<i>Cossyphus</i>	425
<i>Pteraplatea canariensis</i>	29	<i>Cymatogaster</i>	410
<i>valenciennæ</i>	29	<i>Leuciscus</i>	117
<i>Pterengraulis</i>	51	<i>Liparis</i>	400
<i>atherinoides</i>	51	<i>Lyoliparis</i>	400
<i>Pteridium</i>	266	<i>Oligocephalus</i>	291
<i>Pterocephala</i>	31	<i>Zygocetes</i>	177
<i>Pterognathus</i>	458	<i>pulcher, Eques</i>	353
<i>satricus</i>	458	<i>Labrus</i>	426
<i>Pteromyleus</i>	30	<i>Pareques</i>	353
<i>asperimus</i>	30	<i>Pinelomatopon</i>	426
<i>Pteronotus</i>	156	<i>Poecilurichthys</i>	97
<i>Pterophryne</i>	504	<i>pulehra, Tigoma</i>	120
<i>Pterophrinoides</i>	504	<i>pullata, Dicerolene</i>	453
<i>Pteroplatea</i>	29	<i>pullum, Chondrostoma</i>	146
<i>altavala</i>	29	<i>pullus, Cantherines</i>	494
<i>crebrispunctata</i>	29	<i>Monacanthus</i>	494
<i>marmorata</i>	29	<i>Pinelodus</i>	154

List of scientific names—Continued

	Page		Page
pulverentulus, Myloleucus	119-121	punctulatus, Callurus	238
pulvereus, Zygonectes	178	Cottus	383
pulverulenta, Tigoma	120	Hippocampus	244
pulverulentum, Oxygeneum	113	Lampugus	265
puncta, Furcaria	412	Lythrurus	126
puncticeps, Cryptopterus	85	Minnulus	126
punctarens, Gymnothorax	92	Pimelodus	155
punctata, Alutera	495	Poecilichthys	292
Davidia	495	Potomocoitus	383
Gambusia	184, 185	Scarus	433
Girella	338	Sicyases	490
Gorgasia	77	Squalus	12
Lamna	19	Pungtius	238
Muraena	91	brachypoda	239
Muraenophis	92	mainensis	239
Perca	309, 345	pungtius	239
punctulata, Gambusia	185	pungtius brachypoda, Gasterosteus	239
Solea	217	pungtius, Gasterosteus	238, 239
Trigla	407	Pungtius	239
punctatissima, Anguilla	77	pumceus, Hyborhynchus	145
punctatissimus, Canthigaster	500	purpurascens, Lepomis	301
Ophioscion	347	variegata, Salpa	327
Tetrodon	500	purpuratus bouvieri, Salmo	56
punctatum, Cichlasoma	417	Salmo	56
Hoplosternum	160	purpurea, Tigoma	119
Hyperprosopon argenteum	410	purpurescens, Anoplarchus	468
Megalonema	159	archolepis, Anoplarchus	468
Myctophum	169	purpureus, Gadus	209
Opisthognathus	453	Labrus	428
Ostracion	496	Lutjanus	326
Sicydium	447	pusilla, Aphoristia	231
Theraps	417	Bothrocara	477
punctatus, Apomotis	299	Olupea	42
Apriodon	15	Maynea	477
Balistes	492	Perca	295
Blennius	462, 469	Platessa	227
Bryttus	293, 299	pusillus, Coregonus	62
Caranx	270, 273	Enneapterygius	458
Carcharias	15	Leucichthys	62
Cephalopholis fulvus	309	Symphurus	231
Cotylopus	448	putaol, Gymnotus	101
Decapterus	270	putnami, Acipenser	34
Dermatolepis	313	Cottogaster	235
Diodon	501	Euchalarodus	227
Eques	353	Liopsetta	227
Fundulus	174	Pycnomma	438
Gadus	213	semsquamatum	438
Gymnothorax	92	Pygididae	159
Hoplosternum	160	Pygidium	159
Ictalurus	152	fusum	159
Isurus	19	striatum	159
Labrus	416	pygmaea bilineata, Umbra	174
Maurolicus amethystino	73	pygmaea, Eucalia	238
Micristodus	18	Umbra	174
Monacanthus	494	pygmaeus, Gadus	210
Morrhua	211	Gasterosteus	238
Myrophis	82, 83	Leuciscus	174
Opisthognathus	453	Pygosteus	238
Ostracion	496	Pyloctetis limosus	155
Pareques	353	pyramidatus, Cyclopterus	397
Prionotus	407	pyrrhogaster, Chrosomus	113
Prionurus	363	pyrrhomelas, Erogala	129
Profundulus	174	Photogenus	129
Silurus	151, 152	Pythomichthys	90
Squalus	12, 14, 15	sanguineus	90, 91
Stichæus	469		
Tetrodon	499		
Trachinus	310	quadracus, Apeltes	239
Upeneus	343	Gasterosteus	239
Xesurus	363	quadrangularis, Selene	358
puncticeps, Cryptopterenchelys	85	quadratus, Porocottus	388
Cryptopterus	85	Zeus	358
puncticulata, Gambusia	185	quadracornis, Acanthonotus	497
puncticulatus, Apogonichthys	293	Acanthostracion	497
Chilomycterus	502	Aspidophorus	393
Holocentrum	238	Cottus	388
punctifer, Crotalopsis	88	Hypsogonus	393
Hybognathus (Dionda)	144	Oncocottus	388
Trigonophallus	187	Ostracion	497
punctiferus, Menephorus	309	quadrifidens, Pleuronectes	228
punctillatus, Gobius	440	quadrifasciatus, Chasmodes	462
punctipinnis, Ayresia	413	Pholis	462
Dermatostethus	240	quadrifilis, Pathypterois	166
puncto-mediorum, Antennarius	505	Porocottus	388
punctulata, Gambusia punctata	185	quadrifidus, Pomacentrus	416
Microperca	294	quadrilaterale, Prosopium	65
Nivicola	292	quadrilateralis, Coregonus	65

List of scientific names—Continued

	Page		Page
quadriflineatus, <i>Hæmulon</i>	331	radiatus, <i>Lophus</i>	507
quadriloba, <i>Raja</i>	31	Ogcocephalus.....	507
quadriporus, <i>Gobius</i>	439	Sparus.....	426
<i>Rhinogobius</i>	439	radiosus, <i>Antennarius</i>	506
quadripunctatus, <i>Scomber</i>	259	radula, <i>Balistes</i>	493
quadrimemus, <i>Exocoetus</i>	201	<i>Melichthys</i>	493
quadriseriatus, <i>Artedius</i>	376	Radulmus.....	379
<i>Icelinus</i>	376	<i>asprellus</i>	379
quadrispinosus, <i>Scarus</i>	434	<i>boleoides</i>	379
quadrituberculatus, <i>Pleuronectes</i>	226	rafinesqui, <i>Acipenser</i> (<i>Huso</i>).....	35
quadrocellatus, <i>Ancyclopsetta</i>	224	<i>Colletia</i>	168
quappella, <i>Etheostoma</i>	291	<i>Cylindrosteus</i>	37
quatius rondeleti, <i>Anthias</i>	326	<i>Nyctophus</i>	168
Quassilabia.....	112	Rafinesquellus.....	290
Quassiremus.....	87	<i>pottsi</i>	290
<i>evionthas</i>	87	rafinesqui, <i>Acipenser</i> (<i>Huso</i>).....	35
<i>nothochir</i>	87	<i>Cylindrosteus</i>	37
quatuordecimlaminatus, <i>Echeneis</i>	449	rahleghanus, <i>Squalus</i>	20
quensel, <i>Pleuronectes</i>	228	Raja.....	24
quercinum, <i>Macrostoma</i>	166	<i>americana</i>	25
quercinus, <i>Notoscopelus</i>	166	<i>aquila</i>	31
Querimana.....	253	<i>brostris</i>	32
<i>curema</i>	253	<i>chantenay</i>	25
<i>gaimardianus</i>	254	<i>cooperi</i>	26
<i>gyrans</i>	253	<i>desmarestia</i>	25
<i>hospes</i>	254	<i>diaphanes</i>	25
<i>incilis</i>	253	<i>eglanteria</i>	25
<i>setosus</i>	254	<i>erinacea</i>	25
<i>thoburni</i>	253	<i>fimbriata</i>	32
<i>trichodon</i>	254	<i>flagellum</i>	30
querna, <i>Azevia</i>	221	<i>granulata</i>	25
<i>Cyclopsetta</i>	221	<i>guttata</i>	28
quiebra, <i>Lichia</i>	277	<i>inornata inermis</i>	26
quiescens, <i>Poecilichthys</i>	294	<i>interrupta</i>	26
<i>Prionotus</i>	408	<i>jordani</i>	26
<i>Uramdea</i>	385	<i>maclura</i>	29
Quiletula.....	444	<i>manata</i>	32
<i>y-cauda</i>	444	<i>micrura</i>	29
quinctifer, <i>Cirololops</i>	88	<i>marinari</i>	30
quinnat, <i>Salmo</i>	55	<i>obtusa</i>	26
quinquarius, <i>Pentanemus</i>	256	<i>ocellata</i>	25
<i>Polynemus</i>	256	<i>quadriloba</i>	31
quinquecinctus, <i>Pomacanthus</i>	360	<i>radiata</i>	25
quinquefasciata, <i>Promicrops</i>	312	<i>rosispinus</i>	26
quinquefasciatus, <i>Serranus</i>	312	<i>scabrata</i>	25
quinquehneatum, <i>Hæmulon</i>	331	rai, <i>Brama</i>	265
quitzeensis, <i>Platycephalus</i>	182	<i>Sparus</i>	265
<i>Zoogoneticus</i>	182	Raizero.....	327
quoyi, <i>Cestracion</i>	10	<i>aratus</i>	327
		Raja.....	24
Rabdophorus.....	359	<i>abyssicola</i>	26
Rabida.....	155	<i>ackleyi</i>	25
<i>eleuthera</i>	156	<i>aleutica</i>	26
<i>exilis</i>	156	<i>altavella</i>	29
<i>funnebris</i>	156	<i>binoculata</i>	26
<i>furiosa</i>	156	<i>centroura</i>	28
<i>gilberti</i>	156	<i>diaphanes</i>	25
<i>insignis</i>	156	<i>eglanteria</i>	25
<i>leptacantha</i>	155	<i>equatorialis</i>	26
<i>miura</i>	156	<i>erinacea</i>	25
<i>nocturna</i>	156	<i>fulla</i>	25
Rabirubia.....	327	<i>giorna</i>	31
<i>genizara</i>	426	<i>granulata</i>	25
<i>inermis</i>	327	<i>inornata</i>	26
rabirubia, <i>Anthias</i>	327	<i>interrupta</i>	26
Rabula.....	91, 93	<i>jamaicensis</i>	29
<i>davisi</i>	91	<i>lævis</i>	25
<i>longicauda</i>	91	<i>microtrachys</i>	25
<i>marmorea</i>	91	<i>mobular</i>	31
<i>panamensis</i>	91	<i>montereyensis</i>	26
<i>raciovii</i> , <i>Heterophallus</i>	184	<i>ornata</i>	26
Rachycentridæ.....	281	<i>parmifera</i>	26
Rachycentron.....	281	<i>pastinaca</i>	28
<i>canadus</i>	281	<i>percellens</i>	23
Radcliffella.....	446	<i>plutonia</i>	26
radiale, <i>Diplectrum</i>	318	<i>radiata</i>	25
radialis, <i>Serranus</i>	318	<i>rhina</i>	26
radians, <i>Scarus</i>	432	<i>rosispinus</i>	26
<i>Serranus</i>	318	<i>say</i>	26
<i>Sparisoma</i>	432	<i>scabrata</i>	25
radiata, <i>Iridio</i>	426	<i>sentia</i>	25
<i>Raja</i>	25	<i>stabuliforis</i>	25
<i>Raja</i>	25	<i>stellulata</i>	26
radiato, <i>Turdus oculo</i>	426	<i>trachura</i>	26
radiatus, <i>Halosaurus</i>	74	<i>tuberculata</i>	28
<i>Labrus</i>	426	Rajidae.....	24

List of scientific names—Continued

	Page		Page
raja, Squatina	23	Reinhardtius	221
raleighana, Harriotta	33	hippoglossoides	221
rammelsbergii, Mugil	253	remifer, Isopisthus	353
ramosum, Pristipoma	334	remiger, Myctophum	169
ramosus, Rhoniscus	334	remigera, Bothrocara	477
Ramularia	224	Remilegia	449
dendrica	224	australis	449
ranchovii, Heterophallus	184	Remora	449
Xiphophorus	193	remora	449
helleri	193	remora, Echeeneis	449
ranula, Liparis	402	Remora	449
ranulus, Careproctus	402	Remorina	449
Ranzania	504	albescens	449
makua	504	remoroides, Echeeneis	449
truncata	504	Remoropsis	449
raphidoma, Belone	196	brachyptera	449
Tylosurus	196	remotus, Carcharias	16
Raro	95	Serranus	312
rarus, Rhinoscopelus	168	Reniceps	17
Scopelus	168	tiburo	17
rasocio, Scorpena	370	repandus, Serranus	315
rashleighanus, Squalus	20	reticularis, Antennarius	505
rastralis, Anchoviella	50	reticulata, Amla	31
Stolephorus	50	Poecilia	191
rastrelliger, Physiculus	212	Spatularia	36
Pteropodus	369	Thalassophryne	488
Sebastichthys	369	reticulatus, Bryttus	299
Rastrinus	378	Catostomus	105
scutiger	378	Diodon	502
Rataboura	89	Eriscion	355
Rathbunella	452	Esor	173
allenii	452	Haliutichthys	507
hypoplecta	452	Lebistes	191
rathbuni, Fundulus	179	Lepidogaster	488
Mulloides	344	Lycodes	474
Polycurrus	351	Monocheir	229
Fontinus	372	Notropis	133
Upeneus	344	Otolithus	355
Zonoscion	351	Pseudoxiphophorus	187
Zygocetes	179	Tristropis	315
rauchi, Acipenser (Huso)	35	retifer, Scylliorhinus	11
Ichthyobus	102	retifera, Muræna	90
raucus, Sargus	338	retiferum, Scyllium	11
rava, Pteroplatea	29	retractus, Calliodon	431
ravenelli, Esor	173	Cryptotomus	431
Pomotis	302	retrocurrens, Hæmulon	329
recalva, Montereya	391	retropinnis, Catostomus	107
recalvus, Blennicottus	391	Microdesmus	473
rectangulare, Acara	421	Ophichthys	87
rectangularis, Astatheos	421	Ophichthys	87
rectifranum, Eupomacentrus	414	rotosella, Amla	295
Pomacentrus	414	Apogon	295
recurvirostrata, Sayris	197	retrospinus, Holocentrum	236
reddingi, Orthoprists	335	Plectroptops	236
redi, Orthogoriscus	503	retzii, Orthogoriscus	503
refulgens, Oligoplites	278	rex, Catostomus	108
regale, Myctophum	168	Etheostoma	283
Nannobranchium	168	Percina	283
regalis, Cynoscion	355	rex-mullorum, Apogon	295
Enchelyopus	213	rex-salmonorum, Trachipterus	216
Johnius	355	Trachypterus	216
Salmo	56	rhabdophora, Brachyrhaphis	186
Scomber	257	Gambusia	186
Scomberomorus	257	Rhacocilius	411
regani, Chirostoma	250	toxotes	411
Gambusia	186	Rhamdella	156
Rhamdia	158	Rhamdia	156
regis, Atherinops	252	barbata	158
Colpichthys	252	baronis-mulleri	157
regius, Blennius	213	brachycephala	157
Hybognathus	143	bransfordi	157
Lampris	216	cabrerae	157
Ophichthys	87	depressa	158
Urophycis	213	godmani	157
Zeus	216	guatemalensis	157
regulus, Psenes	268	heteracantha	158
Sebastes	363	hypselura	157
reighardi, Leucichthys	63	laticauda	157
Reighardina	9	managuensis	157
unicolor	9	microptera	157
reinhardi, Careproctus	402	motaguensis	157
Liparis	402	nicaraguensis	157
reinhardtii, Motella	208	oxace	157
reinhardtii, Corynolophus	509	parryi	158
Himantolophus	509	petenensis	158
		polycaula	158

List of scientific names—Continued

	Page		Page
<i>Rhamdia regani</i>	158	<i>Rhinogobius</i>	439
<i>rogersi</i>	158	<i>boleosoma</i>	439
<i>sacrilicii</i>	158	<i>claytoni</i>	439
<i>salvini</i>	157	<i>curacao</i>	439
<i>underwoodi</i>	158	<i>eigenmanni</i>	439
<i>wagneri</i>	157	<i>enceomus</i>	439
<i>Rhamphichthys brevirostris</i>	101	<i>garmani</i>	439
<i>Rhamphoberyx</i>	235	<i>glaucofrænulum</i>	439
<i>leucopus</i>	235	<i>manglicola</i>	439
<i>poecilopus</i>	235	<i>mowbrayi</i>	439
<i>Rhamphocottidae</i>	393	<i>parvus</i>	439
<i>Rhamphocottus</i>	393	<i>quadriporus</i>	439
<i>richardsoni</i>	393	<i>shufeldti</i>	439
<i>Rhagma</i>	322	<i>similis</i>	439
<i>thaumasium</i>	322	<i>stigmaturus</i>	439
<i>Rhegmatidae</i>	322	<i>tortuga</i>	439
<i>Rhegnopteri</i>	256	<i>Rhinoliparis</i>	405
<i>Rhencus</i>	333	<i>attenautus</i>	405
<i>panamensis</i>	334	<i>barbulifer</i>	405
<i>Rheocrypta</i>	285	<i>Rhinonemus</i>	209
<i>copelandi</i>	285	<i>Rhinoptera</i>	31
<i>rhessodon</i> , <i>Arbacia</i>	490	<i>bonasus</i>	31
<i>Gobiesox</i>	490	<i>encenadae</i>	31
<i>Rhina</i>	22	<i>lalandi</i>	31
<i>rhina</i> , <i>Antimora</i>	211	<i>steindachneri</i>	31
<i>Raja</i>	26	<i>vespertilio</i>	31
<i>Rhincodon</i>	18	<i>Rhinopteridae</i>	31
<i>typicus</i>	18	<i>Rhinoscion</i>	348
<i>Rhineodon</i>	18	<i>saturnus</i>	348
<i>typicus</i>	18	<i>Rhinoscopelus</i>	168
<i>Rhineodontidae</i>	18	<i>andreae</i>	168
<i>Rhinesomus</i>	496	<i>coccol</i>	169
<i>triqueter</i>	496	<i>oceanicus</i>	169
<i>Rhinichthys</i>	139	<i>rarus</i>	168
(<i>Apocope</i>) <i>nevadensis</i>	141	<i>Rhinotriads</i>	13
<i>velifer</i>	141	<i>henlei</i>	14
<i>arenatus</i>	140	<i>rhizophora</i> , <i>Fundulus</i>	175
<i>atronasus</i>	140	<i>rhodochloris</i> , <i>Sebastichthys</i>	367
<i>badius</i>	140	<i>Sebastomus</i>	367
<i>bowersi</i>	139	<i>rhodopus</i> , <i>Trachinotus</i>	276
<i>cataractae</i>	139	<i>rhodorus</i> , <i>Ascelichthys</i>	392
<i>croceus</i>	140	<i>rhodospilus</i> , <i>Gobiesox</i>	489
<i>dulcis</i>	139	<i>rhodoterus</i> , <i>Holconotus</i>	410
<i>henshawi</i>	141	<i>Rhodymenichthys</i>	468
<i>lunatus</i>	140	<i>dolichogaster</i>	468
<i>luteus</i>	139	<i>rhomalea</i> , <i>Gnathypops</i>	452
<i>marmoratus</i>	139	<i>rhomaleus</i> , <i>Opistognathus</i>	453
<i>maxillosus</i>	139	<i>Squalius</i>	119
<i>meleagris</i>	140	<i>rhombea</i> , <i>Moharra</i>	342
<i>obtusius</i>	140	<i>rhombeus</i> , <i>Gerres</i>	342
<i>ocella</i>	139	<i>Rhombochirus</i>	450
<i>simus</i>	140	<i>osteocheir</i>	450
<i>transmontanus</i>	139	<i>rhombidals</i> , <i>Lebias</i>	181
<i>rhiniichthyoides</i> , <i>Tigoma</i>	141	<i>Otolithus</i>	354
<i>Rhinobatidae</i>	23	<i>Perca</i>	338
<i>Rhinobatos</i>	23	<i>Salma</i>	338
<i>glaucostigma</i>	24	<i>rhomboides</i> , <i>Chetodon</i>	277
<i>horkeli</i>	23	<i>rhomboides</i> , <i>Hemicaranx</i>	271
<i>lentiginosus</i>	23	<i>Lagodon</i>	337
<i>leucorhynchus</i>	24	<i>Sparus</i>	337
<i>percellens</i>	23	<i>Trachinotus</i>	227
<i>productus</i>	24	<i>Rhomboldichthys</i>	217
<i>spinus</i>	24	<i>leopardinus</i>	218
<i>stellio</i>	24	<i>spinus</i>	217
<i>Rhinobatus</i>	23	<i>Rhomboplites</i>	327
<i>columna</i>	23	<i>aurobens</i>	328
<i>electricus</i>	23	<i>Rhomboides</i>	362
<i>glaucostictus</i>	23	<i>Rhombus</i>	266
<i>glaucostigma</i>	24	<i>bahianus</i>	217
<i>horkeli</i>	23	<i>lævis cornubiensis</i>	228
<i>lentiginosus</i>	23	<i>ocellatus</i>	217
<i>leucorhynchus</i>	24	<i>polyspilos</i>	223
<i>marcgravi</i>	23	<i>soleiformis</i>	218
<i>producta</i>	24	<i>Rhoniciscus</i>	334
<i>rhinobatus</i>	23	<i>branioki</i>	334
<i>sen</i>	23	<i>croco</i>	334
<i>spinus</i>	24	<i>ramosus</i>	334
<i>stellio</i>	23	<i>sinuosus</i>	335
<i>undulatus</i>	23	<i>rhothea</i> , <i>Cottus</i>	383
<i>Rhinoberyx</i>	235	<i>Uranidea</i>	383
<i>chryseus</i>	235	<i>rhotheus</i> , <i>Leucosomus</i>	117
<i>Rhinocameridae</i>	33	<i>Rhotheca</i>	289
<i>Rhinocygnus</i>	21	(<i>Rhotheca</i>) <i>blennius</i> , <i>Etheostoma</i>	287
<i>Rhinodon</i>	18	<i>rhotheca</i> , <i>Thoburnia</i>	108
<i>Rhinogoblops</i>	440	<i>rhothecus</i> , <i>Castostomus</i>	108
<i>nichosii</i>	440	<i>rhyncheus</i> , <i>Actipenser</i>	35

List of scientific names—Continued

	Page		Page
Rhynchias	451	Rivulus harti	180
septipinnis	451	heyeri	180
Rhynchichthys	235	isthmensis	180
brachyrhynchus	235	marmoratus	179
Rhynchoeratas	511	tenuis	180
brevirostris	511	roanoka, Alvordius	284
leucorhinus	511	Hadropterus	284
rostratus	511	robalito, Centropomus	306
Rhypticus	316	Robalo	306
maculatus	317	prieto	306
nigripinnis	317	roberti, Exocoetus	201
nigromaculatus	316	Hemirhamphus	198
pitulosus	316	Hyporhamphus	198
subbifrenatus	316	robertsi, Anchoviella	49
xanti	316	Ceratobatis	32
Rhythidostomus	102	Siphostoma	241
richi, Onus	214	Stolephorus	49
richardi, Mesoprion	327	Syngnathus	241
ricei, Cottopsis	386	robertsoni, Astatheros	420
Cottus	386	Cichlasoma	420
richardi, Caranx	272	robini, Hypostomus	160
Hemirhamphus	198	robusta, Agosia	140
Salmo	55	Apocope	140
richardsoni, Acipenser (Huso)	35	Gila	114
Astronesthes	71	Hermosilla	340
Catostomus	103	Moniana	132
Chauliodus	71	Perca	311
Corvina	352	robustum, Megalonema	159
Cottus	385	Moxostoma	111
Hybopsis heterodon	136	robustus, Benthocometes	482
Rhamphocottus	393	Clypeocottus	381
Trachidermis	383	Hippoglossoides	222
richardsoni, Coregonus	64	Ictalurus	152
Richardsonius	117	Mylopharodon	113
balteatus	118	Neobythites	482
lateralis	118	Notropis	132
microdon	118	Ptychocheilus	111
suslawi	118	Zoogoneticus	182
thermophilus	118	Roeus	307
Richia	293	comes	355
brevispina	293	saxatilis	307
Richiella	293	striatus	307
riolata, Bothrocara	477	rochei, Auxis	259
rietus, Xenomystax	81	Scomber	259
riddlei, Torus	188	rodo, Pristipoma	333
rigidus, Pimelodus	159	Roeboidea	98
rijersmaei, Ocyurus	327	guatemalensis	98
rim, Scomber	273	macrolepis	95
rimator, Bathystoma	329, 331	occidentalis	98
Hemulon	331	salvadoris	98
rimensis, Rusciculus	391	Rogenia	41
Rimicola	490	alba	41
beryllinus	490	rogersi, Pimelodus	158
eigenmanni	490	Rhamdia	158
(Rimicola) beryllinus, Göttesox	490	Urolophus	30
rimosus, Etropus	220	Roncador	348
Rineloricaria	162	stearnsi	348
lima	162	roncador, Umbrina	350
ringens, Atopochnus	465	ronchus, Bairdiella	346
Balistes	493	Corvina	346
Phenodon	71	rondeleti, Anthias quartus	326
Sudis	165	Carcharodon	20
Xanthichthys	493	Rondeletia	171
riparium, Holocentrum	236	bicolor	171
risso, Alpinmaris	163	Rondeletidae	171
rossoanus, Notacanthus	76	rondeletti, Exocoetus	200
Rissola	485	Exonantes	200
marginata	485	Orthogoriscus	508
ritteri, Lampanyctus	167	Sargus	338
Pleuronichthys	225	Scombresox	197
Xeneretmus	376	Xiphias	264
Xenopyxis	356	Ronquilius	452
rivalis, Salmo	60	jordani	452
riverendi, Cyprinodon	181	roosevelti, Salmo	59
Trifarous	181	rosacea, Mycteroperca	314
riwolana, Seriola	280	rosaceus, Cymatogaster	410
riwolanus, Zonichthys	280	Epinephelus	314
riularis, Salmo	58	Mesoprion	326
riulatus, Curritus	358	Paraliparis	404
Diodon	502	Sebastes	367
Rivulus	179	Sebastomus	367
brunneus	180	Zalemibus	410
chucunaque, sucubi	180	rosei, Clevelandia	445
chucunaque	180	Hemirhamphus	198
cylindraceus	179	Hyporhamphus	198
flabellicauda	180	Troglichthys	195
godmanni	180	rosarium, Acipenser (Huso)	35

List of scientific names—Continued

	Page		Page
rosei, Salmo.....	59	rubrifrons, Alburnus.....	124
roseipinnis, Lythurus.....	128	Erinemus.....	138
Notropis.....	128	Nocomus.....	138
roseus, Cryptotomus.....	431	Notropis.....	124, 128
Hydrophlox.....	127	Zygonectes.....	177
Luxilus.....	127	rubrilateralis, Cocusius plumbeus.....	116
Merulmus.....	407	rubripinna, Cyprinella.....	131
Prionotus.....	407	rubripinne, Sparsoma.....	433
Rosacola alexandri.....	365	rubripinnis, Argyræus.....	128
babeoeki.....	365	Scarus.....	433
miniatu.....	365	rubrirostris, Catetys.....	480
pinniger.....	365	rubrivinctus, Hispanicus.....	368
rosipes, Novaculichthys.....	430	Sebastichthys.....	368
Xyrichthys.....	430	rubrofluvialis, Cyprinodon.....	181
rosispinis, Raia.....	26	Cyprinodon bovinus.....	181
Raja.....	26	rubro-punctatus, Salaria.....	463
rossi, Salmo.....	60	rubropunctatus, Scartichthys.....	463
Salvelmus.....	60	rubrum, Chorististum.....	315
rostrata, Albula.....	40	rudis, Centrochromis.....	415
Aldrovandia.....	75	Glyphidodon.....	415
Anguilla.....	77	rufa, Morone.....	307
Loricaria.....	161, 162	Perca.....	236
Muraena.....	77	rufilineatus, Nothonotus.....	290
Platessa.....	227	Poecilichthys.....	290
Sturisoma.....	161	rufipinnis, Exocoetus.....	201
rostratum, Astatheros.....	421	Exonantes.....	201
Holocentrum.....	236	rufum, Acutomentum.....	366
rostratus, Apterionotus.....	101	rufus, Balistes.....	493
Canthigaster.....	500	Bodianus.....	425
Gymnothorax.....	91	Centropistes.....	318
Halosaurus.....	75	Holocentrus.....	236
Heros.....	421	Labrus.....	425
Heterostichus.....	458	Onos.....	208
Holocentrus.....	235	Sebastodes.....	366
Notacanthus.....	76	rugifer, Ophichthys.....	88
Phycus.....	213	rugosus, Platichthys.....	226
Polyacanthonotus.....	76	rupertianus, Aclpenser.....	35
Rhynchoceratus.....	511	rupestre, Etheostoma (Rhothoea).....	290
Scarus.....	435	Nanostoma.....	290
Squalus.....	19	rupestris, Ambloplites.....	303
Stenarchus.....	101	Arbacia.....	491
Tetrodon.....	500	Bodianus.....	303
Zeus.....	275	Coryphæna.....	205
rotengulus, Leuciscus.....	117	Coryphænoides.....	205
rothrocki, Notogrammus.....	469	Gadus.....	210
Porodinus.....	470	Gobiesox.....	491
rousseau, Syngnathus.....	241	Sebastichthys.....	368
rubellio, Eximia.....	391	Sebastomus.....	368
Greeleya.....	391	Serranus.....	313
rubellus, Alburnus.....	123	Xiphister.....	469
rubens, Tautoga.....	424	Rupiscartes.....	463
rubber, Apogon.....	295	atlanticus.....	463
Cephapholis fulvus.....	309	rupiscartes, Moxostoma.....	111
Coregonus.....	163	Scartomyzon.....	111
Elaphotoxon.....	271	ruppelli, Monacanthus.....	494
Epinephelus.....	315	ruptus, Julis.....	427
Gadus.....	214	Ruscariops.....	378
Gymnocephalus.....	309	creaseri.....	378
parvus, Sebastes.....	368	Ruscarius.....	378
Rutilus.....	126	manyi.....	378
Scomber.....	271	ruscarius, Gurnardius.....	406
Sebastes.....	368	Prionotus.....	406
ruberrimus, Astryanax.....	97	Rusciculus.....	391
Gunnellus.....	468	rimensis.....	391
Sebastodes.....	368	Rusulus.....	390
Sebastopyr.....	368	saburæ.....	390
rubescens, Algansea.....	116	russula atlantica, Scorpaena.....	370
Auchenopterus.....	457	russula, Scorpaena.....	371
Conger.....	79	ruthenus, Aclpenser.....	34
Exocoetus.....	201	rutilla, Moniana.....	131
Exonantes.....	201	Rutilus amblops.....	133
Steinergia.....	266	anomalus.....	146
rubicundus, Aclpenser.....	35	columbianus.....	122
Glyphisodon.....	415	compressus.....	123, 128
Hypsypops.....	415	melanurus.....	110
rubiginosus, Sicyases.....	490	oregonensis.....	122
Rubio volador.....	407	plagyrrus.....	128
rubio, Prionotus.....	407	ruber.....	126
rubra, Garmannia.....	445	storenianus.....	137
Liopropoma.....	315	rutteri, Neoliparis.....	399
Sciæna.....	235	Ruvettus.....	261
rubre, Liopropoma.....	315	pretiosus.....	261
rubri-cauda, Pomotis.....	299	terminckii.....	261
rubricroceus, Hybopsis.....	126, 127	Rypticus.....	316
Hydrophlox.....	127	arenatus.....	316

List of scientific names—Continued

	Page		Page
<i>Rrypticus bicolor</i>	316	<i>Salmo canadensis</i>	60
<i>coriaceus</i>	316	<i>canis</i>	54
<i>maculatus</i>	316	<i>callaris</i>	61
<i>aponaceus</i>	316	<i>campbelli</i>	61
<i>xanti</i>	316	<i>canadensis</i>	60
<i>sabalo</i> , <i>Arges</i>	169	<i>canis</i>	54
<i>sabanensis</i> , <i>Hippoglossina</i>	223	<i>carinatus</i>	56
<i>sabina</i> , <i>Trygon</i>	29	<i>clarkii</i>	56
<i>sabina</i> , <i>Hybopsis</i>	135	<i>declivifrons</i>	57
<i>Notropis</i>	135	<i>jordani</i>	57
<i>sabinus</i> , <i>Amphotistius</i>	29	<i>tahoensis</i>	57
<i>saburus</i> , <i>Chasmodes</i>	462	<i>clupeaformis</i>	64
<i>Rusulus</i>	390	<i>confinis</i>	59
<i>Saccopharyngida</i>	95	<i>confluentus</i>	55
<i>Saccopharynx</i>	95	<i>consuetus</i>	54
<i>ampullaceus</i>	95	<i>cooperi</i>	55
<i>chordatus</i>	95	(<i>Coregonus</i>) <i>harengus</i>	62
<i>flagellum</i>	95	<i>lucidus</i>	62
<i>saccostoma</i> , <i>Lasiognathus</i>	510	<i>sapidissimus</i>	64
<i>sacrifici</i> , <i>Rhamdia</i>	158	<i>tullibee</i>	64
<i>sadina</i> , <i>Clupea</i>	46	<i>crenatus</i>	57
<i>Etrumeus</i>	46	<i>curius</i>	61
<i>Safole</i>	305	<i>declivifrons</i>	57
<i>taeniura</i>	305	<i>dermatinus</i>	54
<i>sagamius</i> , <i>Ebisus</i>	373	<i>edentulus</i>	100
<i>Sagenichthys</i>	354	<i>eperlanus</i>	67
<i>mordax</i>	354	<i>eremogenes</i>	58
<i>sagitta</i> , <i>Amblyopus</i>	448	<i>erythrorhynchus</i>	61
<i>Poecilichthys</i>	293	<i>evermanni</i>	59
<i>Torrentaria</i>	293	<i>fario</i>	55
<i>Tylosurus</i>	195	<i>foetens</i>	163
<i>Tyntilastes</i>	448	<i>fontinalis</i>	60
<i>sagittula</i> , <i>Euctenogobius</i>	441	<i>gairdneri beardsliei</i>	57
<i>Gobionellus</i>	441	<i>crenatus</i>	57
<i>Saguaraca</i>	235	<i>gilberti</i>	58
<i>saida</i> , <i>Boreogadus</i>	209	<i>shasta</i>	58
<i>Gadus</i>	209	<i>stonei</i>	58
<i>saira</i> , <i>Cololabis</i>	198	<i>gairdneri</i>	58
<i>Salar</i>	55	<i>gibber</i>	54
<i>lewisii</i>	55	<i>gibbsii</i>	57
<i>virginalis</i>	57	<i>gilberti</i>	58
<i>salar</i> , <i>Salmo</i>	55, 56	<i>gloveri</i>	56
<i>Salapia</i>	460	<i>gorbuscha</i>	54
<i>Salarias alticus</i>	463	<i>grömlandicus</i>	66
<i>atlanticus</i>	463	<i>harengus</i>	61
<i>chiostictus</i>	463	<i>hearnei</i>	60
<i>margaritaceus</i>	463	<i>henshawi</i>	57
<i>rubro-punctatus</i>	463	<i>hudsonicus</i>	60
<i>salarias rubropunctatus</i>	463	<i>immaculatus</i>	60
<i>textilis</i>	463	<i>irideus</i>	58
<i>vomerinus</i>	463	<i>iridia</i>	58
<i>salarias rubropunctatus</i> , <i>Salarias</i>	463	<i>japonicus</i>	54
<i>Salarichthys</i>	463	<i>jordani</i>	57
<i>textilis</i>	463	<i>kennerlyi</i>	54, 55
<i>Salema aurata</i>	339	<i>keta vel kayko</i>	54
<i>Salema</i>	337	<i>kisutch</i>	54, 55
<i>rhomboidalis</i>	338	<i>laevigatus</i>	61
<i>tridens</i>	338	<i>lagocephalus</i>	54
<i>saliens</i> , <i>Blennius</i>	463	<i>lactor</i>	64
<i>Oligopilites</i>	277	<i>lavaretus</i>	64
<i>Scomber</i>	277	<i>lewisii</i>	56
<i>Thunnus</i>	259	<i>lordii</i>	61
<i>salin</i> , <i>Sparus</i>	338	<i>lycodon</i>	55
<i>salix</i> , <i>Atherinichthys</i>	248	<i>macdonaldi</i>	57
<i>salix</i> , <i>Algansea</i>	116	<i>macdonaldi</i>	61
<i>Ceratichthys</i>	116	(<i>Mallotus</i>) <i>pacificus</i>	66
<i>salley</i> , <i>Archomenidia</i>	248	<i>malma</i>	61
<i>Atherinichthys</i>	248	<i>marogravi</i>	100
<i>Salmo</i>	55	<i>marstoni</i>	61
<i>adironiacus</i>	59	<i>mas</i>	56
<i>agassizii</i>	60	<i>masoni</i>	53
<i>agua-bonita</i>	59	<i>melampterus</i>	55
<i>alpes</i>	60	<i>miltschutch</i>	54
<i>alleganensis</i>	59	<i>miltschutch</i>	54
<i>amethystinus</i>	59	<i>miltschutch</i>	54
<i>ansonii</i>	55	<i>mülleri</i>	170
<i>angularum</i>	58	<i>mykiss</i>	56
<i>arabatch</i>	55	<i>agua-bonita</i>	59
<i>arcuatus</i>	66	<i>macdonaldi</i>	57
<i>arcturus</i>	61	<i>myops</i>	163
<i>bairdii</i>	61	<i>namaycush</i>	59
<i>bathocetor</i>	57	<i>naresi</i>	61
<i>beardslei</i>	57	<i>nerka</i>	55
<i>bouvieri</i>	56	<i>nelsoni</i>	59
<i>brevicauda</i>	56	<i>nerka</i>	55
		<i>newberryi</i>	58

List of scientific names—Continued

	Page		Page
<i>Salmo nigrescens</i>	60	<i>salmandrus</i> , <i>Alburnops</i>	132
<i>nitidus</i>	60	<i>Hudsonius</i>	132
<i>nummifer</i>	61	<i>salvadoris</i> , <i>Rhomboides</i>	98
<i>omiscornaycus</i>	214, 215	<i>salvatoris</i> , <i>Poecilia</i>	192
<i>omul</i>	61	<i>Salvelini</i>	60
<i>oquassa</i>	61	<i>alpinus</i>	60
<i>orientalis</i>	55	<i>Salvelinus</i>	60
(<i>Osmerus</i>) <i>olidus</i>	68	<i>agassizii</i>	60
<i>otsego</i>	64	<i>arcturus</i>	61
<i>ouananiche</i>	56	<i>aureolus</i>	60
<i>pallidus</i>	59	<i>fontinalis</i>	60
<i>parki</i>	61	<i>malma</i>	61
<i>paucidens</i>	55	<i>spectabilis</i>	61
<i>penshinensis</i>	56	<i>marstoni</i>	61
<i>pleuriticus</i>	57	<i>naresi</i>	61
<i>pluvius</i>	61	<i>oquassa</i>	61
<i>proteus</i>	54	<i>rossii</i>	60
<i>purpuratus</i>	56	<i>spectabilis</i>	61
<i>bouvieri</i>	56	<i>stagnalis</i>	60
<i>quinnat</i>	55	<i>timagamensis</i>	60
<i>regalis</i>	56	<i>salvelinus</i> , <i>Salmo</i>	60
<i>richardi</i>	55	<i>salvini</i> , <i>Agonostomus</i>	254
<i>rivalis</i>	60	<i>Cotylopus</i>	448
<i>rivularis</i>	58	<i>Heros</i>	418
<i>roosevelti</i>	59	<i>Parapetenis</i>	418
<i>rosel</i>	59	<i>Pimelodus</i>	167
<i>rossii</i>	60	<i>Rhamdia</i>	167
<i>salar</i>	55, 56	<i>Sicydium</i>	448
<i>ouananiche</i>	56	<i>sancta-helenæ</i> , <i>Caranx</i>	270
<i>salvelinus</i>	60	<i>Decapterus</i>	270
<i>sanguinolentus</i>	55	<i>Gymnothorax</i>	92
<i>saurus</i>	163	<i>Muraena</i>	92
<i>scouleri</i>	54	<i>sancta-luceæ</i> , <i>Corvula</i>	345
<i>sebago</i>	56	<i>Vacuqua</i>	345
<i>shasta</i>	58	<i>sancta-marthæ</i> , <i>Vomer</i>	275
<i>siscowet</i>	60	<i>sancta-petri</i> , <i>Vomer</i>	275
<i>siskawitz</i>	60	<i>sancta-rosæ</i> , <i>Ulvicola</i>	467
<i>smaragdus</i>	58	<i>sancta-eustatii</i> , <i>Callionymus</i>	450
<i>socialis</i>	66	<i>sancti-laurentii</i> , <i>Engyophrys</i>	220
<i>spectabilis</i>	61	<i>sancti-martini</i> , <i>Gobiesox</i>	490
<i>spilurus</i>	57	<i>sancti-pauli</i> , <i>Holocentrum</i>	236
<i>stagnalis</i>	60	<i>Holocentrus</i>	236
<i>stomias</i>	57	<i>Sander</i>	282
<i>stonei</i>	58	<i>sandvicensis</i> , <i>Gobius</i>	440
<i>striatus</i>	54	<i>Pimelepterus</i>	339
<i>symmetricus</i>	59	<i>sandwichensis</i> , <i>Balistes</i>	493, 494
<i>tahoensis</i>	57	<i>sanguifluus</i> , <i>Poecilichthys</i>	290
<i>tapduma</i>	55	<i>sanguineus</i> , <i>Antennarius</i>	505
<i>thymallus</i>	66	<i>Apodichthys</i>	467
<i>toma</i>	59	<i>Holocentrus</i>	363
<i>truncatus</i>	58	<i>Pythonichthys</i>	90, 91
<i>trutta</i>	55	<i>Sicyases</i>	490
<i>tschawytseha</i>	55	<i>sanguinolentus</i> , <i>Pomotis</i>	300
<i>tschawytsehaformis</i>	54	<i>Salmo</i>	55
<i>tsuppitch</i>	55	<i>santa-ana</i> , <i>Gasterosteus</i>	238
<i>tudes</i>	61	<i>santa-ana</i> , <i>Notolepidomyzon</i>	104
<i>tumbil</i>	164	<i>Pantosteus</i>	104
<i>umbra</i>	60	<i>santamarie</i> , <i>Momana</i>	131
<i>ursinus</i>	60	<i>Notropis</i>	131
<i>utah</i>	56	<i>sapidissima</i>	42
<i>virginalis</i>	57	<i>Clupea</i>	42
<i>warreni</i>	55	<i>sapidissimus</i> , <i>Salmo</i> (<i>Coregonus</i>).....	64
<i>whitei</i>	59	<i>saponaceus</i> , <i>Anthias</i>	316
<i>salmoides</i> , <i>Labrus</i>	297, 298	<i>Rypiticus</i>	316
<i>Micropterus</i>	297	<i>sara</i> , <i>Cybtium</i>	256
<i>salmonea</i> , <i>Ericara</i>	53, 54	<i>saragus</i> , <i>Lepodus</i>	265
<i>Lepomis</i>	298	<i>Saranus apiarius</i>	308
<i>Perca</i>	282	<i>Sarchurus</i>	36
<i>salmoneus</i> , <i>Esox</i>	163, 172	<i>argenteus</i>	36
<i>Mugil</i>	40	<i>vittatus</i>	36
<i>Scombrocottus</i>	373	<i>Sarcidium</i>	142
<i>salmonicolor</i> , <i>Merulinus</i>	407	<i>scopiferum</i>	142
<i>Salmonidæ</i>	54	<i>Sarcura</i>	23
<i>Salmonoidel</i>	54	<i>Sarda</i>	259
<i>Salmo perca</i>	214	<i>chiliensis</i>	259
<i>pellucida</i>	215	<i>lineolata</i>	259
<i>Salmo perca</i>	214	<i>orientalis</i>	259
<i>Salpa purpurascens variegata</i>	327	<i>sarda</i>	259
<i>saltans</i> , <i>Chorinemus</i>	277	<i>velox</i>	259
<i>saltator</i> , <i>Hemiramphus</i>	199	<i>sarda</i> , <i>Sarda</i>	259
<i>Scomberoides</i>	277	<i>Scomber</i>	259
<i>saltatorius</i> , <i>Alticus</i>	463	<i>Sardina</i>	43
<i>saltatrix</i> , <i>Gasterosteus</i>	280	<i>pseudohispanica</i>	43
<i>Perca</i>	280	<i>sardina</i> , <i>Atherina</i>	247
<i>Pomatomus</i>	280	<i>Harengula</i>	43

List of scientific names—Continued

	Page		Page
sardina, <i>Melaniris</i>	249	saxicola, <i>Pleuronectes</i>	228
Menidia.....	247	Sebastichthys.....	367
Thyrina.....	249	say, <i>Ampholis</i>	28
Sardinella.....	43	Raja.....	28
anchovia.....	43	sayanus, <i>Aphredoderus</i>	215
aurita.....	43	Scolopsis.....	215
Sardinia.....	43	Sayris.....	197
anchovia.....	43	bimaculatus.....	197
cerulea.....	43	hians.....	197
pseudohispanica.....	43	recurvirostra.....	197
sargoides, <i>Chaetodon</i>	415	serrata.....	197
Sargosomus.....	412	scaber, <i>Antennarius</i>	505
fluviatilis.....	412	Centropomus.....	306
Sargus.....	338	Chironectes.....	505
ambassis.....	336	Hexagrammus.....	374
argenteus.....	338	scabra, <i>Echidna</i>	94
aries.....	338	Lophius aculeatus.....	505
caribaeus.....	338	Trinectes.....	229
caudimacula.....	338	scabrata, <i>Raja</i>	25
flavolineatus.....	338	Raja.....	25
holbrookii.....	338	scabriceps, <i>Cylindrosteus</i>	37
humeralimaculatus.....	338	Paranotrops.....	125
ovis.....	338	Photogenis.....	125
raucus.....	338	scabripinnis, <i>Astyanax</i>	97
rondeletii.....	338	Tetragonopterus.....	97
sargus.....	338	Scafaria.....	264
tridens.....	338	Scaphirhynchops.....	35
variegatus.....	338	Scaphirhynchus.....	35
vitula.....	338	platyrhynchus.....	35
sargus, <i>Diplodus</i>	338	rafinesquei.....	35
Sargus.....	338	scaphiurus, <i>Opisthognathus</i>	452
Sparus.....	338	scaphopsis, <i>Ocelorhynchus</i>	207
Sarothrodus.....	359	Macrurus (<i>Ocelorhynchus</i>).....	207
amplexocollis.....	360	Scaphorhynchus.....	35
ateniatus.....	360	scapularis, <i>Pristipoma</i>	332
maculo-cinctus.....	360	scapularis, <i>Anistremus</i>	332
nigrirostris.....	360	Strongylura.....	195
Sarritor.....	395	Tylosurus.....	195
frenatus.....	395	Scaridae.....	433
leptorhynchus.....	395	Scarcina.....	262
sarritor, <i>Prionotus</i>	407	argyrea.....	262
Satanoperca.....	424	Scartella.....	461
(Satanoperca) <i>crassilabris</i> , <i>Geophagus</i>	424	brevipinnis.....	461
satrilius, <i>Neodinus</i>	453	microstoma.....	461
Pterognathus.....	453	Scartes.....	463
satis parva et glabra, <i>Pleuronectes</i>	217	scartes, <i>Fundulus</i>	177
saturatus, <i>Ambloplites</i>	343	Scartichthys.....	463
Rhinoscion.....	343	rubropunctatus.....	463
Saurenhelys.....	82	Scartomyzon.....	111
cancrivora.....	82	cervinus.....	111
Saurida.....	164	rupiscartes.....	111
mediterranea.....	163	Scarus.....	432, 433
nebulosa.....	164	abildgardii.....	431
Saurus.....	163	acutus.....	434
anolis.....	164	alternans.....	434
argenteus.....	277	amplus.....	432
brevirostris.....	163	aracanga.....	433, 434
griseus.....	163	atomarius.....	431
intermedius.....	164	aurofrenatus.....	432
longirostris.....	163	bollmani.....	433
lucoiceps.....	163	brachialis.....	433
mexicanus.....	163	caeruleus.....	434
myops.....	163	(Catesby).....	432
spixianus.....	163	chloris.....	432
truncatus.....	163	chrysopterus.....	432
saurus, <i>Elops</i>	39	circumnotatus.....	433
Esox.....	197	coccineus.....	432
Oligoplites.....	277	coelestinus.....	435
Salmo.....	163	compressus.....	434
Scomber.....	277	croicensis.....	433, 434
Scomberesox.....	197	cuzamilae.....	434
Synodus.....	163	dentiens.....	431
saururii, <i>Brama</i>	265	diadema.....	433
Taractes.....	265	distinctus.....	432
savanna, <i>Muraena</i>	78	emarginatum.....	433
Muraenesox.....	78	emblematicus.....	435
saxatilis, <i>Catulus</i>	11	erythrinoides.....	432
Chaetodon.....	414	evermanni.....	434
Glyphidodon.....	415	flavescens.....	433
Johnius.....	350	flavomarginatus.....	434
Mentidarrhus.....	350	flavo-marginatus.....	434
Perca.....	307	frondosus.....	433
Poecilichthys.....	287	gibbus.....	433
Roccus.....	307	gnathodus.....	434
saxicola, <i>Eosebastes</i>	367	guacamaia.....	435
Gymnothorax ocellatus.....	93	holocyaneos.....	434

List of scientific names—Continued

	Page		Page
<i>Scarus hoplomystax</i>	432	<i>Sciæna ensifera</i>	345
humeralis.....	433	ericymba.....	346
insulæ-santæ-cruis.....	434	fusca.....	352
lacrimosus.....	432	gigas.....	352
lateralis.....	432	grisea.....	352
maschalespilos.....	433	istia.....	346
melanotis.....	432	imberbis.....	348
miniofenatus.....	432	imiceps.....	348
novei.....	434	lineata.....	307
nuchalis.....	434	maculata.....	436
obtusus.....	434	multofasciata.....	349
oxybrachius.....	432	oscitans.....	347
pepo.....	433	oseula.....	352
perrico.....	435	plumieri.....	333
pleianus.....	435	rubra.....	235
punctulatus.....	433	sciæra.....	348
quadrispinosus.....	434	tahmel.....	339
radians.....	432	undecimalis.....	305
rostratus.....	435	<i>Sciænidae</i>	99, 344
rubripinnis.....	433	<i>Sciænops</i>	348
spinidens.....	431, 432	ocellata.....	348
squalidus.....	433	sciæra, <i>Sciæna</i>	348
strigatus.....	433	Serraria.....	284
superbus.....	434	scierus, <i>Hadropterus</i>	284
tæniopterus.....	433	Ophioscion.....	348
trilobatus.....	434	serrula, <i>Hadropterus</i>	284
trispinosus.....	434	scintillans, <i>Zastomias</i>	70
truncatus.....	433	Scitaliscus.....	473
turchesius.....	435	scitula, <i>Anchovia</i>	49
vaigiensis.....	431	Anchoviella.....	49
vetula.....	433, 434	scituluopsis, <i>Synodus</i>	164
virginalis.....	433	scitulus, <i>Merulinus</i>	407
virens.....	433	Prionotus.....	407
viridis.....	432	scurus, <i>Diplectrum</i>	318
seaurus, <i>Lycodopsis</i>	472	Hæmulon.....	330
scepticus, <i>Minnilus</i>	123	Sparus.....	329, 330
Notropis.....	123	<i>Sclerodermi</i>	491
Triglops.....	380	<i>Sclerognathus</i>	102, 103
<i>Schedophilopsis</i>	269	cyprinus.....	102
spinosus.....	269	cyprinella.....	102, 103
<i>Schedophilus</i>	268	meridionalis.....	104
heathi.....	268	<i>Sclerotis</i>	300
medusophagus.....	268	miniatus.....	300
<i>Schilbeodes</i>	155	scoffieldi, <i>Anchoviella</i>	49
gyrinus.....	155	Stolephorus.....	49
Schipa.....	34	<i>Scolecocoma</i>	9
Schistorus.....	310	<i>Scoliodon</i>	14
Schizophallus.....	184	lalandi.....	15
schmardæ, <i>Dasyatis</i>	28	laticaudus.....	14
Pastinachus.....	27	longurio.....	15
schmidti, <i>Edriolychnus</i>	508	porosus.....	15
schmidti, <i>Hoplunnis</i>	81	terra-novæ.....	14, 15
schneideri, <i>Ophichthys</i>	88	<i>scolopacea</i> , <i>Nemichthys</i>	83
Pocilia.....	190	<i>scolopaceus</i> , <i>Nemichthys</i>	83
schœpfi, <i>Cylichthys</i>	502	<i>scolopax</i> , <i>Balistes</i>	245, 246
Diodon.....	502	Centriscus.....	245
schœpfi, <i>Balistes</i>	495	Macrorhamphosus.....	246
Ceratacanthus.....	495	<i>Scolopsis sayanus</i>	215
scholaris, <i>Thyrsites</i>	261	<i>Scomber</i>	257
schomburgkii, <i>Pempheris</i>	323	alalunga.....	260
schoneveldii, <i>Cataphractus</i>	395	albacores.....	260
schrankii, <i>Hæmulon</i>	330	alletteratus.....	258
schypha, <i>Antaceus</i>	34	auratus.....	258
<i>Sciadeichthys</i>	148	australasicus.....	258
emphysetus.....	148	balantiophthalmus.....	271
proops.....	148	bisus.....	259
troscheli.....	148	boops.....	270
<i>Sciades pictus</i>	148	carangus.....	273
troscheli.....	148	chrysurus.....	276
(<i>Sciades</i>) <i>emphysetus</i> , <i>Bagrus</i>	148	colias.....	257, 258
sciadicus, <i>Fundulus</i>	177	crumenophthalmus.....	270, 271
Zygonectes.....	177	crysos.....	272
<i>Sciadonus</i>	483	dekayi.....	258
pedicellaris.....	483	diego.....	258
<i>Sciæna</i>	348	fasciatus.....	279
aluta.....	346	germo.....	260
aquila.....	61	gladius.....	263
argentea.....	334	gracilis.....	258
argyrea.....	341	grex.....	258
armata.....	273	gunneri.....	217
(<i>Bairdiella</i>) <i>bedotis</i>	346	hippos.....	273
caprodes.....	282, 283	ignobilis.....	273
cirrhusa.....	350	japonicus.....	258
coro.....	333	lacertus.....	258
croker.....	349	latus.....	276
edwardi.....	353	macrophthalmus.....	258

List of scientific names—Continued

	Page		Page
<i>Scomber maculatus</i>	257, 258	<i>Scorpena calcarata</i>	371
<i>mediterraneus</i>	259	<i>colesi</i>	370
<i>pelagicus</i>	264	<i>cristulata</i>	372
<i>pelamides</i>	258	<i>dactyloptera</i>	370
<i>pelamis</i>	258	<i>dactylopterus</i>	370
<i>pelamitus</i>	259	<i>echinata</i>	372
<i>pelamys</i>	259	<i>grandicornis</i>	371
<i>plumieri</i>	271	<i>guttata</i>	371
<i>pneumatophorus</i>	257, 258	<i>histrio</i>	371
<i>quadripuntatus</i>	258	<i>inermis</i>	371
<i>regalis</i>	257	<i>massiliensis</i>	308
<i>rim</i>	273	<i>mystes</i>	371
<i>rochei</i>	258, 259	<i>nuttingi</i>	371
<i>ruber</i>	271	<i>ocephalalis</i>	371
<i>saliens</i>	277	<i>pannosa</i>	371
<i>sarda</i>	259	<i>picta</i>	370
<i>saurus</i>	277	<i>plumieri</i>	370
<i>scombrus</i>	257	<i>polylepis</i>	370
<i>sloani</i>	280	<i>porcus</i>	370
<i>speciosus</i>	273	<i>rascacio</i>	370
<i>thazard</i>	259	<i>russula</i>	371
<i>thynnus</i>	259	<i>atlantica</i>	370
<i>trachurus</i>	270	<i>(Sebastoplus) sierra</i>	372
<i>undulatus</i>	258	<i>sonora</i>	371
<i>vernalis</i>	257	<i>stearnsi</i>	371
<i>zonatus</i>	277	<i>tridecimspinosa</i>	370
<i>Scomberesocidae</i>	197	<i>Scorpenichthyidae</i>	382
<i>Scomberesox</i>	197	<i>Scorpenichthys</i>	382
<i>brevirostris</i>	198	<i>lateralis</i>	377
<i>camperi</i>	197	<i>marmoratus</i>	382
<i>equirostrum</i>	197	<i>Scorpenidae</i>	363
<i>forsteri</i>	197	<i>Scorpenodes</i>	370
<i>rondeletii</i>	197	<i>xyris</i>	370
<i>saurus</i>	197	<i>Scorpidae</i>	358
<i>scutellatum</i>	197	<i>scorpioides, Acanthocottus</i>	386
<i>storeri</i>	197	<i>Cottus</i>	386
<i>scomberus, Esox</i>	173	<i>Scorpius californiensis</i>	358
<i>Scomberoides</i>	260	<i>scorpius, Acanthocottus</i>	386
<i>saltator</i>	277	<i>Cottus</i>	386
<i>scomberoides, Coryphæna</i>	265	<i>scouleri, Salmo</i>	54
<i>Scomberomorus</i>	257	<i>scovelli, Siphostoma</i>	242
<i>concolor</i>	257	<i>Syrticus</i>	242
<i>maculatus</i>	257	<i>scovelli, Etheostoma</i>	293
<i>plumieri</i>	257	<i>scripps, Otophidium</i>	485
<i>regalis</i>	257	<i>scripta, Balistes</i>	495
<i>sierra</i>	257	<i>Osbeckia</i>	495
<i>Scombridae</i>	257	<i>scriptus, Balistes</i>	495
<i>Scombriformes</i>	256	<i>Gymnothorax</i>	92
<i>Scombrocottus</i>	373	<i>Pseudoscopelus</i>	452
<i>salmonus</i>	373	<i>scrutator, Isesthes</i>	462
<i>Scombroptidae</i>	297	<i>scuderi, Hæmulon</i>	330
<i>Scombrus alalonga</i>	260	<i>scutata, Echeuis</i>	449
<i>scombrus, Scomber</i>	257	<i>scutator, Belone</i>	196
<i>Scopelogys</i>	170	<i>scutatus, Elurichthys</i>	148
<i>dispar</i>	170	<i>Bagre</i>	148
<i>tristis</i>	170	<i>scutellatum, Scomberesox</i>	197
<i>Scopelus caninianus</i>	169	<i>Scutia</i>	95
<i>Scopelus</i>	169	<i>scuticaris, Bascanichthys</i>	86
<i>affine</i>	169	<i>Sphagebranchus</i>	86
<i>andree</i>	168	<i>scutiger, Icelus</i>	378
<i>arcticus</i>	170	<i>Rastrinus</i>	378
<i>benoiti</i>	169	<i>scutum, Achirus</i>	229
<i>cocoi</i>	168, 169	<i>Solea</i>	229
<i>crassiceps</i>	233	<i>Scylliorhinus</i>	11
<i>engraulis</i>	168	<i>scylla, Exocoetus</i>	201
<i>glacialis</i>	170	<i>Hybopsis</i>	135
<i>gracilis</i>	169	<i>Scylliorhinae</i>	11
<i>hygomi</i>	169	<i>Scylliorhinus</i>	11
<i>madelrensis</i>	167	<i>catulus (canicula)</i>	13
<i>metapoclampus</i>	168	<i>cephalus</i>	11
<i>mizolepis</i>	233	<i>indicus</i>	12
<i>muelleri</i>	170	<i>profundorum</i>	12
<i>rarus</i>	168	<i>retifer</i>	11
<i>scopifer, Phenacobius</i>	142	<i>Scyllum</i>	11
<i>scopiferum, Sarcidium</i>	142	<i>capense</i>	11
<i>scopiferus, Notropis</i>	132	<i>laticeps</i>	11
<i>scops, Gnathypops</i>	453	<i>retiferum</i>	11
<i>Scorpena</i>	370	<i>scyllum, Triakis</i>	14
<i>agassizi</i>	371	<i>Scymnus rostratus</i>	21
<i>albifimbria</i>	371	<i>Scyris</i>	274
<i>albofasciata</i>	371	<i>analis</i>	274
<i>americana</i>	393	<i>indica</i>	274
<i>americanus</i>	393	<i>Scytalichthys</i>	89
<i>atlantica</i>	370	<i>miurus</i>	89
<i>bergii</i>	372	<i>Scytalina</i>	473
<i>brasilensis</i>	371	<i>cerdale</i>	473
<i>bufo</i>	370	<i>Scytalinidae</i>	473

List of scientific names—Continued

	Page		Page
Scytalinus.....	473	Sebastodes semicinctus.....	367
Scytalophis.....	87, 88	serranoides.....	365
magnoculus.....	87	swifti.....	367
sebagi, Salmo.....	56	wilsoni.....	366
Sebastes.....	363	Sebastolobus.....	364
auriculatus.....	368	alascanus.....	364
bougainvillei.....	370	altivelis.....	364
caurinus.....	369	Sebastomus.....	367, 368
elongatus.....	368	ayresii.....	367
fasciatus.....	363, 369	chlorostictus.....	367
glaucus.....	364	constellatus.....	367
helvomaculatus.....	367	eos.....	368
imperialis.....	370	gilhi.....	368
macrochir.....	364	rhodochloris.....	367
macrocephalus.....	372	rosaceus.....	367
marinus.....	363	rupestris.....	368
melanops.....	364	umbrosus.....	367
minutus.....	370	Sebastopsis.....	370
nebulosus.....	369	xyris.....	370
nematophthalmus.....	322	Sebastopyr.....	368
nigrocinctus.....	370	ruberrimus.....	368
ovalis.....	365	Sebastosomus.....	364
paucispinis.....	364	aleuticus.....	365
regulus.....	363	ciliatus.....	365
rosaceus.....	367	flavidus.....	364
ruber.....	368	melanops.....	365
parvus.....	368	mystinus.....	365
viviparus.....	364	pinniger.....	365
Sebastichthys.....	369, 370	serranoides.....	365
alutus.....	366	simulans.....	365
atrovirens.....	365	sco, Luxilus.....	115
aurora.....	366	Sectator.....	340
carinatus.....	369	ocyurus.....	340
chlorostictus.....	369	sectatrix, Kyphosus.....	340
chrysomelas.....	369	Perca.....	340
constellatus.....	367	marina.....	340
diploproa.....	367	sector, Serrivomer.....	84
entomelas.....	366	secundodorsalis, Thunnus.....	260
fasciolaris.....	369	secundus, Caranx.....	272
introniger.....	366	sedentarius, Chaetodon.....	360
levis.....	368	seemani, Arius.....	150
maliger.....	369	Galeichthys.....	150
miniatus.....	365	Selache.....	20
mystinus.....	364, 365	Selachii.....	10
nigrocinctus.....	370	selachops, Apterichthys.....	84
proriger.....	366	Sphagebranchus.....	84
var. brevispinis.....	366	Selachostomi.....	36
rastralliger.....	369	Selanonius.....	19
rhodochloris.....	367	walkeri.....	19
rubrivinctus.....	368	Selar.....	270
rupestris.....	368	Selma.....	367, 339
saxicola.....	367	aurata.....	339
serriceps.....	369	Selenaspis.....	148
sinensis.....	368	dowii.....	148
umbrosus.....	367	Selene.....	275
veixillaris.....	369	argentea.....	275
zacentrus.....	368	cerstedii.....	275
Sebastocarus.....	369	quadrangularis.....	358
serriceps.....	369	selene, Carpiodes.....	103
Sebastodes.....	168, 364, 370	Luxilus.....	132
aereus.....	367	Selenichthys.....	216
aleutianus.....	365	seleniris, Trachipterus.....	216
alexandri.....	365	selenops, Hiodon.....	41
atrurubens.....	365	Hiodon.....	41
ayresii.....	367	sellaris, Acanthocottus.....	388
babcocki.....	365	Alvordius.....	284
clavilatus.....	366	Hadropterus.....	284
crameri.....	367	Porocottus.....	388
deani.....	367	sellicauda, Apogon.....	295
eigenmanni.....	366	Epinephelus.....	311
emphaeus.....	367	sellifer, Halichoeres.....	428
eos.....	368	Iridio.....	428
flavidus.....	364	Sema.....	409
gilberti.....	369	signifer.....	409
gilhi.....	368	semiararmatus, Gasterosteus.....	237
goodell.....	364	semicinctus, Eosebastes.....	367
hopkinsi.....	366	Gillellus.....	455
jordani.....	364	Iridio.....	427
melanops.....	365	Julis.....	427
melanostomus.....	366	semicoronata, Seriola.....	279
nigrocinctus.....	364	semifasciata, Pileoma.....	282
ovalis.....	366	semifasciatum, Pileoma.....	283
paucispinis.....	364	Triakis.....	14
rosaceus.....	364	semifasciatus, Gobius.....	441
ruberrimus.....	368	Serranus.....	317
rufus.....	366	semiloricatus, Gasterosteus.....	237

List of scientific names—Continued

	Page		Page
semiluna, Sparus.....	327	serra, Alepidosaurus.....	171
semilunata, Diana.....	264	Gomenion.....	280
seminolis, Fundulus.....	178	Serrana.....	353
Zygonectes.....	178	hispanis.....	353
seminuda, Albulas.....	40	Serranidae.....	317
Gila.....	114	serranoides, Sebastosomus.....	365
Gymneleotris.....	438	Serranus acanthophorus.....	317
seminudus, Electris.....	438	aculeatus (oculatus).....	328
Enypnias.....	445	acutirostris.....	315
Gobius.....	445	aquidens.....	319
Lycias.....	475	albomaculatus.....	317
Lycodes.....	475	altivelis.....	310
Plecotomus.....	160	angustifrons.....	311
semiradiatus, Lepidosteus.....	36	arara.....	311
semiruber, Labrus.....	425	armatus.....	313
semiscaber, Cottopsis.....	384	asperus.....	310
Cottus.....	384	bivittatus.....	318
semiscabra centropleura, Uranidea.....	383	bonaci.....	314
semispinosus, Caranx.....	270	brunneus.....	314
semisquamatum, Pycnomma.....	438	bulleri.....	319
Semitapicus.....	100	calopteryx.....	319
Semotilus.....	116, 117	camelopardalis.....	315
stromaculatus.....	117	capuena.....	331
thoreauanus.....	117	carsuna.....	309
biguttatus.....	137	catus.....	311
cephalus.....	117	colonus.....	320
diplemius.....	125	conspersus.....	311
diplemia.....	117	coronatus.....	308
dorsalis.....	117	couchi.....	308
hammondi.....	117	courtadei.....	310
macrocephalus.....	117	creolus.....	320
notatus.....	177	cyclopomatus.....	314
olivaceus.....	177	decimalis.....	314
speciosus.....	117	dimidiatus.....	314
thoreauanus.....	117	dubius.....	309
senegalensis, Vomer.....	275	emarginatus.....	315
semilis, Gambusia.....	185	erythrogaster.....	312
senta, Raja.....	25	falcatus.....	315
sephen, Raja.....	28	fascicularis.....	318, 319
septemfasciatum, Cichlosoma.....	421	felinus.....	315
septemfasciatum, Astatheros.....	421	fimbriatus.....	311
septentrionalis, Gaidropsarus.....	208	flavescens.....	319
Motella.....	208	fureifer.....	320
Perca.....	307	fuscus.....	315
sepiopinnis, Ammodytes.....	451	galeus.....	312
Rhynchias.....	451	guasa.....	312
Serranus bonaci.....	314	gigas.....	310
serena, Dioda.....	144	humeralis.....	317
sergeanti, Osmerus.....	67	impetiginosus.....	310
Seriola.....	269, 276, 279	mermis.....	313
bipinnulata.....	279	interstitialis.....	314
bonariensis.....	280	irradians.....	319
boschii.....	280	italara.....	312
carolinensis.....	279	jacome.....	319
coronata.....	280	labriformis.....	311
declivis.....	280	lamprurus.....	321
dorsalis.....	279	latepictus.....	314
dubia.....	280	luceopercanus.....	319
dumerili.....	279	maculato-fasciatus.....	317
dussumieri.....	278	maculosus.....	311
falcata.....	280	margaritifera.....	311
gigas.....	279	marginalis.....	311
jalandi.....	279	mentzeli.....	311
ligulata.....	280	morio.....	312
mazatlanana.....	279	morhua.....	310
pinnulata.....	278	mystacinus.....	310
proxima.....	279	nigriculus.....	310
rivoliiana.....	280	nigritus.....	312
semicoronata.....	279	niveatus.....	310, 311
stearnsi.....	279	oculatus.....	328
succincta.....	278	ocyurus.....	318
zonata.....	279	olax.....	314
Seriolichthys.....	279	ouatallii.....	309
Seriolides.....	278	panamensis.....	308
Seriolophus.....	280	(Paralabrax) deweegen.....	319
carangoides.....	333	petrosus.....	313
Seriophus.....	333	phæton.....	309
politus.....	333	phoebe.....	319
serotinus, Accipenser.....	35	pixanga.....	310
serpens, A. dipenser.....	37	præsiglitor.....	319
Serpens marinus maculosus.....	261, 262	psittacinus.....	312
serpens, Gempylus.....	475	quaquefasciatus.....	318
Lycodes.....	475	radialis.....	318
Ophidurus.....	85	remotus.....	315
serpentina, Muraena.....	77	repandus.....	315
serpentinus, Blennius.....	470, 471	rupestris.....	313
Dendrothys.....	77		

List of scientific names—Continued

	Page		Page
<i>Serranus semifasciatus</i>	317	<i>Sicyases fasciatus</i>	490
<i>stadthouderi</i>	311	<i>haeres</i>	490
<i>tamniops</i>	309	<i>punctulatus</i>	490
<i>tigris</i>	315	<i>rubiginosus</i>	490
<i>tinca</i>	315	<i>sanguineus</i>	490
<i>undulosus</i>	315	<i>Sicydium</i>	447
<i>varius</i>	310	<i>antillarum</i>	447
<i>Seriaria</i>	284	<i>buscki</i>	447
<i>sciara</i>	284	<i>caguitæ</i>	447
<i>serrula</i>	284	<i>gymnogaster</i>	447
<i>serrata</i> , <i>Fistularia</i>	245	<i>plumieri</i>	447
<i>Sayris</i>	197	<i>punctatum</i>	447
<i>serrato-granulata</i> , <i>Perca</i>	281	<i>salvini</i>	448
<i>serratum</i> , <i>Hæmulon</i>	329	<i>siragus</i>	447
<i>serratus</i> , <i>Gasterosteus</i>	237	<i>vincente</i>	447
<i>Nautocrates</i>	278	<i>Sicyogaster</i>	488
<i>serriceps</i> , <i>Sebastichthys</i>	369	<i>Sidara</i>	93
<i>Sebastocarus</i>	369	<i>castanea</i>	92
<i>serrifer</i> , <i>Conodon</i>	333	<i>chlevastes</i>	93
<i>Serrivomer</i>	84	<i>verrilli</i>	93
<i>beanii</i>	84	<i>Siderea</i>	93
<i>sector</i>	84	<i>picta</i>	93
<i>serrula</i> , <i>Anisotremus</i>	333	<i>sidera</i> , <i>Muraena</i>	93
<i>Chalinura</i>	204	<i>siderius</i> , <i>Hybognathus</i>	140
<i>Hadropterus sciurus</i>	284	<i>Hyborhynchus</i> (<i>Hybognathus</i>).....	140
<i>Priacanthus</i>	323	<i>sieboldii</i> , <i>Heros</i>	423
<i>Pristipoma</i>	333	<i>Sierra</i>	257
<i>Pseudopriacanthus</i>	323	<i>cavalla</i>	257
<i>Serraria</i>	284	<i>sierra</i> , <i>Doryrhamphus</i>	243
<i>sessilicauda</i> , <i>Monolene</i>	220	<i>Pontinus</i>	372
<i>setapinnis</i> , <i>Vomer</i>	275	<i>Scomberomorus</i>	257
<i>Zeus</i>	274, 275	<i>Scorpaena</i> (<i>Sebastopus</i>).....	372
<i>setapinnis cubensis</i> , <i>Vomer</i>	275	<i>sierrita</i> , <i>Tylosurus</i>	196
<i>Setarches</i>	372	<i>sigalutes</i> , <i>Gilbertidia</i>	393
<i>guntheri</i>	372	<i>Gilbertina</i>	392, 393
<i>parmatus</i>	372	<i>Sigmistes</i>	391
<i>setifer</i> , <i>Argyrosus</i>	275	<i>caullas</i>	391
<i>Monacanthus</i>	494	<i>Sigmops</i>	72
<i>setiger</i> , <i>Dasycottus</i>	392	<i>stigmaticus</i>	72
<i>setigerus</i> , <i>Lophiomus</i>	504	<i>Sigmurus</i>	348
<i>Lophius</i>	504	<i>vermicularis</i>	348
<i>setipinnis</i> , <i>Argyreus</i>	275	<i>Signalosa</i>	46
<i>setosus</i> , <i>Mugil</i>	254	<i>atchafalaya</i>	46
<i>Querimana</i>	254	<i>vanhuyningi</i>	47
<i>Tetraodon</i>	500	<i>mexicana</i>	47
<i>seu conger</i> , etc., <i>Muraena</i>	94	<i>campi</i>	47
<i>severus</i> , <i>Heros</i>	416	<i>petenensis</i>	47
<i>sexcornutus</i> , <i>Ostracion</i>	497	<i>signata</i> , <i>Cyclothone</i>	72
<i>sexdecim-lamellata</i> , <i>Echeneis</i>	449	<i>signatus</i> , <i>Bathymaster</i>	451
<i>sexfasciatus</i> , <i>Cichlosoma</i>	421	<i>Hypoprius</i>	17
<i>Hæmulon</i>	329	<i>Microgobius</i>	443
<i>sexfasciatus</i> , <i>Astatheros</i>	421	<i>signifer</i> , <i>Brytus</i>	299
<i>Hæmulon</i>	329	<i>Centristhmus</i>	319, 320
<i>sex-maculatus</i> , <i>Diodon</i>	501	<i>Chatoëssus</i>	44
<i>sexradiatus</i> , <i>Polynemus</i>	409	<i>Coregonus</i>	66
<i>shasta</i> , <i>Cottus</i>	383	<i>Monacanthus</i>	494
<i>Salmo</i>	58	<i>Neotriopus</i>	423
<i>gairdneri</i>	58	<i>Sema</i>	409
<i>shavianus</i> , <i>Cetorhinus</i>	20	<i>Stypodon</i>	113
<i>shufeldti</i> , <i>Gobius</i>	439	<i>Tyhmallus</i>	66
<i>Mancalia</i>	508	<i>silenus</i> , <i>Zaprora</i>	452
<i>Rhinogobius</i>	439	<i>Siluræ</i>	147
<i>Typhlosaras</i>	508	<i>Silurus bagre</i>	147
<i>shumardi</i> , <i>Alburnops</i>	134	<i>belis</i>	149
<i>Hadropterus</i>	285	<i>catus</i>	153, 159
<i>Hypopsis</i>	134	<i>clarias</i>	245, 246
<i>Imostoma</i>	285	<i>cornutus</i>	153
<i>siatis</i> , <i>Argentina</i>	68	<i>enpreus</i>	148
<i>Corvula</i>	345	<i>felis</i>	153
<i>Noturus</i>	155	<i>fuscus</i>	159
<i>Vacuocqua</i>	345	<i>galeatus</i>	155
<i>sibbaldi</i> , <i>Syngnathus</i>	343	<i>gyrinus</i>	143
<i>Siboma</i>	120	<i>hertzbergi</i>	155
<i>atraria</i>	119	<i>lunus</i>	155
<i>conformis</i>	120	<i>lividus</i>	153
<i>crassicauda</i>	120	<i>marinus</i>	147, 148
<i>sicana</i> , <i>Cerna</i>	312	<i>melas</i>	154
<i>sicciferum</i> , <i>Holocentrum</i>	236	<i>militaris</i>	159
<i>sicculum</i> , <i>Chirostoma</i>	251	<i>nebulosus</i>	155
<i>sicculus</i> , <i>Labridesthes</i>	251	<i>olivaris</i>	155
<i>scula</i> , <i>Pimelopterus boscii</i>	349	<i>punctatus</i>	151, 152
<i>sculus</i> , <i>Lampugus</i>	265	<i>viscosus</i>	155
<i>Sicya</i>	447	<i>xanthocephalus</i>	154
<i>Sicyases</i>	490	<i>var. fuscus</i>	153
<i>carneus</i>	490	<i>Silus</i>	68
<i>erythroptus</i>	490	<i>ascanti</i>	68

List of scientific names—Continued

	Page		Page
<i>slus</i> , Argentina.....	68	<i>skib</i> , Pomatomus.....	280
<i>slma</i> , Carenitra.....	401	<i>Skiffa</i>	183
<i>Careproctus</i>	401	<i>bilineata</i>	183
<i>Electris</i>	436	<i>lerna</i>	183
<i>Simenichelyidae</i>	78	<i>multipunctata</i>	183
<i>Simenichelys</i>	78	<i>variegata</i>	183
<i>parasiticus</i>	78	<i>sloani</i> , Chauliodus.....	71
<i>similis</i> , <i>Amphistichus</i>	411	<i>Leicobatus</i>	29
<i>Fundulus</i>	177	<i>Scomber</i>	260
<i>Hæmulon</i>	330	<i>Urobatis</i>	29
<i>Hydrargyra</i>	175, 177	<i>sloani</i> vermiculatus, <i>Urobatis</i>	29
<i>Iridio</i>	427	<i>sluteri</i> , <i>Brannerella</i>	464
<i>Rhinogobius</i>	439	<i>Smaragdus</i>	440
<i>simillima</i> , <i>Palometa</i>	266	<i>costalesi</i>	441
<i>similimus</i> , <i>Gerrus</i>	342	<i>stigmaticus</i>	441
<i>Poronotus</i>	266	<i>valenciennesi</i>	441
<i>simotera</i> , <i>Ulocentra</i>	286	<i>smaragdus</i> , <i>Electris</i>	437
<i>simoterum</i> , <i>Hystoma</i>	266	<i>Eretelis</i>	437
<i>simplex</i> , <i>Apilurus</i>	261	<i>Gobius</i>	440, 441
<i>Oreanidus</i>	339	<i>Gobionellus</i>	441
<i>Pseudoscarus</i>	435	<i>Salmo</i>	58
<i>Tetragonurus</i>	261	<i>Smaris</i>	340
<i>simplicidens</i> , <i>Girella</i>	339	<i>lineatus</i>	342
<i>Incisidens</i>	339	<i>martincus</i>	340
<i>simpsoni</i> , <i>Ictalurus</i>	152	<i>smaris</i> , <i>Sparus</i>	340
<i>simula</i> , <i>Chalinura</i>	204	<i>Smeeticus</i>	316
<i>simulus</i> , <i>Hemiplites</i>	304	<i>bicolor</i>	316
<i>Sebastosomus</i>	365	<i>smiridus</i> , <i>Merluccius</i>	214
<i>Synodus</i>	164	<i>smithi</i> , <i>Ceratichthys</i>	145
<i>simulus</i> , <i>Ophoscion</i>	348	<i>Lepidosteus</i>	37
<i>sinus</i> , <i>Alburnellus</i>	135	<i>smithi</i> , <i>Chiola</i>	145
<i>Careproctus</i>	401	<i>Cyprinus</i> (<i>Abramis</i>).....	41
<i>Catetix</i>	480	<i>Halientichthys</i>	507
<i>Crescatus</i>	98	<i>snyderi</i> , <i>Catostomus</i>	106
<i>Hycoptis</i>	135	<i>Dialarchus</i>	391
<i>Manticichthys</i>	351	<i>Gnathypops</i>	453
<i>Rhinichthys</i>	140	<i>Hyporhamphus</i>	199
<i>sinaloa</i> , <i>Paralichthys</i>	224	<i>Oligocottus</i>	391
<i>Siphostoma</i>	241	<i>Palometa</i>	267
<i>Syngnathus</i>	241	<i>Peprilus</i>	267
<i>Umbra</i>	350	<i>Sobaco</i>	492
<i>sinensis</i> , <i>Hispanicus</i>	368	<i>sobaco</i> , <i>Balistes</i>	492
<i>Sebastichthys</i>	368	<i>Canthidermis</i>	492
<i>Trygon</i>	28	<i>sobra</i> , <i>Mesoprion</i>	326
<i>sinuosus</i> , <i>Pomadasy</i>	335	<i>socialis</i> , <i>Salmo</i>	66
<i>Rhoniciscus</i>	335	<i>socius</i> , <i>Alburnus</i>	124
<i>Siphagonus barbata</i>	394	<i>Notropis</i>	124
<i>barbatus</i>	394	<i>socorroense</i> , <i>Thalassoma</i>	429
<i>Siphoteles</i>	119, 120, 122	<i>socorroensis</i> , <i>Thalassoma</i>	429
<i>bicolor</i>	122	<i>sogo</i> , <i>Holocentrus</i>	235
<i>columbianus</i>	122	<i>solar</i> , <i>Chaetodon</i>	362
<i>formosus</i>	122	<i>solandri</i> , <i>Acanthocybium</i>	256
<i>mobavensis</i>	123	<i>Cybium</i>	256
<i>obesus</i>	122	<i>solaris</i> , <i>Orthogoriscus</i>	503
<i>olivaceus</i>	122	<i>Solea browni</i>	229
<i>oregonensis</i>	122	<i>fischeri</i>	230
<i>vittatus</i>	122	<i>fonsecensis</i>	230
<i>Siphostoma</i>	240	<i>gronovi</i>	229
<i>arctum</i>	242	<i>klunzingeri</i>	229
<i>auliscus</i>	240	<i>lunata</i>	217
<i>barbata</i>	240	<i>mazatlan</i>	230
<i>brevicaudum</i>	241	<i>panamensis</i>	230
<i>carinatum</i>	240	<i>pilosa</i>	230
<i>crinigerum</i>	241	<i>punctata</i>	217
<i>dendriticum</i>	242	<i>scutum</i>	229
<i>exile</i>	243	<i>soleiformis</i> , <i>Rhombus</i>	218
<i>florida</i>	241	<i>Solenostomus</i>	245
<i>mackayi</i>	240	<i>Soleotalpa</i>	230
<i>poei</i>	241	<i>unicolor</i>	230
<i>robertsi</i>	241	<i>solis</i> , <i>Lepomis</i>	300
<i>scovelli</i>	242	<i>Pomotis</i>	300
<i>sinaloa</i>	241	<i>Somniodide</i>	21
<i>starksi</i>	242	<i>Somniosus</i>	21
<i>torrei</i>	242	<i>brevipinna</i>	21, 22
<i>zatropis</i>	243	<i>microcephalus</i>	22
<i>stragus</i> , <i>Sicydium</i>	447	<i>somnolentus</i> , <i>Electris</i>	436
<i>Siremba guentheri</i>	483	<i>Labotes</i>	323
<i>messieri</i>	480	<i>sonnerati</i> , <i>Opisthognathus</i>	452
<i>nigripinnis</i>	480	<i>sonntagii</i> , <i>Antennarius</i>	506
<i>sisco</i> , <i>Argyrosomus</i>	62	<i>Chironectes</i>	506
<i>huronius</i> , <i>Leucichthys</i>	62	<i>sonora</i> , <i>Atherinopsis</i>	252
<i>Leucichthys</i>	62	<i>Scorpena</i>	371
<i>siscowet</i> , <i>Cristivomer</i>	60	<i>sonoriensis</i> , <i>Catostomus</i>	106
<i>Salmo</i>	60	<i>Girardinus</i>	189
<i>siskawitz</i> , <i>Salmo</i>	60	<i>soporator</i> , <i>Bathygobius</i>	440
<i>sluslawi</i> , <i>Leuciscus</i>	118	<i>Gobius</i>	440
<i>Richardsonius</i>	118	<i>sordida</i> , <i>Muraena</i>	94

List of scientific names—Continued

	Page		Page
sordidus, Chaetodon	414	speciosus, Pomotis	301
Oitharichthys	218	Scomber	273
Psettichthys	218	Semotilus	117
Verilus	328	spectabilis, Pœcilichthys	291
spadicea, Dioda	144	Salmo	61
Lampetra	9	Salvelinus	61
spadiceus, Centrurrophis	87	malma	61
Tetrapleurodon	9	spectrum, Careproctus	402
spallanzani, Isurus	19	Lophius	505
Leptocephalus	79	Osmerus	67
Sparactodon	280	mordax	67
malma	280	spectrunculus, Hybopsis	134
Sparidae	336	speculiger, Exocoetus	201
Sparsoma	431, 438	speculigera, Lampadena	167
abildgaardii	432	spekeus, Amblyopsis	195
atomarium	431	spenleri, Spheroides	498
aurofratatum	432	Tetrodon	255
brachiale	432	spet, Esoc	498
chrysotermum	432	Spharoides tuberculatus	84
cyanolepis	432	Sphagebranchus	84
distinctum	432	anguiformis	84, 91
flavescens	433	kendallii	84
frondosum	433	ophionetus	84
hoplomystax	432	rostratus	84
lorito	432	scuticaris	86
maschalespilos	433	selachops	84
niphobles	432	teres	86
oxybrachium	432	(Sphagebranchus) anguiformis, Ophichthys	84
radians	432	sphenops cuneata, Mollienesia	193
rubripinna	433	sphenops, Mollienesia	192
strigatum	433	Pœcilia	192
viride	432	vandepolli, Mollienesia	193
xystrodon	431	Spheroide tubercule	498
Sparisomidae	431	Spheroides	498
sparoides, Labrus	304, 305	asterias	498
Pomoxis	305	harperi	498
Sparus	418	les	498
abildgaardii	432	(Spheroides) eulepidotus, Tetrodon	498
annularis	338	Sphceroides	498
argenteus	337	angusticeps	498
argyrops	336	asterias	498
atlanticus	310	eulepidotus	498
aureo-ruber	432	harperi	498
aureus	302	lobatus	498
bajonado	336	maculatus	498
brachysomus	337	marmoratus	498
castaneola	265	nepheus	498
caxis	325	spengleri	498
chromis	413	tuberculatus	498
chrysomelanurus	311	Sphyraena	255
chrysops	336	acus	196, 197
chrysurus	327	argentea	255
cruentatus	308	aureoviridis	305
falcatus	425	barracuda	255
niger	265	becuna	255
ovicephalus	338	borealis	255
pagrus	337	ensis	255
probatocephalus	338	guachanco	255
radiatus	426	guntheri	255
raii	265	lucasana	255
rhomboides	337	picudilla	255
salin	338	sphyraena	255
sargus	338	picuda	255
scirurus	329, 330	viridensis	255
semiluna	327	vulgaris	255
smaris	340	sphyraena, Argentina	68
surinamensis	424	Chirostoma	251
synagrus	327	Esoc	255
tetracanthus	325	Sphyraena	255
vermicularis	327	sphyraenarum, Echeus	449
virginicus	332	Sphyraenidae	225
vittatus	332	Sphyraenops	297
xanthurus	336	bairdiana	297
Spathula	36	biardianus	297
spathula, Polyodon	36	sphyraenops, Lestidiops	166
Squalus	36	Sphyrichthys	18
Spathularia	36	Sphyra	18
reticulata	36	zygæna	18
spatula, Atractosteus	38	Sphyridæ	17
Icelus	379	Spicara	340
Lepisosteus	38	flexuosa	340
Spatularia reticulata	36	martinica	340
speciosa, Gambusia	185	spicifer, Hippichthys	243
speciosus, Gnathanodon	273	Syngnathus	243
Gnathodon	273	spillmani, Alvordius	283

List of scientific names—Continued

	Page		Page
spilonotopterus, Cypselurus.....	202	Squalidae.....	21
Exocoetus.....	202	qualidus, Scarus.....	433
spilonota, Poecilia.....	192	squalipeta, Echeeneis.....	449
spilonotus, Coregonus.....	65	Squalus.....	117
Monacanthus.....	493	alicæ.....	119
Prosopium.....	65	copei.....	119
spilopterus, Citharichthys.....	218, 219	crucius.....	119
Photogenis.....	129	galnæ.....	118
spilopus, Exocoetus.....	201	hyalope.....	117
spilota, Uranidea.....	385	lemmon.....	119
spilopterygius, Balistes.....	492	photogenis.....	123
spilotum, Etheostoma nanganæ.....	285	pronger.....	120
spilotus, Hypohomus.....	285	rhomaleus.....	119
spilura, Mollenesia.....	192	Squaloidæ.....	21
spilurus, Archocentrus.....	420	Squalus.....	13, 21
Curimatus.....	100	acanthias.....	21
Heros.....	420	acronotus.....	16
Poecilia.....	192	americanus.....	19
Salmø.....	57	arcticus.....	14
Spinachia.....	239	argus.....	12
spinachia.....	239	borealis.....	21, 22
vulgaris.....	239	brucus.....	22
spinachia, Gasterosteus.....	239	caeruleus.....	17
Spinachia.....	239	camcula.....	11
Spinachinae.....	239	cans.....	13
Spinax (Acanthias) suckleyi.....	21	carcharias.....	15, 20
Spinax fabricii.....	21	(Carcharias) terra-novæ.....	14
hulianus.....	21	cetaceus.....	20
spinax, Squalus.....	21	cirratus.....	12
spinescens, Auliscops.....	239	commersoni.....	15
spinecauda, Acanthenchelys.....	89	cornubicus.....	19
spinicephalum, Exoglossum.....	146	edentulus.....	31
Spinicephalus.....	145	elephas.....	20
fibulatus.....	145	ferox.....	19
spiniceps, Photocorynus.....	509	galeus.....	13, 14
spindens, Scarus.....	431, 432	glacialis.....	22
spinifer, Acanthonus.....	482	glaucus.....	17
Engraulis.....	50	griseus.....	11
spinifera, Anchoviella.....	50	gunneri.....	20
spiniger, Icelus.....	378	hurundinaceus.....	17
Spinivomer.....	84	homanus.....	20
goodii.....	84	isodus.....	20
spinosa, Acanthemblemaria.....	465	littoralis.....	19
Malthopsis.....	507	longimanus.....	15
Thysanocara.....	160	macrodus.....	19
spinosissimum immaculatum, Cichlasoma.....	420	maximus.....	20
spinosissimus, Archocentrus.....	420	microcephalus.....	21, 22
Aspidophorus.....	394	monensis.....	19
Diodon.....	501	mustelus.....	13, 14
Heros (Cichlasoma).....	420	nasus.....	19
spinosus, Aneistrus.....	160	norwegianus.....	22
Calycilepidotus.....	381	obscurus.....	16
Centronotus.....	281	obtusus.....	15
Cychechthys.....	502	pelegrinus.....	20
Cyclopterus.....	398	pennanti.....	19
Diodon.....	502	platyodon.....	15
Eumicrotremus.....	398	porosus.....	15
Notacanthus.....	76	pristis.....	12, 14, 15
Platophrys.....	217	punctatus.....	12
Polypnus.....	74	punctulatus.....	20
Rhinobatos.....	24	rashleighanus.....	19
Rhinobatus.....	24	rostratus.....	36
Rhomboidichthys.....	217	spathula.....	21
Schedophilopris.....	269	spinax.....	22
Squalus.....	22	spinosus.....	22
Trachinotus.....	276	squatina.....	11
spinulosa, Chalinura.....	204	stellaris.....	21
Malthopsis.....	507	suckleyi.....	16, 17
spinulosus, Gasterosteus.....	237	tiburo.....	18
Spirinchus.....	67	vulpes.....	18
thaleichthys.....	67	vulpinus.....	18
sprhngulus, Leuciscus.....	128	zygæna.....	285
spixianus, Saurus.....	163	squamata, Swainia.....	119
spixii, Arius.....	150	Tigoma.....	285
Pinelodus.....	150	squamatus, Etheostoma (Hadi opterus).....	292
Platysomus.....	275	squamiceps, Claricola.....	292
Vomer.....	275	Etheostoma.....	186
splendens, Beryx.....	234	squamulentus, Ceratichthys.....	138
Exocoetus.....	200	Couesius.....	224
splendida, Petenia.....	423	Paralichthys.....	353
splendatum, Pristipoma.....	332	Squampennes.....	355
splendatus, Amstotremus.....	332	squampinnis, Cynoscion.....	342
spongicola, Evermannichthys.....	446, 447	Garres.....	355
Garmanna.....	446	Otolithus.....	36
spongiosa, Hakeutea.....	508	squamosa viridis, Aous maxima.....	272
Spratelloides bryoporus.....	41	squamosus, Trachurus.....	

List of scientific names—Continued

	Page		Page
squamulosus, <i>Chaetodon</i>	361	stellatus, <i>Fario</i>	56
Squatina.....	22	Mustellus.....	13
angelus.....	22	Platichthys.....	226
californica.....	22	Pleuronectes.....	226
dumeril.....	22	Tetraodon.....	497, 73
fimbriata.....	22	Valenciennellus.....	273
lævis.....	22	Xurel.....	361
lewis.....	22	stelleri, <i>Cottus</i>	368
oculata.....	22	<i>Cyclopterus</i>	374
raja.....	23	<i>Hexagrammos</i>	373, 374
squatina.....	22	<i>Liparops</i>	368
vulgaris.....	22	<i>Myoxocephalus</i>	367
Squatina, <i>Squalus</i>	22	Stellerna.....	364
Squatina.....	22	<i>xyosterna</i>	347
Squatinae.....	22	Stellicarens.....	347
Squatinoidei.....	22	<i>zestocarus</i>	347
squeteague, <i>Labrus</i>	355	Stellifer.....	346
<i>maculatus</i> , <i>Labrus</i>	355	<i>coloniensis</i>	347
<i>stabuliforis</i> , <i>Raja</i>	25	<i>erycymba</i>	346
<i>stadthouderi</i> , <i>Serranus</i>	311	<i>illecebrosus</i>	347
<i>stagnalis</i> , <i>Salvelinus</i>	60	<i>lanceolatus</i>	346
<i>stahli</i> , <i>Auchenistius</i>	457	<i>melanocheir</i>	347
<i>Monosira</i>	344	<i>microps</i>	346
<i>stanleyi</i> , <i>Coregonus</i>	65	<i>mindii</i>	347
<i>Frosopium</i>	65	<i>osotans</i>	347
<i>starksi</i> , <i>Memidia</i>	247	<i>stellifer</i>	346
<i>Siphostoma</i>	242	<i>zestocarus</i>	347
<i>Stolephorus</i>	48	stellifer, <i>Bodianus</i>	346
<i>Syngnathus</i>	242	<i>Stellifer</i>	346
Starksia.....	456	<i>Xenisma</i>	179
<i>cremnobates</i>	456	stellifera, <i>Corvina</i>	346
<i>holderi</i>	456	<i>Xenisma</i>	179
starr, <i>Pomadasys</i>	334	stelliferoides, <i>Bassogigas</i>	482
Stathmonotus.....	465	<i>Neobythites</i>	482
<i>hemphilli</i>	465	Stelliferus.....	346
<i>tekla</i>	466	<i>illecebrosus</i>	347
stearnsi, <i>Blennius</i>	460	<i>zestocarus</i>	347
<i>Colotrigia</i>	404	stello, <i>Perca</i>	310
stearnsi, <i>Corvina</i>	348	<i>Rhinobatus</i>	23
<i>Lutjanus</i>	325	<i>Rhinobatus</i>	26
<i>Prionotus</i>	406	stellulata, <i>Raja</i>	336
<i>Roncador</i>	348	Stenesthes.....	168
<i>Scorpaena</i>	371	(<i>Stenobranchius leucopsarum</i> , <i>Myctophum</i>	61
<i>Seriola</i>	279	Stenodus.....	61
Stegastes.....	415	<i>mackenzii</i>	498
<i>bairdii</i>	415	Stenometopus.....	336
<i>chrysurus</i>	415	Stenotoma.....	336
<i>dorsalis</i>	416	Stenotomus.....	336
<i>imbricatus</i>	416	<i>caprinus</i>	336
<i>niveatus</i>	416	<i>chrysops</i>	336
Steindachnerella.....	203	<i>versicolor</i>	336
steindachneri, <i>Argo</i>	265	Stephanoberyoidæ.....	232
<i>Arius</i>	151	Stephanoberyx.....	232
<i>Oichlasoma</i>	419	<i>gillii</i>	232
<i>Diabasis</i>	330	<i>monæ</i>	494
<i>Hæmulon</i>	330	Stephanolepis.....	407
<i>Ophioblennius</i>	464	stephanophrys, <i>Prionotus</i>	308
<i>Parapetenia</i>	419	Stereolepis.....	308
<i>Rhinoptera</i>	31	<i>californicus</i>	308
<i>Tachysurus</i>	151	<i>gigas</i>	33
<i>Thalassoma</i>	429	Sterletæ.....	33
Steindachneria.....	303	Sterletus, <i>Averuncus</i>	395
<i>argentea</i>	303	<i>Cerathichthys</i>	139
Steindachneridion.....	203	<i>Maechtybopsis</i>	139
Steinegeria.....	266	Sternarchus.....	101
<i>rubescens</i>	266	<i>rostratus</i>	101
Steinegeriidæ.....	266	<i>viridescens</i>	101
stegidolepis, <i>Macrourus</i>	205	Sternas.....	380
<i>Macrurus</i> (<i>Macrourus</i>).....	379	<i>xenesthesus</i>	380
Stelgidonotus.....	379	sternula, <i>Gasteropeleus</i>	73
<i>latifrons</i>	379	Sternoptichidæ.....	73
Stelgis.....	365	Sternoptix.....	73
<i>vulvus</i>	365	<i>diaphana</i>	73
Stelgistrus.....	379	<i>obscura</i>	73
<i>beringiana</i>	379	Sternoptyx.....	73
Stelgistrum beringianum.....	346	<i>gardani</i>	266
Stelliferæ, les.....	11	<i>mediterraneus</i>	74
<i>stellaris</i> , <i>Squalus</i>	507	<i>obscura</i>	73
<i>stellata</i> , <i>Haleutæa</i>	34	<i>olfersi</i>	74
<i>stellatus</i> , <i>Acipenser</i>	286	Sternopygus.....	101
<i>Apogonichthys</i>	286	<i>dariensis</i>	101
<i>Astrapogon</i>	309	<i>humboldti</i>	101
<i>Bodianus</i>	309	<i>lineatus</i>	101
<i>Caranx</i>	273	<i>tumifrons</i>	101
<i>Cephalopholis</i>	309		

List of scientific names—Continued

	Page		Page
<i>Sternotremia</i>	215	<i>stomias</i> , <i>Atheresthes</i>	221
<i>isolepis</i>	215	<i>Bathytroctes</i>	52
<i>stevensi</i> , <i>Thaleichthys</i>	66	<i>Chasmistes</i>	108
<i>Stichæide</i>	469	<i>Narcetes</i>	52
<i>Stichæus</i>	469	<i>Platysomichthys</i>	221
<i>punctatus</i>	469	<i>Salmo</i>	57
<i>stigma</i> , <i>Gymnelis</i>	477	<i>Trigllops</i>	314
<i>Ophidium</i>	477	<i>Stomiidae</i>	69
<i>stigmæa</i> , <i>Ulocentra</i>	287	<i>Stomiatoidei</i>	69
<i>stigmæum</i> , <i>Boleosoma</i>	287	<i>Stomocatus</i>	105
<i>stigmæus</i> , <i>Citharichthys</i>	219	<i>Stomodon bilinearis</i>	214
<i>stigmanotus</i> , <i>Ahynnodontophus</i>	94	<i>stonei</i> , <i>Hydrophlox</i>	127
<i>stigmaticus</i> , <i>Ceraticthys</i>	137	<i>Notopis</i>	127
<i>Gobionellus</i>	441	<i>Salmo</i>	58
<i>stigmatistium</i> , <i>Lepophidium</i>	455	<i>gairdneri</i>	58
<i>Leptophidium</i>	455	<i>storeri</i> , <i>Acipenser</i>	34
<i>stigmatiscus</i> , <i>Sigmops</i>	72	<i>Aledon</i>	603
<i>Smaragdus</i>	441	<i>Cetomimus</i>	172
<i>stigmatura</i> , <i>Algansea</i>	116	<i>Leuciscus</i>	117
<i>Bollmannia</i>	443	<i>Scomberesox</i>	197
<i>Erogala</i>	129	<i>storeria</i> , <i>Cichla</i>	304
<i>stigmaturus</i> , <i>Gobius</i>	439	<i>storerianus</i> , <i>Erimemus</i>	137
<i>Photogenis</i>	129	<i>Rutilus</i>	137
<i>Rhinogobius</i>	439	<i>stormi</i> , <i>Cephalocassis</i>	149
<i>stilbe</i> , <i>Zalocys</i>	277	<i>stoutii</i> , <i>Bdelostoma</i>	8
<i>Stilbiscus</i>	89	<i>Polistotrema</i>	8
<i>edwardsi</i>	89	<i>strabo</i> , <i>Icelinus</i>	376
<i>stilbus</i> , <i>Leuroglossus</i>	68	<i>Ophioscion</i>	347
<i>Nototropis</i>	125	<i>straminea</i> , <i>Hybopsis delicosa</i>	134
<i>Notropis</i>	125	<i>stramineus</i> , <i>Hybognathus</i>	133
<i>stampsoni</i> , <i>Triglopsis</i>	359	<i>Hybopsis</i>	134
<i>stipes</i> , <i>Atherina</i>	246	<i>Strephon</i>	159
<i>Hepsetia</i>	246	<i>striata</i> , <i>Alausa</i>	43
<i>Joturus</i>	254	<i>Argentina</i>	68
<i>stirurus</i> , <i>Chloroscombrus</i>	276	<i>Perca</i>	331
<i>Stizostedion</i>	282	<i>striato</i> , <i>Perca marina capite</i>	330
<i>glaucum</i>	282	<i>strutulus</i> , <i>Brycon</i>	98
<i>vitreum</i>	282	<i>Chalcinopsis</i>	98
<i>Stizostethium</i>	282	<i>stratum</i> , <i>Bathystoma</i>	331
<i>Stoasodon</i>	30	<i>Pygidium</i>	159
<i>narnari</i>	30	<i>stratus</i> , <i>Acanthurus</i>	363
<i>Stolephorus</i>	47	<i>Anthias</i>	311, 318
<i>astilbe</i>	49	<i>Blennius</i>	462
<i>branchiomelas</i>	49	<i>Bodianus</i>	325
<i>brevirostrum</i>	50	<i>Centropristes</i>	317
<i>cultratus</i>	48	<i>Chatodon</i>	359
<i>curtus</i>	48	<i>Ctenochatus</i>	363
<i>epsetus</i>	47	<i>Diodon maculato</i>	502
<i>eurystole</i>	47	<i>Epinephelus</i>	311, 318
<i>exiguus</i>	47	<i>Holocentrus</i>	235
<i>garmani</i>	51	<i>Hypsoblennius</i>	462
<i>gilberti</i>	51	<i>Labrus</i>	317
<i>guanensis</i>	49	<i>Luciocharax</i>	100
<i>hiulcus</i>	48	<i>Monacanthus</i>	494
<i>ischanus</i>	48	<i>Roccus</i>	307
<i>lucidus</i>	48	<i>Salmo</i>	54
<i>lyolepis</i>	49	<i>strigata</i> , <i>Trigla</i>	407
<i>miarchus</i>	47	<i>strigatum</i> , <i>Sparisoma</i>	433
<i>mundoculus</i>	50	<i>strigatus</i> , <i>Atemnarius</i>	505
<i>naso</i>	49	<i>Eucenogobius</i>	441
<i>opercularis</i>	51	<i>Gobionellus</i>	441
<i>per fasciatus</i>	47	<i>Holacanthus</i>	361
<i>pertheatus</i>	47	<i>Prionotus</i>	407
<i>rastralis</i>	50	<i>Scarus</i>	433
<i>robertsi</i>	47	<i>Xiphophorus</i>	193
<i>rostralis</i>	47	<i>helleri</i>	193
<i>seofieldi</i>	49	<i>strigilata</i> , <i>Loricaria</i>	182
<i>spunifer</i>	47	<i>strigosus</i> , <i>Acanthurus</i>	363
<i>starksi</i>	48	<i>Anarhichas</i>	472
<i>surinamensis</i>	50	<i>stromini</i> , <i>Zeus</i>	217
<i>stolifera</i> , <i>Clupea</i>	44	<i>Stromateidae</i>	266
<i>Dussumiera</i>	45, 46	<i>Stromateiformes</i>	266
<i>Jenkinsia</i>	46	<i>Stromateus cryptosus</i>	267
<i>Lile</i>	44	<i>longipinnis</i>	266
<i>stolzmanni</i> , <i>Belone</i>	196	<i>medius</i>	267
<i>Ersicion</i>	356	<i>palometa</i>	266, 267
<i>Otolithus</i>	356	<i>paru</i>	266
<i>Strongylura</i>	196	<i>triacanthus</i>	267
<i>stomata</i> , <i>Hippoglossina</i>	223	<i>strombi</i> , <i>Apogonichthys</i>	296
<i>Stomas</i>	69	<i>Strongylura</i>	185
<i>affinis</i>	69	<i>angusticeps</i>	196
<i>atriventer</i>	69	<i>argalus</i>	196
<i>oculubrimus</i>	69	<i>caudimaculata</i>	195
<i>ferox</i>	69	<i>diplostenia</i>	196
<i>hexagonatus</i>	69	<i>euryops</i>	196

List of scientific names—Continued

	Page		Page
<i>Strongylura exilis</i>	196	<i>superbus</i> , <i>Scarus</i>	434
<i>gualanensis</i>	196	<i>superciliosus</i> , <i>Hexagrammos</i>	374
<i>marina</i>	196	<i>Hyborhynchus</i>	145
<i>microps</i>	196	<i>Labrax</i>	373
<i>notata</i>	195	<i>supernas</i> , <i>Leucichthys</i>	63
<i>pterrura</i>	196	<i>surinamensis</i> , <i>Anisotremus</i>	332
<i>scapularis</i>	195	<i>Batrachoides</i>	486
<i>stolzmanni</i>	196	<i>Batrachus</i>	486
<i>tumucu</i>	195	<i>Galeichthys</i>	150
<i>strongylura</i> , <i>Belone</i>	195	<i>Hexanematichthys</i>	150
<i>strumosus</i> , <i>Gobieso</i>	489	<i>Holocentrus</i>	323
<i>Sturio</i>	33, 34	<i>Lobotes</i>	323
<i>vulgaris</i>	33	<i>Lutjanus</i>	332
<i>sturio</i> , <i>Acipenser</i>	33, 34	<i>Pleuronectes</i>	217
<i>Sturisma</i>	161	<i>Poecilia</i>	190
<i>rostrata</i>	161	<i>Sparus</i>	424
<i>Stygicola</i>	479	<i>Stolephorus</i>	50
<i>dentatus</i>	479	<i>susana</i> , <i>Boleosoma</i>	288
<i>Stygoenes</i>	159	<i>suspectus</i> , <i>Phycocetes</i>	478
<i>humboldtii</i>	159	<i>sutor</i> , <i>Blepharis</i>	274
<i>Stylephoridae</i>	216	<i>swaini</i> , <i>Poecilichthys</i>	291
<i>Stylephorus</i>	216	<i>Swamia</i>	285
<i>chordatus</i>	216	<i>squamata</i>	285
<i>Styleror</i>	86	<i>swampina</i> , <i>Hydrargyra</i>	175
<i>stylifer</i> , <i>Hippocampus</i>	244	<i>swani</i> , <i>Bothragonus</i>	396
<i>Stylophthalmdae</i>	71	<i>Hypsagonus</i>	396
<i>Stylophthalmus</i>	71	<i>swannanoa</i> , <i>Etheostoma</i>	289
<i>paradoxus</i>	71	<i>Nanostoma</i>	289
<i>Stypodon</i>	113	<i>swifti</i> , <i>Eosebastes</i>	367
<i>signifer</i>	113	<i>Sebastodes</i>	367
<i>suavis</i> , <i>Cyprinella</i>	131	<i>Syacium</i>	218
<i>subaequalis</i> , <i>Corvina</i>	345	<i>latifrons</i>	218
<i>Vacuqua</i>	345	<i>micrurum</i>	218
<i>subarcuata</i> , <i>Zygæna</i>	18	<i>ovale</i>	218
<i>subarcuatum</i> , <i>Hæmulon</i>	330	<i>papillosum</i>	218
<i>subarmatus</i> , <i>Acanthurus</i>	362	<i>Symbranchia</i>	76
<i>subbifrenatus</i> , <i>Rhypticus</i>	316	<i>Symbranchus</i>	76
<i>subbifurcata</i> , <i>Uvaria</i>	470	<i>symmetricus</i> , <i>Apomotis</i>	299
<i>subbifurcatus</i> , <i>Pholis</i>	470	<i>Caranx</i>	270
<i>subcaerulea</i> , <i>Amia</i>	38	<i>Hesperoleucus</i>	121
<i>subditus</i> , <i>Hesperoleucus</i>	122	<i>Lepomis</i>	299
<i>subfucus</i> , <i>Labrus</i>	424	<i>Pogonichthys</i>	121
<i>subfurcatus</i> , <i>Bermudichthys</i>	430	<i>Salmo</i>	59
<i>subligarius</i> , <i>Centropristis</i>	320	<i>Trachurus</i>	270
<i>subniger</i> , <i>Chiasmodon</i>	452	<i>Symmetrurus</i>	116
<i>suborbitalis</i> , <i>Holocentrum</i>	236	<i>Symbranchus vittatus</i>	76
<i>suborbitalis</i> , <i>Adioryx</i>	236	<i>Symphurus</i>	230
<i>Macrurus</i>	203	<i>stramentatus</i>	231
<i>Nematonurus</i>	203	<i>stricaudus</i>	231
<i>Plectromus</i>	233	<i>diomedianus</i>	231
<i>subosseus</i> , <i>Nannocampus</i>	244	<i>elongatus</i>	231
<i>subterranea</i> , <i>Lucifuga</i>	479	<i>fasciolaris</i>	231
<i>subterraneus</i> , <i>Lucifuga</i>	479	<i>leel</i>	231
<i>Typlichthys</i>	194, 195	<i>marginatus</i>	231
<i>subtruncata</i> , <i>Belone</i>	195	<i>nigrescens</i>	230
<i>subtruncatus</i> , <i>Homoprius</i>	345	<i>piger</i>	230
<i>subulatus</i> , <i>Oreynus</i>	260	<i>plagusa</i>	231
<i>Thunnus</i>	260	<i>plagusia</i>	231
<i>succincta</i> , <i>Seriola</i>	278	<i>pusillus</i>	231
<i>sucetta</i> , <i>Cyprinus</i>	109	<i>williamsi</i>	231
<i>Erismyzon</i>	109	<i>synagris</i> , <i>Lutjanus</i>	3, 30, 327
<i>suckleyi</i> , <i>Spinax</i> (<i>Acanthias</i>).....	21	<i>synagrus</i> , <i>Sparus</i>	327
<i>Squalus</i>	21	<i>Synanceia cervus</i>	381
<i>sucklii</i> , <i>Catostomus</i>	105	<i>Synanceia cervus</i>	381
<i>sucubti</i> , <i>Rivulus chucunaque</i>	180	<i>Synapobranchidae</i>	78
<i>Sudis</i>	165	<i>Synapobranchus</i>	78
<i>Sudis</i>	165	<i>bathybius</i>	78
<i>hyalina</i>	165	<i>kaupu</i>	78
<i>intermedius</i>	165	<i>pinnatus</i>	78
<i>ringens</i>	165	<i>Synapteretmus</i>	166
<i>suensoni</i> , <i>Chlorhinus</i>	83	<i>Synbranchidae</i>	76
<i>suerti</i> , <i>Coryphæna</i>	264	<i>Synbranchus</i>	76
<i>suertii</i> , <i>Meletta</i>	42	<i>fuliginosus</i>	76
<i>Sufflamen</i>	492	<i>immaculatus</i>	76
<i>verres</i>	492	<i>marmoratus</i>	76
<i>sufflamen</i> , <i>Balistes</i>	492	<i>transversalis</i>	76
<i>Canthidermis</i>	492	<i>Synchiridae</i>	393
<i>sugillatus</i> , <i>Ophisurus</i>	88	<i>Synchirus</i>	393
<i>Suillus</i>	425	<i>gilli</i>	393
<i>Suillus</i> , <i>Lachnolaimus</i>	425	<i>Synechogianis</i>	151
<i>sulef</i> , <i>Muraenoides</i>	467	<i>beadlei</i>	151, 152
<i>sulcatus</i> , <i>Coryphæna</i>	207	<i>Synentognathi</i>	195
<i>Herpetoichthys</i>	87	<i>Syngnathidae</i>	240
<i>Trachonurus</i>	207	<i>Syngnathinae</i>	240
<i>sumichrasti</i> , <i>Citharichthys</i>	219	<i>Syngnathus</i>	240
<i>sundiacus</i> , <i>Bagrus</i>	149	<i>abboti</i>	240

List of scientific names—Continued

	Page		Page
<i>Syngnathus acus</i>	240	<i>tabacarius</i> , <i>Centropristes</i>	319
<i>aequoreus</i>	243	<i>tacheté</i> , <i>le Diodon</i>	501
<i>affinis</i>	242	<i>tænia</i> , <i>Bassozetus</i>	481
<i>arctus</i>	242	<i>Bathyonus</i>	481
<i>arundinaceus</i>	240	<i>Blennius</i>	468
<i>ascendens</i>	241	<i>Clinostomus</i>	119
<i>auliscus</i>	240	<i>tæmata</i> , <i>Atherina</i>	246
<i>bairdianus</i>	242	<i>tæmatum</i> , <i>Hæmulon</i>	332
<i>brachycephalus</i>	241	<i>tænatus</i> , <i>Anisotremus</i>	332
<i>barbaræ</i>	240	<i>Chirocentron</i>	45
<i>brachycephalus</i>	241	<i>Evoxymetopon</i>	262
<i>bervicaudus</i>	241	<i>Pseudoxiphophorus bimaculatus</i>	188
<i>brevirostris</i>	240	<i>Tropidichthys</i>	500
<i>californiensis</i>	240	<i>Tæniocoonger</i>	82
<i>carinatus</i>	240	<i>digueti</i>	82
<i>crinigerum</i>	241	<i>tæniops</i> , <i>Cephalopholis</i>	309
<i>dekayi</i>	242	<i>Serranus</i>	309
<i>dendriticus</i>	242	<i>tæniopterus</i> , <i>Balistes</i>	491
<i>dimidiatus</i>	240	<i>Cottus</i>	388
<i>elcapitanense</i>	242	<i>Perissias</i>	220
<i>elcapitensis</i>	242	<i>Platophrys</i>	220
<i>elucens</i>	241	<i>Pseudoscarus</i>	433
<i>ethon</i>	241	<i>Scarus</i>	433
<i>fasciatus</i>	242	<i>Tæniomembras</i>	246
<i>fistulatus</i>	241	<i>Tæniophis</i>	91
<i>flaviostris</i>	241	<i>westphali</i>	91
<i>floridae</i>	241	<i>Tæniosomi</i>	216
<i>fuscus</i>	242	<i>Tæniotoca</i>	411
<i>griseolineatus</i>	240	<i>lateralis</i>	411
<i>hippocampus</i>	244	<i>tæniura</i> , <i>Safole</i>	305
<i>jonesi</i>	241	<i>tæniurus</i> , <i>Dules</i>	305
<i>leptorhynchus</i>	240	<i>Labrus</i>	430
<i>linea</i>	241	<i>Poecilurichthys</i>	97
<i>louisianæ</i>	242	<i>Tachysurus gulosis</i>	151
<i>mackayi</i>	241	<i>jordani</i>	150
<i>marmoreus</i>	241	<i>lentiginosus</i>	149
<i>milbertianus</i>	242	<i>longicephalus</i>	150
<i>mindil</i>	242	<i>Tachysurus</i>	150
<i>ophidion</i>	243	<i>chinensis</i>	150
<i>peckianus</i>	242	<i>emmelane</i>	151
<i>peckii</i>	242	<i>evermanni</i>	151
<i>pelagicus</i>	210 242	<i>liropus</i>	151
<i>picturatus</i>	241	<i>steindachneri</i>	151
<i>poeyi</i>	241	<i>tahoensis</i> , <i>Catostomus</i>	107
<i>robertsi</i>	241	<i>Salmo</i>	51
<i>rousseau</i>	241	<i>clarkii</i>	57
<i>sibbaldi</i>	243	<i>taiasica</i> , <i>Gobius</i>	442
<i>sinaloæ</i>	241	<i>Talsmania</i>	53
<i>spicifer</i>	243	<i>æquatoris</i>	53
<i>starksi</i>	242	<i>antillarum</i>	53
<i>tenulis</i>	241	<i>tang</i> , <i>Mugil</i>	253
<i>torrei</i>	242	<i>tanneri</i> , <i>Hyperchoristus</i>	70
<i>tweddlii</i>	242	<i>tapdisma</i> , <i>Salmo</i>	55
<i>viridescens</i>	242	<i>Taractes</i>	265
<i>Synodontidae</i>	163	<i>asper</i>	265
<i>Synodus</i>	163	<i>longipinnis</i>	265
<i>acutus</i>	164	<i>saussurii</i>	265
<i>argenteus</i>	40	<i>Tarandichthys</i>	375
<i>evermanni</i>	164	<i>cavifrons</i>	375
<i>foetens</i>	163, 172	<i>filamentosus</i>	376
<i>intermedius</i>	164	<i>tenuis</i>	376
<i>jenkinsi</i>	163	<i>tarasorum</i> , <i>Algansea</i>	116
<i>lacerta</i>	163	<i>Tarletonbeania</i>	170
<i>lacertinus</i>	164	<i>crenularis</i>	170
<i>lucioceps</i>	163	<i>tenua</i>	170
<i>malabaricus</i>	99	<i>Tarpon</i>	39
<i>poeyi</i>	164	<i>atlanticus</i>	39
<i>saurus</i>	163	<i>Tarpum</i>	39
<i>scituluopsis</i>	164	<i>tarsodes</i> , <i>Bryostemma</i>	466
<i>simulans</i>	163	<i>tassellatum</i> , <i>Boleosoma</i>	287
<i>synodus</i>	163	<i>tassellatus</i> , <i>Labrus</i>	424
<i>synodys</i> , <i>Esox</i>	163	<i>tau</i> , <i>Batrachoides</i>	486
<i>Synodus</i>	163	<i>Gadus</i>	486, 487
<i>Sypterus</i>	240	<i>Opsanus</i>	487
<i>Syriotes</i>	242	<i>pardus</i> , <i>Batrachus</i>	487
<i>affinis</i>	242	<i>Tauridea</i>	383, 386
<i>bairdianus</i>	242	<i>taurinus</i> , <i>Aspicottus</i>	381
<i>exilis</i>	243	<i>Enophrys</i>	381
<i>fuscus</i>	242	<i>taurocephalus</i> , <i>Alburnops</i>	145
<i>louisianæ</i>	242	<i>taurus</i> , <i>A.budefduf</i>	415
<i>scovelli</i>	242	<i>Carcharias</i>	15
<i>Syrraxis</i>	27	<i>Carpiodes</i>	102
<i>Syrrhina</i>	23	<i>Glyphidodon</i>	415
<i>syrtensium</i> , <i>Argentina</i>	68	<i>Tautoga</i>	424
<i>tabacaria</i> , <i>Fistularia</i>	245	<i>alia</i>	424
<i>Prionodes</i>	319	<i>coerulea</i>	424
		<i>fusca</i>	424

List of scientific names—Continued

	Page		Page
<i>Tautoga niger</i>	424	<i>testudineus</i> , <i>Arothron</i>	499
<i>nigra</i>	424	<i>Cheilichthys</i>	499
<i>onitis</i>	424	<i>Tetraodon</i>	499
<i>rubens</i>	424	<i>Tetrabanchus</i>	75
<i>tautoga</i> , <i>Labrus</i>	424	<i>tetracantha cincta</i> , <i>Parapetenia</i>	419
<i>Tautoglabrus</i>	424	<i>grisea</i> , <i>Parapetenia</i>	419
<i>adpersus</i>	424	<i>lata</i> , <i>Parapetenia</i>	419
<i>taylori</i> , <i>Arius</i>	151	<i>Parapetenia</i>	419
<i>Asemathichthys</i>	390	<i>torralbasi</i> , <i>Parapetenia</i>	419
<i>Ophidium</i>	485	<i>tetracanthus</i> , <i>Cantrarchus</i>	419
<i>Otophidium</i>	485	<i>cinctus</i> , <i>Heros</i>	419
<i>teapae</i> , <i>Cichlasoma</i>	417	<i>griseus</i> , <i>Heros</i>	419
<i>Theraps</i>	417	<i>latus</i> , <i>Heros</i>	419
<i>Tectospondyli</i>	21	<i>Sparus</i>	325
<i>Tekla</i>	456	<i>torralbasi</i> , <i>Heros</i>	419
<i>tekla</i> , <i>Stathmonotus</i>	466	<i>Tetragonopterus</i>	359
<i>Teleostei</i>	39	<i>gus</i>	95
<i>telescopium</i> , <i>Pomatomus</i>	296	<i>brevimanus</i>	95
<i>Telescopi</i>	296	<i>fischeri</i>	97
<i>telescopus arcuansanus</i> , <i>Notropis</i>	124	<i>fulgens</i>	95
<i>telescopus</i> , <i>Netropis</i>	124	<i>(Hemibrycon) polyodon</i>	97
<i>Photogenis</i>	124	<i>humilis</i>	95
<i>Telipomis</i>	293	<i>macrophthalmus</i>	97
<i>Temeculina</i>	119, 120	<i>mexicanus</i>	95
<i>temminckii</i> , <i>Carcharias</i>	15	<i>microphthalmus</i>	95
<i>temminckii</i> , <i>Ruvettus</i>	261	<i>nitidus</i>	95
<i>temnistia</i>	380	<i>ovstedti</i>	95
<i>Temnodon</i>	280	<i>petenensis</i>	97
<i>templei</i> , <i>Pomadasys</i>	344	<i>scabripinnis</i>	97
<i>tenebrosus</i> , <i>Alepocephalus</i>	52	<i>Tetragonuridae</i>	269
<i>Antennarius</i>	605	<i>Tetragonurus</i>	269
<i>Caranx</i>	273	<i>atlanticus</i>	269
<i>Xurel</i>	273	<i>cuvieri</i>	269
<i>tenellus</i> , <i>Fundulus</i>	177	<i>simplex</i>	261
<i>Hyborhynchus</i>	145	<i>Tetranarc</i>	27
<i>tentabunda</i> , <i>Trigla</i>	409	<i>californica</i>	27
<i>tentaculata</i> , <i>Venefica</i>	82	<i>occidentalis</i>	27
<i>tenua</i> , <i>Tarletonbeania</i>	170	<i>tetranemus</i> , <i>Extrarus</i>	139
<i>tenuis</i> , <i>Cichlasoma</i>	419	<i>Hybopsis</i>	139
<i>Moxostoma</i>	109	<i>Tetraodon</i>	499
<i>tenuicauda</i> , <i>Macrurus</i>	206	<i>ammocryptus</i>	499
<i>Macrurus</i>	206	<i>hispidus</i>	500
<i>tenuiculum</i> , <i>Myctophum</i>	170	<i>laevigatus</i>	497
<i>tenuifilis</i> , <i>Antennarius</i>	505	<i>lineatus</i>	499
<i>tenuirostris</i> , <i>Anguilla</i>	77	<i>mola</i>	503
<i>tenuis</i> , <i>Anthias</i>	322	<i>perspicillaris</i>	500
<i>Atherinopsis</i>	251	<i>polius</i>	499
<i>Cottus</i>	384	<i>pluviatilis</i>	499
<i>Cottus</i>	180	<i>reticularis</i>	499
<i>Cynodontichthys</i>	213	<i>rostratus</i>	500
<i>Gadus</i>	376	<i>setosus</i>	500
<i>Icelinus</i>	251	<i>stellatus</i>	497, 499
<i>Leuresthes</i>	419	<i>testudineus</i>	499
<i>Parapetenia</i>	192	<i>trichocephalus</i>	499
<i>Poecilia</i>	180	<i>Tetraodontidae</i>	497
<i>Rivulus</i>	241	<i>Tetrapleurodon</i>	9
<i>Syngnathus</i>	376	<i>spadiceus</i>	9
<i>Tarandichthys</i>	384	<i>tetrapturorum</i> , <i>Echeneis</i>	450
<i>Uranidea</i>	213	<i>Tetrapturus amplus</i>	263
<i>Urophycis</i>	422	<i>indicus</i>	263
<i>teporatus</i> , <i>Heros</i>	237	<i>mazara</i>	263
<i>teraculeatus</i> , <i>Gasterosteus</i>	99	<i>mitsukurii</i>	263
<i>tereira</i> , <i>Macrondon</i>	46	<i>tetraspilus</i> , <i>Upeneus</i>	344
<i>teres</i> , <i>Alosa</i>	105	<i>Tetrodon angusticeps</i>	498
<i>Cyprinus</i>	99	<i>annulatus</i>	499
<i>Macrondon</i>	86	<i>capistratus</i>	500
<i>Sphagebranchus</i>	109	<i>(Cheilichthys) pachygaster</i>	499
<i>Teretulus</i>	111	<i>curvus</i>	497
<i>cervinus</i>	142	<i>fluvialis</i>	499
<i>teretulus hosternus</i> , <i>Phenacobius</i>	142	<i>formosus</i>	500
<i>teretulus</i> , <i>Phenacobius</i>	40	<i>furthii</i>	499
<i>tergatus</i> , <i>Hiodon</i>	474	<i>geometricus</i>	499
<i>terre-novae</i> , <i>Lycodes</i>	14	<i>heraldi</i>	499
<i>Scoliodon</i>	14	<i>hispidus maculatus</i>	498
<i>Squalus (Carcharias)</i>	14	<i>implutus</i>	500
<i>Triglops</i>	180	<i>laevigatus</i>	497
<i>ommatistius</i>	417	<i>laterna</i>	500
<i>terrabe</i> , <i>Theraps</i>	186	<i>le</i>	498
<i>terrabensis</i> , <i>Brachyrhaphis</i>	186	<i>lineolatus</i>	497
<i>Gambusia</i>	290	<i>marmoratus</i>	498
<i>tesselata</i> , <i>Etheostoma</i>	231	<i>mathematicus</i>	497
<i>Plagusia</i>	289	<i>mola</i>	503
<i>tessellatus</i> , <i>Hadropterus</i>	424	<i>nepheus</i>	498
<i>Labrus</i>	290	<i>ornatus</i>	500
<i>Nothonotus</i>	459	<i>oxyrhynchus</i>	500
<i>le tar</i> , <i>Lepadogaster</i>	459		

List of scientific names—Continued

	Page		Page
Tetrodon pachycephalus.....	497	Thoburnia.....	107
psittacus.....	500	rhodocæ.....	108
punctatissimus.....	500	Tholichthys.....	359
punctatus.....	499	ossetus.....	359
rostratus.....	500	thompsoni, Ama.....	38
spengleri.....	498	Carpoides.....	103
(Spheroides) eulepidotus.....	498	Gnatholepis.....	443
truncatus.....	504	Lepidosteus.....	37
turgidus.....	498	Triglopsis.....	389
Tetronarce.....	27	thomsoni, Cottunculus.....	392
tetrophthalma, Lioglossina.....	223	Cottus.....	392
tetrophthalmus, Anableps.....	194	thoracatus, Callichthys (Holosternum).....	180
Lioglossina.....	223	Thoracostei.....	237
Teuthis.....	362	thoreauanus, Semotilus.....	117
australis.....	362	Semotilus atromaculatus.....	117
crestonis.....	363	Thorichthys.....	422
hepatus.....	362	acutus.....	422
texana, Anguilla.....	77	affinis.....	422
Cyprinella.....	130	aureus.....	422
Erogia.....	130	callolepis.....	422
texanus, Catostomus.....	108	elliotti.....	422
Xyrauchen.....	108	helleri.....	422
texensis, Dionda.....	144	thrissa, Clupea.....	44
textilis, Salarias.....	464	thrissina, Clupea.....	44
Salarichthys.....	464	Harengula.....	62
thalassinum, Moxostoma.....	110	Thrissomimus.....	91
Napostoma.....	289	Thronodontus.....	91
thalassinus, Cynoscion.....	355	reticulata tessellata.....	259
Doratonotus.....	429	Thunnidae.....	258
Gobius.....	444	thunnina, Thynnus.....	259
Microgobius.....	444	Thynnus.....	260
Myioleucus.....	122	alisoni.....	260
Nothonotus.....	289	coretta.....	260
Otolithus.....	355	sahensis.....	259
Ptychostomus.....	110	secundodorsalis.....	260
Thalassoma.....	428	subulatus.....	260
bifasciatum.....	429	Thymallidae.....	66
grammaticum.....	429	thymalloides, Coregonus.....	66
lucasanum.....	429	Thymallus.....	66
nitidum.....	429	montanus.....	66
socoeroense.....	429	ontariensis.....	66
socoeroensis.....	429	signifer.....	66
steindachneri.....	429	tricolor.....	66
virens.....	429	thymallus, Salmo.....	66
Thalassophryne.....	488	Thynnus.....	258
dowi.....	488	affinis.....	258
maculosa.....	488	atlanticus.....	260
reticulata.....	488	balteatus.....	260
Thaleichthys.....	66	brasiliensis.....	258
albatrossis.....	67	brevipinnis.....	258
pacificus.....	66	coretta.....	260
stevensi.....	66	leachianus.....	258
thaleichthys, Osmerus.....	67	pacificus.....	260
Spirinchus.....	67	pompius.....	278
thausasium, Rhegma.....	322	thunnina.....	258
Thecopterus.....	392	vagrans.....	258
aleuticus.....	392	thynnus, Scomber.....	259
Theragra.....	209	Thyrsoidea concolor.....	93
chalcogramma.....	209	Thyrina.....	248
fucensis.....	209	balsana.....	249
Theraps.....	417	chagresi.....	249
balteatus.....	418	crystallina.....	248
eigenmanni.....	417	evermanni.....	248
frontalis.....	417	guatemalensis.....	249
gadovi.....	417	guila.....	248
gibbiceps.....	417	mekki.....	249
irregularis.....	417	sardina.....	249
lentiginosus.....	417	Thyrmops.....	249
mulleri.....	417	pachylepis.....	249
mojarra.....	417	Thyrus.....	220
nebulifer.....	417	pellucidus.....	220
nicaraguensis.....	417	Thyriscus.....	379
nigritus.....	417	anoplus.....	379
teapæ.....	417	Thyrsoidea maculipinnis.....	92
terræ.....	417	thyrsoidea, Lemnisoma.....	261
zonatus.....	417	Thyrsoidea concolor.....	92, 93
thermalis, Pœcilia.....	192	militaris.....	92
thermophilus, Richardsonius.....	118	Thyrstes acanthoderma.....	261
Therobromus.....	68	niger.....	261
callochmi.....	68	scholaris.....	261
theta, Daphnus.....	168	Thyristops violaceus.....	260
thoburni, Alcides.....	375	Thysanocara.....	160
Mugil.....	253	spinosa.....	160
Parietellus.....	375	Tiaroga.....	143
Querimana.....	253	cobitis.....	143

List of scientific names—Continued

	Page		Page
tiburo, Reniceps.....	17	torsk, Gadus.....	214
Squalus.....	17	tortuga, Ctenogobius.....	439
Zygæna.....	17	Holocentrus.....	286
Tiburon.....	17	Rhinogobius.....	439
Tigoma.....	119	torvus, Cottunculus.....	392
atralia.....	119	Monomitopus.....	480
bicolor.....	120	tota, Perca.....	310
crassa.....	120	toweri, Goodea.....	184
egregia.....	118	townsendi, Angelichthys.....	361
gibbosa.....	119	Gobius.....	444
gracilis.....	119	Myctophum.....	167
humboldtii.....	119	Nyctimaster.....	167
intermedia.....	119	toxotes, Rhachochilus.....	411
lineata.....	119	Toxus.....	188
nigrescens.....	120	cubensis.....	188
orcuttii.....	120	riddlei.....	188
pulchella.....	120	Trachocorystes.....	159
pulchra.....	120	amblops.....	159
pulverentula.....	120	typus.....	159
purpurea.....	119	Trachichthyidae.....	224
squamata.....	119	Trachichthys pretiosus.....	224
rhinichthyoides.....	141	Trachidermis richardsoni.....	383
tigrina, Ichthyophis.....	94	Trachinocephalus.....	163
tigrinus, Galeocerdo.....	14	myops.....	163
Holocentrus.....	319	Trachinotus.....	276
Myrichthys.....	85	argenteus.....	277
Prionodes.....	319	balloni.....	276
tigris, Antennarius.....	506	carolinus.....	277
Carcharias.....	19	culveri.....	277
Chironectes.....	506	cupreus.....	277
Isurus.....	19	falcatus.....	276
Mycteroperca.....	315	fuscus.....	276
Serranus.....	315	goodii.....	277
tilelli, Hemilepidotus.....	381	kennedyi.....	277
timagamiensis, Salvelinus.....	60	paloma.....	277
timlei, Raja.....	27	palometa.....	276
timpanogenesis, Hybopsis.....	119	pampanus.....	276, 277
Timucu.....	195, 199	rhodopus.....	277
timucu, Belone.....	195	spinosus.....	276
Esox.....	195	Trachinus adscensionis.....	310
Strongylura.....	195	cirrhosus.....	332
tinca, Serranus.....	315	gasteropetecus.....	337
tinella, Algansea.....	115	osbecki.....	310
Leuciscus.....	115	punctatus.....	310
tippecanoe, Etheostoma.....	292	trichodon.....	357
Nivicola.....	292	Trachipteridae.....	216
Tirus.....	163	Trachipterus.....	216
marmoratus.....	163	arcticus.....	216
tlauhucensis, Evarra.....	142	tex-salmonorum.....	216
Todichthys.....	410	trachyurus.....	216
agassizi.....	410	Trachonurus.....	207
toe-roe, Otolithus.....	354	sulcatus.....	207
toma, Salmo.....	59	trachura, Gasterosteus.....	237
tomcod, Gadus.....	210	Raja.....	26
Microgadus.....	210	Trachurops.....	270
tomcodus fuscus, Gadus.....	210	brachychira.....	271
luteus, Gadus.....	210	brachychirus.....	271
mixtus, Gadus.....	210	crumenophthalma.....	271
pruinosis, Gadus.....	210	Trachurus.....	270
Tomleodon.....	490	aliciolus.....	279
Tomotichla.....	423	boops.....	272
geddesi.....	423	europæus.....	270
underwoodi.....	423	fasciatus.....	279
topeka, Codoma.....	143	japonicus.....	270
Notropis.....	143	lathamii.....	270
Torpedinidae.....	27	linnaii.....	270
torpedinus, Trygonobatus.....	29	squamosus.....	272
Torpedo.....	27	symmetricus.....	270
bancrofti.....	27	trachurus.....	270
brasiliensis.....	27	trachurus, Cottus.....	381
californica.....	27	japonicus, Caranx.....	270
nobiliana.....	27	Scomber.....	270
occidentalis.....	27	Trachurus.....	270
pictus.....	27	Trachynotus fasciatus.....	276
torquatus, Labrus bifasciatus.....	429	glacioides.....	276
torralbasi, Glaridichthys.....	183	nasutus.....	276
Heros tetracanthus.....	419	rhodopus.....	276
Parapetenia tetracantha.....	419	trachypoma, Myripristis.....	234
torrei, Dasyatis.....	28	trachypomus, Ostichthys.....	234
Pastinachus.....	28	Trachypterus.....	216
Siphostoma.....	242	bogmarus.....	216
Syngnathus.....	242	rex-salmonorum.....	216
Torrentaria.....	293	selenis.....	216
fasciata.....	293	vogmarus.....	216
sagitta.....	293	Trachyrincus.....	203
torridus, Lutjanus.....	326	helolepis.....	203

List of scientific names—Continued

	Page		Page
trachyrincus, Lepidolepsus.....	203	tridentatus, Entosphenus.....	10
trachyurus, Trachipterus.....	216	Petromyzon.....	10
tractus, Acanthurus.....	363	tridentiger, Alogambusia.....	188
trahira, Erythrinus.....	99	Gambusia.....	188
Macrodon.....	99	tridigitatus, Dactyloscopus.....	454
tranquebar, Aspidophoroides.....	397	Trifarcus.....	180
transluceus, Gobius.....	440	felinianus.....	181
transmonta, Columbia.....	215	riverendi.....	181
transmontanus Asipenser.....	34	trifasciata, Hydrargyra.....	176
Rhynchithys.....	130	Lepomis.....	298
transversalis, Synbranchus.....	76	trifurca, Perca.....	318
transversum, Poecilosoma.....	201	Trigla carolina.....	407, 408
traski, Hysteroecarpus.....	412	evolvans.....	407
Trematopsis.....	503	fasciata.....	407
willughbei.....	503	palmipes.....	407
willughben.....	503	strigata.....	407
tremebundus, Elassodiscus.....	402	tentabunda.....	409
Triacanthidae.....	491	volitans.....	408, 409
triacanthus, Argyriosus.....	275	Triglide.....	406
Naucleus.....	278	Triglops.....	379
Poronotus.....	267	beani.....	380
Stromateus.....	396	metopias.....	380
Xeneretmus.....	396	ommatistius.....	379
Xenochirus.....	396	terre-novæ.....	380
Triacis.....	14	pingell.....	379
triacis, Orthonopias.....	391, 392	pleurostictus.....	379
Trienodon.....	14	septicus.....	380
obesus.....	14	terre-novæ.....	380
triagramma, Heros.....	418	xenostethus.....	380
Triakis.....	14	Triglopsis.....	389
scyllium.....	14	ontariensis.....	389
semifasciatum.....	14	stumpsoni.....	389
triangularis, Balistes.....	492	Triglopsis thompsoni.....	389
triangulum, Chaetodon.....	359	Trigonobatus.....	28
Lutjanus.....	348	vulgaris.....	28
Tribranchus.....	77	Trigonophallus.....	187
tribulus, Prionotus.....	408	punctifer.....	187
Trichodon.....	256	Trigonorhinus alveata.....	24
octofilis.....	256	trigonus, Lactophrys.....	497
opercularis.....	256	Ostracion.....	497
trichiapterus, Pteraclis.....	266	trigrammus, Chirus.....	374
Trichituridae.....	262	trilineatum, Pristipoma.....	332
Trichlurus.....	262	trilobatus, Lutjanus.....	317
argenteus.....	262	Scarus.....	434
lepturus.....	262	Triloburus.....	318
nitens.....	262	philadelphicus.....	318
trichoccephalus, Cheilichthys.....	499	trilobus, Blepsnas.....	382
Tetraodon.....	499	trimaculata, Parapetania.....	420
Trichocycilus.....	501	trimaculatum, Heros.....	420
erinaceus.....	501	Trinectes.....	229
Trichoderma.....	494	scabra.....	229
Trichodiodon.....	501	trinitatis, Anchovia.....	50
pilosus.....	501	Anchoviella.....	50
Trichodon.....	357	trinstegus, Acanthurus.....	362
japonicus.....	357	Chaetodon.....	362
lineatus.....	357	triourus, Bodianus.....	323
trichodon.....	357	tripes, Neolotus.....	261
trichodon, Mugil.....	254	Tripteronotus.....	64
Querimana.....	254	hautin.....	64
Trachinus.....	357	Tripterygium carminale.....	458
Trichodon.....	357	tripterygius, Cottus.....	393
Trichodontidae.....	357	tripunctulatus, Maurolicus.....	73
Trichonotus.....	39	Valenciennellus.....	73
Trichopsetta.....	220	triquetar, Ostracion.....	496
ventralis.....	220	Rhinosomus.....	496
triehoistia, Codoma.....	130	triserialis, Murenophis.....	88
Erogala.....	130	Ophichthus.....	88
trichurus, Macrourus.....	206	triseriata, Platyrrhina.....	24
Macrurus.....	206	triseriatus, Platyrrhinoidis.....	24
tricolor, Chaetodon.....	361	trispinatus, Moxostoma.....	105
Engraulis.....	48	Trisotropis.....	313
Holacanthus.....	361	aguaji.....	314
Thymallus.....	66	bonaci.....	314
Tricropterus.....	273	boulengeri.....	313
tricornis, Ostracium.....	497	chlorostomus.....	314
trieuspidadus, Hyporhamphus.....	198	dimidiatus.....	314
trieuspis, Cottus.....	389	interstitialis.....	314
Gymnocanthus.....	389	jordan.....	314
tridecemlineatus, Esox.....	173	microlepis.....	314
tridecemspinosa, Metzelaaria.....	370	reticulatus.....	315
tridecmspinosa, Scorpæna.....	370	stomias.....	314
tridens, Lutjanus.....	318	venenosus.....	313
Pocilia.....	190	apua.....	318
(Acropocilia).....	190	xanthostictus.....	314
Salema.....	338	xenarchus.....	314
Sargus.....	338	trispinosa, Corvina.....	346

List of scientific names—Continued

	Page		Page
trispinosus, <i>Odontopyxis</i>	396	turgidus, <i>Tetrodon</i>	498
<i>Scarus</i>	434	turneri, <i>Lycodes</i>	476
tristis, <i>Moniana</i>	132	turrubarensis, <i>Gambusia</i>	189
tristechus, <i>Atractosteus</i>	38	<i>Poecilopsis</i>	189
<i>Esox</i>	38	tuscumbia, <i>Etheostoma</i>	293
trivittatus, <i>Grammistes</i>	331	<i>Psychromaster</i>	293
Troglichthys.....	195	tuyrensis, <i>Astatheros</i>	421
<i>rosæ</i>	195	<i>Cichlasoma</i>	421
Trogloglanis.....	155	<i>Loricaria</i>	162
<i>pattersoni</i>	155	tweediei, <i>Syngnathus</i>	242
Trompa.....	434	Tylosurus.....	196
troosti, <i>Lepidosteus</i>	37	<i>acus</i>	197
tropica, <i>Echeneis</i>	449	<i>cantrami</i>	197
tropicus, <i>Atractosteus</i>	38	<i>caribbeus</i>	197
<i>Phtheichthys</i>	449	<i>euryops</i>	196
<i>Platypræculus</i>	192	<i>fodiator</i>	196
Tropichthys.....	500	<i>galeatus</i>	197
<i>tæniatus</i>	500	<i>gladius</i>	196
Tropidimius.....	328	<i>guanensis</i>	196
<i>dentatus</i>	328	<i>pacificus</i>	197
Tropidodus.....	10	<i>pterurus</i>	196
troscheli, <i>Herichthys</i>	423	<i>raphidoma</i>	196
<i>Heros</i>	423	<i>sagitta</i>	195
<i>Scades</i>	148	<i>scapularis</i>	195
<i>Scadeichthys</i>	148	<i>sierita</i>	196
troscheli, <i>Abudefduf</i>	415	Tympanomum.....	503
<i>Glyphidodon</i>	415	<i>planci</i>	503
troubridgei, <i>Holoconotus</i>	409	Tyntlastes.....	448
troubridgei, <i>Homalopomus</i>	214	<i>brevis</i>	448
truculentus, <i>Chlorophthalmus</i>	165	<i>sagitta</i>	448
truncata, <i>Belone</i>	196	Typhlichthys.....	194
<i>Mathea</i>	507	<i>osborni</i>	195
<i>Ranzania</i>	504	<i>subterraneus</i>	195
truncatus, <i>Blennius</i>	461	<i>wyandotte</i>	195
<i>Dinectus</i>	35	Typhlogobius.....	447
<i>Salmo</i>	58	<i>californiensis</i>	447
<i>Saurus</i>	163	Typhlosaras.....	508
<i>Scarus</i>	433	<i>shufeldti</i>	508
<i>Tetrodon</i>	504	typicus, <i>Ophiocion</i>	347
Trutta.....	55, 56	<i>Plagyrius</i>	128
trutta, <i>Salmo</i>	55	<i>Rhineodon</i>	18
Trutta.....	55	typhus, <i>Achrophichthys</i>	89
<i>corpore variegato</i>	55	<i>Carangichthys</i>	273
Trycherodon.....	115	tyrannus, <i>Anguilla</i>	77
<i>megalops</i>	115	<i>brevicaudata</i> , <i>Brevoortia</i>	44
Trygon.....	28	<i>Brevoortia</i>	44
<i>gymnura</i>	28	<i>Clupea</i>	44
<i>longa</i>	28	uhleri, <i>Citharichthys</i>	219
<i>osteoticta</i>	29	Ulama.....	341
<i>sabina</i>	29	<i>lefyoi</i>	341
<i>sinensis</i>	28	<i>meeki</i>	341
Trygonobatus torpedinus.....	29	<i>producta</i>	341
tschawytscha, <i>Oncorhynchus</i>	55	Ulea.....	393
<i>Salmo</i>	55	<i>marmorata</i>	393
tschawytschiformis, <i>Salmo</i>	54	Ulcina.....	397
tschudi, <i>Discopyge</i>	27	<i>guntheri</i>	397
tsilcoosensis, <i>Catostomus</i>	106	<i>olriki</i>	397
tsuppitch, <i>Fario</i>	57	Ulocentra.....	286
<i>Salmo</i>	55	<i>blennius</i>	287
tuberculata, <i>Raja</i>	28	<i>gulberti</i>	286
tuberculatus, <i>Catostomus</i>	109	<i>histrio</i>	286
<i>Phæthonichthys</i>	264	<i>meadia</i>	286
<i>Sphæroides</i>	498	<i>phlox</i>	287
tubercule, le <i>Sphéroide</i>	498	<i>sumotera</i>	286
tuberculée, le <i>Sphéroide</i>	498	<i>stigmæa</i>	287
tubularis, <i>Eleotris</i>	437	<i>verecunda</i>	287
tuckeri, <i>Monacanthus</i>	495	(Ulocentra) <i>histrio</i> , <i>Etheostoma</i>	286
tudes, <i>Gobiesox</i>	489	ulochr, <i>Paraliparis</i>	405
<i>Platysqualus</i>	18	ulvæ, <i>Xiphistes</i>	469
<i>Salmo</i>	61	Ulvaria.....	470
<i>Zygana</i>	17, 18	<i>subbifurcata</i>	470
tuditatus, <i>Hybopsis</i>	145	Ulvicola.....	467
tullibee, <i>Argyrosomus</i>	64	<i>sanctæ rosæ</i>	467
<i>Coregonus</i>	61	umatilla, <i>Agosia</i>	141
<i>Leucichthys</i>	64	<i>Agosiope</i>	141
<i>Salmo (Coregonus)</i>	64	Umbla.....	60
tumidus, <i>Carpiodes</i>	103	Umbra.....	174
<i>Chironectes</i>	504	<i>delicatissima</i>	172
<i>Lophius</i>	504	<i>hmi</i>	174
tumifrons, <i>Sternopygus</i>	101	<i>minor</i>	255
tumicata, <i>Liparis</i>	400	<i>pygmæa</i>	174
tumicatus, <i>Caraliparis</i>	400	<i>bilineata</i>	174
turchesius, <i>Scarus</i>	435	umbratilis, <i>Alburnus</i>	125
Turdus cinereus peltatus.....	341	<i>Gambusia</i>	190
<i>flavus</i>	425	<i>Lythurus</i>	125
<i>oculo-radiato</i>	428	<i>Xenophallus</i>	190
<i>pinnis branchialis carens</i>	325	Umbridae.....	174

List of scientific names—Continued

	Page		Page
<i>umbrier</i> , <i>Cichlaurus</i>	417	<i>unicolor</i> <i>Reighardina</i>	230
<i>Notropis</i>	131	<i>Soleotarpa</i>	230
<i>Urobatus</i>	30	<i>vitulinus</i> , <i>Hypoplectrus</i>	321
<i>Urolophus</i>	30	<i>unicornis</i> , <i>Citharichthys</i>	219
<i>umbrierum</i> , <i>Cichlasoma</i>	417	<i>Unicornis pascis bahamensis</i>	495
<i>Umbrina</i>	350	<i>unicornus</i> , <i>Balistes</i>	496
<i>analis</i>	350	<i>unifasciatus</i> , <i>Hemirhamphus</i>	198
<i>arenata</i>	351	<i>Hyporhamphus</i>	198
<i>broussonnetii</i>	350	<i>unilineatus</i> , <i>Hemigrammus</i>	98
<i>coroides</i>	350	<i>Poecilurichthys</i>	338
<i>dorsalis</i>	351	<i>unimaculata</i> , <i>Perca</i>	275
<i>elongata</i>	351	<i>unimaculatus</i> , <i>Argyreus</i>	470
<i>furnieri</i>	349	<i>Clinus</i>	477
<i>gracilicirrhus</i>	350	<i>Gymnelis viridis</i>	424
<i>gracilis</i>	351	<i>uninotatus</i> , <i>Ctenolabrus</i>	188
<i>januaria</i>	351	<i>Girardinus</i>	188
<i>littoralis</i>	351	<i>Girardichthys</i>	327
<i>martinicensis</i>	351	<i>Mesoprion</i>	306
<i>nanus</i>	351	<i>unionensis</i> , <i>Centropomus</i>	124
<i>panamensis</i>	351	<i>universitatis</i> , <i>Notropis</i>	467
<i>phalena</i>	350	<i>univittatus</i> , <i>Apodichthys</i>	343
<i>roncador</i>	350	<i>Upeneus</i>	343
<i>sinaloæ</i>	350	<i>balteatus</i>	343
<i>undulata</i>	350	<i>dentatus</i>	343
<i>xanti</i>	350	<i>flavovittatus</i>	343
<i>umbrosa</i> , <i>Cyprinella</i>	130	<i>grandisquamis</i>	344
<i>Narcine</i>	27	<i>maculatus</i>	343
<i>umbrosus</i> , <i>Eques acuminatus</i>	353	<i>martinicus</i>	343
<i>Esox</i>	173	<i>parvus</i>	343
<i>Gymnothorax</i>	60	<i>punctatus</i>	344
<i>Pareques</i>	353	<i>rathbun</i>	344
<i>Platichthys</i>	226	<i>tetrasplius</i>	343
<i>Psettichthys</i>	226	<i>xanthogrammus</i>	162
<i>Sebastichthys</i>	367	<i>uracantha</i> , <i>Loricaria</i>	211
<i>Sebastomus</i>	367	<i>Uraleptus</i>	87
<i>Umbrula</i>	351	<i>Uranichthys</i>	87
<i>elongata</i>	351	<i>brachycephalus</i>	383
<i>littoralis</i>	114	<i>Uranidea</i>	384
<i>umpque</i> , <i>Ptychocheilus</i>	39	<i>greeneri</i>	385
<i>Unbarana</i>	379	<i>boyl</i>	385
<i>uncinatus</i> , <i>Icelus</i>	377	<i>kumlieni</i>	384
<i>uncinatus</i> , <i>Artediellus</i>	377	<i>marginata</i>	385
<i>Cottus</i>	108	<i>microstoma</i>	386
<i>uncompaghe</i> , <i>Xyrauchen</i>	330	<i>pollicaris</i>	385
<i>undecimale</i> , <i>Hemulon</i>	303	<i>quiescens</i>	383
<i>undecimalis</i> , <i>Centropomus</i>	305	<i>rhothea</i>	383
<i>Centropomus lucioperca</i>	305	<i>semiscabra centroleura</i>	385
<i>Sciæna</i>	423	<i>spilota</i>	384
<i>undecim-radiatus</i> , <i>Centropomus</i>	158	<i>tenuis</i>	384
<i>underwoodi</i> , <i>Herichthys</i>	423	<i>wheeleri</i>	285
<i>Rhamdia</i>	349	<i>uranidea</i> , <i>Cottogaster</i>	285
<i>Tomocichla</i>	350	<i>Etheostoma</i> (<i>Cottogaster</i>).....	142
<i>undulata</i> , <i>Perca</i>	350	<i>uranops</i> , <i>Phenacobius</i>	454
<i>Umbrina</i>	350	<i>Uranoscopidae</i>	454
<i>undulatus</i> , <i>Menticirrhus</i>	349	<i>Uranoscopus anoplos</i>	454
<i>Micropogon</i>	497	<i>y-græcum</i>	508
<i>Ostracion</i>	23	<i>uranoscopus</i> , <i>Ceratus</i>	158
<i>Rhinobatus</i>	258	<i>Mancalis</i>	309
<i>Scomber</i>	315	<i>urichi</i> , <i>Cæcorhamdia</i>	29
<i>undulosus</i> , <i>Serranus</i>	477	<i>Uripheston</i>	30
<i>unera</i> , <i>Ophidium</i>	76	<i>Urobatis</i>	30
<i>Unibranchiptera</i>	76	<i>concentricus</i>	29
<i>grisea</i>	76	<i>halleri</i>	30
<i>immaculata</i>	76	<i>nebulosus</i>	29
<i>lineata</i>	76	<i>sloani</i>	29
<i>marmorata</i>	321	<i>vermiculatus</i>	30
<i>unicolor aberrans</i> , <i>Hypoplectrus</i>	321	<i>umbrier</i>	30
<i>accensus</i> , <i>Hypoplectrus</i>	321	<i>vermiculatus</i>	467
<i>affinis</i> , <i>Hypoplectrus</i>	9	<i>Urocentrus</i>	81
<i>Ammocetes</i>	496	<i>Uroconger</i>	81
<i>Balistes monocoeros</i>	321	<i>varidens</i>	81
<i>bovinus</i> , <i>Hypoplectrus</i>	321	<i>vicinus</i>	80
<i>Erythrus</i>	360	<i>Urolophus aspidurus</i>	30
<i>Chæstodon</i>	321	<i>asterias</i>	30
<i>chlorurus</i> , <i>Hypoplectrus</i>	321	<i>goodii</i>	30
<i>crocotus</i> , <i>Hypoplectrus</i>	321	<i>halleri</i>	30
<i>gemma</i> , <i>Hypoplectrus</i>	321	<i>nebulosus</i>	30
<i>gummigutta</i> , <i>Hypoplectrus</i>	321	<i>rogersi</i>	30
<i>guttavarius</i> , <i>Hypoplectrus</i>	321	<i>umbrier</i>	478
<i>Hypoplectrus</i>	321	<i>Uronectes</i>	213
<i>indigo</i> , <i>Hypoplectrus</i>	90	<i>Urophycis</i>	213
<i>Muraena</i>	321	<i>chesteri</i>	213
<i>negricans</i> , <i>Hypoplectrus</i>	321	<i>chuss</i>	213
<i>Perca</i>	321	<i>floridanus</i>	213
<i>pinnivarius</i> , <i>Hypoplectrus</i>	321	<i>regius</i>	213
<i>puella</i> , <i>Hypoplectrus</i>	321	<i>tenuis</i>	213

List of scientific names—Continued

	Page		Page
urophthalmus, Heros.....	418	veliger, Histiochnus.....	456
Urosetta.....	223	velox, Carcharhinus.....	16
Uropterygius.....	94	Chilola.....	145
necturus.....	94, 95	Euleptorhamphus.....	198
urophthalmus, Parapetenis.....	418	Sarda.....	259
Urotrygon.....	30	venadorum, Mycteroperca.....	314
aspidurus.....	30	Venefica.....	82
goodei.....	30	ocella.....	82
mundus.....	30	procera.....	82
Uroxis.....	28	tentaculata.....	82
ursinus, Salmo.....	60	venenosa, Perca.....	313
urus, Bubalichthys.....	102	venosa, Meletta.....	42
Carpoides.....	102	venosus, Aluterus.....	495
Ictiobus.....	102	Trisotrops.....	313
ustus, Callyodon.....	431	ventralis, Bathypterois.....	166
Cryptotomus.....	431	Brosomphycus.....	480
utah, Salmo.....	56	Citharichthys.....	220
utawana, Cetostomus.....	105	Cottus.....	389
uter, Catulus.....	12	Eucrotus.....	267
Cephaloscyllium.....	12	Novaculichthys.....	430
		Ogilbia.....	480
vacca, Carpiodes.....	103	Trichopsetta.....	220
Damalichthys.....	412	Xyrichtys.....	430
Vacuocua.....	345	ventricosus, Aplocheilichthys.....	398
macrops.....	345	Blepsius.....	380, 381
sancta-lucea.....	345	Ceratichthys.....	141
sialis.....	345	Cyclopterus.....	398
subaequalis.....	345	venusta, Cyprinella.....	130
vafer, Myrophis.....	83	Emmoeleia.....	428
vagans, Katsuwonus.....	258	Erogala.....	130
Thynnus.....	258	Lima.....	175
vagrans, Chirostoma.....	248	venustus, Dinemus.....	234
Kirtlandia.....	248	Hesperoleucus.....	122
lacinata, Memidia.....	248	Pseudopodus.....	428
vahli, Lycodes.....	474	Xyrichtys.....	430
vaigensis, Scarus.....	431	vera-crucis, Bardella.....	346
vallanti, Pteroplatea.....	29	Nector.....	346
Vaillantia.....	288	verany, Cybium.....	256
camura.....	288	Verasper.....	223
chlorosoma.....	288	moseri.....	223
valenciennesi, Pteroplatea.....	29	verde, Pudiano.....	426
valenciennesi, Erotelis.....	437	verecunda, Ulocentra.....	287
Moxostoma.....	109	verecundum, Etheostoma.....	287
Smaragdus.....	441	Lepidon.....	211
Valenciennesellus.....	73	Verhilde.....	328
stellatus.....	73	Verilus.....	328
tripunctulatus.....	73	sordidus.....	328
vampyrus, Cephalopterus.....	32	Verma.....	84
Vandellius.....	262	kendalli.....	85
vandepolli arubensis, Pechia.....	193	vermelho, Pudiano.....	426
Mollinesia sphenops.....	193	vermicularis, Corvina.....	328
Foecilia.....	193	Sigmurus.....	348
vandoisulus, Clinostomus.....	121	Sparus.....	327
Leuciscus.....	121	vermiculatus, Esox.....	173
vanhyningi, Signalosa atchafalaya.....	47	Exocoetus.....	202
varia, Perca.....	317	Urobasia.....	29
variabilis, Dioplites.....	268	sloani.....	29
Perca.....	365	Xyrichtys.....	430
Pomacentrus.....	413	vermiformis, Neoconger.....	81
variata, Etheostoma.....	289	Oxyurus.....	79
variatus, Characodon.....	183	vernalis, Clupea.....	42
Platycephalus.....	193	Gobio.....	137
Poecilichthys.....	289	Hydrargyra.....	176
varidens, Uroconger.....	81	Hyodon.....	40
variegata, Batrachoides.....	487	Scomber.....	237
Loricaria.....	161	vernulus, Batrachoides.....	487
Muraena.....	93	verreuxi, Conger.....	79
Salpa purpurascens.....	327	verres, Balistes.....	492
Skimia.....	153	Lutjanus.....	425
variegatus, Cyprinodon.....	181	Sufflamen.....	492
Limnurgus.....	182	verrilli, Gymnothorax.....	93
Orbis levis.....	499	Sidera.....	93
Pachylabrus.....	411	verrilli, Brosomphycus.....	480
Sargus.....	338	Lycenchelys.....	475
varius, Malacoptenus.....	459	Lycodes.....	475
Monacanthus.....	494	Ogilbia.....	480
Myxodes.....	459	verrucosa, Ora.....	394
Serranus.....	310	verrucosus, Antennarius.....	506
velatus, Ptychostomus.....	109	Brachyopsis.....	394
velcana, Atherina.....	246	Cottus.....	387
velifer, Carpiodes.....	103	Diodon.....	502
Cetostomus.....	86	Myoxocephalus.....	387
Letharchus.....	141	Verrugato.....	323
Rhynchichthys (Apocope).....	141	pacificus.....	324
vellifera, Apocope.....	141	Verruculus.....	492
Mollinesia.....	192	polylepis.....	492

List of scientific names—Continued

	Page		Page
versicolor, Abramis.....	115	virgata, Brachistoma.....	7
Girardinus.....	191	virginalis, Salmo.....	57
Labrus.....	336	Searus.....	433
Lima.....	191	virginicus, Anisotremus.....	332
Malacopterus.....	460	Polynemus.....	256
Myxodes.....	460	Sparus.....	322
Phytichthys.....	469	viride, Sparisoma.....	432
Poecilichthys.....	291	vindensis, Sphyræna.....	255
Stenotomus.....	336	vindescens, Atherina.....	247
Xiphistes.....	469	Fundulus.....	176
versipunctatus, Gymnothorax.....	92	Osmerus.....	67
vertagus, Gadus.....	210	Synnathus.....	242
verticalis, Echeineis.....	449	viridipallidus, Gobius.....	446
Pleuromichthys.....	225	vindis, Amla.....	38
verus, Conger.....	79	bahamensis, Piscis.....	433
vespertilio, Lophius.....	507	Centropomus.....	306
Ogcocephalus.....	507	Diacope.....	324
Rhinoptera.....	31	Gymnelis.....	477
verus, Carcharodon.....	20	Lutianus.....	324
vetula, Balistes.....	491	Ophidium.....	477
Searus.....	433, 434	Searus.....	432
vetulus, Parophrys.....	225	unimaculatus, Gymnelis.....	477
vevillare, Boleosoma.....	288	viscosus, Cottus.....	385
vevillaris, Pteropodus.....	369	Silurus.....	155
Sebastichthys.....	369	vitrea, Ioa.....	288
vevillarium, Holocentrum.....	236	Perca.....	282
vevillarius, Holocentrus.....	236	vitreum, Stizostedion.....	282
Vexillifer.....	486	vitreus, Poecilichthys.....	288
viana, Murenophis.....	92	vitta, Xyrnichthys.....	430
vianalis, Icelus.....	378	vittata, Aztecula.....	143
vicianus, Gymnothorax.....	92	Channomuræna.....	95
Uroconger.....	81	Clupea.....	41
victoria, Moxostoma.....	109	Codoma.....	143
Vieja.....	433, 434	Echeineis.....	449
Vigil.....	288	Gambusia.....	186
clarus.....	288	Hemitremia.....	121
pellucidus.....	288	Inermia.....	340
vivax.....	288	Lepidomeda.....	147
vigil, Ioa.....	288	Lima.....	191
vigilax, Ceratichthys.....	144, 145	Poecilia.....	191
Villarius.....	152	vittatus, Ammodytes.....	451
prisci.....	153	Catostomus.....	109
villosa, Clupea.....	66	Cyprinus.....	140
Fistularia.....	245	Esox.....	173
villosus, Cottus.....	393	Gobiesox.....	489
Hemirhamphus.....	393	Ichthyophis.....	95
Mallotus.....	66	Leuciscus.....	123
vincente, Sicydium.....	447	Proceros.....	36
vinciguerra, Exocoetus.....	200	Sarchirus.....	36
Exonatus.....	200	Siphateles.....	122
Vindiguerra.....	73	Sparus.....	332
attenuata.....	73	Symbranchus.....	76
luetia.....	73	vitula, Sargus.....	338
nimbaria.....	73	vitulinum, Plectropoma.....	321
vincta, Xurel.....	272	vitulinus, Hypoplectrus unicolor.....	321
vinctipes, Nonostoma.....	289	vitulus, Carpiodes.....	102
vinctus, Blennius.....	461	vivanet, Bodianus.....	325
Caranx.....	272	vivanus, Anthias.....	322
Fundulus.....	176	Hemianthias.....	322
violentus, Lethotremus.....	398	Lutianus.....	326
viola, Antimora.....	211	Mesoprion.....	326
Eques.....	353	vivax, Ammocrypta.....	288
Haloporphyrus.....	211	Chola.....	145
Pareques.....	353	Vigil.....	288
violaceus, Apodichthys.....	466	Vivero.....	312
Cebidichthys.....	466	vivipara, Poecilia.....	190
Escolar.....	260	viviparus, Lophius.....	504
Thyrstopsis.....	260	Sebastes.....	364
virens, Gadus.....	209	viamingii, Coryphæna.....	265
Pollachius.....	209	Vogmarus.....	216
Searus.....	433	vogmarus, Trachypterus.....	216
Thalassoma.....	429	volador, Exocoetus.....	200
virescens, Apodichthys.....	467	Istiophorus.....	203
Clupea.....	42	Rubio.....	407
Eigenmannia.....	101	volans, Dactylopterus.....	409
Eriscon.....	358	Exocoetus.....	200
Gymnothorax.....	92	Trigla.....	409
Otolithus.....	256	volucella, Hybopsis.....	133
Pantosteus.....	105	volucellus, Hybognathus.....	138
Sternarchus.....	101	Vomer.....	274
virgata, Coryphæna.....	265	brasiliensis.....	275
Nivicola.....	292	brownii.....	275
virgatus, Delolepis.....	472	cayennensis.....	275
Poecilichthys.....	292	columbianus.....	275
virgatus, Gobiesox.....	489	cubæ.....	275

List of scientific names—Continued

	Page		Page
Vomer curtus.....	275	williamsoni elmontanus, Coregonus.....	65
declivifrons.....	275	Prosopium.....	65
dominicensis.....	275	williamsoni, Coregonus.....	65
domalis.....	275	Ctenodax.....	269
gabonensis.....	275	Gasterosteus.....	237, 238
goreensis.....	275	Prosopium.....	65
martinicensis.....	275	willoughbyi, Acrotus.....	269
novae-boracensis.....	275	willoughbyi, Trematopsis.....	503
sanctae-mari-thae.....	275	willoughbyi, Balistes.....	493
sanctae-petii.....	275	Canthidermis.....	493
senegaletis.....	275	Trematopsis.....	503
setapinnis.....	275	wilsoni, Acontmentum.....	366
cubensis.....	275	Cottus.....	365
spixii.....	275	Oceanomyzon.....	9
vomer, Argyreolus.....	275	Phenagonistes.....	95
Zeus.....	275	Sebastodes.....	266
vomerius, Atherina.....	250	winehelli, Hybopsis.....	133
Chirostema.....	250	woolmani, Paralichthys.....	224
vomerinus, Anarrhichas.....	472	wrayi, Gambusia.....	185
Salaria.....	464	wrighti, Etheostoma.....	235
vorax, Mesoprion.....	323	Isiophorus.....	233
Pseudorhombus.....	224	wurdemanni, Gobius.....	441
Ptychocheilus.....	114	wyandotte, Typhlichthys.....	195
vulgaris, Acanthias.....	21		
Ameiurus.....	153	xenoccephalus, Hybopsis.....	127
Conger.....	79	Hydrophlox.....	137
Gunnellus.....	467	xenura, Erogala.....	130
Lumpus.....	398	xenurus, Minnilus.....	130
Molva.....	212	xalscone, Xenendum.....	184
Pagrus.....	337	xaniurus, Catulus.....	12
Pimelodus.....	153	Fristiurus.....	12
Pomotis.....	302	Xanthichthys.....	493
Sphyræna.....	255	lineopunctatus.....	493
Spinachia.....	239	ringsens.....	493
Squatina.....	22	xanthocephalus, Silurus.....	154
Trigonobatus.....	25	xanthogrammus, Upeneus.....	343
vulnerata, Apocope.....	141	xanthomelas, Anguilla.....	77
vulneratus, Nethonotus.....	230	xanthops, Odontoseion.....	244
Poecilichthys.....	290	xanthopteron, Hæmulon.....	330
Vulpecula.....	18	xanthopterus, Catostomus.....	107
marina.....	18	xanthosticta, Mycteroperca bonaci.....	314
vulpecula, Galeus.....	13	xanthostictus, Tristropis bonaci.....	314
vulpeculus, Pimelodus.....	154	xanthostigma, Citharichthys.....	218
Vulpes bahamensis.....	39	xanthostigmus, Citharichthys.....	218
vulpes, Albula.....	39	xanthulum, Cynoscion.....	255
Esox.....	39	xanthulus, Eriscion.....	356
Pimelodus.....	152	Leostomus.....	249
Squalus.....	18	Pomacentrus.....	413
vulpinus, Alopias.....	18	Sparus.....	336
Squalus.....	18	xanti, Labrisomus.....	459
Vulsiculus.....	406	Rypticus.....	316
imberbis.....	406	Umbrina.....	350
vulsus, Agonus.....	395	Xenichthys.....	324
Stelgis.....	395	Xantocles.....	375
vultharius, Acipenser.....	34	frenatus.....	375
		xantops, Odontoseion.....	334
wabashensis, Anguilla.....	77	xantus, Lepidopus.....	262
wachna, Gadus.....	210	xenarcha, Mycteroperca.....	214
wagneri, Pimelodus.....	157	Xenarchi.....	215
Rhamdia.....	157	xenarchus, Tristropis.....	314
walker, Selanopus.....	19	Xenatherina.....	249
warreni, Boleichthys.....	294	hsa.....	249
Salmo.....	55	renauchen, Galeichthys.....	150
watanga, Erimystax.....	138	Xenendum.....	183
Hybopsis.....	138	calante.....	184
webbi, Nemabrama.....	234	multipunctatum.....	183
webbi, Blennophis.....	464	xalscone.....	184
Embiotoca.....	411	Xeneretmus.....	396
Ophioblennius.....	464	infrapinnatus.....	396
weberi, Xenogobius.....	442	lelops.....	396
welaka, Notropis.....	124	ritteri.....	396
welshi, Otophidiium.....	486	triacanthus.....	396
wheatlandi, Gasterosteus.....	238	Xenichthyidae.....	324
wheeleri, Uranidea.....	384	Xenichthys.....	324
whipplei alabamæ, Etheostoma.....	292	californiensis.....	324
whipli alabamæ, Etheostoma.....	292	xanti.....	324
Boleichthys.....	292	xenops.....	324
Claricola.....	292	xenicus, Fundulus.....	180
Cyprinella.....	129	Xenisma.....	179
Erogala.....	129	catenatum.....	179
whitei, Balsadichthys.....	183	stellifer.....	179
Goodea.....	183	xenisma, Gurnardus.....	406
Salmo.....	59	Prionotus.....	406
wilderi, Lampetra.....	9	Xenistius.....	324
williamsi, Symphurus.....	231	californiensis.....	324

List of scientific names—Continued

	Page		Page
Xenoberyces.....	232	Xurel melampyus.....	272
Xenocara.....	160	stellatus.....	273
chagresi.....	160	tenebrosus.....	273
Xenochirus.....	396	vinota.....	272
alascanus.....	396	xyosterna, Stellerina.....	394
latifrons.....	396	xyosternus, Brachyopsis.....	394
pentacanthus.....	396	Xyrauchen.....	108
triacanthus.....	396	texanus.....	108
Xenodermichthys.....	53	uncompbagre.....	108
copei.....	53	Xyrichthys.....	430
nodulosus.....	53	binghami.....	430
xenodon, Calotomus.....	431	infirmus.....	430
Leptoscarus.....	431	jessæ.....	429
Xenognathus.....	52	martinicensis.....	430
profundorum.....	52	modestus.....	430
Xenogobius.....	442	mundiceps.....	430
weberi.....	442	pavoninus.....	430
Xenomi.....	172	psittacus.....	430
Xenomystax.....	81	rosipes.....	430
atrarius.....	81	ventralis.....	430
rictus.....	81	venustus.....	430
Xenopallus.....	190	vermiculatus.....	430
umbrahtis.....	190	vitta.....	430
xenops, Xenichthys.....	324	xyris, Scorpæonoides.....	370
xenopterus, Cypselurus.....	202	Sebastopsis.....	370
Exocoetus.....	202	Xyruia.....	429
Xenopterygi.....	458	jessæ.....	429
Xenophthalmichthys.....	69	Xystema.....	341
dane.....	69	havana.....	342
Xenopyxis.....	396	Xyster.....	339
latifrons.....	396	xyster, Zapteryx.....	24
lelops.....	396	Xyterus.....	339
ritteri.....	396	Xytes.....	395
Xenorhynchichthys.....	254	axinophrys.....	395
xenostethus, Sternias.....	380	Xystophorus.....	278
Triglops.....	380	Xystrama.....	296
Xenotis.....	300	pandionis.....	296
aureolus.....	300	Xystreurus.....	223
lythrochloris.....	300	holepis.....	223
megalotis.....	300	xystrodon, Sparisoma.....	431
occidentalis.....	301	Xystroperca.....	315
Xeropes.....	467	pardalis.....	315
fucorum.....	467	Xystroplites.....	301
Xesurus.....	363	gilli.....	302
clarionis.....	363	longimanus.....	302
punctatus.....	363	pallidus.....	302
Xiphias.....	264	Xystrosus.....	142
gladius.....	264	popoche.....	142
imperator.....	264	xysturus, Myrichthys.....	85
rondeletti.....	264	Ophisurus.....	85
Xiphidion.....	369	yalto, Euthynnus.....	289
mucosum.....	469	yalei, Ostracion.....	497
Xiphidium cruxreum.....	469	Yarella.....	72
Xiphidae.....	264	blackfordi.....	72
Xiphiliformes.....	263	yarrowi, Agosia.....	141
Xiphister.....	469	Apocope.....	141
chirus.....	469	y-cauda, Gillichthys.....	444
mucosus.....	469	Quietula.....	444
rupestris.....	469	y-græcum, Astroscopus.....	454
Xiphisteridæ.....	469	Uranoscopus.....	454
Xiphistes.....	469	yucatanæ, Gambusia.....	186
ulvæ.....	469	yuma, Gobiesox.....	490
versicolor.....	469	Yunria.....	137
Xiphophorus.....	193	alta.....	137
bimaculatus, vs.....	187	zacentrus, Hispanicus.....	368
brevis.....	194	Sebastichthys.....	368
gracilis.....	185	zachirus, Errex.....	228
güntheri.....	193	Glyptocephalus.....	228
helleri.....	193	Zadlemus.....	351
brevis.....	194	goodel.....	351
güntheri.....	193	zadockii, Cylindrosteus.....	37
rachovii.....	193	Zalarges.....	73
strigatus.....	193	nimbarius.....	73
jalape.....	193	Zalembius.....	410
montezuma.....	194	rosaceus.....	410
rachovi.....	193	Zalenteus.....	507
strigatus.....	193	elater.....	507
Xiphostomidæ.....	100	Zalocys.....	277
Xiphotheca.....	262	stilbe.....	277
Xurel.....	272	Zalopyr.....	365
fasciatus.....	272	aleutianus.....	365
lata.....	272	atrorubens.....	365
lugubris.....	273	atrovirens.....	365
marginatus.....	272		
medusicola.....	272		

List of scientific names—Continued

	Page		Page
Zalypnus.....	443	zonale arcasanum, Etheostoma.....	289
Zanellidae.....	362	Nanostoma.....	289
Zanellurus.....	263	zonalis, Poeciliichthys.....	289
Zandrus.....	362	zonata, Elasmoma.....	305
centrognathos.....	362	Heterandria.....	191
cornutus.....	362	Seriola.....	279
rauculus, Dermatolepis.....	313	zonatum, Cichlasoma.....	417
zanemus, Ceratichthys.....	138	Elasmoma.....	305
Zanolepididae.....	375	zonatus, Alburnus.....	128
Zanolepis.....	375	Chaetodipterus.....	358
frenatus.....	375	Cocotus.....	128
latipinnis.....	375	Ephippus.....	358
Zanotacanthus.....	76	Esox.....	176
Zaphotias.....	72	Fundulus.....	178
pedalotus.....	72	Scorpaenidae.....	279
Zaprora.....	452	Theraps.....	417
silenus.....	452	Zygocetes.....	177
Zaproridae.....	452	Zonichthys.....	279
Zapteryx.....	24	falcatus.....	280
exasperatus.....	24	fasciatus.....	279
xyster.....	24	mazatlanus.....	279
zapus, Hemilepidotus.....	381	rivolanus.....	280
Zastomias.....	70	zonifer, Clinus.....	459
scintillans.....	70	Erilepis.....	373
zatriopsis, Siphostoma.....	243	Myrionolepis.....	373
zebra, Acanthurus.....	362	Zygocetes.....	178
Arbacia.....	491	zonipsectus, Pomacanthodes.....	361
Fundulus.....	176	zonistius, Cocotus.....	128
Gerres.....	342	Luxilus.....	128
Gobiosoma.....	491	Zonogobius.....	441
Gobius.....	441	zebra.....	441
Hydrargyra.....	179	zonope, Jordania.....	375
Percina.....	283	Zonoscion.....	351
Pleuromma.....	283	rathbuni.....	351
Plancterus.....	179	zonurus, Malacocottus.....	388
Psychrolutes.....	392	Zoogoneticus.....	182
Zonogobius.....	441	diazii.....	182
zebrinus, Fundulus.....	179	dugesii.....	182
Zeldae.....	232	maculatus.....	182
zelotes, Dactyloscopus.....	455	miniatus.....	182
Esiocopus.....	455	quitzeensis.....	182
Hemicaranx.....	271	robustus.....	182
Zenion.....	282	Zophendium.....	140
hololepis.....	282	australe.....	116
zenithus, Argyrosomus.....	64	zopherus, Potamocottus.....	385
Leucichthys.....	64	zophochir, Ophichthys.....	88
Zenopsis.....	232	Ophichthys.....	88
ocellata.....	232	zosterse, Hippocampus.....	244
Zenopidae.....	232	zosterura, Evermannia.....	445
zephyreus, Astroscopus.....	454	zosterurum, Gobiosoma.....	445
Pristis.....	23	zunensis, Leuciscus.....	114
Zesticelus.....	392	Zygæna.....	18
profundorum.....	392	lewini.....	18
Zestidium.....	347	melleus.....	18
colonense.....	347	subarcuata.....	18
illecebrosus.....	347	tudes.....	18
Zestus.....	347	zygæna, Sphyrna.....	18
furthii.....	347	Squalus.....	18
melanocheir.....	347	Zygobatis.....	31
oscutans.....	347	Zygocetes.....	177
zestocarus, Stellicarens.....	347	atriatus.....	184
Stelliferus.....	347	aproguttatus.....	177
zestus, Nebria.....	354	chrysotus.....	177
Zeus capillaris.....	275	cingulatus.....	177
crinitus.....	274	craticula.....	178
gallus.....	274	diaphanus.....	178
geometricus.....	275	menona.....	178
guttatus.....	216	dispar.....	178
imperialis.....	217	escambia.....	178
luna.....	216	extensus.....	179
ocellatus.....	232	funduloides.....	176
opal.....	217	guttatus.....	178
quadratus.....	358	henshalli.....	177
regius.....	216	hieroglyphicus.....	178
rostratus.....	275	murus.....	185
setapinnis.....	275	jenkinsi.....	178
stromii.....	217	lateralis.....	177
vomere.....	275	lumi.....	178
zirahuen, Chirostoma.....	250	lineatus.....	177
Zoarces gronovi.....	473	lineolatus.....	178
labrosus.....	473	luciae.....	178
Zoarcesidae.....	473	macdonaldi.....	177
zoarchus, Lycodes.....	475	manni.....	176

List of scientific names—Continued

	Page		Page
<i>Zygonectes mclapleurus</i>	178	<i>Zygonectes scartes</i>	177
(<i>Micristius</i>) <i>brachypterus</i>	185	<i>sciadicus</i>	177
<i>notatus</i>	177	<i>seminolis</i>	178
<i>notul</i>	178	<i>zonatus</i>	177
<i>pulchellus</i>	177	<i>zonifer</i>	178
<i>pulvereus</i>	178	<i>zygopterus</i> , <i>Galeorhinus</i>	14
<i>rathbuni</i>	179	<i>Zypothyca</i>	261
<i>rubrifrons</i>	177		

